

**FEMS Microbiology Ecology**

July 2010, Volume 73 Issue 1, Pages 121 - 133

<http://dx.doi.org/10.1111/j.1574-6941.2010.00881.x>

© 2010 Federation of European Microbiological Societies.

Published by Blackwell Publishing Ltd. All rights reserved

**Archimer**  
<http://archimer.ifremer.fr>

The definitive version is available at <http://www3.interscience.wiley.com/>

---

## **Marine culturable yeasts in deep-sea hydrothermal vents: species richness and association with fauna**

Gaëtan Burgaud<sup>1,\*</sup>, Danielle Arzur<sup>1</sup>, Lucile Durand<sup>2</sup>, Marie-Anne Cambon-Bonavita<sup>2</sup> & Georges Barbier<sup>1</sup>

<sup>1</sup> Laboratoire Universitaire de Biodiversité et Ecologie Microbienne (EA3882), IFR 148, Université Européenne de Bretagne, Université de Brest, ESMISAB, Technopole Brest-Iroise, Plouzané, France

<sup>2</sup> Ifremer, Centre de Brest, DEEP/LM2E, UMR 6197, Plouzané, France

\*: Corresponding author : G. Burgaud, Tel.: +33 2 98 05 61 48; fax: +33 2 98 05 61 01; email address : [gaetan.burgaud@univ-brest.fr](mailto:gaetan.burgaud@univ-brest.fr)

---

### **Abstract:**

Investigations of the diversity of culturable yeasts at deep-sea hydrothermal sites have suggested possible interactions with endemic fauna. Samples were collected during various oceanographic cruises at the Mid-Atlantic Ridge, South Pacific Basins and East Pacific Rise. Cultures of 32 isolates, mostly associated with animals, were collected. Phylogenetic analyses of 26S rRNA gene sequences revealed that the yeasts belonged to *Ascomycota* and *Basidiomycota* phyla, with the identification of several genera: *Rhodotorula*, *Rhodospiridium*, *Candida*, *Debaryomyces* and *Cryptococcus*. Those genera are usually isolated from deep-sea environments. To our knowledge, this is the first report of yeasts associated with deep-sea hydrothermal animals.

**Keywords:** yeasts • hydrothermal vents • fauna • 26S rRNA gene • FISH

## Introduction

Yeasts are ubiquitous microorganisms that represent a part of the microbiota in all natural ecosystems, such as soils, freshwaters and marine waters from the ocean surface to the deep sea. Marine yeasts are divided into obligate and facultative groups. Obligate marine yeasts are yeasts that have never been isolated from anywhere other than the marine environment, whereas facultative marine yeasts are also known from terrestrial habitats (Kohlmeyer & Kohlmeyer, 1979). Based on these definitions, Kohlmeyer & Kohlmeyer (1979) examined yeasts occurring in marine environments and gathered a list of 176 species isolated from diverse marine habitats. Of those, only 25 were obligate marine yeasts, widely represented by the genera *Metschnikowia*, *Rhodosporidium*, *Candida* and *Torulopsis*.

Hawksworth (2002) hypothesized the existence of 1.5 million fungal species; this estimate is now a commonly used and accepted figure. If this is correct, <5% of the fungi have been described up to now and these almost exclusively from terrestrial environments. In that ecosystem, fungi are known to utilize a wide spectrum of simple and more complex organic compounds. The decomposition activities of fungi are clearly important in relation to the redistribution of elements among organisms and environmental compartments (Gadd, 2007). Bearing in mind these parameters, our hypothesis is that deep sea and especially hydrothermal vents, which remain underexplored habitats for fungi, could be ecological niches hosting specific fungal communities.

Deep-sea hydrothermal vents are localized at seafloor spreading centers called rifts, where seawater seeps into cracked regions caused by the presence of hot basalt and magma.

1  
2  
3 57 Seawater carrying dissolved minerals is then emitted from springs. Two major types of  
4  
5 58 emissions have been found. Warm fluids diffuse at temperatures ranging from 6 to 23°C into  
6  
7 59 seawater at 2-4°C when hot vents called black smokers emit hydrothermal fluid at 270-  
8  
9 60 380°C (Munn, 2003). Thermal gradients in hydrothermal vents are so important that just a  
10  
11 61 few centimeters away, the temperature can fall to 2-4°C allowing mesophilic or psychrophilic  
12  
13 62 organisms as well as thermophilic and hyperthermophilic prokaryotes to grow and interact  
14  
15 63 with all biotic or abiotic components of these ecosystems. Dense animal communities cluster  
16  
17 64 around those hot springs. These communities are supported by the chemolithoautotrophic  
18  
19 65 activities of prokaryotes (Jorgensen and Boetius, 2007).

20  
21 66 The occurrence of fungi (filamentous fungi and yeasts) at deep-sea hydrothermal vents  
22  
23 67 remains an underexplored topic. Over the last years, the interest for the diversity of microbial  
24  
25 68 eukaryotes in these ecosystems emerged using PCR amplification of SSU ribosomal RNA  
26  
27 69 genes and sequence analysis (Edgcomb *et al.* 2002; Lopez-Garcia *et al.*, 2003; 2007). These  
28  
29 70 papers revealed a scarce fungal diversity but some sequences were novel. Only two papers  
30  
31 71 have specifically dealt with fungal diversity at deep-sea hydrothermal vents based on culture-  
32  
33 72 dependent methods (Gadanhó & Sampaio, 2005; Burgaud *et al.*, 2009). Culturable yeasts  
34  
35 73 affiliated to *Ascomycota* and *Basidiomycota* phyla were reported from hydrothermal waters.  
36  
37 74 Some papers assessing fungal diversity at deep-sea vents were also published. Bass *et al.*  
38  
39 75 (2007) reported the presence of sequences affiliated to *Debaryomyces hansenii* and novel  
40  
41 76 sequences closed to *Malassezia furfur* in hydrothermal sediments. Le Calvez *et al.* (2009)  
42  
43 77 reported that fungal diversity from deep-sea vent animals was widely constituted of sequences  
44  
45 78 affiliated to *Chytridiomycota* and *Basidiomycota* phyla. The latter phylum was mostly  
46  
47 79 represented by yeasts with, for example, the *Cryptococcus* and *Filobasidium* genera that form  
48  
49 80 dense clusters.

50  
51 81  
52  
53 82 The occurrence of yeasts in other deep-sea environments has been much more studied.  
54  
55 83 Nagahama *et al.* (2001b) reported that culturable fungal diversity was dominated by  
56  
57 84 ascomycetous yeasts in surface sediments in water depths exceeding 2000 meters (*Candida*,  
58  
59 85 *Debaryomyces*, *Kluyveromyces*, *Saccharomyces* and *Williopsis*). Inversely, diversity was  
60  
61 86 dominated by basidiomycetous yeasts on the subsurface of sediments in water depths  
62  
63 87 exceeding 2000 meters and from deep-sea clams, tubeworms and mussels (*Rhodotorula*,  
64  
65 88 *Sporobolomyces*, *Cryptococcus* and *Pseudozyma*). Recent studies have clearly demonstrated  
66  
67 89 that *Cryptococcus* was the dominant genus sequenced from sediments collected at deep  
68  
69 90 methane cold seeps (Takishita *et al.*, 2006; 2007). Those observations are in agreement with

1  
2  
3 91 Bass *et al.* (2007) who suggest that yeast forms dominate fungal diversity in deep oceans.  
4  
5 92 Several yeasts mostly isolated from deep-sea sediments represented new species in the  
6  
7 93 *Ascomycota* or *Basidiomycota* phyla (Nagahama *et al.*, 1999; 2001a; 2003a; 2003b; 2006a;  
8  
9 94 2008).

10 95  
11  
12 96 In this study, we decided to assess the presence of yeasts at deep-sea hydrothermal vents  
13  
14 97 based on a culture-based approach with an emphasis on yeasts in interactions with the  
15  
16 98 endemic animal fauna thriving in these extreme ecosystems. A recent paper (Gadanhó and  
17  
18 99 Sampaio, 2005) has dealt with the diversity of yeasts in deep-sea vent waters but, to our best  
19  
20 100 knowledge, this is the first report of the culturable yeasts isolated from deep-sea animals.  
21  
22 101 Those interactions with the fauna are discussed based on the cultures obtained from the  
23  
24 102 samples collected during different oceanographic the cruises at Mid-Atlantic Ridge, South-  
25  
26 103 West Pacific Lau Basin and East Pacific Rise.

## 27 104 **Materials and methods**

### 28 29 30 105 *Environmental sampling*

31  
32  
33 106 210 hydrothermal samples were collected during 6 oceanographic cruises at several dates and  
34  
35 107 locations (For hydrothermal vents locations, see Tivey, 2007): (i) BIOLAU in the Lau Basin,  
36  
37 108 South-west Pacific (12/05/1989–27/05/1989; 20°3.0'S, 176°7.8'W; -2620 m); (ii)  
38  
39 109 DIVANAUT2 (19/06/ 1994–01/07/1994) on the MAR at Menez Gwen (37°51'N, 31°31'W; -  
40  
41 110 900 m) and Lucky Strike (37°17'N, 32°16'W; -1650 m) hydrothermal sites; (iii) HERO on the  
42  
43 111 EPR at Elsa site (30/09/1991–04/11/1991; 12°48'N, 103°57'W; -2630 m); (iv) MARVEL  
44  
45 112 (29/08/1997–13/09/1997) on the MAR at Menez Gwen and Lucky Strike sites; (v) EXOMAR  
46  
47 113 (25/07/2005–28/08/2005) on the MAR at Rainbow (36°08'N, 34°00'W, -2300 m), TAG  
48  
49 114 (26°02'N, 44°54'W, -3630 m) and Lost City (30°04'N, 42°12'W, -900 m) sites; (vi)  
50  
51 115 MoMARDREAM-Naut (08/07/2007–19/07/2007) on the MAR at Rainbow site. Depending  
52  
53 116 on cruises, deep-sea sampling was performed using the Deep Submergence Vehicle “Nautile”  
54  
55 117 or the Remote Operated Vehicle (ROV) “Victor 6000” and the N/O “Atalante” and  
56  
57 118 “Pourquoi Pas?” research vessel.

58  
59 119 The deep-sea samples were processed as described by Burgaud *et al* (2009) taking care to  
60  
120 avoid contamination in applying strict sterile sampling conditions.

### 121 122 *Enrichment conditions and isolation*

1  
2  
3 123 The samples were processed directly after the Nautilic or ROV recovery. The collected samples  
4  
5 124 mainly composed of deep-sea hydrothermal vent animals (*Rimicaris exoculata* and  
6  
7 125 *Chorocaris chacei* shrimps and *Bathymodiolus azoricus* mussels) were used to inoculate the  
8  
9 126 GYPS culture medium that led to the best isolation rate during a previous study (Burgaud *et*  
10  
11 127 *al.*, 2009). This medium contained per liter: glucose (Sigma) 1 g, yeast extract (AES) 1 g,  
12  
13 128 peptone (AES) 1 g, starch (Fisher) 1 g, sea salts (Sigma) 30 g. This medium was  
14  
15 129 supplemented per litre with agar 15 g and chloramphenicol (Sigma) 500 mg, pH was also  
16  
17 130 adjusted to 7.5. Cultures were done aerobically at 4°C, 15°C, 25°C (ambient temperature), 35  
18  
19 131 and 45°C (only during EXOMAR) at atmospheric pressure until fungal strains were  
20  
21 132 visualized. During the MoMARDREAM-Naut cruise, some dissections were realized on  
22  
23 133 board on animal samples in order to investigate the yeast location.  
24  
25 134 Each purified strain from our collection (Table 1) has been integrated to the 'Souchothèque de  
26  
27 135 Bretagne' culture collection  
28  
29 136 (<http://www.ifremer.fr/souchotheque/internet/htdocs/generique.php?pagebody=catalogue.php>  
30  
31 137 ) and are available with an accession number associated to their GenBank number.  
32  
33 138

### 34 35 139 ***Physiological characterization and statistical analysis***

36  
37 140 All experiments were done in triplicate. The yeasts were grown in liquid GYPS broth media.  
38  
39 141 The effect of temperature on growth was determined at 5°C, 15°C, 25°C and 35°C at 30 g.L<sup>-1</sup>  
40  
41 142 sea salts. The effect of salinity was analyzed modifying sea salts concentrations in media from  
42  
43 143 0 to 60 g.L<sup>-1</sup> with steps of 15 g.L<sup>-1</sup> at optimal temperature for each strain. Optical densities  
44  
45 144 (OD) were measured at 600 nm with Nanocolor 100D (Macherey-Nagel, Hoerd, France) at  
46  
47 145 17, 22, 25 and 28 hours of growth under each condition of salinity and temperature.  
48  
49 146

### 50 51 147 ***DNA extraction and 26S rDNA sequencing***

52  
53 148 DNA of each strain was extracted using FastDNA Spin Kit (MP Biomedicals, Illkirch,  
54  
55 149 France) specific for fungi and yeasts. Amplifications of the D1/D2 region of 26S rDNA were  
56  
57 150 carried out with rDNA primers ITS5 (5'-GGA AGT AAA AGT CGT AAC AAG-3'), LR6  
58  
59 151 (5'-CGC CAG TTC TGC TTA CC-3'), NL1 (5'-GCA TAT CAA TAA GCG GAG GAA  
60  
152 AAG-3') and NL4 (5'-GGT CCG TGT TTC AAG ACG G-3') as described by Gadanho &  
153  
154 Sampaio (2005). All PCR reactions were performed in 20 µL reaction volumes containing 19  
µL of 1X PCR Buffer (Promega), 2 mM of MgCl<sub>2</sub>, 0.2 mM of each dNTPs (Promega), 0.6

1  
2  
3 155  $\mu\text{M}$  of primers (forward and reverse), 1.25 U of Taq DNA Polymerase (Promega) and  $1\mu\text{L}$  of  
4  
5 156 DNA. The polymerase chain reactions were performed on PTC-200 (Biorad, France). The  
6  
7 157 amplification consisted in an initial denaturation step at  $94^{\circ}\text{C}$  for 2 min, followed by 30  
8  
9 158 iterations of 15 sec at  $94^{\circ}\text{C}$ , 30 sec at  $54^{\circ}\text{C}$ , 1 min at  $72^{\circ}\text{C}$  and a final extension step of 2 min  
10  
11 159 at  $72^{\circ}\text{C}$ . A negative control with sterile distilled water replacing DNA was added. Two kinds  
12  
13 160 of amplification were generated using ITS5-NL4 and NL1-LR6 primers. The amplified DNA  
14  
15 161 fragments were separated by electrophoresis in 0.8% agarose (w/v) gel (Promega) in 0.5X  
16  
17 162 Tris-Borate-EDTA (TBE) Buffer at 90 V for 1h and stained with ethidium bromide. A  
18  
19 163 molecular size marker was used for reference (Lambda DNA/EcoR1+Hind III Markers,  
20  
21 164 Promega). The DNA banding patterns were visualized under UV transillumination and  
22  
23 165 picture files were generated using Gel-Doc 2000 (Biorad, France).

24 166 The sequencing of the D1/D2 region of the 26S rDNA was then realized using NL1 on the  
25  
26 167 ITS5-NL4 fragments and NL4 on the NL1-LR6 fragments. The sequences were obtained by  
27  
28 168 “Big Dye Terminator” technology (Applied Biosystems). This work was done at  
29  
30 169 “Biogenouest” sequencing facility in the “Station Biologique de Roscoff” ([www.sb-roscoff.fr](http://www.sb-roscoff.fr)).  
31  
32 170

### 33 171 *Phylogenetic analyses*

34  
35  
36 172 Sequences were edited and cleaned using Sequencher v 4.8 (Gene Codes). Sequences were  
37  
38 173 then imported to MEGA 4.0 software (Tamura *et al.*, 2007). Each sequence was analyzed in  
39  
40 174 order to find GenBank sequences with close BLAST-N hits (Altschul *et al.*, 1990).  
41  
42 175 Similarities between sequences were assessed using pairwise distance calculation with  
43  
44 176 MEGA 4.0. The sequences were trimmed to ensure that all sequences had the same start and  
45  
46 177 end-point. All the D1/D2 regions of the 26S rDNA sequences were aligned using ClustalW  
47  
48 178 v.1.83 (Thompson *et al.*, 1994). After visual checking and manual curation, an alignment  
49  
50 179 composed of 62 taxa and 579 characters was analysed for the Bayesian estimation of  
51  
52 180 phylogeny using MrBayes v.3.1.2 software (Ronquist and Huelsenbeck, 2003). A two-million  
53  
54 181 generation option has been set to run the Metropolis-coupled Monte Carlo Markov Chain  
55  
56 182 method (*mcmc*). After generation 2 000 000, the standard deviation of split frequencies was  $P$   
57  
58 183 = 0.005997 indicating that a convergence had occurred (P-value of  $< 0.05$ ). The alignment  
59  
60 184 was analysed using MODELTEST v.3.7 (Posada and Crandall, 1998), in order to obtain the  
185  
186 186 more realistic evolutionary model used for phylogenetic analyses (GTR + G model; gamma-  
distribution shape parameter = 0.3978). Phylogeny was then evaluated using two different

1  
2  
3 187 methods: (i) Bayesian inference with MrBayes v.3.1.2 analysis using 2 000 000 generations  
4  
5 188 and the *mcmc* method. The tree search included two *mcmc* searches with four chains (setting  
6  
7 189 default temperature for heating the chains) and a sampling frequency of 100 generations. A  
8  
9 190 'burnin' of 5000 (25% of the 2 000 000 generations/100 sample frequency) was set in order to  
10  
11 191 exclude the first 5000 trees generated. (ii) Maximum likelihood with 100 bootstrap iterations  
12  
13 192 using PHYML (Guindon *et al.*, 2005) and the parameters obtained with MODELTEST v.3.7.  
14  
15 193 The final phylogenetic tree topology was realized using MrBayes v.3.1.2 analysis results.  
16  
17 194 Nodes in the tree show Bayesian posterior probabilities and ML bootstraps respectively.

### 18 195 ***Fluorescent probe design and evaluation***

19  
20 196  
21  
22 197 For the detection of yeasts isolated from deep-sea vent animals by FISH, we designed  
23  
24 198 oligonucleotide probes using the Primrose software ([http://www.bioinformatics-  
25  
26 199 toolkit.org/Primrose/index.html](http://www.bioinformatics-toolkit.org/Primrose/index.html)) as described by Ashelford *et al.* (2002) using a set of high-  
27  
28 200 quality, full-length rRNA sequences of probe target organisms. The PrimRose design tool  
29  
30 201 permitted to produce oligonucleotide probes for the three principal clusters of our collection  
31  
32 202 (Table 3). These probes exhibited no mismatches with the target organisms but exhibited  
33  
34 203 mismatches with the next most similar sequences in the GenBank database proving that the  
35  
36 204 designed probes were *in silico* highly specific. The target sites of newly designed probes were  
37  
38 205 checked for accessibility using the prediction maps based on the 26S rRNA of *Saccharomyces*  
39  
40 206 *cerevisiae* (Inacio *et al.*, 2003). Each probe was in a relative accessible area of the 26S rRNA  
41  
42 207 secondary structure (Fig S1). As it was not possible to test the probes with culture isolates that  
43  
44 208 exhibited zero or one mismatch with the probes, we used an alternative method and tested the  
45  
46 209 probes against all strains from our collection displaying two or more mismatches with the  
47  
48 210 oligonucleotides. All newly designed probes were labelled at the 5' terminus with the  
49  
50 211 fluorescent marker Cy3. All probes were synthesized by (Proligo, France) and stored in sterile  
51  
52 212 distilled water at -20°C. The newly designed probes were checked under *in situ* conditions  
53  
54 213 with target and non-target species. The universal probe Euk516-Fluorescein (5'-  
55  
56 214 ACCAGACTTGCCCTCC-3'; Amann *et al.*, 1995) and the non-Euk516-Cy3 (5'-  
57  
58 215 CCTCCCGTTCAGACCA-3') probes were used as positive and negative control respectively.  
59  
60 216 The average cell brightness was measured using different formamide concentrations from 0 to  
217  
218 80% with steps of 10%. Systematic evaluation of the signal intensities was done by recording  
219  
images of independent visual fields (encompassing at least 100 cells), followed by digital  
image analysis using the daime software (Daims *et al.*, 2006). During this step, the intensities

1  
2  
3 220 of the image pixels analyzed enable determination of single cell fluorescence in relative units  
4  
5 221 (RU).  
6

### 7 222 *Fluorescence In Situ Hybridization*

8  
9  
10 223 *On environmental samples.* Interior branchiostegites of *Rimicaris exoculata* shrimps and  
11 224 byssus of *Bathymodiolus azoricus* mussels were processed for FISH analyses. Following  
12 225 harvest and dissections, animal subsamples were fixed with 4% paraformaldehyde solution in  
13 226 phosphate-buffered saline (PBS) for 3 hours at 4°C in a dark room. After fixation, tissues  
14 227 were washed three times with PBS and stored at -20°C in a storage buffer containing PBS and  
15 228 96% ethanol (1:1).  
16  
17  
18  
19  
20

21  
22 229 *On membrane filters.* The seawater surrounding shrimps (MoMAR08, Rainbow) was sampled  
23 230 in 5 L sterile sampling bags using a peristaltic pump. Immediately after dives, seawater  
24 231 samples for *in situ* hybridizations were mixed with 3 % formaldehyde (final concentration) 2  
25 232 hours at 4°C. Fixed seawater was then filtered on polycarbonate membranes 0.22 µm  
26 233 (Nuclepore®, 47 mm diameter; Whatman, Maidstone, Kent, UK) and rinsed with a PBS 2X -  
27 234 sterile seawater (v:v) buffer. Then filters were dehydrated using ethanol series (50 %, 80 %  
28 235 and absolute, 3 min each). Dried filters were stored at -20°C until hybridization treatments.  
29 236 Three membranes were treated in this study. The filtered volume was 0.8 L for membrane A,  
30 237 1 L for membrane B and 1.5 L for membrane C. The filtered seawater on membranes A and B  
31 238 was from the same sample.  
32  
33  
34  
35  
36  
37  
38  
39

40 239 The samples (environmental samples and membrane filters) were cut in squares and paste  
41 240 with one drop of 0.2% low-gelling point agarose (35-40°C) on slides (Menzel-Glaser,  
42 241 Germany). All slides were then dipped in 0.2% agarose and air dried. Samples were then  
43 242 subjected to dehydration with increasing concentrations of ethanol (50, 80, and 96%, for 3  
44 243 min each). Working solutions of probes had a concentration of 30 ng of DNA per liter. The  
45 244 hybridization buffer, containing 0.9 M NaCl, 20 mM Tris-HCl (pH 7.2), 0.03% SDS, and 0,  
46 245 10, 20, 30, 40, 50, 60, 70 and 80% formamide, and the fluorescent probe were gently mixed  
47 246 in a ratio of 10:1 (vol/vol) to get a final oligonucleotide concentration of 3 ng per liter. For  
48 247 hybridization, slides were placed in sampling tubes and incubated at 46°C in the dark for  
49 248 exactly 3 hours. Following hybridization, the slides were washed with prewarmed washing  
50 249 buffer (20 mM Tris/HCl, 5 mM EDTA (pH 8.0) and 900, 450, 215, 102, 46, 18, 5, 0.6 and 0  
51 250 mM NaCl corresponding respectively to 0, 10, 20, 30, 40, 50, 60, 70 and 80% formamide  
52 251 stringencies) for 20 min at 48°C. Slides were rinsed with double-distilled water, air dried,  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 252 DAPI stained (final concentration 1 µg/ml) and mounted with the antifading reagent Citifluor  
4 253 AF 2 (Citifluor, France) before observations under fluorescent microscope.

## 8 254 **Results**

### 10 255 **Yeast isolation**

11  
12  
13 256 Yeasts were not found in all the studied sites as shown in Table 1. No yeast was isolated from  
14 257 samples collected during HERO (on the East Pacific Rise at Elsa site), DIVANAUT2 and  
15 258 MARVEL (Menez-Gwenn and Lucky Strike) cruises or at TAG site during the EXOMAR  
16 259 oceanographic cruise. The hydrothermal site that yielded the highest number of isolates was  
17 260 clearly Rainbow (29 isolates out of 32 strains). Rainbow is also the site where the highest  
18 261 number of samples was processed (97/210). The yeast collection obtained from deep-sea  
19 262 samples raised thirty-two isolates that could be divided in pigmented yeasts (18) and non-  
20 263 pigmented yeasts (14). Pigmented yeasts consisted widely of red-pigmented yeasts (16),  
21 264 black-pigmented yeast (1) and brown-pigmented yeast (1).

22  
23  
24 265 Regarding yeast isolation versus type of substrate, strains were obtained mostly from  
25 266 hydrothermal shrimps *Rimicaris exoculata* (11), *Chorocaris chacei* (3), *Mirocaris fortunata*  
26 267 (1) and from hydrothermal mussels *Bathymodiolus azoricus* (7). Carbonate colonization  
27 268 modules deployed for 1 year near Rainbow vent yielded a few yeasts (4); sponges led to the  
28 269 isolation of three yeasts. Finally, seawater, gastropods and coral samples permitted to obtain  
29 270 one strain each (Table 1). Those results indicate that yeasts were much more associated with  
30 271 animals rather than mineral substrates. Statistical distribution tests have been performed in  
31 272 order to find out the distribution type of yeasts in hydrothermal sites. The variance to mean  
32 273 ratio ( $s^2/m$ ) was calculated for each site (Cancela da Fonseca, 1966). Values of  $s^2/m$   
33 274 significantly different of 1 corresponds with  $(s^2/m) - 1 > 2(2n/(n - 1)^2)^{1/2}$  and were obtained  
34 275 only for Rainbow site. For this hydrothermal site, an aggregate distribution was observed  
35 276 ( $s^2/m=1.32$ ) indicating that the culturable yeasts isolated were located in specific niches in  
36 277 this hydrothermal site (mainly shrimps and mussels).

37  
38  
39 278  
40 279 During the MoMARDREAM-Naut cruise, dissections of body components were processed  
41 280 for all shrimps (branchiostegites, scaphognathites, exopodites, gills, stomach and digestive  
42 281 tract) and mussels (interior and external faces of shells) to investigate the localization of  
43 282 yeasts in deep-sea animals. For shrimps, a large majority of strains were grown from the inner  
44 283 side of the branchiostegites that can be divided in 3 different compartments: (a) an antero-

1  
2  
3 284 ventral area, which was relatively clear; (b) a posterior area, which always remained light  
4  
5 285 beige; (c) an antero-dorsal area with an intensely rusty coloration (for schematic views, see  
6  
7 286 Zbinden *et al.*, 2004; Corbari *et al.*, 2008). Yeast isolates resulted from this study were all  
8  
9 287 cultivated from the antero-dorsal area characterized by high amounts of minerals and a dense  
10  
11 288 bacterial mat.  
12

13  
14 290 The yeasts were also isolated from *Bathymodiolus azoricus* (7) during the MoMADREAM-  
15  
16 291 Naut oceanographic cruise (Table 1). Most of them (6) were cultivated from external face of  
17  
18 292 the mussel shells and more precisely from the byssus that is a network of filaments allowing  
19  
20 293 attachment to rocks. This tangle gathers a lot of particles and organic matter in decomposition  
21  
22 294 (personal observation). Only one yeast was isolated from the interior of a mussel (Mo32).  
23

### 24 295 25 296 **Physiological analysis**

26 297  
27  
28 298 Three categories of strains were identified (Table 2) based on the definition of halotolerant  
29  
30 299 and halophilic microorganisms (Margesin & Schinner, 2001; Kushner, 1978). Non halophiles  
31  
32 300 are strains with maximal growth without sea salts and a decreasing growth rate with increased  
33  
34 301 sea salts concentration in media. Halotolerant yeasts are strains able to grow in the absence as  
35  
36 302 well as in the presence of salt. Halophiles required salt for an optimal growth. Regarding  
37  
38 303 halophily, optimal salinities, optimal temperatures and OD measurement, 9 physiological  
39  
40 304 groups were defined. Most of the isolated strains were non halophiles (23 strains) and  
41  
42 305 halotolerant (2 strains, maximal OD at 30 g/l sea salts) growing efficiently at an optimal  
43  
44 306 temperature of 25°C. Four strains had poor maximal growth at 25°C including 1 non  
45  
46 307 halophile, 2 halotolerant (maximal OD at 30 and 60 g/l sea salts) and 1 halophile (maximal  
47  
48 308 OD at 30 g/l sea salts). Three strains had maximal and efficient growth at 35°C, including 1  
49  
50 309 non halophile, 1 halotolerant (maximal OD at 45 g/l sea salts) and 1 halophile (maximal OD  
51  
52 310 at 30 g/l sea salts).

### 53 311 54 312 **Identification**

55 313  
56 314 For species identification, a sequence analysis of the D1/D2 domain of the 26S rRNA gene  
57  
58 315 was done (Fig 1). A total of 12 phylotypes was found among the collection of yeasts isolated  
59  
60 316 from deep-sea hydrothermal vents. Eleven phylotypes could be assigned to a known yeast  
317 species and one represents a new yeast species.

1  
2  
3 318  
4  
5 319 Within *Basidiomycota*, the *Sporidiobolales* order was the dominant cluster composed of 16  
6  
7 320 strains. A majority of strains (Ex2, Ex3, Ex4, Ex5, Ex6, Ex7, Ex9, Ex11, Ex12, Mo32, Mo35  
8  
9 321 and Mo37) was identified as *Rhodotorula mucilaginosa* (100% similarity). A large majority  
10  
11 322 of *R. mucilaginosa* was isolated from deep-sea shrimps (14) and the others from deep-sea  
12  
13 323 mussels (2). As member of the *Sporidiobolales* order, isolates affiliated to *Rhodospiridium*  
14  
15 324 *diobovatum* were also isolated (Mo24, Mo33 and Mo38) with 100% similarity. These 3  
16  
17 325 strains were isolated respectively from *Rimicaris exoculata* exuviae in decomposition on  
18  
19 326 smocker rocks, *Bathymodiolus azoricus* and a sponge. One strain isolated from *R. exoculata*  
20  
21 327 was identified as *Sporobolomyces roseus* based on 26S rRNA genes (Mo22) with 100%  
22  
23 328 similarity with the reference strain. Four strains (Mo26, Mo27, Mo28 and Mo29) were  
24  
25 329 affiliated to the *Filobasidiales* order and identified as *Cryptococcus uzbekistanensis* (100%  
26  
27 330 similarity). These four strains were all isolated from a carbonate colonization module. Finally,  
28  
29 331 one isolate (Mo36) from *B. azoricus* mussel was identified as *Leucosporidium scottii* in the  
30  
31 332 *Leucosporidiales* order.

30 333 The *Ascomycota* phylum gathered 9 strains belonging to the *Saccharomycetales* order. Within  
31  
32 334 this order, 4 strains (Mo20, Mo21, Mo40 and Bio2) isolated respectively from *R. exoculata*,  
33  
34 335 *Mirocaris fortunata*, a deep-sea coral and the gills of the gastropod *Ifremeria nautilei* were  
35  
36 336 identified as *Debaryomyces hansenii* (100% similarity). *Candida atlantica* isolates were  
37  
38 337 found in *R. exoculata* exuviae in decomposition (Mo25) and *B. azoricus* (Mo31). One strain  
39  
40 338 isolated from a deep-sea sponge (Ex15) was identified as *Pichia guilliermondii* (100%  
41  
42 339 similarity). Finally, among the *Saccharomycetales* order, one strain was identified as *Candida*  
43  
44 340 *viswanathii* (Bio1) with 100% similarity. One halophilic strain (Mo39) isolated from a deep-  
45  
46 341 sea coral represents a new species in the *Candida* genus and thus was identified as *Candida*  
47  
48 342 sp. This strain has 95% similarity with the reference sequence of *Candida atmosphaerica* (23  
49  
50 343 mismatches on 505 bp). Mo30 isolated from *Bathymodiolus azoricus* was identified as  
51  
52 344 *Phaeotheca triangularis* (mitosporic *Ascomycota*) with 100% similarity. In the *Dothideales*  
53  
54 345 order, one strain (Mo34) isolated from *Bathymodiolus azoricus* was identified as *Hortaea*  
55  
56 346 *werneckii* with 99.98% similarity (one mismatch on 560bp).

54 347 Sequencing of the 26S rRNA genes indicated the presence of *Ascomycota* and *Basidiomycota*  
55  
56 348 in our culture collection. In term of quantity, the phylum *Basidiomycota* (21) was two times  
57  
58 349 higher than the *Ascomycota* (11). In term of species richness, ascomycetous yeasts belonged  
59  
60 350 to 7 different clusters while basidiomycetous yeasts belonged to 5 clusters.

351

## 352 **Fluorescence *in situ* hybridizations**

353 We processed numerous assays to detect fungi on deep-sea hydrothermal vent animal samples  
354 using different existing fluorescent probes from different databases. The Euk516-Cy3 probe  
355 gave positive results on pure cultures but strong background fluorescence on hydrothermal  
356 samples led to the renouncement of its use. The probe MY1574 targeting *Eumycota*  
357 organisms (Baschien *et al.*, 2008) showed very weak fluorescence on pure cultures. Thus, we  
358 decided to design our own probes (Table 3) based on our culture collection that was divided  
359 in 3 main clusters: **MitoFilo** (*Cryptococcus* / Mitosporic *Filobasidiales* order), **MitoSporidio**  
360 (*Rhodotorula*, *Rhodosporidium* / Mitosporic *Sporidiobolales* order) and **Sacch**  
361 (*Debaryomyces*, *Pichia* / *Saccharomycetales* order). The probes designed revealed a strong  
362 specificity for the target organisms. The optimal conditions for the *in situ* hybridization  
363 protocol use stringent conditions of 20% formamide (Fig S1).

364 Our aim was to check the applicability of the FISH method to the *in situ* detection of yeasts in  
365 deep-sea hydrothermal fauna samples. Hydrothermal body components of endemic shrimps  
366 (*Rimicaris exoculata*) and mussels (*Bathymodiolus azoricus*) were fixed for FISH  
367 experiments directly after dissection. The pieces of shrimps and mussels that gave the higher  
368 number of fungi isolation (interior branchiostegites of shrimps and byssus of mussels) were  
369 analyzed for yeast cell fluorescence. Although shrimp and mussel samples from Rainbow site  
370 led to the highest rate of isolation, no FISH signal was ever observed. The FISH detection  
371 limit of  $10^3$ - $10^4$  target cells per ml is relatively high (Daims *et al.*, 2005) and thus, the absence  
372 of FISH signals does not necessarily mean that the target organisms were not present in the  
373 samples.

374 To test this hypothesis, several volumes of water were concentrated on polycarbonate  
375 membrane filters to yield sufficient cells for FISH experiments with these new probes.  
376 Membrane filters were embedded in low gelling-point agarose to minimize cell loss. Yeast  
377 cells could be visualized in a low quantity on these membrane filters (Fig 2). Such results are  
378 another evidence of the yeast cells presence in hydrothermal vents but at low concentration.  
379 Using FISH on membrane filters, yeast cells detected were affiliated to 3 genera:  
380 *Rhodosporidium*, *Rhodotorula* and *Cryptococcus*.

## 382 **Discussion**

383

1  
2  
3 384 ***Occurrence of yeasts in deep-sea hydrothermal vents***  
4

5  
6 385 In this study, the main aim was to isolate yeast strains from deep-sea hydrothermal endemic  
7  
8 386 fauna knowing that yeasts can be isolated from seawater surrounding hydrothermal fauna  
9  
10 387 (Gadanhó and Sampaio, 2005). Yeast isolation was successful even if the retrieved species  
11  
12 388 richness was relatively low. Thirty-two strains were isolated mostly from *Rimicaris exoculata*  
13  
14 389 shrimps. The association with shrimps is probably favorable for yeasts that could benefit from  
15  
16 390 nutrients due to the water circulation in the gill chamber. Most of our strains were isolated  
17  
18 391 from the Rainbow hydrothermal site which confirms previous results (Gadanhó and Sampaio,  
19  
20 392 2005). The Rainbow hydrothermal field hosted in ultramafic rocks is a unique vent enriched  
21  
22 393 in CH<sub>4</sub>, H<sub>2</sub>, CO, Fe and depleted in H<sub>2</sub>S (Charlou *et al.*, 2002). The high yeast isolation ratio  
23  
24 394 may indicate that yeasts thrive in hydrothermal sites depleted in H<sub>2</sub>S. The isolation rate of  
25  
26 395 non-pigmented yeasts on sulfur-free media significantly higher than those on sulfur-based  
27  
28 396 media in a previous study (Gadanhó & Sampaio, 2005) support such hypothesis.

29 397  
30 398 Several yeasts were also isolated from mussels and more precisely from the byssus  
31  
32 399 constituted of filaments with a high concentration of minerals and organic matter. These  
33  
34 400 yeasts may have a role in the decomposition of organic material entrapped in mussel byssi in  
35  
36 401 deep-sea vents. These results seem promising as they confirm the data obtained in previous  
37  
38 402 studies and suggest that yeasts may interact with deep-sea hydrothermal vent fauna.

39 403  
40 404 ***Pattern of the culturable yeast communities***

41 405 *New species.*

42  
43 406 The yeast that was firstly isolated from stomach of a marine fish was described as *D. hansenii*  
44  
45 407 and deposited in the Centraalbureau voor Schimmelcultures (CBS 5307) database. In a recent  
46  
47 408 paper, based on the intergenic spacer (IGS) region of the rRNA gene, this strain was re-  
48  
49 409 evaluated as *Candida sp.* (Nguyen *et al.*, 2009). This strain is identical to another one isolated  
50  
51 410 from deep-sea hydrothermal vent waters and annotated MARY089 (Gadanhó and Sampaio,  
52  
53 411 2005). These two strains isolated from different marine environments were finally reported as  
54  
55 412 a single new undescribed species within the *Candida* genus. In our collection, strain Mo39,  
56  
57 413 isolated from deep-sea coral near Rainbow hydrothermal vents (Table 1), has the same 26S  
58  
59 414 rRNA gene sequence as CBS 5307 and MARY089. Mo39 is halophilic and thus supposed to  
60  
415 be an autochthonous marine yeast species. This new ecotype can be characterized as an  
416  
obligate marine yeast and its complete description is currently under progress.

1  
2  
3 417  
4  
5 418 *Known species*  
6  
7 419 Two strains (Mo25 and Mo31) isolated from *Rimicaris exoculata* and *Bathymodiolus azoricus*  
8  
9 420 samples were identified as *Candida atlantica*. This result seems in keeping with previous  
10  
11 421 published reports that have isolated this species from coastal seawater in the South of  
12  
13 422 Portugal (Gadanho *et al.*, 2003) and in deep-sea hydrothermal vent waters (Gadanho and  
14  
15 423 Sampaio, 2005). The very first *C. atlantica* strain was isolated from shrimp eggs in the North  
16  
17 424 Atlantic Ocean (Siepmann and Höhnk, 1962). *C. atlantica* seemed to be a marine obligate  
18  
19 425 yeast and some interactions with shrimps seemed to occur. Our physiological analysis has  
20  
21 426 revealed that Mo25 and Mo31 were non-halophiles, which does not mean that they are unable  
22  
23 427 to grow in marine environments. They may have a role in deep-sea environments in  
24  
25 428 interaction with endemic crustaceans even if they are not in optimal growth conditions. One  
26  
27 429 isolate (Bio1) isolated from seawater surrounding mussels at Lau Basin in the South-West  
28  
29 430 Pacific was clearly identified as *Candida viswanathii*. Kohlmeyer & Kohlmeyer (1979)  
30  
31 431 characterized this yeast as marine facultative. More recently, *C. viswanathii* was isolated from  
32  
33 432 a shrimp (*Peneaus braziliensis*) in the Gulf of Mexico. Its synonym, *Candida lodderae* was  
34  
35 433 recently reported in deep-sea hydrothermal vent waters at Rainbow site (Gadanho and  
36  
37 434 Sampaio, 2005) and characterized as the most abundant yeast.

38  
39 435  
40  
41 436 *Leucosporidium scottii* isolates (Mo36) were retrieved only in the oceanic regions close to  
42  
43 437 Antarctica and are known to be psychrophilic and probably autochthonous marine species  
44  
45 438 (Lachance and Starmer, 1998). Such strains known for their presence in cold polar marine  
46  
47 439 environments could be another evidence that confirms the hypothesis of global exchanges  
48  
49 440 from polar environments to deep-sea vents based on results from bacteria (Maruyama *et al.*,  
50  
51 441 2000) and filamentous fungi (Burgaud *et al.*, 2009). *Hortaea werneckii* (Mo34) was  
52  
53 442 characterized as halophilic in our physiological study. This is not surprising as this black  
54  
55 443 yeast-like fungus was characterized as halophilic or extremely halotolerant in different studies  
56  
57 444 (Gunde-Cimerman *et al.*, 2000; Kogej *et al.*, 2005) where it was frequently isolated from  
58  
59 445 hypersaline waters of solar salterns. In a molecular survey, *H. werneckii* was identified (based  
60  
446 on internal transcribed spacers and 5.8 S rRNA gene) in deep-sea methane seep sediments at a  
447 depth of 2965 meters (Lai *et al.*, 2007). *Phaeotheca triangularis* (Mo30) was also frequently  
448 isolated from salted environments (Gunde-Cimerman *et al.*, 2000) and characterized as  
449 halophile. This confirmed previous results on *P. triangularis* showing a better growth with  
450 5% additional salts (Zalar *et al.*, 1999). In our study, Mo30 was characterized as halotolerant

1  
2  
3 451 with 4.5% sea salts optimal concentration and thus hypothesized as marine adapted yeast.  
4  
5 452 This is the first report about the presence of *Phaeotheca triangularis* at deep-sea vents.  
6  
7 453 Mo22 is described as *Sporobolomyces roseus*. The genus *Sporobolomyces* is composed of  
8  
9 454 strains mainly isolated from the pylophane (Bai *et al.*, 2002). However, a previous study has  
10  
11 455 proved that strains of the genus *Sporobolomyces* are frequently isolated from marine  
12  
13 456 ecosystems and the frequency of isolation increases when distance from the coastline and  
14  
15 457 depth increase (Hernandez-Saavedra *et al.*, 1992). Moreover, yeasts from this genus were  
16  
17 458 found in benthic invertebrates collected from deep-sea floor in the Pacific Ocean (Nagahama  
18  
19 459 *et al.*, 2001b). Our strain was isolated from a deep-sea hydrothermal shrimp in the Atlantic  
20  
21 460 Ocean and characterized as halotolerant with an optimal salinity of 6% sea salts. This may  
22  
23 461 indicate that yeasts of this genus are also able to live in deep-sea vents and interact with  
24  
25 462 endemic crustaceans.  
26  
27 463

26 464 A previous study of yeasts in oceanic environments (Fell, 1976) reported that yeast  
27  
28 465 communities appeared to be constituted of ubiquitous and endemic species. Typical  
29  
30 466 ubiquitous strains were the ascomycetous yeast *Debaryomyces hansenii* and the  
31  
32 467 basidiomycetous ones *Cryptococcus* and *Rhodotorula*. Kohlmeyer and Kohlmeyer (1979)  
33  
34 468 confirmed this statement and characterized these genera mainly as facultative marine yeasts.  
35  
36 469 Some of these results, especially for *Rhodotorula* yeasts showing a strong ubiquity, were  
37  
38 470 confirmed based on their presence in several habitats such as deep-sea vents (Gadanhó and  
39  
40 471 Sampaio, 2005), deep-sea sediments (Nagahama *et al.*, 2001b), coastal waters (Gadanhó *et*  
41  
42 472 *al.*, 2003; 2004) and oligotrophic lakes (Libkind *et al.*, 2003). Our results confirm their  
43  
44 473 ubiquity and indicate that these strains seem to be allochthonous. Strain Ex15 identified as  
45  
46 474 *Pichia guilliermondii* has also been characterized as non halophile and may be another  
47  
48 475 allochthonous yeast strain as reported by Kohlmeyer and Kohlmeyer (1979).  
49  
50 476

49 477 The members of the genus *Rhodospiridium* have been characterized as non halophiles (Mo24  
50  
51 478 and Mo33) and halotolerant (Mo38). Based on previous reports, this genus seemed to be  
52  
53 479 restricted to marine environments (Gadanhó and Sampaio, 2005). *R. diobovatum* in deep-sea  
54  
55 480 vents seemed to be able to colonize different substrates (shrimps, mussels and sponges). The  
56  
57 481 isolation of a strain from shrimp exuviae in decomposition may indicate a role as a recycler of  
58  
59 482 organic material and so a probable implication in carbon cycle in deep-sea environments.  
60  
61 483

61 484 ***Adaptation to marine conditions***

1  
2  
3 485 The isolation of culturable yeasts led to an old question about marine yeasts “Are there any  
4  
5 486 indigenous marine yeasts ?” (Kohlmeyer & Kohlmeyer, 1979) and to the resulting question  
6  
7 487 “Which are the indigenous species ?”. Based on our results, one can suggest that halophilic  
8  
9 488 strains are marine indigenous yeasts and that others, halotolerant and non-halophiles, are  
10  
11 489 ubiquitous terrestrial strains present in deep-sea waters due to sedimentation or other natural  
12  
13 490 or anthropogenic phenomena. But almost all yeast species can grow well in media with NaCl  
14  
15 491 concentrations exceeding those normally present in the sea (Kohlmeyer & Kohlmeyer, 1979).  
16  
17 492 Few yeast species with a physiological dependence on NaCl or other seawater components  
18  
19 493 have been reported (Nagahama, 2006b). Thus, our results appeared in good agreement with  
20  
21 494 such statements. Only 2 strains described as halophiles (Mo34 and Mo39) in our study can be  
22  
23 495 described as obligate marine yeasts.

### 24 496 *FISH observations*

25 497  
26  
27 498 FISH using labeled oligonucleotide probes targeting rRNA has been used as a powerful  
28  
29 499 technique for assessing both microbial identity and spatial distributions *in situ* in complex  
30  
31 500 environmental contexts (Yang *et al.*, 2008). Our results indicate a very low-level of yeasts at  
32  
33 501 deep-sea vents. As a first conclusion, regarding diversity and quantification (added to  
34  
35 502 previous results of Gadanho and Sampaio, 2005), it seems that yeasts at deep-sea vents  
36  
37 503 represent a minor community that might not be major actors in biogeochemical cycles.  
38  
39 504 However, fluorescent signals are correlated to the cellular content of ribosomes and  
40  
41 505 consequently to the microbial growth rates. Recently, the detection limits of conventional  
42  
43 506 FISH with Cy3-labeled probe EUB338 were found to be approximately 370 16S rRNA  
44  
45 507 molecules per cell for *Escherichia coli* hybridized on glass microscope slides and 1,400 16S  
46  
47 508 rRNA copies per *E. coli* cell in environmental samples (Hoshino *et al.*, 2008). So, in addition  
48  
49 509 to a low concentration of yeast cells, low detection of yeasts may be caused by low ribosome  
50  
51 510 content of most yeasts in the deep-sea environment due to low-level metabolic activities of  
52  
53 511 yeasts living under extreme environmental abiotic factors (high hydrostatic pressure, low  
54  
55 512 temperatures,...). Our attempts to cultivate the yeast strains resulted from this study under  
56  
57 513 elevated hydrostatic pressure have been successful, but ribosomal activities were lower under  
58  
59 514 high hydrostatic pressure than at atmospheric pressure. Such results may account for the low  
60  
515 fungal detection using FISH (unpublished data). Consequently, care must be taken when  
516  
dealing with diversity and biomass estimations when using FISH alone.



1  
2  
3 517 The quantification of yeasts using FISH has been impossible due to a non homogeneous  
4  
5 518 repartition of microorganisms on filters. Moreover, bacterial and yeast cells were only visible  
6  
7 519 in some regions of the filters without minerals due to strong autofluorescence. However we  
8  
9 520 can say that yeast concentrations are really low, as shown by the only few cells visualized  
10  
11 521 after filtration of seawater surrounding shrimps. This result is in keeping with the relatively  
12  
13 522 low diversity revealed by Gadanho and Sampaio (2005) ranging from 0 to 10 cfu/L for pink  
14  
15 523 yeasts and from 0 to 6000 cfu/L for non-pigmented yeasts. To better analyze the fungal  
16  
17 524 presence in deep-sea animals, one could work with phylum-specific probes on histological  
18  
19 525 sections of animals and use the CARD-FISH (Amann & Fuchs, 2008) or the DOPE-FISH  
20  
21 526 (Stoecker *et al.*, 2010) methods to amplify probe signals.

22 527 These data raise emerging questions regarding the ecological role of such microorganisms in  
23  
24 528 deep-sea vents and about the old question of the ubiquity or endemism of those strains. Yeasts  
25  
26 529 at deep-sea vents may be facultative parasites or opportunistic pathogens of endemic deep-sea  
27  
28 530 animals as it has already been hypothesized in previous works (Van Dover *et al.*, 2007;  
29  
30 531 Burgaud *et al.*, 2009). However, a role in the decomposition of abundant organic material may  
31  
32 532 occur.

33 533 Considering all the results obtained, we can say that yeasts may seem to interact with deep-  
34  
35 534 sea hydrothermal endemic fauna even if the density is low. These yeasts are mainly composed  
36  
37 535 of ubiquitous species but obligate marine yeasts have also been harvested. However, the  
38  
39 536 results obtained using *in situ* hybridization have allowed us to visualize these ubiquitous species  
40  
41 537 showing that they are able to live and grow in deep-sea hydrothermal vents. Yeasts associated  
42  
43 538 with endemic animals in deep-sea vents may be exposed to favorable conditions and could  
44  
45 539 benefit from a stable source of nutrients (Nagahama *et al.*, 2001b). Yeasts were reported from  
46  
47 540 dead and healthy individuals which may also indicate their facultative saprophytism and so  
48  
49 541 emphasize the wide role of fungi in the decomposition of organic matter from terrestrial  
50  
51 542 environments to deep-sea hydrothermal vents. Even if yeasts were isolated from animal body  
52  
53 543 components, they were not visualized using FISH. To better understand the interaction with  
54  
55 544 animals and fungi in deep-sea vents, we need to work on tissues as in Van Dover *et al.* (2007)  
56  
57 545 and also with probes specific to fungal phyla (*Ascomycota*, *Basidiomycota* and  
58  
59 546 *Chytridiomycota*). In conclusion, several questions regarding the role of yeasts in deep-sea  
60  
547 hydrothermal vents and the endemism or ubiquity of the isolated yeasts remain a difficult task  
548  
549 548 without clear answers. Their culture under high hydrostatic pressures would be an interesting  
549 study to better characterize their lifestyle and role at deep-sea vents.

1  
2  
3 5504  
5 551 **Acknowledgements**6  
7 552

8  
9 553 We thank the chief scientists of the BIOLAU, DIVANAUT2, HERO, MARVEL, EXOMAR  
10 554 and MoMARDREAM-Naut cruises, the pilots and support crews of the oceanographic vessels  
11 555 and Deep Submergence Vehicles of Ifremer. We greatly acknowledge Dr Jerome Mounier for  
12 556 his valuable help and comments on FISH and Jerome Lepioufle for valuable advice on the  
13 557 manuscript. We thank all the members of the GDR Ecchis for their advice and suggestions.  
14 558 We would like to thank the editor and the anonymous reviewers who have provided helpful  
15 559 comments on the refinement of this manuscript. We also thank ANR Deep-Oases, Ifremer,  
16 560 Region Bretagne and French Research Ministry for their financial support. We finally thank  
17 561 Françoise Gaill and the CHEMECO/European Science Foundation EURODEEP for  
18 562 discussions and financial support.

19  
20  
21  
22  
23  
24  
25  
26 563  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

564 **References**

- 565 Altschul S, Gish W, Miller W, Myers E & Lipman D (1990) Basic local alignment search tool. *J Mol Bio* **215**:  
566 403–410.
- 567
- 568 Amann RI, Ludwig W & Schleifer K (1995) Phylogenetic identification and *in situ* detection of individual  
569 microbial cells without cultivation. *Microbiol Rev*, **59**: 143-169.
- 570
- 571 Amann RI & Fuchs BM (2008) Single-cell identification in microbial communities by improved fluorescence in  
572 situ hybridization techniques. *Nat Rev Microbiol*, **6**: 339-348.
- 573
- 574 Ashelford KE, Weightman AJ, & Fry JC (2002) PRIMROSE: a computer program for generating and estimating  
575 the phylogenetic range of 16S rRNA oligonucleotide probes and primers in conjunction with the RDP-II  
576 database. *Nucleic Acids Res* **30**: 3481-3489.
- 577
- 578 Bai F, Zhao JH, Takashima M, Jia JH, Boekhout T & Nakase T (2002) Reclassification of the *Sporobolomyces*  
579 *roseus* and *Sporidiobolus pararoseus* complexes, with the description of *Sporobolomyces phaffii* sp. nov. *Int J*  
580 *Syst Evol Microbiol* **52**: 2309-2314.
- 581
- 582 Baschien C, Manz W, Neu T, Marvanova L & Szewzyk U (2008) *In Situ* Detection of Freshwater Fungi in an  
583 Alpine Stream by New Taxon-Specific Fluorescence In Situ Hybridization Probes. *Appl Environ Microb* **74**:  
584 6427-6436.
- 585
- 586 Bass D, Howe A, Brown N, Barton H, Demidova M, Michelle H, Li L, Sanders H, Watkinson SC, Willcock S &  
587 Richards TA (2007) Yeast forms dominate fungal diversity in the deep oceans. *Proc R Soc B* **274**: 3069-3077.
- 588
- 589 Burgaud G, Le Calvez T, Arzur D, Vandenkoornhuysse, P & Barbier G (2009) Diversity of culturable marine  
590 filamentous fungi from deep-sea hydrothermal vents. *Environ Microbiol* **11**: 1588-1600.
- 591
- 592 Cancela da Fonseca JP (1966) L'outil statistique en biologie du sol. III. Indices d'intérêt écologique. *Revue*  
593 *d'Ecologie et de Biologie du Sol* **3**: 381–407.
- 594
- 595 Charlou JL, Donval JP, Fouquet Y, Jean-Baptiste P & Holm N (2002) Geochemistry of high H<sub>2</sub> and CH<sub>4</sub> vent  
596 fluids issuing from ultramafic rocks at the Rainbow hydrothermal field (36j14VN, MAR). *Chem Geol*, **191**: 345-  
597 359.
- 598
- 599 Corbari L, Zbinden M, Cambon-Bonavita MA, Gaill F & Compère P (2008) Bacterial symbionts and mineral  
600 deposits in the branchial chamber of the hydrothermal vent shrimp *Rimicaris exoculata*: Relationship to moult  
601 cycle. *Aquat Biol*, **1**: 225-238.
- 602
- 603 Daims H, Stoecker K & Wagner M (2005) Fluorescence *In situ* Hybridisation for the Detection of Prokaryotes.  
604 In: *Advanced Methods in Molecular Microbial Ecology*. BIOS Scientific Publishers, Abingdon, UK. pp. 213-  
605 239.
- 606
- 607 Daims H, Lückner S, & Wagner M (2006) *daime*, a novel image analysis program for microbial ecology and  
608 biofilm research. *Environ Microbiol* **8**: 200-213.
- 609
- 610 Edgcomb VP, Kysela DT, Teske A, & de Vera Gomez A (2002) Benthic eukaryotic diversity in the Guaymas  
611 Basin hydrothermal vent environment. *Proc Natl Acad Sci USA* **99**: 7658-7662.
- 612
- 613 Fell JW (1976) Yeasts in oceanic regions. In: Jones EBG (ed) *Recent advances in aquatic mycology*. Elec,  
London, pp 93-124.
- 614
- 615 Gadanho M, Almeida JM, & Sampaio JP (2003) Assessment of yeast diversity in a marine environment in the  
616 south of Portugal by microsatellite-primed PCR. *Antonie van Leeuwenhoek* **84**: 217-227.
- 617
- 618 Gadanho M & Sampaio JP (2004) Application of temperature gradient gel electrophoresis to the study of yeast  
619 diversity in the estuary of the Tagus river, Portugal. *FEMS Yeast Res* **5**: 253-261.
- 620
- 621 Gadanho M & Sampaio J (2005) Occurrence and Diversity of Yeasts in the Mid-Atlantic Ridge Hydrothermal  
Fields Near the Azores Archipelago. *Microb Ecol* **50**: 408-417.

- 1  
2  
3 622  
4 623 Gadd GM (2007) Geomycology: biogeochemical transformations of rocks, minerals and radionuclides by fungi,  
5 624 bioweathering and bioremediation. *Mycol Res* **111**: 3-49.  
6 625  
7 626 Guindon S, Lethiec F, Duroux P & Gascuel O (2005) PHYML online – a web server for fast maximum  
8 627 likelihoodbased phylogenetic inference. *Nucleic Acids Res* **33**: 557–559.  
9 628  
10 629 Gunde-Cimerman N, Zalar P, de Hoog S & Plemenitas A (2000) Hypersaline waters in salterns - natural  
11 630 ecological niches for halophilic black yeasts. *FEMS Microbiol Ecol* **32**: 235-240.  
12 631  
13 632 Hawksworth DL (2002) The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycol Res*  
14 633 **105**: 1422-1432.  
15 634  
16 635 Hernandez-Saavedra NY, Hernandez- Saavedra D & Ochoa JL (1992) Distribution of *Sporobolomyces* (Kluyver  
17 636 et van Niel) genus in the western coast of Baja California Sur, Mexico. *Syst Appl Microbiol* **15**: 319-322.  
18 637  
19 638 Hoshino T, Yilmaz LS, Noguera DR, Daims H & Wagner M (2008) Quantification of Target Molecules Needed  
20 639 To Detect Microorganisms by Fluorescence In Situ Hybridization (FISH) and Catalyzed Reporter Deposition-  
21 640 FISH. *Appl Environ Microb* **74**: 5068-5077.  
22 641  
23 642 Joergensen BB, & Boetius A (2007) Feast and famine-microbial life in the deep-sea bed. *Nat Rev Microbiol* **5**:  
24 643 770-781.  
25 644  
26 645 Kogej T, Ramos J, Plemenitas A, & Gunde-Cimerman N (2005) The halophilic fungus *Hortaea werneckii* and  
27 646 the halotolerant fungus *Aureobasidium pullulans* maintain low intracellular cation concentrations in hypersaline  
28 647 environments. *Appl Environ Microb* **71**: 6600-6605.  
29 648  
30 649 Kohlmeyer J & Kohlmeyer E (1979) Marine Mycology: The Higher Fungi. New-York, USA; Academic Press.  
31 650  
32  
33 651 Inacio J, Behrens S, Fuchs BM, Fonseca A, Spencer-Martins I & Amann R (2003) In situ accessibility of  
34 652 *Saccharomyces cerevisiae* 26S rRNA to Cy3-labeled oligonucleotide probes comprising the D1 and D2  
35 653 domains. *Appl Environ Microb* **69**: 2899-2905.  
36 654  
37 655 Kushner DJ (1978) Life in high salt and solute concentrations. In: Kushner DJ (ed) *Microbial life in extreme*  
38 *environments*. Academic Press, London, pp 317-368.  
39 656  
40 657 Lachance MA & Starmer WT (1998) Ecology and yeasts. In: Kurtzman CP, Fell JW (eds) *The yeasts, a*  
41 *taxonomic study*, 4th edn. Elsevier, Amsterdam, The Netherlands, pp 21-30.  
42 658  
43 659 Lai X, Cao L, Tan H, Fang S, Huang Y & Zhou S (2007) Fungal communities from methane hydrate-bearing  
44 660 deep-sea marine sediments in South China Sea. *ISME J* **1**: 756-762.  
45 661  
46 662 Le Calvez T, Burgaud G, Mahe S, Barbier G & Vandenkoornhuysse P (2009) Fungal Diversity in Deep Sea  
47 663 Hydrothermal Ecosystems. *Appl Environ Microb* **75**: 6415-6421.  
48 664  
49 665 Libkind D, Brizzio S, Ruffini A, Gadanho M, van Broock M & Sampaio JP (2003) Molecular characterization of  
50 666 carotenogenic yeasts from aquatic environments in Patagonia, Argentina. *Antonie van Leeuwenhoek* **84**: 313-  
51 667 322.  
52 668  
53 669 Lopez-Garcia P, Philippe H, Gail F & Moreira D (2003) Autochthonous eukaryotic diversity in hydrothermal  
54 670 sediment and experimental microcolonizers at the Mid-Atlantic Ridge. *Proc Natl Acad Sci USA* **100**: 697-702.  
55 671  
56 672 Lopez-Garcia P, Vereshchaka A & Moreira D (2007) Eukaryotic diversity associated with carbonates and fluid-  
57 673 seawater interface in Lost City hydrothermal field. *Environ Microbiol* **9**: 546-554.  
58 674  
59 675 Margesin R & Schinner F (2001) Potential of halotolerant and halophilic microorganisms for biotechnology.  
60 676 *Extremophiles* **5**: 73-83.  
677  
678 Maruyama A, Honda D, Yamamoto H, Kitamura K & Higashihara T (2000) Phylogenetic analysis of  
psychrophilic bacteria from the Japan Trench, including a description of the deep-sea species *Psychrobacter*

- 1  
2  
3 679 *pacificensis* sp. nov. *Int J Syst Evol Microbiol* **50**: 835-846.  
4 680  
5 681 Munn, CB (2003) *Marine Microbiology – Ecology and Applications*. Oxford, UK: Bios-Garland Scientific.  
6 682  
7 683 Nagahama T, Hamamoto M, Nakase T & Horikoshi K (1999) *Kluyveromyces nonfermentans* sp. nov., a new  
8 684 yeast species isolated from the deep sea. *Int J Syst Evol Microbiol* **49**: 1899-1905.  
9 685  
10 686 Nagahama T, Hamamoto M, Nakase T & Horikoshi K (2001a) *Rhodotorula lamellibrachii* sp. nov., a new yeast  
11 687 species from a tubeworm collected at the deep-sea floor in Sagami Bay and its phylogenetic analysis. *Antonie*  
12 688 *van Leeuwenhoek* **80**: 317-323.  
13 689  
14 690 Nagahama T, Hamamoto M, Nakase T, Takami H & Horikoshi K (2001b) Distribution and identification of red  
15 691 yeasts in deep-sea environments around the northwest Pacific Ocean. *Antonie van Leeuwenhoek* **80**: 101-110.  
16 692  
17 693 Nagahama T, Hamamoto M, Nakase T, Takaki Y & Horikoshi K (2003a) *Cryptococcus surugaensis* sp. nov., a  
18 694 novel yeast species from sediment collected on the floor of Suruga Bay. *Int J Syst Evol Microbiol* **53**: 2095-  
19 695 2098.  
20 696  
21 697 Nagahama T, Hamamoto M, Nakase T & Horikoshi K (2003b) *Rhodotorula benthica* sp. nov. and *Rhodotorula*  
22 698 *calyptogenae* sp. nov., novel yeast species from animals collected from the deep-sea floor, and *Rhodotorula*  
23 699 *lysiniophila* sp. nov., which is related phylogenetically. *Int J Syst Evol Microbiol* **53**: 897-903.  
24 700  
25 701 Nagahama T, Hamamoto M & Horikoshi K (2006a) *Rhodotorula pacifica* sp. nov., a novel yeast species from  
26 702 sediment collected on the deep-sea floor of the north-west Pacific Ocean. *Int J Syst Evol Microbiol* **56**: 295-299.  
27 703  
28 704 Nagahama T (2006b) Yeast biodiversity in freshwater, marine and deep-sea environments. *Biodiversity and*  
29 705 *Ecophysiology of Yeasts*.  
30 706  
31 707 Nagahama T, Abdel-Wahab MA, Nogi Y, Miyazaki M, Uematsu K, Hamamoto M & Horikoshi K (2008)  
32 708 *Dipodascus tetrasporus* sp. nov., an ascosporeogenous yeast isolated from deep-sea sediments in the Japan  
33 709 Trench. *Int J Syst Evol Microbiol* **58**: 1040-1046.  
34 710  
35 711 Nguyen H, Gaillardin C & Neuvéglise C (2009) Differentiation of *Debaryomyces hansenii* and *Candida famata*  
36 712 by rRNA gene intergenic spacer fingerprinting and reassessment of phylogenetic relationships among *D.*  
37 713 *hansenii*, *C. famata*, *D. fabryi*, *C. flarerii* (= *D. subglobosus*) and *D. prosopidis*: description of *D. vietnamensis* sp.  
38 714 nov. closely related to *D. nepalensis*. *FEMS Yeast Res* **9**: 641-662.  
39 715  
40 716 Posada D & Crandall K (1998) Applications note. MODELTEST: testing the model of DNA substitution.  
41 717 *Bioinformatics* **14**: 817-818.  
42 718  
43 719 Ronquist F & Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models.  
44 720 *Bioinformatics* **19**: 1572-1574  
45 721  
46 722 Siepmann R & Höhnk W (1962) Über Hefen und einige Pilze (Fungi imp., Hyphales) aus dem Nordatlantik.  
47 723 *Veroeff Inst für Meeresforschung in Bremerhaven* **8**: 79-97.  
48 724  
49 725 Stoecker K, Dorninger C, Daims H & Wagner M (2010) Double Labeling of Oligonucleotide Probes for  
50 726 Fluorescence *In Situ* Hybridization (DOPE-FISH) Improves Signal Intensity and Increases rRNA Accessibility.  
51 727 *Appl Environ Microb* **76**: 922-926.  
52 728  
53 729 Takishita K, Tsuchiya M, Reimer JD & Maruyama T (2006) Molecular evidence demonstrating the  
54 730 basidiomycetous fungus *Cryptococcus curvatus* is the dominant microbial eukaryote in sediment at the  
55 731 Kuroshima Knoll methane seep. *Extremophiles* **10**: 165-169.  
56 732  
57 733 Takishita K, Yubuki N, Kakizoe N, Inagaki Y & Maruyama T (2007) Diversity of microbial eukaryotes in  
58 734 sediment at a deep-sea methane cold seep: surveys of ribosomal DNA libraries from raw sediment samples and  
59 735 two enrichment cultures. *Extremophiles* **11**: 563-576.  
60 736  
737 Tamura K, Dudley J, Nei M & Kumar, S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA)  
738 software, version 4.0. *Mol Biol Evol* **24**: 1596-1599.  
739

- 1  
2  
3 740 Tivey, MK (2007) Generation of seafloor hydrothermal vent fluids and associated mineral deposits.  
4 741 *Oceanography* **20**: 50-65.  
5 742
- 6 743 Thompson JD, Higgins DG & Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive  
7 744 multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix  
8 745 choice. *Nucleic Acids Res* **22**: 4673-4680.  
9 746
- 10 747 Van Dover CL, Ward ME, Scott JL, Underdown J, Anderson B, Gustafson C, Whalen M & Carnegie RA (2007)  
11 748 A fungal epizootic in mussels at a deep-sea hydrothermal vent. *Mar Ecol* **28**: 54-62.  
12 749
- 13 750 Zalar P, de Hoog GS & Gunde-Cimerman N (1999) Ecology of halotolerant dothideaceous black yeasts. *Stud*  
14 751 *Mycol* **43**: 38-48.  
15 752
- 16 753 Zbinden M, Le Bris N, Gaill F & Compere P (2004) Distribution of bacteria and associated minerals in the gill  
17 754 chamber of the vent shrimp *Rimicaris exoculata* and related biogeochemical processes. *Mar Ecol Progr Ser* **284**:  
18 755 237-251.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

756 **Tables and Figures**

757

758 Table 1. Culture collection of yeasts from deep-sea hydrothermal vents.

759

760

761

762

763

For Peer Review

	<b>Location (Depth)</b>	<b>Sample processed (type)</b>	<b>Strain</b>
3	764		
4	765	South Pacific West;	B2E07: Seawater surrounding mussels
5	766	(Lau Basin; -2620m)	B9E07: Gastropod ( <i>Ifremeria nautili</i> ) gills
6	767	Mid-Atlantic Ridge	EX6E01 to EX6E04: <i>Rimicaris exoculata</i>
7	768	(Rainbow; -2300m)	EX6E05: <i>Chorocaris chacei</i>
8	769		MoPR1: <i>Rimicaris exoculata</i>
9	770		MoPR1: <i>Mirocaris fortunata</i>
10	771		MoPR2: <i>Rimicaris exoculata</i>
11	772		MoPR3: Sloughs of shrimp on smocker rocks
12	773		MoPR5: Colonization module TRAC (Carbonates)
13	774		MoPR6: <i>Bathymodiolus azoricus</i>
14	775		MoPR8: <i>Rimicaris exoculata</i>
15	776		MoPR9: Sponge
16	777		MoPR9: Coral
17	778	Mid-Atlantic Ridge	EX18E02: Siliceous sponge
18	779	(Lost-City; -700m)	Ex15

780

781

782

783

784

785

786

787

788

789

790

791



1  
2  
3 792 Table 2. Physiological analysis of the yeast collection. This table shows distribution of  
4 793 halotolerant and halophilic strains of the collection depending on their optimal salinities (g/l  
5 794 sea salts), optimal temperatures (°C) and maximal optical densities of cultures on GYPS broth  
6 795 medium (120 rpm on a rotary shaker) measured at 600nm at 4 different incubation times (17h,  
7 796 22h, 25h and 28h).  
8  
9

10 797  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

798

		Low OD (<1.1)	High OD (>2.0)	
	Optimum	25°C	25°C	35°C
<b>Non halophile</b>	0-15 g/l	Mo25	Bio1, Bio2, Ex2, Ex3, Ex4, Ex5, Ex6, Ex7, Ex9, Ex11, Ex12, Mo20, Mo21, Mo24, Mo26, Mo27, Mo28, Mo29, Mo31, Mo32, Mo33, Mo35, Mo40	Ex15
<b>Halotolerant</b>	30 g/l	Mo36	Mo37, Mo38	
	45 g/l			Mo30
	60 g/l	Mo22		
<b>Halophile</b>	30 g/l	Mo34		Mo39

799

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

800 Table 3. Yeast oligonucleotide probes and their sequences, target organisms and binding  
801 positions on the 26S rRNA.

802  
803  
804  
805

For Peer Review

Probe	Hybridization stringency (% formamide)	rRNA subunit, binding position <sup>(a)</sup> and relative probe accessibility <sup>(b)</sup>	Probe sequence (5'-3')	Target organisms (Genus/Species)
Sacch	20	26S; 162-177 ; 44 to 66%	GGCATCTCATCGCACG	<i>Debaryomyces</i> <i>Pichia</i>
MitoFilo	10	26S; 397-412 ; 60%	ACACCGCAGAACGGCT	Members of the genus <i>Cryptococcus</i> <sup>(c)</sup>
MitoSporidio	20	26S; 164-179 ; 44 to 66%	TGGGCGTCCGCACCAT	Members of the genera <i>Rhodotorula</i> and <i>Rhodospidium</i> <sup>(d)</sup>

(a) Nucleotide position according to *Saccharomyces cerevisiae* 26Sr RNA between NL1 and NL4 primers.


(b) According to Inacio *et al.*, 2003.

(c) *Cryptococcus saitoi*, *C. randhawii*, *C. uzbekistanensis*, *C. adeliensis*, *C. vishniacii*, *C. socialis*, *C. friedmannii* and *C. uniguttulatus*.

(d) *Rhodotorula mucilaginosa*, *R. glutinis*, *R. graminis*, *R. dairenensis*, *Rhodospidium babjevae* and *R. diobovatum*.

1  
2  
3 Figure 1: Phylogenetic tree of deep-sea yeast isolates (coloured terminals) and close relatives  
4 obtained by analysis of the D1/D2 domain of the 26S rRNA gene. Topology was built using  
5 MrBayes v.3.1.2 from a ClustalW 1.83 alignment. Node support values are given in the  
6 following order: MrBayes posterior probabilities/PHYML 100 bootstraps. Black squares  
7 represent nodes supported by an excess of 0.95 posterior probabilities and 95% bootstraps.  
8 *Mucor flavus* (EU071390) belonging to the *Zygomycota* phylum was used as outgroup. All  
9 sequences are listed with their GenBank accession numbers.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review




Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 2: Fluorescence *in situ* hybridization with specific oligonucleotide probes on membrane filters. (a, b and c) Membrane filter labelled with DAPI and hybridized using MitoSporidio probe indicating the presence of bacteria and yeast cells (blue). Yeasts belonging to *Rhodotorula* and *Rhodospiridium* genera are visualized in pink (composite of blue and red). (d) Membrane filter labelled with DAPI and hybridized with MitoFilo indicating the presence of yeasts belonging to *Cryptococcus* genera. White arrows indicate the yeast cells.

For Peer Review







1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




1  
2  
3 Figure S1 : Target sites of the fluorescent oligonucleotide probes designed on a model of the  
4 *Saccharomyces cerevisiae* 26S rRNA secondary structure in which the D1 and D2 domains  
5 (delimited by NL1 and NL4) are enlarged (Inacio *et al.*, 2003). Each probe was evaluated  
6 without formamide in order to check whether the probe binds to the ribosomes of the target  
7 cells. The optimal hybridization conditions were determined in a series of FISH experiments  
8 with increasing formamide concentrations for a probe target and a non-target organism : (i)  
9 Sacch probe, *Debaryomyces hansenii* (Target) and *Candida atlantica* (Non-Target) with two  
10 mismatches ; (ii) MitoSporidio probe, *Rhodospiridium diobovatum* (Target) and  
11 *Cryptococcus uzbekistanensis* (Non-Target) with five mismatches and (iii) MitoFilo probe,  
12 *Cryptococcus uzbekistanensis* (Target) and *Rhodospiridium diobovatum* (Non-Target) with  
13 seven mismatches. Relative probe accessibility was determined for each probe : MitoFilo,  
14 about 60% ; MitoSporidio and Sacch, 44 to 66%.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review




Peer Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Phylogenetic tree of deep-sea yeast isolates (coloured terminals) and close relatives obtained by analysis of the D1/D2 domain of the 26S rRNA gene. Topology was built using MrBayes v.3.1.2 from a ClustalW 1.83 alignment. Node support values are given in the following order: MrBayes posterior probabilities/PHYML 100 bootstraps. Black squares represent nodes supported by an excess of 0.95 posterior probabilities and 95% bootstraps. *Mucor flavus* (EU071390) belonging to the Zygomycota phylum was used as outgroup. All sequences are listed with their GenBank accession numbers.

392x327mm (450 x 450 DPI)



Fluorescence in situ hybridization with specific oligonucleotide probes on membrane filters. (a, b and c) Membrane filter labelled with DAPI and hybridized using MitoSporidio probe indicating the presence of bacteria and yeast cells (blue). Yeasts belonging to *Rhodotorula* and *Rhodosporidium* genera are visualized in pink (composite of blue and red). (d) Membrane filter labelled with DAPI and hybridized with MitoFilo indicating the presence of yeasts belonging to *Cryptococcus* genera.

White arrows indicate the yeast cells.

255x231mm (600 x 600 DPI)




Figure S1 : Target sites of the fluorescent oligonucleotide probes designed on a model of the *Saccharomyces cerevisiae* 26S rRNA secondary structure in which the D1 and D2 domains (delimited by NL1 and NL4) are enlarged (Inacio et al., 2003). Each probe was evaluated without formamide in order to check whether the probe binds to the ribosomes of the target cells. The optimal hybridization conditions were determined in a series of FISH experiments with increasing formamide concentrations for a probe target and a non-target organism : (i) Sacch probe, *Debaryomyces hansenii* (Target) and *Candida atlantica* (Non-Target) with two mismatches ; (ii) MitoSporidio probe, *Rhodospiridium diobovatum* (Target) and *Cryptococcus uzbekistanensis* (Non-Target) with five mismatches and (iii) MitoFilo probe, *Cryptococcus uzbekistanensis* (Target) and *Rhodospiridium diobovatum* (Non-Target) with seven mismatches. Relative probe accessibility was determined for each probe : MitoFilo, about 60% ; MitoSporidio and Sacch, 44 to 66%.  
237x157mm (600 x 600 DPI)