

*This is a pre-copy-editing, author-produced PDF of an article accepted for publication in ICES Journal of Marine Science following peer review. The definitive publisher-authenticated version is available online at:*

<http://dx.doi.org/10.1093/icesjms/fst030>

---

## Discarded fish in European waters: general patterns and contrasts

Sebastian S. Uhlmann<sup>1,\*</sup>, Aloysius T. M. van Helmond<sup>1</sup>, Elísabet Kemp Stefánsdóttir<sup>2</sup>, Sigríður Sigurðardóttir<sup>2</sup>, John Haralabous<sup>3</sup>, Jose Maria Bellido<sup>4</sup>, A. Carbonell<sup>4</sup>, Tom Catchpole<sup>5</sup>, Dimitrios Damalas<sup>6</sup>, Laurence Fauconnet<sup>7</sup>, Jordan Feekings<sup>8</sup>, Teresa Garcia<sup>4</sup>, Niels Madsen<sup>8</sup>, Sandra Mallold<sup>4</sup>, Sveinn Margeirsson<sup>2</sup>, Andreas Palialexis<sup>3</sup>, Lisa Readdy<sup>5</sup>, Julio Valeiras<sup>4</sup>, Vassiliki Vassilopoulou<sup>3</sup> and Marie-Joëlle Rochet<sup>7</sup>

<sup>1</sup> IMARES Wageningen UR, Haringkade 1, 1976 CP IJmuiden, The Netherlands

<sup>2</sup> Matís, Icelandic Food and Biotech R&D, Vínlandsleið 12, 113 Reykjavík, Iceland

<sup>3</sup> Institute of Marine Biological Resources, Hellenic Centre for Marine Research, Agios Kosmas, Hellinikon, 16610 Athens, Greece

<sup>4</sup> Instituto Español de Oceanografía, Centro Oceanográfico de Murcia, c/Varadero, 1, 30740 San Pedro del Pinatar, Murcia, Spain

<sup>5</sup> Cefas, Pakefield Road, Lowestoft, Suffolk NR33 0HT, UK

<sup>6</sup> European Commission, Joint Research Centre, Institute for the Protection and Security of the Citizen, Maritime Affairs Unit, via E. Fermi 2749, I-21027 Ispra (VA), Italy

<sup>7</sup> IFREMER, Rue de l'Île d'Yeu, 44311 Nantes, France

<sup>8</sup> Technical University of Denmark, National Institute of Aquatic Resources, North Sea Science Park, PO Box 101, DK-9850 Hirtshals, Denmark

\*: Corresponding author : Sebastian S. Uhlmann, tel: +31 317 480133 ; fax: +31 317 487326 ; email address : [sebastian.uhlmann@wur.nl](mailto:sebastian.uhlmann@wur.nl)

---

### Abstract:

To reduce the practice of discarding commercially fished organisms, several measures such as a discard ban and extra allowances on top of landings quotas ("catch quota") have been proposed by the European Commission. However, for their development and successful implementation, an understanding of discard patterns on a European scale is needed. In this study, we present an international synthesis of discard data collected on board commercial, towed-gear equipped vessels operating under six different national flags spanning from the Baltic to the Mediterranean Seas mainly between 2003 and 2008. We considered discarded species of commercial value such as Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), European hake (*Merluccius merluccius*), and European plaice (*Pleuronectes platessa*). Comparisons of discard per unit effort rates expressed as numbers per hour of fishing revealed that in the Mediterranean Sea minimum size-regulated species such as hake are generally discarded in much lower numbers than elsewhere. For most species examined, variability in discard rates across regions was greater than across fisheries, suggesting that a region-by-region approach to discard reduction would be more relevant. The high uncertainty in discard rate estimates suggests that current sampling regimes should be either expanded or complemented by other data sources, if they are to be used for setting catch quotas.

**Keywords:** Bycatch ; Common Fisheries Policy reform ; Data Collection Framework ; discard reduction ; Europe ; monitoring

50 1. Introduction

52 Discarding unwanted catch at sea in response to regulatory and/or market forces during  
commercial fishing is generally considered to be a waste of natural resources. It evades the  
54 eyes and often goes unrecorded. But, knowing how much is lost is important, for at least  
three reasons: firstly, discards might make up a large part of the total catch, possibly  
56 exceeding the amount of landings; secondly, stock viability and productivity may be  
compromised if large, and unregistered numbers of organisms are removed periodically on  
58 top of the registered landings (Crowder and Murawski, 1998; Punt *et al.*, 2006); thirdly,  
quantification of the magnitude of discarding is the first step in a framework to resolve it  
60 (Kennelly and Broadhurst, 2002).

62 In Europe, estimating the amount of discards is legislated via the Data Collection Framework  
(‘DCF’; EEC, 2000). As part of nationally-adopted onboard observer programmes, trained  
64 personnel collect the biomass, length, age and species compositions of discards from their  
most important commercial fisheries (EEC, 2009), with the main aim to feed these data into  
66 stock assessments. This is done via at-sea sampling (ICES, 2011), and all the data are stored  
and administered by the respective national authorities. Although various analyses of these  
68 data have been done, many studies were restricted to regional fisheries (e.g. Stratoudakis *et al.*,  
1999; Viana *et al.*, 2011; Feekings *et al.*, 2012; Madsen *et al.*, 2013). However,  
70 synthesizing discard data from as many different fisheries, regions and countries as possible  
is required to facilitate European-wide management approaches. So far, such a synthesis was  
72 hampered by i) the diversity of procedures in collecting and processing data, ii) the disparate  
intensities of sampling compared to total fishing effort across countries, iii) the lack of a  
74 common data exchange format and storage facility, and iv) national regulations which  
precluded sharing of detailed commercial catch data (STECF, 2006, 2008; Hinz *et al.*, 2013).

76 Considering that a reduction of discards is set to be a cornerstone of the European Common  
78 Fisheries Policy (CFP) reform (EEC, 2011), a comprehensive pan-European synthesis of  
discard data across species, fishing regions and fleets is important. This may aid the  
80 decision-making process by providing input to questions such as on what level discard-  
reduction initiatives need to be implemented: species, fisheries, or region-based (i.e. fishing  
82 ground). An important component of the CFP-reform proposal is a landing obligation, or  
discard ban, prohibiting the at-sea disposal of some commercially-valuable species from 2014  
84 onwards (Article 15; EEC, 2011; EEC, 2012a). Alternatively, catch quota could substitute the  
current landings quota (EEC, 2011). In either case, the complete catch would need to be  
86 accounted for. Shifting from a landings to a catch quota management system would require  
that catch quotas are set based on reliable estimates of discarded amounts and/or proportions.  
88 However, discard rates of a given species are likely to fluctuate within a fishery (e.g.  
Feekings et al., 2012; Poos et al., 2013) and/or across different fisheries, seasons, years and  
90 regions (Stratoudakis et al., 1999; Borges et al., 2005; Borges et al., 2006). The starting point  
for designing mitigation measures and management plans to reduce discards is to describe  
92 and characterise these patterns.

94 In this study, onboard observer data from discard-intensive fisheries using towed gears from  
Denmark, England, France, Greece, The Netherlands, and Spain were compiled. These data  
96 were used to describe species-specific discard patterns among and between fisheries and  
regions. Owing to logistical and financial constraints, only a fraction of operations carried  
98 out by a fleet can be monitored, which will render extrapolations across the entire population  
of operations uncertain (Depestele et al., 2011). Extrapolations require the use of raising or  
100 auxiliary variables such as landings or fishing effort. Following ICES (2011) this could be  
done “*according to sampling theory [where] the standard raising procedure within a given*

102 *stratum (e.g. quarter and area) should be: i) samples are raised to haul level based on*  
104 *sampled proportion; ii) sampled hauls are raised to trip level based on the proportion of*  
106 *hauls sampled; and iii) sampled trips are raised to métier level based on the proportion of*  
108 *trips sampled*'. But, the availability and quality of raising variables is not uniform and varies  
110 across countries (ICES, 2007), so that no single raising procedure can be recommended at the  
European level (ICES, 2011). For example, the total number of trips within a stratum may  
not be known, or may be either over- or underestimated due to the switching of gears  
throughout a trip or depending on post-stratification methods (ICES, 2010). To circumvent  
these issues, discard estimates at the level of sampled trips are presented here.

112 To allow for an integration and comparison of discard data from various fisheries and national  
sampling programmes, an index has to be defined that takes into account the unit of fishing  
114 effort (i.e. DPUE, Discards per Unit of Effort; Rochet and Trenkel, 2005). Fishing effort  
measured as the hours spent actually fishing is a commonly-used effort descriptor among EU  
116 member states for towed gears. A DPUE index of abundance, hereafter called 'discard rate',  
can be a useful tool for policy makers to identify discard-intensive fisheries and improve  
118 discard management by developing mitigation strategies. Another useful measure, is the ratio  
between discards and catch (discards and landings). Thus, in this study, we combined discard  
120 data from six different countries and several different regions (spanning from the Baltic to the  
Mediterranean Seas) to compare discard rates of commercially-valuable species such as  
122 Atlantic cod (*Gadus morhua*); haddock (*Melanogrammus aeglefinus*); European hake  
(*Merluccius merluccius*); and European plaice (*Pleuronectes platessa*). The aim was to  
124 contrast discard rates and ratios between fisheries or regions. We compared the coefficients  
of variation of discard rates and ratios across fisheries for a given region and across regions  
126 for a given fishery. If discard patterns were found to be more homogeneous across regions  
than fisheries, a fisheries-by-fisheries approach to discard reduction might be more relevant.

128

## 130 2. Material and methods

### 132 2.1 Dataset

A dataset was built from pre-processed and aggregated trip-level information that was  
134 provided by each partner detailing the mean ( $\pm$  standard deviation) number of  
discarded/landed species per hour from sampled trips per métier, fishing region, sub-region;  
136 together with the corresponding number of sampled trips from towed gears. Thereby, fishing  
activity was linked to the European level 5 métier definition, requiring data at the level of  
138 fishing ground (hereafter ‘region’), gear type, and target species assemblage (e.g. demersal  
fish – hereafter ‘fish’, small pelagic fish, cephalopods and fish, crustaceans, crustaceans and  
140 fish; FAO, 1980; EEC, 2008; ICES, 2009). Hereafter the term ‘fishery’ is used to designate a  
gear type and target species assemblage combination. All biological data such as the numbers  
142 and weights (where available) of discarded and landed species were summarized by region,  
sub-area per region (i.e. ICES Divisions or FAO areas of the Mediterranean Sea), métier and  
144 vessel flag country (hereafter country) together with technical information (average trip  
duration, fleet size and fishing effort). ICES Division ‘IIIa’ was subdivided into Skagerrak  
146 and Kattegat to reflect the stock classifications used by ICES. A summary of a detailed  
comparison of each of the national discard sampling programmes is provided in Table 1.

148

Biological data were collected on a haul-by-haul basis and, for the majority of samples,  
150 consisted of landings and discard observations of commercially-valuable species (including  
invertebrates such as crustaceans, molluscs and cephalopods). Numbers discarded, numbers  
152 landed (when these were registered), and lengths (cm) were recorded. For the purpose of our  
study, numbers rather than weights were used, because species weights of catch and discards

154 were not recorded in all national sampling programmes owing to the challenge of obtaining  
accurate weight measurements at sea. Although length-weight relationships may have  
156 allowed for transformations of available numbers-at-length into weights, this approach was  
not chosen, because it would have implied the mixing of measurements (available from  $n=5$   
158 partners; Table 1) with estimated weights (theoretically available from  $n=2$  partners, Table 1)  
when combining data from different countries. All numbers were raised to the haul level (if a  
160 sub-sample was measured; based on the proportion between the total and sampled fraction)  
and subsequently to the trip level (based on either the proportion of sampled fishing  
162 operations or fishing time; see Table 1 and ICES, 2011 for details)). These raised numbers of  
landings and discards per species per sampled trip were standardized by sampled fishing time  
164 (i.e. tow duration, in hours) to derive a discard rate (i.e. DPUE), as the numbers landed or  
discarded per hour per sampled trip. The ratio between discards and catch (discards +  
166 landings) rates was used as the discard ratio. From all sampled trips, an average and a  
standard deviation was then calculated for discard rates and ratios as follows.

168

## 2.2 Estimation of discard rates and ratios and their variability

170 To compare species-specific discard rates and ratios (at the level of sampled trips) across  
regions and fisheries, means and standard deviations across countries and sub-areas within  
172 regions were combined. The most appropriate auxiliary variables, such as total fishing effort,  
were not available in comparable units at the required level of aggregation and desired quality  
174 from all countries. Therefore, discard rates were weighted by national sampling effort (i.e.  
number of observed trips) under the assumption that sampling effort was proportional to a  
176 fleet's activity. Thereby, mean numbers of discarded or landed species per hour and trip were  
combined for a given fishery and region as:

178

$$M = \sum_{i \in I, k \in K} \frac{n_{i,k} m_{i,k}}{N} \quad (\text{Equation 1})$$

180 Where  $M$  is the mean number of a discarded or landed species per given fishery and region  
 and  $N$  is the total number of sampled trips per given fishery and region.  $I$  is the set of all sub-  
 182 areas within the region and  $K$  is the set of all countries.  $n_{i,k}$  is the number of sampled trips in  
 sub-area  $i$ , by country  $k$ , for the specified métier; and  $m_{i,k}$  is the mean number of a discarded  
 184 or landed species in sub-area  $i$ , by country  $k$ , for the specified fishery.

186 From the standard deviation that was associated with each mean number of a discarded or  
 landed species per hour, the variance  $V$  was calculated per species, fishery and region as  
 188 follows, whereby  $v_{i,k}$  is the variance for sub-area  $i$ , by country  $k$ , for the specified fishery.

$$V = \sum_{i \in I, k \in K} \frac{v_{i,k}(n_{i,k} - 1) + (m_{i,k} - M)^2 n_{i,k}}{N - 1} \quad (\text{Equation 2})$$

190  
 In  $n=97$  cases, standard deviations ( $SD$ , square root of the variance) of discard rates were  
 192 larger than the mean ( $M$ ). Available length-frequency distributions (Helmond and Uhlmann,  
 2011) were graphically examined and found to be positively skewed, which implies that a  
 194 log-normal distribution would describe the data more appropriately than a normal distribution  
 (Limpert et al., 2001). Accordingly, geometric means ( $GM$ ) and the multiplicative standard  
 196 deviation ( $GSD$ ) were calculated from the combined means ( $M$ ) and standard deviations  
 following Limpert et al. (2001):

198

$$GM = \frac{M}{\sqrt{1 + \left(\frac{SD}{M}\right)^2}} \quad (\text{Equation 3})$$

$$GSD = \exp\left(\sqrt{\log\left(1 + \left(\frac{SD}{M}\right)^2\right)}\right) \quad (\text{Equation 4})$$

200

Differences of discard and landings rates (i.e. per unit effort) between fisheries and/or regions  
 202 are illustrated in bar plots with inferential error bars (Cumming et al., 2007) calculated as:

$$GSE = GSD \frac{1}{\sqrt{N}} \quad (\text{Equation 5})$$

204

The inferential error bars show a confidence interval ( $GM/GSE$ ;  $GM*GSE$ ) for the median of  
 206 discarded or landed numbers. ‘Discard’ or landing rate’ hereafter refers to the geometric  
 mean of discarded or landed numbers per hour. Statistical significance at  $p < 0.05$  was  
 208 inferred when the gap between error bars was of the same size as the error bar itself with  $>10$   
 sampled trips. For fewer trips a greater gap is needed for a similar significant difference.

210

As a measure of the variability of discard rates and ratios across fisheries or regions, we  
 212 computed the coefficient of variation for discards rates and ratios by fisheries and region. To  
 calculate the respective CVs, the average and the standard deviation of discard rates and  
 214 ratios for a given fishery (across regions) or for a given region (across fisheries) were taken.

All calculations were done using the statistical software R (R Development Core Team,  
 216 2005), with the aid of the ‘combinevar’ function from the package ‘fishmethods’ (Nelson,  
 2012).

218



### 2.3 Comparison of discard rates and ratios

220 The comparisons of discard rates and ratios were done specifically for towed-gear fisheries  
that operated under different national flags. These included otter- (OTB) and beam-trawlers  
222 (TBB) targeting crustaceans (CRU) or demersal fish ('fish', DEF; Table 2). Pelagic fisheries  
which require specific sampling procedures were not considered in this study. To make  
224 meaningful i) inter-region (across fishing regions) and ii) inter-fishery (across fisheries)  
comparisons of species-specific discard rates in the following section, we selected non-  
226 pelagic, minimum-landing-size (MLS)-regulated species which were listed in the CFP-reform  
proposal, and were commonly discarded from the above-mentioned fisheries in a number of  
228 different regions, namely: cod (MLS= 35 cm in all regions except Skagerrak/Kattegat, where  
MLS was decreased to 30 cm in 2008 and in the Baltic Sea where it was increased to 38 cm  
230 in 2003); haddock (30 cm in all regions apart from Skagerrak/ Kattegat, where it is 27 cm);  
hake (27 cm in all regions apart from Skagerrak/Kattegat, 30 cm; and Mediterranean Sea, 20  
232 cm); and plaice (27 cm). Acknowledging the different species composition of discards in the  
Mediterranean Sea, for this region the following list was nominated in accordance with the  
234 above criteria: bogue (*Boops boops*; 10 cm according to national legislation in Greece); red  
mullet (*Mullet barbatus barbatus*; 11 cm); and deep-water rose shrimp (*Parapenaeus*  
236 *longirostris*, 2 cm carapace length).

238

## 240 3. Results

### 242 3.1 Dataset

National discard sampling programmes are not standardized at the European level and exhibit  
244 differences in the way vessels are selected for observation, the level of detail that is recorded

during biological sampling (e.g. species numbers, weights, age, and maturity) and what units  
246 of ratio estimators are used to scale up measured numbers (Table 1). Notwithstanding the  
above, sampling effort and landings and discard rates were compiled for 15 towed-gear  
248 fisheries and 11 major European fishing regions (22 ICES Divisions, and five Mediterranean  
geographic sub-areas (GSA); see Helmond and Uhlmann, 2011 for details). Among these  
250 classified fisheries, there were differences in fleet size, fishing effort, and sampling effort  
between countries (Table 2). Apart from one Greek fishery, generally <1% of the number of  
252 days spent at sea were observed in any fishery (Table 2).

### 254 3.2 Comparison of discard rates and ratios

Discard rates varied from <5 up to >300 individuals per hour based on observations between  
256 4 and 776 sampled trips (Figure 1). Observations from <4 trips were not included to avoid  
using non-representative values which in turn will increase the overall variance. The  
258 variability in sampling effort is reflected in the precision of the estimates (Figure 1). With  
<10 observations the uncertainty is large, and even with many samples some discard rates are  
260 difficult to estimate precisely owing to the large variability in discarding patterns (e.g. plaice  
discards by beam trawlers in the North Sea and Eastern Channel have a low precision, even  
262 though 100 trips were observed; Figure 1d).

264 Discard rates of cod and haddock (Figure 1a,b) were generally lower than those of hake and  
plaice (Figure 1c,d). Some of the Mediterranean species such as red mullet and deep-water  
266 rose shrimp exhibited the lowest rates (Figure 1e,f). In general, there were distinct patterns  
when comparing species-specific discard rates across fisheries and regions (Figure 1). For  
268 example, discard rates of Atlantic cod were found to be homogenous across fisheries, but  
were higher in the Skagerrak than in other areas (Table 3; Figure 1a). For haddock,  
270 differences of discard rates between regions were larger than between fisheries (Table 3;

Figure 1b). Hake discard rates were relatively low and similar between different fisheries and  
272 regions, except for bottom-otter trawlers targeting fish in the Celtic Sea or crustaceans in the  
Bay of Biscay (Table 3; Figure 1c). For plaice the differences of discard rates between  
274 fisheries, seemed to be of the same order of magnitude than between regions (Table 3; Figure  
1d). Notably, discard rates of plaice differed greatly between beam and otter trawls in the  
276 North Sea, but were much more homogenous across fisheries in the Irish Sea (Table 3; Figure  
1d). In general, otter trawlers targeting crustaceans were observed to discard the majority of  
278 the cod, hake, and plaice compared to those targeting fish (Figure 1a-d).

280 Both discard rates and ratios were lower in the Mediterranean Sea than in other regions  
(Tables 3 and 4; Figure 1e-g). In the Mediterranean Sea, landings rates largely exceeded  
282 those of discard rates (Figure 1c, e-f), except for bogue (Figure 1g). Discard ratios of hake  
were more homogenous than discard rates (Tables 3 and 4). The discard ratios of hake varied  
284 more in the Mediterranean Sea than in the Celtic Sea, where hake discards exceeded landings,  
even though it is a target species by the fleet operating there (Table 4; Figure 1c).

286

#### 288 4. Discussion

290 Our study highlights the variability of species-specific discard rates at a European scale. A  
stark contrast was observed between rates in the Mediterranean Sea and the other fishing  
292 regions. Further, we found that discard rates were more homogeneous across fisheries than  
regions, suggesting that discard management measures may be devised at a regional level; for  
294 example, by removing quota and catch composition rules (e.g. EEC, 2012b) and incentivising  
the use of more selective gears. In any case, differences in discard rates between species will  
296 also require species-specific approaches to discard reduction such as improvements to gear

selectivity parameters.

298

The low level of discarding of MLS-regulated species among Mediterranean otter-trawl  
300 fisheries may be a consequence of smaller MLS (e.g. hake), a lack of MLS-compliance and  
the absence of over-quota discards in a quota-independent management system of Greek  
302 demersal trawl fisheries (Catchpole et al., 2013; Damalas and Vassilopoulou, 2013).

Although undersized hake for example are being caught by demersal otter trawlers, the  
304 proportion (in weight) of discarded individuals is small (Damalas and Vassilopoulou, 2013).

The fast-growing, small-sized, and highly diverse fish fauna (Stergiou et al., 1997) together  
306 with the existence of local markets for small fish and the low probability of prosecution for  
retaining undersized fish (Damalas and Vassilopoulou, 2013) may be further reasons why a  
308 tendency to retain most of the catch exists in this area.

310 Apart from removing quotas and catch composition rules, incentives to increase the use of  
more selective gears may be another option to reduce discards. One of the more selective  
312 gears and fishing methods in our study, where the majority of the target catch was landed,  
were Danish seines catching cod in the Baltic Sea and plaice in the North Sea (Figure 1b,d).

314 Scottish seines seem equally selective for other target species such as megrim  
(*Lepidorhombus whiffiagonis*; Borges et al., 2006). Some gears and methods have become  
316 more selective in recent years (beyond the period investigated here) in some areas (e.g.  
Kattegat and Skagerrak); and their uptake throughout the fishing community was partly  
318 promoted by incentives such as an increased quota share, access rights and more fishing days  
(Madsen and Valentinsson, 2010).

320

A shortcoming of the current DCF, which complicated the inter-national synthesis of discard  
322 data, was the difficulty to agree upon common métier definitions. For example, target species

assemblage of a level-5-métier could be defined either before the commencement of a trip or  
324 after a trip's completion (i.e. by determining its landings compositions). If we had followed  
the latter rule, it would have resulted in such a large number of métiers, at least among some  
326 countries, that it would have rendered an analysis of combined data meaningless. Alternative  
sampling units other than métiers may be considered for the selection of a sampling frame as  
328 part of at-sea monitoring programmes, for example vessels (ICES, 2012). This will also  
facilitate the standardization of discard sampling approaches (ICES, 2011). Another  
330 shortcoming, which hampered our analysis, was the inability to combine both raw data of  
fishing effort and catch statistics, partly due to the requirements of a data harmonization  
332 software for species weights which were not routinely collected in all programmes  
(Anonymous, 2009; ICES, 2010, 2011) and partly due to confidentiality concerns of releasing  
334 detailed, non-aggregated data to a third party (ICES, 2009); the latter is an issue which has  
hampered also other scientific analyses (Hinz et al., 2013). The lack of recording a species'  
336 sub-sampled and total weight in some sampling programmes precluded the use of the COST  
software (Anonymous, 2009; ICES, 2010).

338

Data incompatibility and confidentiality were also the reasons, why we ended up contrasting  
340 aggregated data at the sampled trip as opposed to the fleet level. However, some inferences  
from patterns at the trip to the fleet level are possible. For example, the greater variability in  
342 discard rates between regions than fisheries may be a consequence of the region-specific  
quota and landings regulations, if acting as the main drivers of discarding (Catchpole et al.,  
344 2013). For example, the main reason for discarding cod by Danish otter trawlers in the Baltic  
Sea were catches below MLS, whereas in the North Sea and Eastern Channel cod discards  
346 were also driven by lack of sufficient quotas (Catchpole et al., 2013). Regional differences in  
MLS regulations may also be associated with higher discard rates of hake from bottom-otter  
348 trawlers in the Celtic Sea (MLS=27 cm), compared with lower rates by the same fishery in

the Mediterranean Sea (MLS=20 cm; Figure 1).

350

Nevertheless, the interpretation of differences between discard rates based on the available  
352 dataset is difficult for two reasons: firstly, not all species are caught and discarded in  
significant amounts in all regions, thus for each region we did not necessarily have data on  
354 the same species from all countries. Secondly, an additional problem is that the specific  
reason as to why a species is discarded can often be difficult to disentangle; especially if  
356 similar drivers such as quota and MLS regulations exists in different regions or target species  
vary throughout seasons and fisheries. For example, we have almost exclusively considered  
358 CFP-reform-listed fish as opposed to invertebrate crustacean species (other than deep-water  
rose shrimp) in our analysis. Thereby, we essentially mix comparisons of discard rates of  
360 non-target with those of target species. For bottom otter trawlers targeting crustaceans,  
discarded fish typically exceeded their landings rates during those sampled trips, whereas for  
362 those targeting fish the opposite patterns was eminent (Figure 1 a-d) Furthermore, the exact  
reasons why some fish with an associated landings quota were discarded above MLS can only  
364 be inferred (Catchpole et al., 2013); unless fishers (or observers, for example in the US  
Northeast Fisheries observer programme; Wigley et al., 2012) note why they chose to discard  
366 some fish over others (e.g. lack of quota, low market prize, or poor quality). Such reasons  
together with a plethora of likely other biological, technical, environmental and socio-  
368 economic factors will contribute to fluctuating discard rates between species (Borges et al.,  
2006), regions (Stratoudakis et al., 1999; Eliassen et al., 2013), gears and years (Borges et al.,  
370 2005), among others.

372 Introducing a discard ban or landing obligation in combination with catch limits across 27  
Member States, 11 fishing regions, 27 species, and approximately 84 000 registered vessels  
374 (EEC, 2011; Eurostat, 2012) may compromise the profitability of some discard-intensive

fisheries at least in the short-term. A discard ban in isolation would increase costs and  
376 decrease income if the catch includes significant proportions of unwanted organisms (Condie  
et al., unpubl. manuscript). But, if the benefits of non-compliance still outweigh the costs of  
378 sanctions (Batsleer et al., 2013), there may be little incentive for those with increased costs to  
comply with the desired outcome of reduced discards. Thus, the introduction of a discard ban  
380 will also require ancillary management measures such as catch quotas to stimulate more  
selective fishing practices (Condie et al., 2013). For the allocation of catch quotas it will be  
382 important, as the European Commission noted, that these “need to reflect as much as possible  
the actual fishing patterns of vessels and their likely catch composition” (EEC, 2012c). This  
384 study provides at a European scale a first portrayal of the fishing and discarding pattern for  
some of the considered species, fisheries and regions.

386

Our analysis of patterns in discard rates and ratios are based on measured numbers-at-length  
388 as opposed to length-weight-relationship-estimated weights. If weights were used, patterns  
may have differed depending on the proportion of small and light-weight individuals in  
390 discarded fractions. For example, 100 discarded cod would have translated into a much  
greater weight than 100 discarded bogue or plaice, owing to differences in MLS (e.g. cod, <  
392 38 cm in the Baltic Sea versus bogue, < 10 cm in the Mediterranean or plaice, < 27 cm) and  
their body morphology (flat versus round shapes).

394

Our analysis is based on the assumption that all the sampling programmes considered here  
396 have a similar degree of bias. Such bias may be associated with the selection of vessels on a  
voluntary basis, deployment of observers, and their sampling procedures. Deployment and  
398 observer bias (Benoît and Allard, 2009) are inherent to sampling programmes and difficult, if  
not impossible, to quantify. However, some of the sampling programmes used in this study  
400 were evaluated based on surrogate measures, such as comparing the relative biomass of

marketable fish between observed and unobserved trips gleaned from logbooks (Tsagarakis et al., 2008); the representativeness of sampled trips versus total effort in time and space (ICES, 2011); or selecting vessels for sampling from randomly-generated lists and where sampling effort was allocated in proportion to the fisheries' annual fishing effort in the preceding year (Catchpole et al., 2011). Despite these shortcomings, on-board observer programmes remain the most complete source of information on all components of the catch by fishing vessels.

The variability across samples resulted in wide confidence intervals for many discard rate estimates. If discard estimates are to be used in the future to set species-specific catch quotas within reasonable confidence limits, observations from a much greater number of fishing trips will be needed to more precisely estimate discard amounts. Alternative, innovative sampling techniques (e.g. self-sampling, Uhlmann et al., 2011; vessel monitoring by satellite systems, VMS, Hintzen et al., 2012; and closed-circuit TV, CCTV, Kindt-Larsen et al., 2011) may be necessary to overcome the high costs of observers and resulting small sample sizes. Otherwise, the number of species for which target precision levels can be achieved will remain small.

Onboard observer programmes, in their complexity require, like any other scientific survey, uniform sampling standards, or at least their detailed description (Cotter and Pilling, 2007, ICES, 2011) to allow for the inter-national integration of data. These programmes need to be continuously adapted because of perpetual changes in fishing activities. Despite some institutional inertia, the national efforts and the international coordination have allowed significant progress to be made. This study contributes to further improvements.

424

426 Acknowledgements



428 This study was carried out within the MariFish-funded BADMINTON project (Bycatch and  
Discards: management indicators, trends and location; <http://83.212.243.10/badminton.html>).  
430 The funders had no role in study design, data collection and analysis, decision to publish, or  
preparation of the manuscript. Our thanks go out to all the skippers who were happy to take  
432 observers onboard and likewise to all observers who diligently collected data.

434

#### References

436

Anonymous. 2009. Common tool for raising and estimating properties of statistical estimates  
438 derived from the Data Collection Regulation. Studies and Pilot projects for carrying out the  
common fisheries policy. Project No SI2.467814, Brussels. 118 pp.  
440 <http://wwz.ifremer.fr/cost>.

442 Batsleer J., Poos J.J., Marchal P., Vermard Y. , Rijnsdorp A.D. 2013. Mixed fisheries  
management: Protecting the weakest link. Marine Ecology Progress Series, doi:  
444 10.3354/meps10203.

446 Benoît, H. P., and Allard, J. 2009. Can the data from at-sea observer surveys be used to make  
general inferences about catch composition and discards? Canadian Journal of Fisheries and  
448 Aquatic Sciences, 66: 2025–2039.

450 Borges, L., Rogan, E., and Officer, R. 2005. Discarding by the demersal fishery in the waters  
around Ireland. Fisheries Research, 76: 1–13.

452

454 Borges, L., Zuur, A. F., Rogan, E., and Officer, R. 2006. Modelling discard ogives from Irish demersal fisheries. *ICES Journal of Marine Science*, 63: 1086–1095.

456 Catchpole, T. L., Enever, R., Maxwell, D. L., Armstrong, M. J., Reese, A., and Revill, A. S. 2011. Constructing indices to detect temporal trends in discarding. *Fisheries Research*, 107: 458 94–99.

460 Catchpole, T. L., Rochet, M. J., Feekings, J., Madsen, N., Nikolic, N., Palialexis, A., Vassilopoulou, V., Valeiras, J., and Garcia, T. 2013. Using inferred drivers of discarding 462 behaviour to develop a fishery specific mitigation framework. Submitted manuscript (ICESJMS-2012-447) to *ICES Journal of Marine Science* ‘Bycatch and Discards’ symposium 464 issue.

466 Condie, H. M., Grant, A., and Catchpole, T. L. 2013. Incentivising selective fishing under a policy to ban discards; lessons from European and global fisheries. *Marine Policy*. 468

Condie, H. M., Grant, A., and Catchpole, T. L. unpublished manuscript. Does banning 470 discards in an otter trawler fishery create incentives for more selective fishing? *Fisheries Research*. 472

Cotter, A. J. R., and Pilling, G. M. 2007. Landings, logbooks and observer surveys: 474 improving the protocols for sampling commercial fisheries. *Fish and Fisheries*, 8: 123–152.

476 Crowder, L. B., and Murawski, S. A. 1998. Fisheries bycatch: Implications for management. *Fisheries*, 23: 8–17. 478

Cumming, G., Fidler, F., and Vaux, D. L. 2007. Error bars in experimental biology. *Journal of Cell Biology*, 177: 7–11.

Damalas, D., and Vassilopoulou, V. 2013. Slack regulation compliance in the Mediterranean fisheries: a paradigm from the Greek Aegean Sea demersal trawl fishery, modelling discard ogives. *Fisheries Management and Ecology*, 20: 21–33.

Depestele, J., Vandemaele, S., Vanhee, W., Polet, H., Torrele, E., Leirs, H., and Vincx, M. 2011. Quantifying causes of discard variability: an indispensable assistance to discard estimation and a paramount need for policy measures. *ICES Journal of Marine Science*, 68: 1719–1725.

EEC. 2000. Council Regulation (EC) No 1543/2000 of 29 June 2000 establishing a Community framework for the collection and management of the data needed to conduct the common fisheries policy. Brussels. *Official Journal of the European Union*, L 176: 1–16.

EEC. 2008. Council regulation (EC) No 199/2008 of 25 February 2008 concerning the establishment of a community framework for the collection, management and use of data in the fisheries sector and support for scientific advice regarding the Common Fisheries Policy. *Official Journal of the European Union*, L 60: 1–12.

EEC. 2009. Council Regulation (EC) No 409/2009 of 18 December 2009 adopting a multiannual Community programme for the collection, management and use of data in the fisheries sector for the period 2011-2013. *Official Journal of the European Union*, L 41: 8–71.

504

EEC. 2011. Proposal of 13 July 2011 for a regulation of the European Parliament and of the  
506 Council on the Common Fisheries Policy. 2011/0195. Brussels. 87 pp.

508 EEC. 2012a. Proposal of 29 August 2012 for a regulation of the European parliament and of  
the council on certain technical and control measures in the Skagerrak and amending  
510 Regulation (EC) No 850/98 and Regulation (EC) No 1342/2008. Brussels. 18 pp.

512 EEC. 2012b. Fisheries: commission acts to reduce discards in the west of Scotland. Press  
release of 23 February 2012. Directorate-General for Maritime Affairs and Fisheries,  
514 Brussels.

516 EEC. 2012c. CFP reform – the discard ban. Information paper.  
[http://ec.europa.eu/fisheries/reform/docs/discards\\_en.pdf](http://ec.europa.eu/fisheries/reform/docs/discards_en.pdf).

518

Eliassen, S., Papadopoulou, N., Vassilopoulou, V., and Catchpole, T. (2013) Socio-economic  
520 and institutional incentives influencing fishers behaviour in relation to fishing practices and  
discard. Submitted manuscript (ICES-JMS-2012-435) to ICES Journal of Marine Science  
522 ‘Bycatch and Discards’ symposium issue.

524 Eurostat. 2012. <http://epp.eurostat.ec.europa.eu/portal/page/portal/eurostat/home/>.

526 FAO. 1980. International standard statistical classification of fishing gear.  
<ftp://ftp.fao.org/fi/document/cwp/handbook/annex/AnnexM1fishinggear.pdf>.

528

Feekings, J., Bartolino, V., Madsen, N., and Catchpole, T. 2012. Fishery discards: Factors  
530 affecting their variability within a demersal trawl fishery. PLoS One, 7: e36409.

- 532 Helmond, A.T.M., and Uhlmann, S.S. (eds) 2011. Report on an overview of existing discard  
data by métier, area, and member state. MariFish, London. 300 pp.  
534 <http://83.212.243.10/badminton.html>.
- 536 Hintzen, N. T., Bastardie, F., Beare, D., Piet, G. J., Ulrich, C., Deporte, N., Egekvist, J., et al.  
2012. VMStools: Open-source software for the processing, analysis and visualisation of  
538 fisheries logbook and VMS data. Fisheries Research, 115-116: 31–43.
- 540 Hinz, H., Murray, L. G., Lambert, G. I., Hiddink, J. G., and Kaiser, M. J. (2013).  
Confidentiality over fishing effort data threatens science and management progress. Fish and  
542 Fisheries. Article first published online : 22 May 2012, doi: 10.1111/j.1467-  
2979.2012.00475.x  
544
- ICES. 2007. Report of the Working Group on Discard Raising Procedures, 6-9 February  
546 2007, San Sebastian, Spain. ICES CM 2007 ACFM:06. Copenhagen.
- 548 ICES. 2009. Definition of a standard data-exchange format for sampling, landings, and effort  
data from commercial fisheries. ICES Cooperative Research Report No. 296, Copenhagen. 44  
550 pp.
- 552 ICES. 2010. Report of the ICES Workshop on ecosystem indicators of discarding (WKEID),  
28 September – 1 October 2010, Copenhagen. 60 pp.  
554
- ICES. 2011. Report of the Study Group on Practical Implementation of Discard Sampling  
556 plans (SGPIDS). ICES Advisory Science Committee. ICES CM 2011/ACOM:50,

Copenhagen. 110 pp.

558

ICES. 2012. Report of the Planning Group on Commercial Catches, Discards And Biological  
560 Sampling (PGCCDBS). 30th January – 3rd February, 2012, Rome. 163 pp.

562 Kennelly, S. J., and Broadhurst, M. K. 2002. By-catch begone: changes in the philosophy of  
fishing technology. *Fish and Fisheries*, 3: 340–355.

564

Kindt-Larsen, L., Kirkegaard, E., and Dalskov, J. 2011. Fully documented fishery: a tool to  
566 support a catch quota management system. *ICES Journal of Marine Science*, 68: 1606–1610.

568 Limpert, E., Stahel, W.A., and Abbt, M. 2001. Log-normal distributions across the sciences –  
keys and clues. *BioScience* 51, 341–352.

570

Madsen, N., Feekings, J., and Lewy, P. 2013. Discarding of plaice (*Pleuronectes platessa*) in  
572 the Danish North Sea trawl fishery. *Journal of Sea Research*, 75: 129–134.

574 Madsen, N., and Valentinsson, D. 2010. Use of selective devices in trawls to support recovery  
of the Kattegat cod stock: a review of experiments and experience. *ICES Journal of Marine*  
576 *Science*, 67: 2042–2050.

578 Nelson, G. A. 2011. Package ‘fishmethods’. March 27, 2012 Massachusetts Division of  
Marine Fisheries. <http://cran.r-project.org>, accessed on 12/06/2012.

580

Poos, J. J., Aarts, G., Vandemaele, S., Willems, W., Bolle, L. J., and van Helmond, A. T. M.  
582 2013. Estimating spatial and temporal variability of juvenile North Sea plaice from

opportunistic data. *Journal of Sea Research*, 75: 118–128.

584

Punt, A. E., Smith, D. C., Tuck, G. N., and Methot, R. D. 2006. Including discard data in fisheries stock assessments: two case studies from south-eastern Australia. *Fisheries Research*, 79: 239–250.

588

R Development Core Team. 2005. *R: a Language and Environment for Statistical Computing*.

590

R Foundation for Statistical Computing, Vienna, Austria. <http://cran.r-project.org>.

592

Rochet, M.-J., and Trenkel, V. 2005. Factors for the variability of discards: assumptions and field evidence. *Canadian Journal of Fisheries and Aquatic Sciences*, 62: 224–235.

594

STECF. 2006. Commission staff working paper report of the scientific, technical, and

596

economic committee for fisheries discards from community vessels: opinion expressed during a plenary meeting held in Ispra from 6-10 November 2006. 56 pp.

598

STECF. 2008. Commission staff working paper report of the scientific, technical, and

600

economic committee for fisheries discards from community vessels: opinion expressed during a plenary meeting of 14-18 April 2008 in Hamburg. 206 pp.

602

Stergiou, K. I., Christou, E. D., Georgopoulos, D., Zenetos, A., and Souvermesoglou, C.

604

1997. Hellenic Seas: physics, chemistry, biology and fisheries. *Oceanography and Marine Biology: An Annual Review*, 35: 415–538.

606

Stratoudakis, Y., Fryer, R. J., Cook, R. M., and Pierce, G. J. 1999. Fish discarded from

608

Scottish demersal vessels: Estimators of total discards and annual estimates for targeted

gadoids. *ICES Journal of Marine Science*, 56: 592–605.

610

Tsagarakis, K., Machias, A., Giannoulaki, M., Somarakis, S., and Karakassis, I. 2008.

612

Seasonal and temporal trends in metrics of fish community for otter-trawl discards in a Mediterranean ecosystem. *ICES Journal of Marine Science*, 65: 539–550.

614

Uhlmann, S. S., Bierman, S. M., and Helmond, A. T. M. van 2011. A method of detecting

616

patterns in mean lengths of samples of discarded fish, applied to the self-sampling

programme of the Dutch bottom-trawl fishery. *ICES Journal of Marine Science*, 68: 1712–

618

1718.

620

Viana, M., Graham, N., Wilson, J. G., and Jackson, A. L. 2011. Fishery discards in the Irish Sea exhibit temporal oscillations and trends reflecting underlying processes at an annual

622

scale. *ICES Journal of Marine Science*, 68: 221–227.

624

Wigley, S. E., Blaylock, J., Rago, P. J., Murray, K. T., Nies, T. A., Seagraves, R. J., Potts, D., Drew, K. 2012. Standardized Bycatch Reporting Methodology 3-year Review Report - 2011

626

Part 2. US Department of Commerce, Northeast Fisheries Science Centre Reference

Document 12-27; 226 pp. Available from: National Marine Fisheries Service, 166 Water

628

Street, Woods Hole, MA 02543-1026, or online at

<http://www.nefsc.noaa.gov/nefsc/publications/>



630 Tables

632 Table 1. Sampling allocation schemes, species identification and measurement procedures, and  
raising units of national discard sampling programmes part of the European Data Collection

634 Framework (DCF).

636	Programme	Allocation <sup>a</sup>	Identification <sup>b</sup>	Measurement <sup>c</sup>	Raising unit <sup>d</sup>
	Denmark				
638	All DCF-fisheries	Random	Partial	Numbers/weights	Fishing operation
	Spain				
640	Otter trawl (Med. Sea)	Opportunistic	Partial	Numbers/weights	Fishing operation
	Otter trawl (Atlantic)	Random	Partial	Numbers/weights	Fishing operation
642	France				
	All DCF-fisheries	Opportunistic	All	Numbers/weights	Fishing operation
644	England				
	All DCF-fisheries	Random	All	Numbers	Fishing operation
646	Greece				
	Otter trawl	Random	All	Numbers/weights	Fishing operation
648	Netherlands				
	Beam trawl	Opportunistic	All	Numbers	Fishing time

650 <sup>a</sup> Allocation of sampling effort. For example, how the units of the sampling frame (e.g. vessels,  
trips) were chosen: by a (stratified) random, opportunistic/cooperative design (ICES, 2011).

652 <sup>b</sup> Identification of either all or selected (partial) species within a catch sample.

<sup>c</sup> Measurement includes numbers and/or weights of discarded or landed species.

654 <sup>d</sup> Sampling unit includes the estimator used to raise species numbers/weights from haul to trip level.

656 Table 2. List of discard-intensive, towed-gear fisheries for which data were provided by country;  
together with an indication of the range of fishing and sampling effort within a given period:  
number of registered vessels, annual total and % observed fishing effort (days at sea, D.A.S.).

658

Fishery	Country	Period	No. vessels	Total D.A.S.	% observed D.A.S.
Otter trawl for crustaceans					
	Denmark	2003-08	221-350	15 719-28 152	0.29-0.55
	France	2003-08	390-504	104 310-161 280	0.11-0.26
	England	2002-08	NA	4 179-5 161	0.19-1.29
Otter trawl for fish					
	Denmark	2003-08	476-809	27 706-57 687	0.22-0.71
	Spain <sup>a</sup>	2003-07	167-210	109 683-294 673	0.05-0.12
	Spain <sup>b</sup>	2003-08	182-188	23 512-34 664	0.12-0.19
	Greece	2003-06	5-12	378-2 545	4.37-34.56
	Greece <sup>c</sup>	2003-08	326-336	53 624-59 552	0.06-0.22
	France	2003-08	1 530-1 832	550 800-616 600	0.05-0.17
	England	2002-08	NA	31 612-50 578	0.17-0.51
Beam trawl for fish					
	Denmark	1997-2008	2-17	313-2 111	0.00-5.16
	France	2003-05	42-79	15 120-27 876	0.09-0.15
	Netherlands	2003-08	99-139	14 210-21 027	0.17-0.30
	England	2002-08	NA	30 929-49 384	0.15-0.47

660 <sup>a</sup> Fishery active in North-East Atlantic ICES Divisions: VIIb; VIIc; VIIj; VIIk; VIIg; VIIh; VIIc;  
and IXa.

662 <sup>b</sup> Fishery active in the Western Mediterranean Sea: GSA3701.

664 <sup>c</sup> Different otter trawl fleets in the Greek part of the Mediterranean Sea were considered as a single  
fishery.

Table 3. Coefficients of variation (%) of discard rates, where applicable, for selected species calculated across fisheries for a given region (inter-fishery) and across regions for a given fishery (inter-region).

	Atlantic cod	Haddock	European hake	European plaice	Red mullet	Deep-water rose shrimp	Bogue
670	Inter-fishery						
	Baltic Sea	14					
672	Celtic Sea		84	83			
	Irish Sea			14			
674	Mediterranean		70		80	109	121
	North Sea	62	77	188			
676	Skagerrak	15	48				
	Inter-region						
678	Otter trawls (crustaceans)	53	63	104	114		
680	Otter trawls (fish)	43	79	126	120		
682	Beam trawls (fish)		53		62		
684							

686 Table 4. Coefficients of variation (%) of discard ratios, where applicable, for selected species, calculated across fisheries for a given region (inter-  
688 fishery) and across regions for a given fishery (inter-region).

688		Atlantic cod	Haddock	European hake	European plaice	Red mullet	Deep-water rose shrimp	Bogue
690	Inter-fishery							
	Baltic Sea	69						
692	Celtic Sea		25	3				
	Irish Sea				9			
694	Mediterranean			60		76	183	71
	North Sea	29	40		73			
696	Skagerrak	9	57					
	Inter-region							
698	Otter trawls (crustaceans)	22	35	<1	13			
700	Otter trawls (fish)	43	28	63	19			
702	Beam trawls (fish)		65		6			

704

## Figures


706

Figure 1. Discard and landings rates (with inferential error bars) of commercially-valuable  
708 species across fisheries for a given region (inter-fishery, top row) and across regions for a  
given fishery (inter-region, bottom row of plots): (a) Atlantic cod; (b) haddock; (c) European  
710 hake; and (d) European plaice, when combined across countries and ICES Divisions; and (e)  
red mullet; (f) deep-water rose shrimp; and (g) bogue when combined across countries fishing  
712 in the Mediterranean Sea. To improve visibility of bar plots, the y-axis scaling was broken  
where large differences between landings and discard rates existed. The number above each  
714 bar represent the number of observed trips (if  $\geq 4$ ).


716

718





# (a) Atlantic cod



# (b) Haddock




# (c) European hake







# (d) European plaice



**(e) Red mullet**



**(f) Deep-water rose shrimp**



**(g) Bogue**

