


Rapport interne LPO/11-05

UMR 6523 Laboratoire de Physique des Océans 	DELAYED MODE QUALITY CONTROL OF OVIDE ARGO DATA FLOAT WMO 6900490	
Date : 23 juin 2011	Auteurs : Lagadec Catherine Thierry Virginie	Archivage : LPO

Liste de diffusion :

LPO


Carole Despinoy (ODE/LPO)

DELAYED MODE QUALITY CONTROL OF OVIDE ARGO DATA FLOAT WMO 6900490

C. Lagadec - V. Thierry

23 juin 2011

Float WMO 6900490



1 Presentation and DMQC summary

Number	Deployment (cycle OD) cycle OD	Last cycle 43
Provor WMO 6900490	02/07/08 17h33	
CTS2 04-S2-77	N 59.688 W 39.598	
Date of control	Float status	Last cycle
April 2011	DEAD	07/09/09
Coriolis transmission		23/06/11

TAB. 1: Status of the float

Warning : Note that all the figures are plotted with the latest QC flag values (the modifications mentioned table 2 are taken into account).

1.1 QC flag checks and interesting profiles

Cycle	Para- meter	Vertical level	Old flag	New flag	Comments	Coriolis transmission
all cycles (except 0D)	PSAL	level 1 (where PRES inf. 7)	1	4	untrustable data	21/04/11

TAB. 2: Float 6900490. Summary of the modifications of the real-time QC flags and of the interesting or suspicious data.

The resolution is equal to 10 dbar from the surface to 800 dbar, then 25 dbar from 800 to 2000 dbar. Salinity data between 0 and 6 dbar are suspicious because they are acquired when the pump of the CTD is turned off.

1.2 Salinity correction from the OW method

We cannot see any evidence of a drift or bias in the salinity measurements. We thus conclude that it is not necessary to correct the salinity data. Errors bars are maximum value between 0.01 and those determined from the OW method with parameters from the OW configuration 1.

2 Data

OW CONFIGURATION	1	3	11
CONFIG_MAX_CASTS	250	250	250
MAP_USE_PV	1	1	1
MAP_USE_PV_ELLIPSE	1	1	1
MAP_USE_FACTEUR	1	1	1
MAPSCALE_LONGITUDE_LARGE	3.2	3.2	3.2
MAPSCALE_LONGITUDE_SMALL	0.8	0.8	0.8
MAPSCALE_LATITUDE_LARGE	2	2	2
MAPSCALE_LATITUDE_SMALL	0.5	0.5	0.5
MAPSCALE_PHL_LARGE	0.5	0.5	0.5
MAPSCALE_PHL_SMALL	0.1	0.1	0.1
MAPSCALE_AGE	0.69	0.69	0.69
MAP_P_EXCLUDE	500	500	500
MAP_P_DELTA	250	250	250
Reference data base	CTD only	CTD only	ARGO
Comments		no break point	

TAB. 3: Parameters of the OW method.





FIG. 1: Profiles position and relationship between cycle number, date and color.




FIG. 2: θ/S diagrams. (Left panel) Flags are not taken into account. (Right panel) Quality flags are taken into account.




FIG. 3: Temperature section along the float trajectory. Quality flags are not taken into account.

Float WMO 6900490 – PSAL




FIG. 4: Salinity section along the float trajectory. Quality flags are not taken into account.

Float WMO 6900490 – PRES




FIG. 5: Pression as fonction of cycle number and vertical level index along the float trajectory. Quality flags are taken into account.




FIG. 6: Potential temperature, salinity and potential density sections along the float trajectory (interpolated on standard levels). Quality flags are taken into account.

3 Comparison to the OVIDE 2008 nearest CTD profile




FIG. 7: Comparison of the cycle 0A with the nearest CTD profile done after the float deployment.

4 Cycle 10 - Comparison to the nearest historical CTD profiles





FIG. 8: Flotteur 6900490, cycle 10. Upper panel : Position of the Argo profile (red) and of the nearest CTD profiles (black). The nearest CTD profile in time is in magenta while the nearest CTD profile in space is in blue. Lower panels : Temperature, salinity and potential density as function of pressure for the Argo profile (stars) and for the nearest CTD profile in time (magenta line) and for the nearest CTD profile in space (blue line). The color of the Argo profile represents the QC flag (green for a QC=1 ; blue for a QC=2 ; orange for a QC=3 and red for a QC=4).

6900490 – Cycle 10



6900490 – Cycle 10 – Date Argo profile 12–Oct–2008
 Dates historicals profiles 30–Aug–1997 (blue) and 14–Aug–2002 (magenta)




FIG. 9: Float 6900490, cycle 10. The Argo profile (stars) is compared to the nearest CTD profiles (black line) and to two specific profiles : the nearest profile in time (magenta) and the nearest profile in space (blue). The color of the Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4). (Upper panels) Temperature (left panel), salinity (middle panel) and potential density (right panel) as function of pressure. (Lower panels) θ/S diagrams.

5 Cycle 10 - Comparaison to the nearest ARGO profiles





FIG. 10: Flotteur 6900490, cycle 10. Upper panel : Position of the analysed Argo profile (red) and of the nearest Argo profiles (black). The nearest Argo profile in time is in magenta while the nearest CTD profile in space is in blue. Lower panels : Temperature, salinity and potential density as function of pressure for the analysed Argo profile (stars) and for the nearest Argo profile in time (magenta line) and for the nearest Argo profile in space (blue line). The color of the analysed Argo profile represents the QC flag (green for a QC=1 ; blue for a QC=2 ; orange for a QC=3 and red for a QC=4).

6900490 – Cycle 10



6900490 – Cycle 10 – Date Argo profile 12–Oct–2008
 Dates historicals profiles 06–Dec–2009 (blue) and 22–Jan–2009 (magenta)




FIG. 11: Float 6900490, cycle 10. The analysed Argo profile (stars) is compared to the nearest Argo profiles (black line) and to two specific profiles : the nearest Argo profile in time (magenta) and the nearest Argo profile in space (blue). The color of the analysed Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4). (Upper panels) Temperature (left panel), salinity (middle panel) and potential density (right panel) as function of pressure. (Lower panels) θ/S diagrams.

6 Cycle 12 - Comparison to the nearest historical CTD profiles





FIG. 12: Flotteur 6900490, cycle 12. Upper panel : Position of the Argo profile (red) and of the nearest CTD profiles (black). The nearest CTD profile in time is in magenta while the nearest CTD profile in space is in blue. Lower panels : Temperature, salinity and potential density as function of pressure for the Argo profile (stars) and for the nearest CTD profile in time (magenta line) and for the nearest CTD profile in space (blue line). The color of the Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4).

6900490 – Cycle 12



6900490 – Cycle 12 – Date Argo profile 01–Nov–2008
 Dates historicals profiles 08–Aug–1997 (blue) and 03–Jul–2005 (magenta)




FIG. 13: Float 6900490, cycle 12. The Argo profile (stars) is compared to the nearest CTD profiles (black line) and to two specific profiles : the nearest profile in time (magenta) and the nearest profile in space (blue). The color of the Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4). (Upper panels) Temperature (left panel), salinity (middle panel) and potential density (right panel) as function of pressure. (Lower panels) θ/S diagrams.

7 Cycle 12 - Comparaison to the nearest ARGO profiles





FIG. 14: Flotteur 6900490, cycle 12. Upper panel : Position of the analysed Argo profile (red) and of the nearest Argo profiles (black). The nearest Argo profile in time is in magenta while the nearest CTD profile in space is in blue. Lower panels : Temperature, salinity and potential density as function of pressure for the analysed Argo profile (stars) and for the nearest Argo profile in time (magenta line) and for the nearest Argo profile in space (blue line). The color of the analysed Argo profile represents the QC flag (green for a QC=1 ; blue for a QC=2 ; orange for a QC=3 and red for a QC=4).

6900490 – Cycle 12



6900490 – Cycle 12 – Date Argo profile 01–Nov–2008
 Dates historical profiles 21–Dec–2005 (blue) and 06–Mar–2008 (magenta)




FIG. 15: Float 6900490, cycle 12. The analysed Argo profile (stars) is compared to the nearest Argo profiles (black line) and to two specific profiles : the nearest Argo profile in time (magenta) and the nearest Argo profile in space (blue). The color of the analysed Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4). (Upper panels) Temperature (left panel), salinity (middle panel) and potential density (right panel) as function of pressure. (Lower panels) θ/S diagrams.

8 Cycle 15 - Comparison to the nearest historical CTD profiles





FIG. 16: Flotteur 6900490, cycle 15. Upper panel : Position of the Argo profile (red) and of the nearest CTD profiles (black). The nearest CTD profile in time is in magenta while the nearest CTD profile in space is in blue. Lower panels : Temperature, salinity and potential density as function of pressure for the Argo profile (stars) and for the nearest CTD profile in time (magenta line) and for the nearest CTD profile in space (blue line). The color of the Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4).

6900490 – Cycle 15



6900490 – Cycle 15 – Date Argo profile 01–Dec–2008
 Dates historicals profiles 15–Aug–1991 (blue) and 15–Jun–2006 (magenta)




FIG. 17: Float 6900490, cycle 15. The Argo profile (stars) is compared to the nearest CTD profiles (black line) and to two specific profiles : the nearest profile in time (magenta) and the nearest profile in space (blue). The color of the Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4). (Upper panels) Temperature (left panel), salinity (middle panel) and potential density (right panel) as function of pressure. (Lower panels) θ/S diagrams.

9 Cycle 15 - Comparaison to the nearest ARGO profiles





FIG. 18: Flotteur 6900490, cycle 15. Upper panel : Position of the analysed Argo profile (red) and of the nearest Argo profiles (black). The nearest Argo profile in time is in magenta while the nearest CTD profile in space is in blue. Lower panels : Temperature, salinity and potential density as function of pressure for the analysed Argo profile (stars) and for the nearest Argo profile in time (magenta line) and for the nearest Argo profile in space (blue line). The color of the analysed Argo profile represents the QC flag (green for a QC=1 ; blue for a QC=2 ; orange for a QC=3 and red for a QC=4).

6900490 – Cycle 15



6900490 – Cycle 15 – Date Argo profile 01–Dec–2008
 Dates historicals profiles 23–Aug–2009 (blue) and 13–Nov–2008 (magenta)




FIG. 19: Float 6900490, cycle 15. The analysed Argo profile (stars) is compared to the nearest Argo profiles (black line) and to two specific profiles : the nearest Argo profile in time (magenta) and the nearest Argo profile in space (blue). The color of the analysed Argo profile represents the QC flag (green for a QC=1; blue for a QC=2; orange for a QC=3 and red for a QC=4). (Upper panels) Temperature (left panel), salinity (middle panel) and potential density (right panel) as function of pressure. (Lower panels) θ/S diagrams.

10 OW method, CONFIGURATION # 1




FIG. 20: Figures from the OW method. (Left) Position of the historical and float data. (Right) Comparison, on various θ levels, between the float data and the historical data interpolated at the float position.




FIG. 21: Figures from the OW method. Comparison of the θ/S diagram of the float with the historical database. (left) raw data; (right) corrected data using the OW correction.




FIG. 22: Figures from the OW method. Salinity anomaly : (left) raw data ; (right) corrected data using the OW correction .




FIG. 23: Correction proposed by the OW method.




FIG. 24: Chosed levels by the OW method.

11 OW method, CONFIGURATION # 3




FIG. 25: Figures from the OW method. (Left) Position of the historical and float data. (Right) Comparison, on various θ levels, between the float data and the historical data interpolated at the float position.




FIG. 26: Figures from the OW method. Comparison of the θ/S diagram of the float with the historical database. (left) raw data; (right) corrected data using the OW correction.




FIG. 27: Figures from the OW method. Salinity anomaly : (left) raw data ; (right) corrected data using the OW correction .




FIG. 28: Correction proposed by the OW method.




FIG. 29: Chosed levels by the OW method.

12 OW method, CONFIGURATION # 11




FIG. 30: Figures from the OW method. (Left) Position of the historical and float data. (Right) Comparison, on various θ levels, between the float data and the historical data interpolated at the float position.




FIG. 31: Figures from the OW method. Comparison of the θ/S diagram of the float with the historical database. (left) raw data; (right) corrected data using the OW correction.




FIG. 32: Figures from the OW method. Salinity anomaly : (left) raw data ; (right) corrected data using the OW correction .




FIG. 33: Correction proposed by the OW method.




FIG. 34: Chosed levels by the OW method.