

Mapping of *Posidonia oceanica* beds around Elba Island (western Mediterranean) with integration of direct and indirect methods

Luigi PIAZZI, Stefano ACUNTO *, Francesco CINELLI

Dipartimento di Scienze dell'Uomo e dell'Ambiente, Università di Pisa, Via A. Volta 6, 56126 Pisa, Italy

Received 18 October 1999; revised 14 February 2000; accepted 14 February 2000

Abstract – Direct and indirect methods were used for mapping *Posidonia oceanica* beds around Elba Island. Side scan sonar and a visual infra red scanner were used to detect the extension of beds. Also, direct observations were adopted to assess the bed density and to calibrate the indirect methods. Samples of *P. oceanica* were collected and phenological variables and macroalgal epiphytes were studied to assess the health of the beds. *P. oceanica* around Elba Island covered an area of 3 680 ha. The largest beds occupied the four main gulfs of the island, characterized by gentle slopes and sandy bottoms. There, the beds showed uniform cover while a patchy structure was common on the rocky and steep bottoms of the western and south-eastern coasts of the island. *P. oceanica* beds at Elba Island demonstrated a good state of health. The epiphytic algal assemblage of the leaves was dominated by species generally found on this habitat, while, on the rhizomes, we observed a high covering of turf-forming species. The methods adopted allowed accurate maps at a scale of 1:10 000. The integration of data obtained both by direct and indirect methods and the precision of the positioning system were particularly important. © 2000 Ifremer/CNRS/IRD/Éditions scientifiques et médicales Elsevier SAS

Posidonia oceanica / seagrass / algae / mapping / airborne teledetection / Mediterranean

Résumé – Cartographie des herbiers à *Posidonia oceanica* de l'île d'Elbe (Méditerranée occidentale) par intégration de méthodes directes et indirectes. Le sonar latéral et la télédétection aérienne ont été utilisés pour la localisation des herbiers, tandis que des observations directes en plongée autonome et au moyen de véhicules sous-marins l'ont été pour identifier certaines images ou pour mesurer la densité des herbiers. Des faisceaux de *P. oceanica* ont été prélevés pour l'étude de la phénologie de la plante et de la communauté macroalgale épiphyte, de manière à évaluer la condition écologique des herbiers. *P. oceanica* autour de l'île d'Elbe couvre une surface de 3 680 ha. La pente limite l'expansion des herbiers, d'où une étendue très basse par rapport à la longueur de la côte (25,3 ha km⁻¹). Les herbiers les plus étendus apparaissent dans les baies principales de l'île : Campo, Lacona, Stella et Procchio. Les herbiers y présentent une couverture uniforme, tandis qu'une structure en taches est commune sur les fonds rocheux et à pente forte, le long des côtes occidentales et sud-orientales. Les paramètres structuraux et phénologiques traduisent une bonne condition écologique des herbiers à *P. oceanica*. De petites aires de régression sont présentes autour des ports principaux et le long de la côte nord-orientale, où les effets d'anciennes mines sont encore visibles. Les méthodes utilisées ont permis la réalisation de cartes très précises à une échelle de 1:10 000. L'intégration des données directes et indirectes et la précision du système de positionnement (DGPS) ont été particulièrement importantes pour la connaissance de l'étendue des herbiers et de leurs situation écologique. Ces résultats constituent un document pour la gestion du système littoral de l'île d'Elbe. © 2000 Ifremer/CNRS/IRD/Éditions scientifiques et médicales Elsevier SAS

Posidonia oceanica / algues / cartographie / sonar latéral / télédétection aérienne / Méditerranée

* Correspondence and reprints: acunto@discat.unipi.it

1. INTRODUCTION

Cartography of the sea bottom plays a key role in coastal management. Knowledge of the distribution of the main marine biocenoses is of fundamental importance in conservation and monitoring programs. Sea bottom surveys have often involved mapping of seagrasses [7, 12, 14, 19, 26, 28, 43]. Seagrass beds are among the most important benthic communities in temperate coastal areas, where they have a fundamental ecological role. They produce organic matter, maintain the sedimentary equilibrium and protect sandy shores from erosion [15, 22, 27, 30]. The regression of seagrass beds observed throughout the world and caused by increasing pollution [42] needs accurate monitoring.

The knowledge of the extension of a bed and the definition of its limits must be considered as only part of the information necessary to manage this resource. It is very important to acquire data on some structural (density) and phenological characteristics which are indicators of the health of the beds and plants

respectively [24, 25]. Also, of interest is the epiphytic community that can give information on the whole system (possible stress can be detected on the epiphytic community before seagrass is damaged) [16, 29].

In the Mediterranean Sea, mapping programs have been carried out to detect the distribution of the endemic marine phanerogam *Posidonia oceanica* (L.) Delile, widely distributed in a depth range from 0 to 40 m. The aim of this work was to implement protective measures and to evaluate the resources to be managed.

Underwater inspections by scuba diving were made in restricted areas [3, 13, 21, 23, 31, 38, 40] while indirect methods were employed in large scale mapping programs. Among the different methods used to map *P. oceanica*, side scan sonar [9, 17, 18, 20, 32, 37] was demonstrated to be the best instrument to map the lower limits of the meadows while aerial photographs [13, 33] and satellite images [8, 37] appeared to be techniques particularly suitable for the shallower layer. Thus, the best results have been obtained with the integration of several methods, chosen on the basis of the characteristics of the area, the maximum depth to be reached and the precision required [5, 6, 10, 34, 37].

In this study we present the results of a mapping program of the *P. oceanica* beds around Elba Island (Italy), realised using both direct and indirect methods. We examine the potential of the techniques adopted, discuss the results in relation to other Mediterranean coasts and give information on the health of the beds.

2. MATERIALS AND METHODS

Elba is the largest island of the Tuscan Archipelago (Western Mediterranean, Italy) (*figure 1*), with a surface area of 223.5 km² and a coast 147 km long.

Mapping of *Posidonia oceanica* meadows was carried out from April to July 1995. Side scan sonar (hereafter SSS) was used to discriminate *P. oceanica* beds from rocky and sandy bottoms. We used two recorders (graphic and digital) and a 'fish' bearing two transducers transmitting an acoustic signal at a frequency variable from 100 to 500 kHz. The return




Fig. 1. Location of Elba Island.




Fig. 2. Distribution of *Posidonia oceanica* beds (in black) around Elba Island. Numbers indicate the following localities. 1: Nasuto Cape; 2: Gulf of Procchio; 3: Gulf of Viticcio; 4: Enfola Cape; 5: Portoferraio; 6: Inferno Bay; 7: Portoazzurro; 8: Gulf of Stella; 9: Gulf of Lacona; 10: Gulf of Campo.

signal was picked up by the 'fish' and transmitted to the recorders on board ship. The 'fish' was dragged at a maximal speed of 5 knots (9.26 km h^{-1}). The ship's position was determined by a differential global positioning system (DGPS Trimble, precision $\pm 1 \text{ m}$). The data of SSS were real time interfaced with the positioning system using the Trimble hydro software. The SSS was used at a frequency of 100 kHz, covering a 200-m-wide band of sea bottom (100 m for each side). Sonograms at 500 kHz were prepared when a higher resolution was necessary. Navigation lines ran parallel to the coast 150 m apart, in order to allow sonograms to overlap. Some lines perpendicular to the coast were also made.

Airborne teledetection was obtained from the visible infrared scanner 201 (VIRS 201) in September 1994 and in April 1995. The VIRS is a multi-

spectral imaging radiometer bearing two separate sensors acquiring data in the visible, near infrared (400–1 000 nm) and thermic infrared (8–12 nm) wave lengths. The number 201 indicates the possibility to detect 20 bands in the VIS- near infrared and one in the thermic infrared.

The image built is a line scanner type; the instrument is able to acquire wide areas through a succession of narrow bands perpendicular to the fly direction. Data are recorded and then analysed by dedicated software for PC.

Direct observations by scuba diving and remote operated vehicle were used to calibrate indirect methods. From the maps we calculated the surface area (ha) of *Posidonia oceanica* beds and the R_1 index equal to the seagrass bottom cover per kilometre of coastline [35].

Moreover, information on the health of the beds was obtained through the study of structural and phenological variables of the beds and of epiphytic flora. This study was carried out in July 1995 at three localities (Inferno Bay, Nasuto Cape, Gulf of Viticcio) (figure 2) randomly chosen from a set of all the possible localities around the island. Shoot density was assessed in five 0.1 m² samples collected by scuba diving at depths ranging from 10 to 20 m. At each locality ten shoots were sampled and, in the laboratory, the following phenological variables were calculated: mean number of leaves, mean leaf length per shoot and Leaf Area Index (mean leaf surface per shoot/2 × mean shoot density) [24, 25, 36]. Macroalgal epiphytes of leaves and rhizomes were identified in the laboratory using a binocular microscope.

3. RESULTS

Seven bionomic maps and seven bathymetric maps at a scale of 1:10 000 were realised (available at *Dipartimento di Scienze dell'Uomo e dell'Ambiente*, University of Pisa). *Posidonia oceanica* stands showed a continuous distribution along the coast of Elba Island (figure 2). The total surface colonised by the seagrass at Elba Island was 3 680 ha, and R₁ was 25.03 ha km⁻¹. The distribution of the seagrass was affected by the morphology of the bottom. The largest beds were in the main gulfs of the Island: Procchio at the north-side, Campo, Lacona and Stella at the south-side. Wide beds were also found between Enfola Cape and Portoferraio and in the north-eastern part of the island. Upper limits occurred at depths ranging between 3 and 10 m, while lower limits were localised between 30 and 40 m.

P. oceanica often colonised rocky bottoms, in particular along the western, north-western and south-eastern coasts of Elba, where meadows showed a patchy distribution. Plants showed a uniform cover on the

sandy bottoms present in the main gulfs, where wide matte structures were also found. Meadows showed eroded edges along north-eastern and south-western coasts and restricted damaged areas near Rio Marina and in the Bays of Portoferraio and Porto Azzurro (figure 2).

The shoot density in the studied beds ranged from 500 ± 55 to 675 ± 95 shoots m⁻². Leaf Area Index varied from 8.4 ± 0.1 to 18.6 ± 9.5. The mean number of leaves ranged between 4.8 ± 0.3 and 6.2 ± 0.5 leaves/shoot while the mean leaf length between 40.8 ± 9.9 and 62.5 ± 4.2 cm (table I).

Twenty-eight macroalgal species were found on leaves (1 Chlorophyta, 9 Fucophyceae, 18 Rhodophyta) and fifty-one on rhizomes (10 Chlorophyta, 7 Fucophyceae, 34 Rhodophyta) (table II). Macroalgal assemblages on leaves were dominated by the encrusting Corallinaceae *Hydrolithon farinosum* (Lamouroux) Penrose and Chamberlain, *H. cruciatum* (Bressan) Chamberlain and *Pneophyllum fragile* Kützing and by the Fucophyceae *Myronema orbiculare* J. Agardh. On the rhizomes the most abundant species were the Rhodophyta *Acrothamnion preissii* (Sonder) Wollaston, *Womersleyella setacea* (Hollenberg) R.E. Norris and *Peyssonnelia* spp.

4. DISCUSSION AND CONCLUSIONS

The distribution of *Posidonia oceanica* beds around Elba Island was strictly linked to the morphology of the bottom. A steep slope often limited the extension of the beds explaining a low value of the seagrass bottom cover compared to the length of the coastline (25.03 ha·km⁻¹) (table III). Despite the short distance from the mainland, Elba meadows appeared more similar to those of other islands of the Tuscan Archipelago [18] than to the meadows occurring along the continental coasts. In fact, the surface area

Table I. Structural and phenological variables of *Posidonia oceanica* measured in three different locations of Elba Island.

| | Inferno Bay | Nasuto Cape | Gulf of Viticcio |
|----------------------------------|-------------|-------------|------------------|
| Density (shoot·m ⁻²) | 500 ± 55 | 675 ± 95 | 613 ± 60 |
| Mean number of leaves per shoot | 6.4 ± 0.2 | 6.2 ± 0.5 | 4.8 ± 0.3 |
| Mean leaf length per shoot | 40.8 ± 9.9 | 62.5 ± 4.2 | 55.2 ± 5.2 |
| Leaf area index | 8.4 ± 0.1 | 18.6 ± 9.5 | 16.2 ± 2.8 |

Table II. Floristic list of macroalgal epiphytes of *Posidonia oceanica*.

| Taxa | Leaves | Rhizomes |
|---|--------|----------|
| CHLOROPHYTA | | |
| <i>Chaetomorpha mediterranea</i> (Kützing) Kützing | | + |
| <i>Cladophora albida</i> (Nees) Kützing | + | + |
| <i>Cladophora coelotrix</i> Kützing | | + |
| <i>Cladophora echinus</i> (Biaioletto) Kützing | | + |
| <i>Cladophora prolifera</i> (Roth) Kützing | | + |
| <i>Cladophora vagabunda</i> (Linnaeus) Hoek | | + |
| <i>Flabellia petiolata</i> (Turra) Nizamuddin | | + |
| <i>Pseudochlorodesmis furcellata</i> (Zanardini) Børgesen | | + |
| <i>Valonia macrophysa</i> Kützing | | + |
| <i>Valonia utricularis</i> Kützing | | + |
| FUCOPHYCEAE | | |
| <i>Arthrocladia villosa</i> (Hudson) Duby | | + |
| <i>Asperococcus compressus</i> Griffiths ex Hooker | + | |
| <i>Cladosiphon cylindricus</i> (Sauvageau) Kylin | + | + |
| <i>Dictyopteris polypodioides</i> (De Candolle) Lamouroux | | + |
| <i>Dictyota linearis</i> (C. Agardh) Greville | + | + |
| <i>Feldmannia caespitula</i> (J. Agardh) Knoepfler-Péguy | + | |
| <i>Giraudia sphaclarioides</i> Derbés et Solier | + | |
| <i>Halopteris filicina</i> (Grateloup) Kützing | | + |
| <i>Myriactula gracilis</i> Van der Ben | + | |
| <i>Myriactula rivulariae</i> (Suhr) J. Feldmann | + | + |
| <i>Myrionema orbiculare</i> J. Agardh | + | |
| <i>Sphaclaria cirrosa</i> (Roth) C. Agardh | + | + |
| RHODOPHYTA | | |
| <i>Acrosorium venulosum</i> (Zanardini) Kylin | | + |
| <i>Acrothamnion preissii</i> (Sonder) Wollaston | + | + |
| <i>Anthithamnion cruciatum</i> (C. Agardh) Nägeli | | + |
| <i>Anthithamnion ogdeniae</i> Abbott | | + |
| <i>Apoglossum gregarium</i> (Dawson) Wynne | + | + |
| <i>Asparagopsis armata</i> Harvey (sporophyte) | | + |
| <i>Botryocladia boergesenii</i> J. Feldmann | | + |
| <i>Ceramium codii</i> (Richards) G. Feldmann | | + |
| <i>Ceramium diaphanum</i> (Lighfoot) Roth | + | + |
| <i>Ceramium flaccidum</i> (Kützing) Ardissonne | + | |
| <i>Chondria mairei</i> G. Feldmann | + | |
| <i>Crouania attenuata</i> (C. Agardh) J. Agardh | | + |
| <i>Dasya corymbifera</i> J. Agardh | | + |
| <i>Dasya ocellata</i> (Grateloup) Harvey | + | |
| <i>Erythrotrichia carnea</i> (Dillwyn) J. Agardh | + | + |

Table II. (Continued)

| Taxa | Leaves | Rhizomes |
|--|--------|----------|
| <i>Eupogodon spinellus</i> (C. Agardh) Kützing | | + |
| <i>Feldmannophycus raissiae</i> (J. et G. Feldmann) Augier et Boudouresque | | + |
| <i>Halopitys incurvus</i> (Hudson) Batters | | + |
| <i>Herposiphonia secunda</i> (C. Agardh) Ambronn | + | + |
| <i>Heterosiphonia crispella</i> (C. Agardh) Wynne | | + |
| <i>Hydrolithon cruciatum</i> (Bressan) Chamberlain | + | |
| <i>Hydrolithon farinosum</i> (Lamouroux) Penrose et Chamberlain | + | |
| <i>Hypoglossum hypoglossoides</i> (Stackhouse) Collins et Harvey | | + |
| <i>Jania rubens</i> Lamouroux | | + |
| <i>Laurencia obtusa</i> (Hudson) Lamouroux | | + |
| <i>Lomentaria chylocladiella</i> Funk | + | |
| <i>Monosporus pedicellatus</i> (Smith) Solier | | + |
| <i>Peyssonnelia bornetii</i> Boudouresque et Denizot | | + |
| <i>Peyssonnelia rubra</i> (Greville) J. Agardh | | + |
| <i>Plenosporium borneri</i> (Smith) Nägeli | + | |
| <i>Plocamium cartilagineum</i> (Linnaeus) Dixon | | + |
| <i>Pneophyllum fragile</i> Kützing | + | |
| <i>Polysiphonia furcellata</i> (C. Agardh) Harvey | | + |
| <i>Polysiphonia scopulorum</i> (Harvey) Womersley | + | + |
| <i>Polysiphonia subulifera</i> (C. Agardh) Harvey | + | + |
| <i>Pterothamnion plumula</i> (Ellis) Nägeli | | + |
| <i>Ptilothamnion pluma</i> (Dillwyn) Thuret | | + |
| <i>Rhodophyllis divaricata</i> (Stackhouse) Papenfuss | | + |
| <i>Seirospora sphaerospora</i> J. Feldmann | + | + |
| <i>Spermothamnion flabellatum</i> Bornet | + | |
| <i>Spyridia filamentosa</i> (Wulfen) Harvey | | + |
| <i>Stylonema alsidii</i> (Zanardini) Drew | + | + |
| <i>Womersleyella setacea</i> (Hollenberg) R.E. Norris | + | + |

(ha) occupied by *P. oceanica* along these coasts was much larger than that occupied along the coasts of the islands (table III). Moreover, the continental meadows showed large damaged areas produced by anthropic activities and densities of leaves were generally low [17]. On the contrary, as observed for the other islands [18], structural and phenological variables demonstrate that Elba seagrass meadows are healthy. Nevertheless, restricted areas of regression were found close to the main harbours and along the

Table III. Surface area (ha) of *Posidonia oceanica* beds in different zones of the north-western coasts of Italy.

| Reference | Zone | Surface area (ha) | Coastline length (km) | Surface area per km of coastline (ha·km ⁻¹) |
|---------------------------|--------------------|-------------------|-----------------------|---|
| Present study | Elba Island | 3 680 | 147 | 25.03 |
| Cinelli and Piazzì, 1990 | Tuscany | 20 277 | 300 | 67.59 |
| Cinelli et al., 1995 | Tuscan Archipelago | 5 043 | 125 | 40.34 |
| Bianchi and Peirano, 1995 | Liguria | 4 800 | 332 | 14.46 |

north-eastern coast of Elba where the effects of ancient mine activity are still visible. This points to the sensitivity of *P. oceanica* to human activities and is why it is regarded as a good biological indicator of water quality of coastal Mediterranean areas [39].

The epiphytic algal assemblage of the leaves was dominated by species generally found in this habitat. While, on the rhizomes, it was characterized by a high covering of the introduced Rhodophyta *Acrothamnion preissii* and *Womersleyella setacea*. The development of these turf-forming species changed the structure of the population of epiphytes, with a decrease in the number of species. This phenomenon was described in the other islands of the Tuscan Archipelago [41] and along the continental coasts of Tuscany [2].

The declaration of a marine national park in Elba Island requires a detailed knowledge of the distribution of marine biocenoses [4, 23]. The maps obtained should be a basis for monitoring of *P. oceanica* beds and a fundamental tool for administrators and decision makers. In particular, data about the extension of beds and their current state of conservation should contribute to adequate coastal management. These objectives can be obtained through an extremely precise cartographic representation at an adequate scale. The methods used in this work revealed a high efficiency that relied upon the integration of data obtained by different methods and the precision of the positioning system. The use of VIRS 201 gave good results, providing a fast acquisition of upper limits of the beds. At the moment, side scan sonar is the best method for large scale acquisition of the extent of the beds and of their deep limits. Moreover, recent studies have suggested the possibility of detecting information about structural and phenological variables [1, 11].

Acknowledgements

Mapping of *Posidonia oceanica* beds around Elba Island was funded by the Italian *Ministero dell'Ambiente, Ispettorato centrale per la difesa del mare*. We are grateful to Fabio Rindi for his help with the classification of algae.

REFERENCES

- [1] Acunto, S., Lyons, A.P., Pouliquen, E., Some anatomical features of *Posidonia oceanica* leaves and their importance to acoustic scattering, *Biol. Mar. Medit.* (2000) in press.
- [2] Airoidi L., Rindi F., Cinelli F., Structure, seasonal dynamics and reproductive phenology of a filamentous turf assemblage on a sediment influenced rocky subtidal shore, *Bot. Mar.* 38 (1995) 227–237.
- [3] Astier, J.M., Cartographie des fonds marins de la région de Toulon par le groupe Ecomair, *Ann. Soc. Sci. Nat. Archéol. Toulon* (1972) 1–15.
- [4] Augier H., Boudouresque C.F., Végétation marine de l'île de Port-Cros (Parc national), VI. Le récif-barrière La Baie de Port-Man et le problème de la régression de l'herbier de Posidonies, *Bull. Mus. Hist. nat. Marseille* 30 (1970) 145–164.
- [5] Augier H., Boudouresque C.F., Végétation marine de l'île de Port-Cros (Parc national), VI. Le récif-barrière de Posidonies, *Bull. Mus. Hist. Nat. Marseille* 30 (1970) 221–228.
- [6] Augier H., Boudouresque C.F., Végétation marine de l'île de Port-Cros (Parc national), XII. Documents pour la carte des peuplements benthiques, *Trav. Sci. Parc. nation. Port-Cros* 2 (1976) 9–22.
- [7] Bay D., Demoulin V., The seagrass beds of Hansa Bay (north coast of Papua New Guinea), *Bull. Soc. Roy. Bot. Belgique* 122 (1989) 3–17.
- [8] Belsher T., Meinesz A., Lefevre J.R., Boudouresque C.F., Simulation of SPOT satellite imagery for charting shallow water benthic communities in the Mediterranean, *Mar. Ecol. P.S.Z.N.I.* 9 (1988) 157–165.

- [9] Bianchi, C.N., Peirano, A., Atlante delle fanerogame marine della Liguria: *Posidonia oceanica* e *Cymodocea nodosa*, ENEA Centro ricerche ambiente marino, La Spezia, 1995, 146 p.
- [10] Boudouresque C.F., Meinesz A., Lefevre J.R., Cartographie des peuplements benthiques marins de la Corse: I : la formation récifale à *Posidonia oceanica* de Saint-Florent, Ann. Inst. océanogr. Paris 61 (1985) 27–38.
- [11] Bozzano, R., Castellano, L., Utilizzo di un sonar ad alta frequenza per la mappatura di praterie di *Posidonia oceanica*, Biol. Mar. Medit. (2000) in press.
- [12] Bulthuis D.A., Distribution of seagrasses in a North Puget Sound estuary: Padilla Bay, Washington, USA, Aquat. Bot. 50 (1995) 99–105.
- [13] Calvo S., Fradà-Orestano C., Abbadessa P., The suitability of a topographical instrument for an integrated approach to the cartography of *Posidonia oceanica* meadows, Oceanol. Acta 16 (1993) 273–278.
- [14] Cambridge M.L., McComb A.J., The loss of seagrass in Cockburn Sound, Western Australia, I. The time course and magnitude of seagrass decline in relation to industrial development, Aquat. Bot. 20 (1984) 229–243.
- [15] Cambridge M.L., Hocking P.J., Annual primary production and nutrient dynamics of the seagrasses *Posidonia sinuosa* and *Posidonia australis* in south-western Australia, Aquat. Bot. 59 (1997) 277–295.
- [16] Casola E., Scardi M., Mazzella L., Fresi E., Structure of the epiphytic community of *Posidonia oceanica* leaves in a shallow meadow, Mar. Ecol. P.S.Z.N.I. 8 (1987) 285–296.
- [17] Cinelli, F., Piazzì, L., Mappatura delle praterie a *Posidonia oceanica* (L.) Delile lungo le coste Toscane, CIBM Livorno Relazione Tecnica interna, 1990, 80 p.
- [18] Cinelli F., Pardi G., Papi I., Benedetti-Cecchi L., Mappatura delle praterie a *Posidonia oceanica* (L.) Delile intorno alle isole minori dell'Arcipelago Toscano, Atti Soc. Tosc. Sc. Nat. 102 (1995) 93–110.
- [19] Clark C.D., Riplet H.T., Green E.P., Edwards A.J., Mumby P.J., Mapping and measurement of tropical coastal environments with hyperspectral and height spatial resolution data, Internat. J. Remote Sens. 18 (1997) 237–242.
- [20] Colantoni P., Gallignani P., Fresi E., Cinelli F., Patterns of *Posidonia oceanica* (L.) Delile beds around the Island of Ischia (Gulf of Naples) and in adjacent waters, Mar. Ecol. P.S.Z.N.I. 3 (1982) 53–74.
- [21] Falconetti C., Meinesz A., Charting the seaward limit of *Posidonia oceanica* meadows and of circalittoral biocoenoses along the coast of Monaco, Oceanol. Acta 12 (1989) 443–447.
- [22] Fonseca M.S., Fisher J.S., A comparison of canopy friction and sediment movement between four species of seagrasses with reference to their ecology and restoration, Mar. Ecol. Prog. Ser. 29 (1986) 15–22.
- [23] Gili J.M., Ros J.D., Study and cartography of the benthic communities of Medes islands (NE Spain), Mar. Ecol. P.S.Z.N.I. 6 (1985) 219–238.
- [24] Giraud G., Essai de classement des herbiers de *Posidonia oceanica* (L.) Delile, Bot. Mar. 20 (1977) 487–491.
- [25] Giraud G., Sur un méthode de mesure et de comptage des structures foliaires de *Posidonia oceanica* (L.) Delile, Bull. Mus. Hist. Nat. Marseille 34 (1979) 33–39.
- [26] Kirkman H., Baseline and monitoring methods for seagrass meadows, J. Env. Manag. 47 (1996) 191–201.
- [27] Kuo, J., Phillips, R.C., Walker, D.I., Kirkman, H., Seagrass Biology, Sciences UWA Publ. Univ. Western Australia, Netherlands, W.A.6907, 1996, 366 p.
- [28] Larkum A.W.D., West R.J., Long term changes of seagrass meadows in Botany Bay, Australia, Aquat. Bot. 37 (1990) 55–70.
- [29] May V., The use of epiphytic algae to indicate environmental changes, Aust. J. Ecol. 7 (1982) 101–102.
- [30] Mazzella L., Zupo V., Reti trofiche e flussi di energia nei sistemi a fanerogame marine, G. Bot. It. 129 (1995) 337–350.
- [31] Meinesz A., Laurent R., Cartographie et état de la limite inférieure de l'herbier de *Posidonia oceanica* dans les Alpes Maritimes - France. Campagne Poseidon, Bot. Mar. 21 (1978) (1976) 513–526.
- [32] Meinesz A., Cuvalier M., Laurent R., Méthodes récentes de cartographie et de surveillance des herbiers de phanérogame marines, Leurs applications sur les côtes françaises de la Méditerranée, Vie et Milieu 31 (1981) 27–34.
- [33] Meinesz A., Simonian M., Cartes de la végétation sous-marine des Alpes Maritimes (Côtes françaises de la Méditerranée) II: La végétation mixte à *Cymodocea nodosa*, *Zostera noltii*, *Caulerpa prolifera* et la limite supérieure de l'herbier de *Posidonia oceanica* entre Juan-les-pins et le Golfe Juan, Annls Inst. Océanogr. Paris 59 (1983) 21–35.
- [34] Meinesz A., Boudouresque C.F., Lefevre J.R., A map of the *Posidonia oceanica* beds of Marina d'Elbu (Corsica, Mediterranean), Mar. Ecol. P.S.Z.N.I. 9 (1988) 243–252.
- [35] Meinesz A., Belsher T., Boudouresque C.F., Lefevre J.R., Première évaluation des potentialités du satellite SPOT pour la cartographie des peuplements benthiques superficiels de Méditerranée Occidentale, Oceanol. Acta 14 (1991) 299–307.
- [36] Panayotidis, P., Contribution à l'étude qualitative et quantitative de l'association *Posidonietum oceanicae* Funk 1927, thèse 3^e cycle Univ. Aix-Marseille 2, 1980, 213 p.
- [37] Pasqualini V., Pergent-Martini C., Clabaut P., Pergent G., Mapping of *Posidonia oceanica* using aerial photographs and side scan sonar: application off the Island of Corsica (France), Estuar. Coast. Shelf Sci. 47 (1998) 359–367.
- [38] Pergent G., Floraison des herbiers à *Posidonia oceanica* dans la région d'Izmir (Turquie), Posidonia Newsletter 1 (1985) 15–22.
- [39] Pergent G., Pergent-Martini C., Boudouresque C.F., Utilisation de l'herbier à *Posidonia oceanica* comme indicateur biologique de la qualité du milieu littoral en Méditerranée: état des connaissances, Mésogée 54 (1995) 3–29.

- [40] Piazzì L., Balestri E., Cinelli F., Mappatura e monitoraggio di una prateria a *Posidonia oceanica* (L.) Delile situata a sud di Livorno (Toscana, Italia), *Inf. Bot. Ital.* 28 (1996a) 67–77.
- [41] Piazzì L., Pardi G., Cinelli F., Ecological aspects and reproductive phenology of *Acrothamnion preissii* (Sonder) Wollaston (Ceramiaceae, Rhodophyta) in the Tuscan Archipelago (Western Mediterranean), *Cryptogamie Algol.* 17 (1996b) 35–43.
- [42] Short F., Willie-Echevarria S., Natural and human-induced disturbance of seagrasses, *Environ. Conserv.* 23 (1996) 17–27.
- [43] Walker D.I., Kendrick G.A., McComb A.J., The distribution of seagrass species in Shark Bay, Western Australia, with notes on their ecology, *Aquat. Bot.* 30 (1988) 305–317.