

Toxicity screening of 13 *Gambierdiscus* strains using neuro-2a and erythrocyte lysis bioassays

Pisapia Francesco ^{1,*}, Holland William C. ², Hardison D. Ransom ², Litaker R. Wayne ², Fraga Santiago ³, Nishimura Tomohiro ⁴, Adachi Masao ⁴, Nguyen-Ngoc Lam ⁵, Séchet Veronique ¹, Amzil Zouher ¹, Herrenknecht Christine ⁶, Hess Philipp ¹

¹ Ifremer, Phycotoxins Laboratory, rue de l'Île d'Yeu, BP 21105, F-44311 Nantes, France

² National Oceanic and Atmospheric Administration, National Ocean Service, National Centers for Coastal Ocean Science, Center for Coastal Fisheries and Habitat Research (CCFHR), 101 Pivers Island Road, Beaufort, NC 28516, USA

³ Instituto Español de Oceanografía (IEO), Centro Oceanográfico de Vigo, Subida a Radio Faro 50, 36390 Vigo, Spain

⁴ LAQUES (Laboratory of Aquatic Environmental Science), Faculty of Agriculture, Kochi University, 200 Otsu, Monobe, Nankoku, Kochi, 783-8502, Japan

⁵ Institute of Oceanography, VAST, Cauda 01, Vinh Nguyen, Nha Trang, Viet Nam

⁶ LUNAM, University of Nantes, MMS EA2160, Pharmacy Faculty, 9 rue Bias, F-44035 Nantes, France

* Corresponding author : Francesco Pisapia, email address : francesco.pisapia@ifremer.fr

Abstract :

Species in the epi-benthic dinoflagellate genus *Gambierdiscus* produce ciguatoxins (CTXs) and maitotoxins (MTXs), which are among the most potent marine toxins known. Consumption of fish contaminated with sufficient quantities of CTXs causes Ciguatera Fish Poisoning (CFP), the largest cause of non-bacterial food poisoning worldwide. Maitotoxins, which can be found in the digestive system of fish, could also contribute to CFP if such tissues are consumed. Recently, an increasing number of *Gambierdiscus* species have been identified; yet, little is known about the variation in toxicity among *Gambierdiscus* strains or species.

This study is the first assessment of relative CTX- and MTX-toxicity of *Gambierdiscus* species from areas as widespread as the North-Eastern Atlantic Ocean, Pacific Ocean and the Mediterranean Sea. A total of 13 strains were screened: (i) seven Pacific strains of *G. australes*, *G. balechii*, *G. caribaeus*, *G. carpenteri*, *G. pacificus*, *G. scabrosus* and one strain of an undetermined species (*Gambierdiscus* sp. Viet Nam), (ii) five strains from the North-Eastern Atlantic Ocean (two *G. australes*, a single *G. excentricus* and two *G. silvae* strains), and (iii) one *G. carolinianus* strain from the Mediterranean Sea. Cell pellets of *Gambierdiscus* were extracted with methanol and the crude extracts partitioned into a CTX-containing dichloromethane fraction and a MTX-containing aqueous methanol fraction. CTX-toxicity was estimated using the neuro-2a cytotoxicity assay, and MTX-toxicity via a human erythrocyte lysis assay.

Different species were grouped into different ratios of CTX- and MTX-toxicity, however, the ratio was not

related to the geographical origin of species (Atlantic, Mediterranean, Pacific). All strains showed MTX-toxicity, ranging from 1.5 to 86 pg MTX equivalents (eq) cell⁻¹. All but one of the strains showed relatively low CTX-toxicity ranging from 0.6 to 50 fg CTX3C eq cell⁻¹. The exception was the highly toxic *G. excentricus* strain from the Canary Islands, which produced 1426 fg CTX3C eq cell⁻¹. As was true for CTX, the highest MTX-toxicity was also found in *G. excentricus*. Thus, the present study confirmed that at least one species from the Atlantic Ocean demonstrates similar toxicity as the most toxic strains from the Pacific, even if the metabolites in fish have so far been shown to be more toxic in the Pacific Ocean.

Highlights

► 13 strains of *Gambierdiscus* have been screened for CTX- and MTX-toxicity. ► *G. excentricus* is the most toxic species found so far in the Atlantic. ► Highest toxicity was observed in the slowest growing strain (*G. excentricus*). ► *G. excentricus* (Atlantic) showed similar CTX-toxicity as *G. polynesiensis*.

Keywords : Ciguatera fish poisoning, *Gambierdiscus*, Ciguatoxins, Maitotoxins, Neuro-2a assay, Erythrocyte lysis assay

1. Introduction

Dinoflagellates in the genera *Gambierdiscus* and *Fukuyoa* produce ciguatoxins (CTXs) and maitotoxins (MTXs), cyclic polyether neurotoxins that rank in the top five most potent natural toxins isolated to date (Fusetani and Kem, 2009).

Ciguatoxins, like brevetoxins, bind voltage-gated sodium channels (VGSCs) at site 5 on the alpha-subunit causing an influx of Na⁺ into affected cells that disrupts cellular function, especially in nerve cells (Benoit et al., 1986; Legrand et al., 1982; Lombet et al., 1987). Ciguatoxins are lipophilic and they could readily accumulate in the marine food chain reaching their highest concentration in fish, as hypothesized by Randall (1958), albeit with considerable lag-time between the bloom of *Gambierdiscus* sp. and CTX-related CFP outbreaks (Chateau-Degat et al., 2005). The genera *Gambierdiscus* and *Fukuyoa* are epibenthic and are found on many substrates including macro-algae, algal turfs, sea grasses and coral rubble (Parsons and Preskitt, 2007; Rains and Parsons, 2015) but they can also be found in near bottom plankton as shown using moored screens (Tester et al., 2014). Algal turfs appear to be very suitable substrates as support for *Gambierdiscus*, even when compared to macrophytes (Leaw et al., 2016). It is commonly assumed that the primary flux occurs from herbivorous grazers of such macro-algae to carnivorous fish (Ledreux et al., 2014), though other vectors such as crustaceans, echinoderms, and bivalves have been implicated (Kelly et al., 1992; Laurent et al., 2008; Roué et al., 2016; Silva et al., 2015). During this accumulation process CTXs are biotransformed, frequently resulting in metabolites of greater toxicities than the algal parent compounds (Lehane and Lewis, 2000). Certain *Gambierdiscus* species also produce other bioactive polyether compounds, such as gambierol (Cuypers et al., 2008; Satake et al., 1993a), gambieric acids (Nagai et al., 1993; Nagai et al., 1992) and gambierone (Rodríguez et al., 2015). The biological activity of gambierone is known to mimic that of CTX3C, although much lower in intensity, whereas the overall toxicity of gambierol and gambieric acids has yet to be characterized. The role, if any, of these three classes of compounds in causing CFP is unknown.

Maitotoxins are amphiphilic molecules that bind non-selective ion channels, causing an influx of Ca²⁺ that significantly raises intracellular Ca²⁺ levels. This is important since Ca²⁺ is one of the major signaling ions in the cell. The increased influx of the ion abnormally activates numerous biochemical pathways, including apoptosis, which disrupt the function of neuronal, muscular and red blood cells (Gusovsky and Daly, 1990; Ogura et al., 1984;

Ohizumi and Kobayashi, 1990). Even though MTXs are more toxic than CTXs when injected intraperitoneally into mice, MTXs are less likely to be involved in causing Ciguatera Fish Poisoning (CFP) because of their low capacity to accumulate in fish flesh and their low oral potency as assessed in mice (Yasumoto et al., 1976). Still, a recent study by Kohli et al. (2014) suggests that MTX could accumulate in carnivorous fish (fed in controlled conditions with *Gambierdiscus*-inoculated herbivorous fish), particularly in their digestive tract and liver, and thus MTXs may potentially contribute to CFP. Also, the large diversity of symptoms of CFP observed in different oceans has been suggested to be related to different CTX profiles (Lewis, 2001) but may also potentially be related to differences in consumer habits, e.g. the consumption of the intestinal parts of fish (Gatti et al., 2008; Hamilton et al., 2010). Consequently, the role of MTXs in contributing to CFP still remains to be clarified, in particular whether such contribution may derive from contamination of fish fillets during dissection of ciguateric fish or only from the consumption of visceral tissues of ciguateric fish.

In addition to uncertainties regarding different toxin profiles and the routes of accumulation little is known about the degree to which toxicity varies among species. One reason this has proven challenging is that the taxonomy has only recently been sufficiently resolved to examine species-specific toxicity (Fraga and Rodríguez, 2014; Fraga et al., 2011; Fraga et al., 2016; Kretzschmar et al., 2016; Litaker et al., 2009; Nishimura et al., 2014; Smith et al., 2016). This taxonomic work includes the separation of the globular *Gambierdiscus* species into the genus *Fukuyoa* (Gómez et al., 2015).

The goal of this study was to characterize the relative toxicity of *Gambierdiscus* strains from the Pacific Ocean, the North-Eastern Atlantic Ocean and the Mediterranean Sea. A total of 13 strains were examined, representing ten known species and one strain for which species annotation is not yet complete (Table 1, section 2.2). Except two strains (CCMP1653 and the strain from Viet Nam), none of the strains studied had previously been shown to produce any known CTXs or MTXs. Hence, this study examined the strains with a targeted cellular bioassay approach to detect activity of hitherto undescribed analogs of CTXs and MTXs.

2. Materials and Methods

2.1. Reference toxins and chemicals

CTX3C was kindly provided by Mireille Chinain (Institut Louis Malardé, Tahiti) and used as the reference standard for the neuroblastoma neuro-2a (N2a) cytotoxicity assay. MTX was purchased from Wako Chemicals USA, Inc. (Richmond, Virginia, USA) and used as the reference standard for the erythrocyte lysis assay (ELA). CTX3C was dissolved and stored in pure MeOH prior to utilization in the N2a assay. MTX was stored in MeOH:H₂O (1:1, v/v), dried and re-dissolved in ELA buffer prior to utilization in the ELA. HPLC grade methanol and dichloromethane were purchased from Sigma Aldrich (St. Louis, Missouri, USA).

Eagle's Minimum Essential Medium (EMEM, ATCC® 30-2003) for culture of N2a cells was purchased from the American Type Culture Collection. The following additives to the N2a medium were purchased from Sigma Aldrich (St. Louis, Missouri, USA): sodium pyruvate, streptomycin, penicillin and fetal bovine serum. N2a assay reagents were also purchased from Sigma Aldrich: trypsin-(ethylenediaminetetraacetic acid) (trypsin-EDTA) and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT). Saponin for the ELA was purchased from Sigma Aldrich (St. Louis, Missouri, USA).

2.2. Laboratory cultures of *Gambierdiscus* spp.

The strains of *Gambierdiscus* which were examined in this study and their location of origin are listed in Table 1. Strains of *G. scabrosus* (Nishimura et al., 2014), *G. excentricus* (Fraga et al., 2011), *G. silvae* (Fraga and Rodríguez, 2014) and *G. balechii* (Fraga et al., 2016) all belong to recently described species. Molecular analysis of *Gambierdiscus* sp. Viet Nam, previously reported as *G. toxicus* Vietnam by Roeder et al. (2010), still needs to be completed, and hence it was not assigned to this species.

Culture experiments were conducted using a semi-continuous batch method at both the Phycotoxins laboratory at the French Research Institute for Exploitation of the Sea (IFREMER), Nantes, France and at the National Oceanic and Atmospheric Administration, Center for Coastal Fisheries and Habitat Research (CCFHR), Beaufort, NC, USA. Cell densities were maintained at levels to ensure the absence of nutrient or CO₂ limitation. At pH > 8.4 cells become progressively more CO₂ limited. Cells of *Gambierdiscus* were harvested

in log phase growth. Slight differences in the experimental protocols necessitated by differences in the equipment available at each location are noted below (sections 2.2.1 and 2.2.2). As a control, *G. pacificus* CCMP1650 was grown in both laboratories to determine if growth rate and toxin values obtained in each laboratory were comparable.

2.2.1. NOAA CCFHR laboratory (Beaufort, NC, USA)

Four strains of *Gambierdiscus* (*G. caribaeus* Bill Hi Gam8, *G. carolinianus* Greece Gam2, *G. carpenteri* Pat Hi Jar7 Gam11 and *G. pacificus* CCMP1650) (Table 1) were grown in 75 cm² tissue culture flasks with vented caps (Falcon®, BD Biosciences, Bedford, MA, USA). Media consisted of 0.2 µm filtered Gulf Stream seawater (at a salinity of 33), vitamins and nutrients were added according to a modified K-medium protocol (Keller and Guillard, 1985; Keller et al., 1987). Phosphate was added in the form of Na₂ β-Glycerophosphoric Acid, 5-Hydrate at twice the concentration used for K-medium preparation. An EDTA-trace metal buffer system was used with the omission of copper as described by Hardison et al. (2012). The media was sterilized via microwave treatment as described in Keller et al. (1988). The culture pH was monitored (Thermo Orion 3 star pH meter, Ross ultra-combination pH electrode) to ensure pH range throughout experiments were between 8.1 and 8.4. This ensured the cells were not CO₂-limited. Cultures were maintained in a Percival Scientific incubator (Boone, IA, USA) at 27°C, under full spectrum lights (Blue Max F20-T12, Full Spectrum Solutions, Mississippi, USA) with an incident photon flux density at 100 µmol photons m⁻² s⁻¹ and a daily light-dark cycle of 12h:12h light:dark (LD). Full spectrum light source was placed in illumination cassettes above the culture flasks. Flasks were randomly placed and the position was changed once a day in order to ensure a homogeneous exposure to light. When the culture reached ~1000-2000 cells mL⁻¹, cells were first retained on a 20 µm sieve, washed with sterile seawater and collected by centrifugation (10 min, 1800 g, 20°C) in 50 mL Falcon tubes. Cell pellets were stored at -20°C until further extraction for toxicity screening.

2.2.2. IFREMER laboratory (Nantes, France)

Ten strains of *Gambierdiscus* were cultured in 75 cm² culture flasks (Corning® CellBIND®, Grosseron SAS, Coueron, France): *G. australes* CCMP1653, VGO1178 and VGO1181, *G. excentricus* VGO791, *G. pacificus* CCMP1650, *G. scabrosus* KW070922_1, *G. silvae* VGO1167 and VGO1180, *G. balechii* VGO917 and *Gambierdiscus* sp. Viet Nam (Table 1).

Media consisted of filtered (0.2 μm) natural Mediterranean seawater (at a salinity of 33) enriched with L1 nutrients with the exception of silica (Guillard and Hargraves, 1993). Cultures were maintained in a growth chamber incubator (Binder KBW240, Binder GmbH, Tuttlingen, Germany) at 25°C, under the same light conditions described above. After three weeks of semi-continuous culture, cells were harvested by centrifugation (20 min, 3000 g, 4°C) in 50 mL Falcon tubes and cell pellets were stored at -20°C until further extraction for toxicity screening.

2.3. Maximum growth rate determination

In order to determine the maximum growth rates during the exponential growth phase, *Gambierdiscus* cells were grown in semi-continuous batch cultures as previously described by Hardison et al. (2012). Briefly, all the cultures were acclimated to the culture conditions specific to each of the two laboratories (sections 2.2.1 and 2.2.2) for several months prior to experimentation. Cultures of *Gambierdiscus* cells were inoculated in 200 mL of culture medium at an initial concentration of ~100-200 cells mL⁻¹ and incubated at randomly determined sites in the incubator which were rotated daily. Cells were kept in the exponential growth phase as follows: cultures were transferred to new medium (1 to 10 dilution) when cell concentration reached ~1000-2000 cells mL⁻¹ and, thus, they never experienced nutrient or CO₂ limitation. An aliquot of culture was taken every 3-4 days during a period of 53-78 days (n = 13-15 samplings, at least three generations) and analyzed for cell concentration (cells mL⁻¹) and mean cellular biovolume (Estimated Spherical Volumes, ESV, μm^3 cell⁻¹) using a Multisizer™ 3 Coulter Counter® (Beckman Coulter, Georgia, USA) particle counter equipped with a 280 μm aperture tube and a 1 mL sample volume. The total volume of cells per liter of culture media (biovolume) was then calculated. Maximum growth rate (μ_{max} , d⁻¹) was the slope calculated by the linear regression of the natural logarithm of the biovolume versus time, after correcting for serial culture dilutions (Sunda and Hardison, 2007). SigmaPlot software (version 12.5) was used to calculate regression slopes and associated relative standard error and R² values. Maximum growth rate (μ_{max} , divisions day⁻¹) was then calculated as follows: μ_{max} (divisions day⁻¹) = μ_{max} (d⁻¹) ln(2)⁻¹.

2.4. Toxin extraction and liquid-liquid partitioning

Cultures of each strain have been grown in three separate flasks. After the cells had been harvested in log phase growth, they were suspended in MeOH (30 mL per 1 million cells)

and disrupted using sonication (CCFHR laboratory) or bead beating (IFREMER laboratory). Sonication was conducted twice for 1 min at 50% of total power (500 W) using a 3 mm diameter probe sonicator (Q-Sonica, Q700, Newtown, Connecticut USA). Grinding with the bead-mill was conducted twice for 30 min at a vibration frequency of 30 Hz using a mixer mill (Retsch MM400, Germany) with glass beads (0.25 g, diameter 250-500 μm) (Serive et al., 2012). Completeness of cell disruption was verified using light microscopy. Crude extracts (CEs) were blown dry under N_2 gas at 40°C. The residue was suspended in $\text{MeOH:H}_2\text{O}$ (3:2, v/v) (25 mL per 1 million cells) and partitioned twice with dichloromethane (DCM) (50 mL per 1 million cells) as previously described by Satake et al. (1993b). The lipophilic CTXs were partitioned into the DCM soluble fraction (DSF) while the amphiphilic MTXs were partitioned into the aqueous methanol (aq. MeOH) soluble fraction (MSF). Once the DSF and MSF fractions were isolated, they were blown dry under N_2 gas at 40°C and stored at -20°C. Just prior to the bioassays, the dried DCM and aq. MeOH residues were re-dissolved in MeOH or $\text{MeOH:H}_2\text{O}$ (1:1, v/v), respectively. An aliquot of the hydrophilic fraction was then evaporated (N_2 gas at 40°C) and stored at -20°C until use in the human erythrocyte lysis assay (ELA). Just prior to running the ELA the dried residue from the MSF fraction was dissolved in ELA buffer.

2.5. Neuroblastoma neuro-2a assay

The neuroblastoma neuro-2a (N2a) cell line is frequently used to estimate levels of CTXs in fish, shellfish or phytoplankton extracts (Pawlowicz et al., 2013). The N2a cytotoxicity assay developed by Manger et al. (1993) and modified by Dickey et al. (1999) was performed at CCFHR laboratory (Beaufort, NC, USA), with some modification (Hardison et al., 2016). Ciguatoxins do not induce N2a cell death, however, when N2a cells are pre-incubated with ouabain (O) and veratridine (V) they become highly sensitive to sodium channel activator toxins. Assays were set up so that the N2a cells are exposed to partially purified cell extracts with and without O and V. If cell death occurs in the samples without O and V it indicates the presence of a non-specific toxic compound other than a sodium channel activator. The details of the assay were as follows.

The N2a cell line was obtained from the American Type Culture Collection (ATCC, CCL 131). Neuro-2a cells were grown and maintained as described by Hardison et al. (2016). The assay was carried out in 96-well flat-bottom CELLCOAT® tissue culture plates with Poly-D-

Lysine coating (Greiner Bio-One, Kremsmünster, Austria). Plates were seeded with 30,000 N2a cells per well and were incubated for 24 h until they were >90% confluent at the bottom of each well. The CTX3C standard, controls and *Gambierdiscus* samples were added next and incubated for 24 h. The standard curve was added in presence of O/V (250 μ M and 25 μ M, respectively) at 50% cell viability to increase sensitivity and specificity to CTXs. The CTX3C standard curve for this assay ranged from 0.001 to 2,000 pg mL^{-1} . A sigmoidal dose-response curve was plotted and an EC_{50} of 1.66 ± 0.16 (SD, $n=12$) pg CTX3C mL^{-1} was calculated using GraphPad Prism 6.0 (Fig. S1) (Hardison et al., 2016). Controls included buffer wells to provide maximum survival estimates and wells with the addition of 1% MeOH (final concentration in well) to identify any cell mortality caused by the presence of MeOH used to dissolve the dried extracts. Half of the sample aliquots (1 μ L additions) from each assay were processed in the presence of O/V (O/V^+) so they were directly comparable to the CTX3C standard curve. The other half was incubated without O/V (O/V^-) to identify non-specific mortality caused by other compounds in the sample. Total well volume was 100 μ L. No more than 500 *Gambierdiscus* cell equivalents were added to each well to avoid matrix effects or non-specific N2a cell death. Each of the three replicate samples was run in duplicate in the N2a assay. Cell viability was assessed after 24 h incubation using the quantitative colorimetric 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay (Mosmann, 1983) using a FLUOstar® Omega microplate reader (BMG Labtech, Germany) at 544 nm. As the cells treated with O/V showed 50% of cell viability relative to control cells in solvent vehicle (1% MeOH in N2a medium), the viability of cells treated with CTX3C standard or algal extracts was estimated relative to O/V^+ wells. Quantitation of CTX3C eq in the samples using the N2a assay was operated within a range from 20% to 80% N2a cell viability with respect to the O/V^+ wells. The limit of quantification (80% N2a cell viability) was 0.197 ± 0.005 (SD, $n=12$) ng CTX3C mL^{-1} . When N2a cell viability was lower than 20%, a ten-fold dilution of the sample (in MeOH) was made. Extracts or strains showing activity with this assay will be referred to as “containing CTXs” or showing “CTX-toxicity”, even though this is a simplification as other compounds may also show sodium specific activity using this model.

2.6. Erythrocyte lysis assay

The erythrocyte lysis assay (ELA), developed by Eschbach et al. (2001) was performed at the CCFHR laboratory (Beaufort, NC, USA), with some modifications (Holland et al., 2013) to

estimate the hemolytic activity of *Gambierdiscus* strains. This assay is based on lysis of erythrocytes due to hemolytic compounds and subsequent photometrical determination of the released hemoglobin. In the context of marine dinoflagellates, the ELA has broadly been used on different red blood cell lines to detect hemolytic compounds from different microalgae such as *Alexandrium* (Tatters et al., 2012), *Karenia* (Tatters et al., 2010), *Ostreopsis cf. ovata* (Nascimento et al., 2012a) and *Gambierdiscus* (Holland et al., 2013).

Human red blood cells (hRBCs) were obtained from the Red Cross (Durham, North Carolina, USA). The hRBCs treated with saponin provided the maximal hemolysis (100% lysis) used to normalize the hemolytic activity of MTX standard or the diluted algal extracts. The hRBCs incubated solely in ELA buffer served as negative control (0% lysis). Details of how the assay was performed can be found in Holland et al. (2013). The 8-point MTX standard curve for this assay ranged from 0.0002 to 2,000 ng mL⁻¹ using purified MTX from Wako Chemicals USA, Inc. (Richmond Virginia, USA). Four replicate wells were used for each of the eight concentrations. A sigmoidal dose-response curve was plotted and an EC₅₀ of 14.2 ± 3.3 (SD, n=4) ng MTX mL⁻¹ was calculated using GraphPad Prism 6.0 (Fig. S2). The minimum hemolytic activity observed was 1.51 ± 0.39 (SD, n=4) % hemolysis when hRBCs were exposed to 1.0 ng MTX mL⁻¹. For each *Gambierdiscus* MSF sample, six dilutions were tested and three replicate wells for each dilution were run. Sigmoidal dose-response curves were plotted and EC₅₀ values (cell eq mL⁻¹) were calculated for each strain using GraphPad Prism 6.0. Quantitation of MTX eq in the samples using the ELA was operated converting EC₅₀ values (cell eq mL⁻¹) for each strain into toxin equivalent per cell (pg MTX eq cell⁻¹) taking into account the EC₅₀ value obtained from the MTX standard curve. Erythrocyte lysis was assessed after 24h incubation at 4°C using a FLUOstar® Omega microplate reader (BMG Labtech, Germany) at 415 nm. Extracts or strains showing activity with this assay will be referred to as “containing MTXs” or showing “MTX-toxicity”, even though this is a simplification as other compounds may also show hemolytic activity using this model.

2.7. Statistical analysis

Statistical analysis was performed using RStudio (Version 0.99.903) (<http://www.rstudio.com>) utilizing the R statistical language version 3.3.1 (<https://www.R-project.org>).

Multiple and linear regression models were obtained using basic functions in R on a data matrix including all qualitative information and quantitative values measured for each replicate on all strains. This corresponded to 42 observations of 7 variables [origin, laboratory, species, μ_{\max} (divisions day⁻¹), biovolume (ESV, μm^3 cell⁻¹), fg CTX3C eq cell⁻¹ (DSF) and pg MTX eq cell⁻¹ (MSF)]. For comparison purpose, the same analyses were performed on the same data matrix with the outlier strain (*G. excentricus* VGO791) excluded, which allowed for showing consistent correlations.

Unsupervised clustering of the strains based on the mean centered and normalized values of biovolume (ESV, μm^3 cell⁻¹), fg CTX3C eq cell⁻¹ (DSF) and pg MTX eq cell⁻¹ (MSF) was performed by 1) calculating the distance matrix between each observation (strain) using the Euclidean distance 2) executing a hierarchical cluster analysis using the Ward's minimum variance method. The result was displayed as a cluster dendrogram, one replicate of *Gambierdiscus* sp. Viet Nam being excluded for this analysis as it presented very dissimilar results to other two replicates.

3. Results

3.1. Maximum growth rates and cellular biovolumes

Maximum growth rates (μ_{\max}) of *Gambierdiscus* in culture ranged from 0.099 to 0.244 divisions day⁻¹, depending on the strain (Table 2). The slowest growing strains were *G. excentricus* VGO791 and *G. balechii* VGO917 (IFREMER laboratory conditions), while the fastest growth was observed for *G. pacificus* CCMP1650 (CCFHR laboratory conditions).

Interestingly, *G. excentricus* VGO791, which was the slowest growing strain, had the largest cellular biovolume (Table 2). *G. pacificus* (CCMP1650), the species with the smallest biovolume (3.8-fold < *G. excentricus*) was the fastest growing species (Table 2). Notwithstanding, overall the correlation was poor between growth rate and cellular volume, e.g. *G. balechii* had a similar growth rate as *G. excentricus* but a substantially smaller cellular biovolume. The overall low growth rates (< 0.5 divisions day⁻¹) observed in this study are consistent with those previously reported in the literature (Kibler et al., 2012; Litaker et al., submitted to *PLoS One* on Feb 2017; Xu et al., 2016; Yoshimatsu et al., 2014). If a comparison could be made, it can be concluded that, in the present study, *Gambierdiscus*

strains behaved as slow-growers when cultured under CCHFR and IFREMER laboratory conditions, with $\mu_{\max} < 0.25$ divisions day⁻¹, which appeared similarly low or somewhat lower than those reported in other studies.

3.2. Screening of DSF toxicity using the neuro-2a assay

Figure S1 shows the sigmoidal dose-response curve of CTX3C standard on the neuro-2a assay (Hardison et al., 2016).

For each strain, both dichloromethane soluble fraction (DSF, a fraction expected to contain CTXs) and the corresponding crude extract (CE) were tested on the neuro-2a assay. All DSFs tested were found to enhance the ouabain/veratridine (O/V) mediated cell mortalities consistent with CTX activation of voltage-gated sodium channels (VGSCs). There was no enhanced mortality without addition of O/V, indicating absence of non-specific toxicity after the initial purification step. In contrast, CEs showed substantial non-specific mortality (cell death in absence of O/V), indicating the presence of bioactive compounds other than VGSC activators such as MTXs or other toxic algal compounds. Thus, the quantitative estimation of CTXs was only possible in DSFs. Results were expressed in CTX3C equivalents (eq) per cell (Fig. 1).

Only the one *G. excentricus* (VGO791) strain examined from the Canary Islands exhibited a high level of CTX-type toxicity, i.e. 1,426 fg CTX3C eq cell⁻¹. The CTX content for all the other strains examined fell into the range of 0.6-40.8 fg CTX3C cell⁻¹ (Fig. 1).

Among the Pacific strains examined in this study, the Vietnamese strain, representing an as of yet undescribed species, showed the highest N2a cytotoxicity followed by the Japanese *G. scabrosus* strain. The two *G. silvae* strains showed low CTX-type toxicity equivalent to that of *G. pacificus*, i.e. around 10 fg CTX3C eq cell⁻¹. The Mediterranean strain of *G. carolinianus* showed background levels of CTX-type activity (< 4 fg CTX3C eq cell⁻¹) similar to the Pacific *G. caribaeus*, *G. carpenteri* and *G. australes* strains. Interestingly, *G. australes* strains originating either from the Pacific or the North-Eastern Atlantic Oceans, showed similar CTX-type activity (Fig. 1).

The strain of *G. pacificus* (CCMP1650) from French Polynesia showed similar levels of CTX-toxicity in cultures from both IFREMER and CCFHR, despite the differences in culture

conditions and extraction procedure (Table 2). Thus, the differences in culture and extraction techniques between the two laboratories appear to not have affected the results.

3.3. Screening of MSF toxicity using the erythrocyte lysis assay

Figure S2 shows the sigmoidal dose-response curve of MTX standard on the human erythrocyte lysis assay (ELA).

Figure 2 shows the MTX-type activity of the 13 strains of *Gambierdiscus* evaluated using the ELA. The strain of *G. excentricus* exhibited the highest hemolytic activity, followed by the Vietnamese strain. The Mediterranean strain of *G. carolinianus* was intermediate and about 5-fold more toxic than the two Atlantic strains of *G. silvae*. The Japanese strain of *G. scabrosus* showed the lowest hemolytic activity among all the strains tested in this study. Also, *G. australes*, *G. caribaeus*, and *G. carpenteri* species showed low MTX-type activity. Interestingly, among the latter ones, *G. australes* strains, originating either from Pacific or the North-Eastern Atlantic Oceans, showed the same MTX-type activity. Overall, though variable, all strains tested showed measurable MTX-toxicity.

As in the evaluation of CTX3C equivalent toxicity, *G. pacificus* (CCMP1650) showed similar levels of MTX-toxicity in cultures from both IFREMER and CCFHR, despite the differences in culture conditions and extraction procedure (Table 2).

3.4 Relationship between CTX and MTX toxicity

The CTX toxin content per unit biovolume varied over four orders of magnitude compared to a 10-fold variation in MTX toxicity per unit biovolume (Fig. 3). A similar pattern was observed when the data were normalized on a per cell basis. The relationship between CTX and MTX for the various species fell into three groups (Fig. 4) with strains of from a given species classifying into the same group.

4. Discussion

CTX-toxicity. The present study showed that *Gambierdiscus* species examined from the Pacific Ocean, North-Eastern Atlantic Ocean and Mediterranean Sea exhibited marked

differences in toxicity ranging from 0.6 fg to > 1400 fg CTX3C equivalents (eq) cell⁻¹ (Fig. 1). The greatest toxicity was exhibited by *G. excentricus* (VGO791) from the Canary Islands. This result is consistent with a previous study reporting *G. excentricus* strains exhibiting between 0.37 - 1.10 pg CTX1B eq cell⁻¹ (Fraga et al., 2011), which would be equivalent to 1.17 - 3.49 pg CTX3C eq cell⁻¹ according to Bottein Dechraoui et al. (2007). The Canary Islands, where these strains were obtained, are a temperate region (North-Eastern Atlantic Ocean) from which Ciguatera Fish Poisoning (CFP) has recently been reported (Boada et al., 2010; Pérez-Arellano et al., 2005). More recently, *G. excentricus* has been found in Brazil, its contribution to CFP in the region has yet to be evaluated (Nascimento et al., 2012b; Nascimento et al., 2015). The observation of CFP in the Canary Islands is important because CFP is typically considered a tropical disease (Lewis, 2001). Thus, *G. polynesiensis* and *G. excentricus* could be considered as primary toxin producing species in the South Pacific and the Eastern Atlantic Oceans, respectively (Chinain et al., 2010; Rhodes et al., 2014; Rhodes et al., 2016).

In contrast, *G. australes*, *G. balechii*, *G. carolinianus*, *G. carpenteri*, *G. pacificus* and *G. silvae* all had toxicities below 50 fg CTX3C eq cell⁻¹ (Fig. 1). In a separate study on Caribbean strains, *G. belizeanus*, *G. caribaeus*, *G. carolinianus*, *G. carpenteri*, *Gambierdiscus* ribotype II, *G. silvae* and *F. ruetzleri* have similarly low toxicities (< 20 fg CTX3C eq cell⁻¹) (Litaker et al., *submitted to PLoS One on Feb 2017*). Further, *G. carolinianus* and *G. carpenteri* strains from the Caribbean, Mediterranean and Pacific, as well as *G. silvae* and *G. australes* strains originating either from the Caribbean, Pacific or the North-Eastern Atlantic Oceans showed similar CTX-type activity, suggesting that each species produce comparable levels of toxin worldwide. The data presented here also indicate that most *Gambierdiscus* species produce relatively low levels of CTXs. This low level of toxin production raises an important question concerning the degree to which these low toxicity species contribute to the overall toxin flux into the food chain relative to the high toxicity species *G. excentricus* and *G. polynesiensis* (Litaker et al., *submitted to PLoS One on Feb 2017*). As these low toxicity species can only be of public health importance if they are able to bloom, it will be important to map abundances of the different *Gambierdiscus* species in the field.

The CTX-toxicity results from this study were consistent with other studies on individual species. The strain of *Gambierdiscus* sp. Viet Nam (previously reported as *G. toxicus* based

on morphology), for example, showed the highest toxicity among all the Pacific strains examined (Fig. 1) and has been shown to produce several CTX analogs (Roeder et al., 2010). Nishimura et al. (2013) similarly detected the CTX-like toxicity of the DSF fraction from the Japanese strain of *G. scabrosus* (species previously reported as *Gambierdiscus* sp. type 1), KW070922_1, using the mouse bioassay (MBA), i.e. 20×10^{-4} MU/1,000 cells. These toxicities are relatively high compared to most other species, but still low compared to *G. excentricus* or *G. polynesiensis*, consistent with the findings in this study.

The three *G. australes* strains in this study were in the same low range of toxicity, independently of their origin (Canary Islands and Hawaii). Of these strains, comparable toxicity data are only available for the Pacific CCMP1653 strain, previously reported as T39. For this particular strain, Babinchak et al. (1986) reported high toxicity of the crude extract using the MBA. Later on, a more specific assay described by Van Dolah et al. (1994) (radio-labelled brevetoxin ($[^3\text{H}]\text{BTX-3}$) displacement assay) was conducted on CCMP1653, showing from no detectable to low Na^+ ion channel activity indicating low toxicity (Sperr and Doucette, 1996). More recently, LC-MS studies conducted by Roeder et al. (2010) on CCMP1653 showed the presence of one CTX analog. Chinain et al. (2010) reported comparatively low toxicity for six *G. australes* strains originating from French Polynesia, ranging from < 0.016 (LOD) to $0.030 \text{ pg CTX3C eq cell}^{-1}$. Rhodes et al. (2010) found intermediate N2a cytotoxicity at sub-pg range ($0.13 \text{ pg CTX3C eq cell}^{-1}$) for the *G. australes* CAWD149 strain (Cook Islands). Nishimura et al. (2013) reported DSF-toxicity (MBA) for a Japanese *G. australes* strain (S080911_1) of 670×10^{-4} MU/1,000 cells, comparable to the highly toxic *G. polynesiensis* species, i.e. $800\text{-}1500 \times 10^{-4}$ MU/1,000 cells (Chinain et al., 1999). Such difference in toxicity between strains suggests that a larger number of strains are needed to assess intraspecific variations in CTX toxicity.

Chinain et al. (2010) observed that the slow growing species examined in their study, *G. polynesiensis* exhibited the highest level of toxicity and hypothesized that slower growing species were more toxic. In this study, *G. excentricus* was the slowest growing species and it also showed by far the highest CTX and MTX toxicity ($1.4 \text{ pg CTX3C eq cell}^{-1}$ and $86 \text{ pg MTX eq cell}^{-1}$) (Figs. 1 and 2, Table 2). A study of *Gambierdiscus* species found in the Caribbean showed an inverse exponential relationship between CTX toxicity on a per-cell and per-biovolume basis consistent with the Chinain et al. (2010) hypothesis (Litaker et al., submitted to *PLoS One* on Feb 2017). The species *G. balechii* however appears an exception

to this rule as it had a comparable growth rate to *G. excentricus* but was substantially less toxic (Figs. 1-3).

MTX-toxicity. The MTX-toxicity of *Gambierdiscus* strains varied more than 50-fold (1.5 - 86 pg MTX eq cell⁻¹). The most maitotoxic species was *G. excentricus* (VGO791; ~80 pg MTX eq cell⁻¹) followed by the Vietnamese strain (*Gambierdiscus* sp. Viet Nam; ~70 pg MTX eq cell⁻¹). The variability was highest for the Vietnamese strain (64% RSD) with one replicate giving a much higher result than the other two, followed by variability for *G. excentricus* (48% RSD). Additional trials with more replication are needed before concluding the toxicity of these species is comparable. The measured MTX-toxicity for *G. excentricus* was lower than the ~600 pg MTX eq cell⁻¹ estimated for the strains VGO790, VGO791 and VGO792 (Fraga et al. 2011) using a modified neuro-2a assay (Caillaud et al., 2010). The MTX content however was estimated using a crude extract whereas MTX-toxicity in the present study was estimated in the aqueous methanol extracts, i.e. after liquid-liquid partitioning. Possible explanations of this discrepancy include differences in the assay applied (N2a vs. ELA) and sources of standard (MTX from different sources). Similarly, the G10DC strain of *G. pacificus*, isolated from Malaysia, was estimated to have a toxicity of 50.2 pg MTX eq cell⁻¹ (Caillaud et al., 2011), 2.5 times more toxic than results from the present study for CCMP1650 strain. The three *G. australes* strains assayed in this study showed low MTX-toxicity (< 5 pg MTX eq cell⁻¹) consistent with the low toxicity of *G. australes* RAV-92 strain of from Raivavae Island (Australes Archipelago) measured by MBA (Chinain et al., 1999). *G. scabrosus* strain (KW070922_1) showed a similarly low hemolytic activity (1.5 pg MTX eq cell⁻¹) consistent with the observed MSF fraction toxicity of 67 x 10⁻⁴ MU/1,000 cells (Nishimura et al. 2013).

Ratio CTX/MTX-toxicity. It should be noted that all the strains examined in this study produced measureable quantities of both CTXs and MTXs (Figs. 1 and 2). The variation among species, however, was not consistent for the two groups of toxins. CTX-toxicity per unit biovolume varied over three orders of magnitude among species whereas the MTX-toxicity varied over one order of magnitude (Fig. 3).

Based on a hierarchical cluster analysis (section 2.7), strains could be classified into three different groups: group I (*G. excentricus* VGO791), group II (*G. australes* CCMP1653, VGO1178, VGO1181, *G. caribaeus* Bill Hi Gam8, *G. carolinianus* Greece Gam2 and *G. carpenteri* Pat Hi Jar7 Gam11), and group III (*G. balechii* VGO917, *G. pacificus*

CCMP1650, *G. silvae* VGO1167, VGO1180, *G. scabrosus* KW070922_1 and *Gambierdiscus* sp. Viet Nam) (Fig. 4). As visually suggested in Figure 3, there is a correlation between CTX and MTX contents (on a per cell basis, in DSF and MSF fractions, respectively): the Pearson correlation coefficient for linear regression was $Rr^2=0.45$ ($P < 0.001$) when including *G. excentricus*, and $Rr^2=0.23$ ($P < 0.01$) when excluding *G. excentricus*. Though the data were limited, different strains of the same species fell into the same grouping indicating that the relationship between CTX and MTX toxicity appears constant for a given species. More data are needed to fully test this hypothesis. It should also be noted that toxicity in this study was assessed using functional assays. Those assays cannot distinguish between production of large amounts of low toxicity CTX or MTX congeners relative to smaller production of high toxicity congeners. *Gambierdiscus* cell extracts that were fractionated and assayed for toxicity using a calcium flux assay indicated that despite strains of a species possessing similar CTX and MTX toxicity, the actual congeners being produced in a given species probably vary (Lewis et al., 2016). How these relative profiles might affect grazing pressure or deter bacterial or fungal infections remains unknown.

Also unidentified is whether the different patterns of CTX and MTX can offer insights into biosynthetic pathways for these studies. Ongoing studies are focusing on growing large-scale cultures of the most toxic strains of *Gambierdiscus* for the purification of the toxic compounds through fractionation and screening using the cellular bioassays.

In summary, it should be noted that only one of the thirteen strains examined has been shown to contain pg amounts of CTX-type toxicity per cell, and this strain of *G. excentricus* needs to be examined in detail for its toxin contents to identify the algal precursor(s) of toxins involved in CFP in the Atlantic Ocean. As this strain of *G. excentricus* exhibits CTX-type toxicity in the same order of magnitude as *G. polynesiensis* from the South Pacific, *Gambierdiscus* species from both oceans should be considered to be a similar potential threat to fish consumers. All strains displayed MTX-toxicity in the pg range ($1.5 - 86 \text{ pg cell}^{-1}$). Also, the variability of CTX and MTX-type toxicities between species and strains appeared to be similar to those previously reported in literature and was equivalent between Atlantic and Pacific strains of *Gambierdiscus*. Several of the findings, including correlation between growth rate and toxicity or variability within and between species, are only indicative of possible trends and more strains should be examined to corroborate the findings.

Acknowledgements

The authors acknowledge Ifremer and the Regional Council of the *Région des Pays de la Loire* for PhD funding of F. Pisapia, and for the contribution through the COSELMAR project (funding and time of P. Hess). The authors also acknowledge Catherine Roullier (MMS laboratory, University of Nantes, France) for conducting statistical analysis. L. Nguyen-Ngoc thanks the NAFOSTED for funding project 106-NN.06-2014.08. S. Fraga thanks MINECO for funding project CICAN. The authors would like to thank Dr. Urban Tillmann of the Alfred Wegener Institute, Bremerhaven, for providing an aliquot of the culture of the *Gambierdiscus* strain from Viet Nam. The work done at the NOAA, National Ocean Service, National Centers for Coastal Ocean Science, Center for Coastal Fisheries was supported by the program funds and NOAA, National Centers for Coastal Ocean Science, Center for Sponsored Coastal Ocean Research, Ecology and Oceanography of Harmful Algal Blooms award, ECOHAB ECO882.

Conflict of interest statement

The authors declare that there is no conflict of interest.

References

- Babinchak, J.A., Jollow, D.J., Voegtline, M.S., Higerd, T.B., 1986. Toxin production by *Gambierdiscus toxicus* isolated from the Florida Keys. *Marine Fisheries Review* 48(4), 53-56.
- Benoit, E., Legrand, A.M., Dubois, J.M., 1986. Effects of ciguatoxin on current and voltage clamped frog myelinated nerve-fiber. *Toxicon* 24(4), 357-364.
- Boada, L.D., Zumbado, M., Luzardo, O.P., Almeida-González, M., Plakas, S.M., Granade, H.R., Abraham, A., Jester, E.L., Dickey, R.W., 2010. Ciguatera Fish Poisoning on the West Africa Coast: an emerging risk in the Canary Islands (Spain). *Toxicon* 56(8), 1516-1519.
- Bottein Dechraoui, M.Y., Wang, Z., Ramsdell, J.S., 2007. Optimization of ciguatoxin extraction method from blood for Pacific ciguatoxin (P-CTX-1). *Toxicon* 49(1), 100-105.
- Bravo, I., Figueroa, R.I., Fraga, S., 2014. Cellular and nuclear morphological variability within a single species of the toxigenic dinoflagellate genus *Gambierdiscus*: relationship to life-cycle processes. *Harmful Algae* 40, 1-8.
- Caillaud, A., de la Iglesia, P., Barber, E., Eixarch, H., Mohammad-Noor, N., Yasumoto, T., Diogène, J., 2011. Monitoring of dissolved ciguatoxin and maitotoxin using solid-phase adsorption toxin tracking devices: application to *Gambierdiscus pacificus* in culture. *Harmful Algae* 10(5), 433-446.
- Chateau-Degat, M.L., Chinain, M., Cerf, N., Gingras, S., Hubert, B., Dewailly, E., 2005. Seawater temperature, *Gambierdiscus* spp. variability and incidence of ciguatera poisoning in French Polynesia. *Harmful Algae* 4(6), 1053-1062.
- Chinain, M., Darius, H.T., Ung, A., Cruchet, P., Wang, Z., Ponton, D., Laurent, D., Pauillac, S., 2010. Growth and toxin production in the ciguatera-causing dinoflagellate *Gambierdiscus polynesiensis* (Dinophyceae) in culture. *Toxicon* 56(5), 739-750.
- Chinain, M., Faust, M.A., Pauillac, S., 1999. Morphology and molecular analyses of three toxic species of *Gambierdiscus* (Dinophyceae): *G. pacificus*, sp. nov., *G. australes*, sp. nov., and *G. polynesiensis*, sp. nov. *Journal of Phycology* 35(6), 1282-1296.
- Cuypers, E., Abdel-Mottaleb, Y., Kopljar, I., Rainier, J.D., Raes, A.L., Snyders, D.J., Tytgat, J., 2008. Gambierol, a toxin produced by the dinoflagellate *Gambierdiscus toxicus*, is a potent blocker of voltage-gated potassium channels. *Toxicon* 51(6), 974-983.
- Dickey, R., Jester, E., Granade, R., Mowdy, D., Moncreiff, C., Rebarchik, D., Robl, M., Musser, S., Poli, M., 1999. Monitoring brevetoxins during a *Gymnodinium breve* red tide: comparison of sodium channel specific cytotoxicity assay and mouse bioassay for determination of neurotoxic shellfish toxins in shellfish extracts. *Natural Toxins* 7(4), 157-165.
- Eschbach, E., Scharsack, J.P., John, U., Medlin, L.K., 2001. Improved erythrocyte lysis assay in microtitre plates for sensitive detection and efficient measurement of haemolytic compounds from ichthyotoxic algae. *J. Appl. Toxicol.* 21(6), 513-519.
- Fraga, S., Rodríguez, F., 2014. Genus *Gambierdiscus* in the Canary Islands (NE Atlantic Ocean) with description of *Gambierdiscus silvae* sp. nov., a new potentially toxic epiphytic benthic dinoflagellate. *Protist* 165(6), 839-853.
- Fraga, S., Rodríguez, F., Caillaud, A., Diogène, J., Raho, N., Zapata, M., 2011. *Gambierdiscus excentricus* sp. nov. (Dinophyceae), a benthic toxic dinoflagellate from the Canary Islands (NE Atlantic Ocean). *Harmful Algae* 11, 10-22.
- Fraga, S., Rodríguez, F., Riobó, P., Bravo, I., 2016. *Gambierdiscus balechii* sp. nov. (Dinophyceae), a new benthic toxic dinoflagellate from the Celebes Sea (SW Pacific Ocean). *Harmful Algae* 58, 93-105.

- Fusetani, N., Kem, W., 2009. Marine toxins: an overview. *Progress in molecular and subcellular biology* 46, 1-44.
- Gatti, C., Oelher, E., Legrand, A.M., 2008. Severe seafood poisoning in French Polynesia: a retrospective analysis of 129 medical files. *Toxicon* 51(5), 746-753.
- Gómez, F., Qiu, D.J., Lopes, R.M., Lin, S.J., 2015. *Fukuyoa paulensis* gen. et sp. nov., a new genus for the globular species of the dinoflagellate *Gambierdiscus* (Dinophyceae). *PLoS One* 10(4), e0119676.
- Guillard, R.R.L., Hargraves, P.E., 1993. *Stichochrysis immobilis* is a diatom, not a chrysophyte. *Phycologia* 32(3), 234-236.
- Gusovsky, F., Daly, J.W., 1990. Maitotoxin: a unique pharmacological tool for research on calcium-dependent mechanisms. *Biochemical Pharmacology* 39(11), 1633-1639.
- Hamilton, B., Whittle, N., Shaw, G., Eaglesham, G., Moore, M.R., Lewis, R.J., 2010. Human fatality associated with Pacific ciguatera contaminated fish. *Toxicon* 56(5), 668-673.
- Hardison, D.R., Holland, W.C., McCall, J.R., Bourdelais, A.J., Baden, D.G., Darius, H.T., Chinain, M., Tester, P.A., Shea, D., Quintana, H.A., Morris, J.A., Jr., Litaker, R.W., 2016. Fluorescent receptor binding assay for detecting ciguaterins in fish. *PLoS One* 11(4), e0153348.
- Hardison, D.R., Sunda, W.G., Litaker, R.W., Shea, D., Tester, P.A., 2012. Nitrogen limitation increases brevetoxins in *Karenia brevis* (Dinophyceae): implications for bloom toxicity. *J. Phycol.* 48(4), 844-858.
- Holland, W.C., Litaker, R.W., Tomas, C.R., Kibler, S.R., Place, A.R., Davenport, E.D., Tester, P.A., 2013. Differences in the toxicity of six *Gambierdiscus* (Dinophyceae) species measured using an *in vitro* human erythrocyte lysis assay. *Toxicon* 65(0), 15-33.
- <http://www.rstudio.com>, RStudio Team (2015). RStudio: integrated Development for R, RStudio, Inc., Boston, MA.
- <https://www.R-project.org>, R Core Team (2016). R: a language and environment for statistical computing, R Foundation for Statistical Computing, Vienna, Austria.
- Keller, M., Guillard, R., 1985. Factors significant to marine dinoflagellate culture. *Toxic Dinoflagellates*. Elsevier, New York, 113-116.
- Keller, M.D., Bellows, W.K., Guillard, R.R., 1988. Microwave treatment for sterilization of phytoplankton culture media. *Journal of Experimental Marine Biology and Ecology* 117(3), 279-283.
- Keller, M.D., Selvin, R.C., Claus, W., Guillard, R.R., 1987. Media for the culture of oceanic ultraphytoplankton1, 2. *J Phycol* 23(4), 633-638.
- Kelly, A.M., Kohler, C.C., Tindall, D.R., 1992. Are crustaceans linked to the ciguatera food chain? *Environmental Biology of Fishes* 33(3), 275-286.
- Kibler, S.R., Litaker, R.W., Holland, W.C., Vandersea, M.W., Tester, P.A., 2012. Growth of eight *Gambierdiscus* (Dinophyceae) species: effects of temperature, salinity and irradiance. *Harmful Algae* 19, 1-14.
- Kohli, G.S., Papiol, G.G., Rhodes, L.L., Harwood, D.T., Selwood, A., Jerrett, A., Murray, S.A., Neilan, B.A., 2014. A feeding study to probe the uptake of maitotoxin by snapper (*Pagrus auratus*). *Harmful Algae* 37, 125-132.
- Kretzschmar, A.L., Verma, A., Harwood, D.T., Hoppenrath, M., Murray, S.A., 2016. Characterisation of *Gambierdiscus lapillus* sp. nov. (Gonyaulacales, Dinophyceae): a new toxic dinoflagellate from the Great Barrier Reef (Australia). *Journal of Phycology*.

Laurent, D., Kerbrat, A.-S., Darius, H.T., Girard, E., Golubic, S., Benoit, E., Sauviat, M.-P., Chinain, M., Molgo, J., Pauillac, S., 2008. Are cyanobacteria involved in Ciguatera Fish Poisoning-like outbreaks in New Caledonia? *Harmful Algae* 7(6), 827-838.

Leaw, C.P., Yong, H.L., Mustapa, N.I., Tan, T.H., Lim, Z.F., Lee, L.K., Lim, P.T., 2016. Habitat complexity affects the benthic harmful dinoflagellate assemblages in the fringing reefs of Malaysia. Session OS08, The 17th International Conference on Harmful Algae (ICHA 2016), Florianópolis, SC, Brazil.

Ledreux, A., Brand, H., Chinain, M., Bottein, M.Y.D., Ramsdell, J.S., 2014. Dynamics of ciguatoxins from *Gambierdiscus polynesiensis* in the benthic herbivore *Mugil cephalus*: trophic transfer implications. *Harmful Algae* 39(0), 165-174.

Legrand, A.M., Galonnier, M., Bagnis, R., 1982. Studies on the mode of action of ciguateric toxins. *Toxicon* 20(1), 311-315.

Lehane, L., Lewis, R.J., 2000. Ciguatera: recent advances but the risk remains. *International Journal of Food Microbiology* 61(2-3), 91-125.

Lewis, R.J., 2001. The changing face of ciguatera. *Toxicon* 39(1), 97-106.

Lewis, R.J., Inserra, M., Vetter, I., Holland, W.C., Hardison, D.R., Tester, P.A., Litaker, R.W., 2016. Rapid extraction and identification of maitotoxin and ciguatoxin-like toxins from Caribbean and Pacific *Gambierdiscus* using a new functional bioassay. *PLoS One* 11(7), e0160006.

Litaker, R.W., Holland, W.C., Hardison, D.R., Pisapia, F., Hess, P., Tester, P., *submitted to PLoS One on Feb 2017*. Relative toxicity of *Gambierdiscus* and *Fukuyoa* species from the Caribbean and Gulf of Mexico. *PLoS One*.

Litaker, R.W., Vandersea, M.W., Faust, M.A., Kibler, S.R., Chinain, M., Holmes, M.J., Holland, W.C., Tester, P.A., 2009. Taxonomy of *Gambierdiscus* including four new species, *Gambierdiscus caribaeus*, *Gambierdiscus carolinianus*, *Gambierdiscus carpenteri* and *Gambierdiscus ruetzleri* (Gonyaulacales, Dinophyceae). *Phycologia* 48(5), 344-390.

Lombet, A., Bidard, J.N., Lazdunski, M., 1987. Ciguatoxin and brevetoxins share a common receptor site on the neuronal voltage-dependent Na⁺ channel. *FEBS Letters* 219(2), 355-359.

Manger, R.L., Leja, L.S., Lee, S.Y., Hungerford, J.M., Wekell, M.M., 1993. Tetrazolium-based cell bioassay for neurotoxins active on voltage-sensitive sodium channels: semiautomated assay for saxitoxins, brevetoxins, and ciguatoxins. *Analytical Biochemistry* 214(1), 190-194.

Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* 65(1-2), 55-63.

Nagai, H., Mikami, Y., Yazawa, K., Gono, T., Yasumoto, T., 1993. Biological activities of novel polyether antifungals, gambieric acids A and B from a marine dinoflagellate *Gambierdiscus toxicus*. *The Journal of Antibiotics (Tokyo)* 46(3), 520-522.

Nagai, H., Murata, M., Torigoe, K., Satake, M., Yasumoto, T., 1992. Gambieric acids, new potent antifungal substances with unprecedented polyether structures from a marine dinoflagellate *Gambierdiscus toxicus*. *Journal of Organic Chemistry* 57(20), 5448-5453.

Nascimento, S.M., Corrêa, E.V., Menezes, M., Varela, D., Paredes, J., Morris, S., 2012a. Growth and toxin profile of *Ostreopsis* cf. *ovata* (Dinophyta) from Rio de Janeiro, Brazil. *Harmful Algae* 13, 1-9.

Nascimento, S.M., Diniz, B., de Alencar, A.G., Meneguelli, A.C., Menezes, M., 2012b. First record of the ciguatera causing genus *Gambierdiscus* in Brazil, *Harmful Algal Blooms*, pp. 8-9.

Nascimento, S.M., Melo, G., Salgueiro, F., Diniz, B.D., Fraga, S., 2015. Morphology of *Gambierdiscus excentricus* (Dinophyceae) with emphasis on sulcal plates. *Phycologia* 54(6), 628-639.

- Nishimura, T., Sato, S., Tawong, W., Sakanari, H., Uehara, K., Shah, M.M., Suda, S., Yasumoto, T., Taira, Y., Yamaguchi, H., Adachi, M., 2013. Genetic diversity and distribution of the ciguatera-causing dinoflagellate *Gambierdiscus* spp. (Dinophyceae) in coastal areas of Japan. *PLoS One* 8(4), e60882.
- Nishimura, T., Sato, S., Tawong, W., Sakanari, H., Yamaguchi, H., Adachi, M., 2014. Morphology of *Gambierdiscus scabrosus* sp. nov. (Gonyaulacales): a new epiphytic toxic dinoflagellate from coastal areas of Japan. *J. Phycol.* 50(3), 506-514.
- Ogura, A., Ohizumi, Y., Yasumoto, T., 1984. Calcium-dependent depolarization induced by a marine toxin, maitotoxin, in a neuronal cell. *Japanese Journal of Pharmacology* 36, P315-P315.
- Ohizumi, Y., Kobayashi, M., 1990. Ca²⁺-dependent excitatory effects of maitotoxin on smooth and cardiac muscle. *ACS Symposium Series* 418, 133-143.
- Parsons, M.L., Preskitt, L.B., 2007. A survey of epiphytic dinoflagellates from the coastal waters of the island of Hawai'i. *Harmful Algae* 6(5), 658-669.
- Pawlowicz, R., Darius, H.T., Cruchet, P., Rossi, F., Caillaud, A., Laurent, D., Chinain, M., 2013. Evaluation of seafood toxicity in the Australes archipelago (French Polynesia) using the neuroblastoma cell-based assay. *Food Additives & Contaminants: Part A* 30(3), 567-586.
- Pérez-Arellano, J.L., Luzardo, O.P., Perez Brito, A., Hernandez Cabrera, M., Zumbado, M., Carranza, C., Angel-Moreno, A., Dickey, R.W., Boada, L.D., 2005. Ciguatera fish poisoning, Canary Islands. *Emerging Infectious Diseases* 11(12), 1981-1982.
- Rains, L.K., Parsons, M.L., 2015. *Gambierdiscus* species exhibit different epiphytic behaviors toward a variety of macroalgal hosts. *Harmful Algae* 49, 29-39.
- Randall, J.E., 1958. A review of ciguatera, tropical fish poisoning with a tentative explanation of its cause. *Bull. Mar. Sci. Gulf Carib.* 8, 236-267.
- Rhodes, L., Harwood, T., Smith, K., Argyle, P., Munday, R., 2014. Production of ciguatoxin and maitotoxin by strains of *Gambierdiscus australes*, *G. pacificus* and *G. polynesiensis* (Dinophyceae) isolated from Rarotonga, Cook Islands. *Harmful Algae* 39(0), 185-190.
- Rhodes, L., Harwood, T., Smith, K., Argyle, P., Munday, R., 2016. Production of ciguatoxin and maitotoxin by strains of *Gambierdiscus australes*, *G. pacificus* and *G. polynesiensis* (Dinophyceae) isolated from Rarotonga, Cook Islands (vol 39, pg 185, 2014). *Harmful Algae* 55, 295-295.
- Rhodes, L.L., Smith, K.F., Munday, R., Selwood, A.I., McNabb, P.S., Holland, P.T., Bottein, M.Y., 2010. Toxic dinoflagellates (Dinophyceae) from Rarotonga, Cook Islands. *Toxicon* 56(5), 751-758.
- Rodríguez, I., Genta-Jouve, G., Alfonso, C., Calabro, K., Alonso, E., Sánchez, J.A., Alfonso, A., Thomas, O.P., Botana, L.M., 2015. Gambierone, a ladder-shaped polyether from the dinoflagellate *Gambierdiscus belizeanus*. *Org. Lett.* 17(10), 2392-2395.
- Roeder, K., Erler, K., Kibler, S., Tester, P., Van The, H., Nguyen-Ngoc, L., Gerds, G., Luckas, B., 2010. Characteristic profiles of ciguatera toxins in different strains of *Gambierdiscus* spp. *Toxicon* 56(5), 731-738.
- Roué, M., Darius, H.T., Picot, S., Ung, A., Viallon, J., Gaertner-Mazouni, N., Sibat, M., Amzil, Z., Chinain, M., 2016. Evidence of the bioaccumulation of ciguatoxins in giant clams (*Tridacna maxima*) exposed to *Gambierdiscus* spp. cells. *Harmful Algae* 57, 78-87.
- Satake, M., Murata, M., Yasumoto, T., 1993a. Gambierol: a new toxic polyether compound isolated from the marine dinoflagellate *Gambierdiscus toxicus*. *Journal of the American Chemical Society* 115(1), 361-362.
- Satake, M., Murata, M., Yasumoto, T., 1993b. The structure of CTX3C, a ciguatoxin congener isolated from cultured *Gambierdiscus toxicus*. *Tetrahedron Letters* 34(12), 1975-1978.

- Serive, B., Kaas, R., Berard, J.B., Pasquet, V., Picot, L., Cadoret, J.P., 2012. Selection and optimisation of a method for efficient metabolites extraction from microalgae. *Bioresour. Technol.* 124, 311-320.
- Silva, M., Rodríguez, I., Barreiro, A., Kaufmann, M., Isabel Neto, A., Hassouani, M., Sabour, B., Alfonso, A., Botana, L.M., Vasconcelos, V., 2015. First report of ciguatera toxins in two starfish species: *Ophidiaster ophidianus* and *Marthasterias glacialis*. *Toxins* 7(9), 3740-3757.
- Smith, K.F., Rhodes, L., Verma, A., Curley, B.G., Harwood, D.T., Kohli, G.S., Solomona, D., Rongo, T., Munday, R., Murray, S.A., 2016. A new *Gambierdiscus* species (Dinophyceae) from Rarotonga, Cook Islands: *Gambierdiscus cheloniae* sp. nov. *Harmful Algae* 60, 45-56.
- Sperr, A.E., Doucette, G.J., 1996. Variation in growth rate and ciguatera toxin production among geographically distinct isolates of *Gambierdiscus toxicus*. Intergovernmental Oceanographic Commission of UNESCO.
- Sunda, W.G., Hardison, D.R., 2007. Ammonium uptake and growth limitation in marine phytoplankton. *Limnol. Oceanogr.* 52(6), 2496-2506.
- Tatters, A.O., Muhlstein, H.I., Tomas, C.R., 2010. The hemolytic activity of *Karenia selliformis* and two clones of *Karenia brevis* throughout a growth cycle. *J. Appl. Phycol.* 22(4), 435-442.
- Tatters, A.O., Van Wagoner, R.M., Wright, J.L.C., Tomas, C.R., 2012. Regulation of spiroimine neurotoxins and hemolytic activity in laboratory cultures of the dinoflagellate *Alexandrium peruvianum* (Balech & Mendiola) Balech & Tangen. *Harmful Algae* 19, 160-168.
- Tester, P.A., Kibler, S.R., Holland, W.C., Usup, G., Vandersea, M.W., Leaw, C.P., Teen, L.P., Larsen, J., Mohammad-Noor, N., Faust, M.A., Litaker, R.W., 2014. Sampling harmful benthic dinoflagellates: comparison of artificial and natural substrate methods. *Harmful Algae* 39(0), 8-25.
- Van Dolah, F.M., Finley, E.L., Haynes, B.L., Doucette, G.J., Moeller, P.D., Ramsdell, J.S., 1994. Development of rapid and sensitive high throughput pharmacologic assays for marine phycotoxins. *Natural Toxins* 2(4), 189-196.
- Vandersea, M.W., Kibler, S.R., Holland, W.C., Tester, P.A., Schultz, T.F., Faust, M.A., Holmes, M.J., Chinain, M., Litaker, R.W., 2012. Development of semi-quantitative PCR assays for the detection and enumeration of *Gambierdiscus* species (Gonyaulacales, Dinophyceae). *Journal of Phycology* 48(4), 902-915.
- Xu, Y., Richlen, M.L., Liefer, J.D., Robertson, A., Kulis, D., Smith, T.B., Parsons, M.L., Anderson, D.M., 2016. Influence of environmental variables on *Gambierdiscus* spp. (Dinophyceae) growth and distribution. *PLoS One* 11(4), e0153197.
- Yasumoto, T., Bagnis, R., Vernoux, J., 1976. Toxicity of the surgeonfishes II. Properties of the principal water-soluble toxin. *Bull. Jpn. Soc. Sci. Fish* 42, 359-365.
- Yoshimatsu, T., Yamaguchi, H., Iwamoto, H., Nishimura, T., Adachi, M., 2014. Effects of temperature, salinity and their interaction on growth of Japanese *Gambierdiscus* spp. (Dinophyceae). *Harmful Algae* 35, 29-37.

Figure captions

Figure 1. Neuro-2a (N2a) cytotoxicity of dichloromethane soluble fractions (DSFs, n=3) of 13 *Gambierdiscus* strains. Results are expressed in fg CTX3C eq cell⁻¹.

Figure 2. Hemolytic activity of aq. MeOH soluble fractions (MSFs, n=3) of 13 *Gambierdiscus* strains evaluated by means of a human erythrocyte lysis assay (ELA). Results are expressed in pg MTX eq cell⁻¹.

Figure 3. Plot of log CTX toxicity (fg CTX3C eq μm^{-3}) versus MTX toxicity (pg MTX eq μm^{-3}).

Figure 4. Dendrogram of a hierarchical cluster analysis of 13 *Gambierdiscus* strains based on the following three variables: CTX-toxicity in the DSF (fg CTX3C eq cell⁻¹), MTX-toxicity in the MSF (pg MTX eq cell⁻¹) and cell biovolume (μm^3 cell⁻¹).

Figure S1. Sigmoidal dose-response curve of CTX3C and CTX1B standards on the neuro-2a (N2a) assay plotted using GraphPad Prism 6.0 (Hardison et al., 2016). Error bars represent the standard deviation (SD, n=12 for CTX3C, n=14 for CTX1B). The CTX standard used in this study was CTX3C only.

Figure S2. Sigmoidal dose-response curve of MTX standard on the human erythrocyte lysis assay (ELA) plotted using GraphPad Prism 6.0. Error bars represent the standard deviation (SD) of four replicates.


Table captions

Table 1. Denomination and origin of *Gambierdiscus* strains examined in this study.

Table 2. Maximum specific growth rates (μ_{max} , divisions day⁻¹) and per-cell CTX- and MTX-toxicity of the *Gambierdiscus* strains cultivated in this study.


Figures

Figure 1. Neuro-2a (N2a) cytotoxicity of dichloromethane soluble fractions (DSFs, n=3) of 13 *Gambierdiscus* strains. Results are expressed in fg CTX3C eq cell⁻¹.




(*) cultured at the CCFHR laboratory (section 2.2.1). The other strains were cultured at the IFREMER laboratory (Nantes, France) (section 2.2.2).

Figure 2. Hemolytic activity of aq. MeOH soluble fractions (MSFs, n=3) of 13 *Gambierdiscus* strains evaluated by means of a human erythrocyte lysis assay (ELA). Results are expressed in pg MTX eq cell⁻¹.




(*) cultured at the CCFHR laboratory (section 2.2.1). The other strains were cultured at the IFREMER laboratory (Nantes, France) (section 2.2.2).

Figure 3. Plot of log CTX toxicity (fg CTX3C eq μm^{-3}) versus MTX toxicity (pg MTX eq μm^{-3}).



(*) cultured at the CCFHR laboratory (section 2.2.1). The other strains were cultured at the IFREMER laboratory (Nantes, France) (section 2.2.2).

Figure 4. Dendrogram of a hierarchical cluster analysis of 13 *Gambierdiscus* strains based on the following three variables: CTX-toxicity in the DSF (fg CTX3C eq cell^{-1}), MTX-toxicity in the MSF (pg MTX eq cell^{-1}) and cell biovolume ($\mu\text{m}^3 \text{cell}^{-1}$).



(*) cultured at the CCFHR laboratory (section 2.2.1). The other strains were cultured at the IFREMER laboratory (Nantes, France) (section 2.2.2).

Tables

Table 1. Denomination and origin of *Gambierdiscus* strains examined in this study.

Location	Species / Strain	Origin	Culture Collection	Reference
Atlantic Ocean	<i>G. australes</i> / VGO1178 * ^x	Punta Hidalgo, Tenerife, Canary Islands	CCVIEO ^(a)	(Fraga and Rodríguez, 2014)
	<i>G. australes</i> / VGO1181 * ^x	Punta Hidalgo, Tenerife, Canary Islands	CCVIEO	Sequencing of LSU rDNA (D1-D3 region) (GENBANK KY549925)
	<i>G. excentricus</i> / VGO791 * ^x	Punta Hidalgo, Tenerife, Canary Islands	CCVIEO	(Fraga et al., 2011)
	<i>G. silvae</i> / VGO1167 * ^x (species formerly known as <i>G. ribotype 1</i>)	Punta Hidalgo, Tenerife, Canary Islands	CCVIEO	(Fraga and Rodríguez, 2014)
	<i>G. silvae</i> / VGO1180 * ^x (species formerly known as <i>G. ribotype 1</i>)	Punta Hidalgo, Tenerife, Canary Islands	CCVIEO	(Fraga and Rodríguez, 2014)
Mediterranean Sea	<i>G. carolinianus</i> / Greece Gam2 * ^x	Crete, Greece	CCFHR ^(b)	Species-specific qPCR assays (Vandersea et al., 2012)
Pacific Ocean	<i>G. australes</i> / CCMP1653 (NOAA 24) * ^x (strain previously reported as T39 strain)	Tern Island, Hawaii	NCMA ^(c)	(Babinchak et al., 1986; Litaker et al., 2009)
	<i>G. balechii</i> / VGO917 * ^x	Manado, Celebes Sea, Indonesia	CCVIEO	(Bravo et al., 2014; Fraga et al., 2016)
	<i>G. caribaeus</i> / Bill Hi Gam8 * ^x	Waikiki Beach, Honolulu, Hawaii	CCFHR	Species-specific qPCR assays (Vandersea et al., 2012)
	<i>G. carpenteri</i> / Pat Hi Jar7 Gam11 * ^x	Waikiki Beach, Honolulu, Hawaii	CCFHR	Species-specific qPCR assays (Vandersea et al., 2012)
	<i>G. pacificus</i> / CCMP1650 (NOAA 9) * ^x	Moorea, Society Islands, French Polynesia	NCMA	(Litaker et al., 2009)
	<i>G. scabrosus</i> / KW070922_1 * ^x (species formerly known as <i>Gambierdiscus</i> sp. type 1)	Kashiwa-jima Island, Otsuki, Kochi, Japan	KU ^(d)	(Nishimura et al., 2013; Nishimura et al., 2014)
	<i>Gambierdiscus</i> sp. / Viet Nam * ^x (strain reported as <i>G. toxicus</i> Vietnam)	Cau Island, Binh Thuan, South China Sea, Viet Nam	VNIO ^(e)	(Roeder et al., 2010)

* = strains cultured at the CCFHR laboratory (Beaufort, NC, USA) (section 2.2.1).

^x = strains cultured at the IFREMER laboratory (Nantes, France) (section 2.2.2).

(a) Culture Collection of Harmful Microalgae of IEO (CCVIEO), Centro de Vigo, Vigo, Spain.

(b) National Oceanographic and Atmospheric Administration (NOAA), Center for Coastal Fisheries Habitat Research (CCFHR), Beaufort, NC, USA.

(c) Provasoli – Guillard National Center for Marine Algae and Microbiota (NCMA), Bigelow Laboratory for Ocean Sciences, East Boothbay, Maine, USA.

(d) Kochi University (KU), Kochi, Japan.

(e) Viet Nam National Institute of Oceanography (VNIO, VAST), Vinh Nguyen, Nha Trang, Viet Nam.

Table 2. Maximum specific growth rates (μ_{\max} , divisions day⁻¹) and per-cell CTX- and MTX-toxicity of the *Gambierdiscus* strains cultivated in this study.

Location	Species / Strain	μ_{\max} (divisions day ⁻¹) \pm RSD	R ²	n	days _{tot}	ESV \pm RSD (n=3) ($\mu\text{m}^3 \text{ cell}^{-1} \times 10^5$)	DSF: fg CTX3C eq cell ⁻¹ \pm SD (n=3)	MSF: pg MTX eq cell ⁻¹ \pm SD (n=3)
Atlantic Ocean	<i>G. australes</i> / VGO1178	ND				1.57 \pm 0.3%	1.4 \pm 0.4	4.3 \pm 1.2
	<i>G. australes</i> / VGO1181	ND				1.24 \pm 1.2%	0.6 \pm 0.3	4.7 \pm 2.8
	<i>G. excentricus</i> / VGO791	0.099 \pm 2.59%	0.993	13	67	2.69 \pm 1.3%	1,426 \pm 55	85.7 \pm 41.5
	<i>G. silvae</i> / VGO1167	ND				0.93 \pm 7.6%	10.3 \pm 2.7	2.2 \pm 0.3
	<i>G. silvae</i> / VGO1180	ND				1.02 \pm 0.7%	12.4 \pm 3.6	3.3 \pm 0.9
Mediterranean Sea	<i>G. carolinianus</i> / Greece Gam2 (*)	0.129 \pm 4.77%	0.971	15	78	1.55 \pm 0.2%	3.3 \pm 0.6	10.6 \pm 0.4
Pacific Ocean	<i>G. australes</i> / CCMP1653	0.149 \pm 4.29%	0.980	13	53	1.37 \pm 0.5%	2.7 \pm 0.9	5.0 \pm 0.5
	<i>G. balechii</i> / VGO917	0.100 \pm 4.46%	0.979	13	53	0.82 \pm 0.5%	3.4 \pm 1.5	19.9 \pm 2.9
	<i>G. caribaeus</i> / Bill Hi Gam8 (*)	0.175 \pm 1.81%	0.996	15	78	1.64 \pm 2.4%	1.6 \pm 1.0	5.3 \pm 1.0
	<i>G. carpenteri</i> / Pat Hi Jar7 Gam11 (*)	0.141 \pm 5.59%	0.964	14	68	2.24 \pm 0.8%	1.4 \pm 0.6	6.3 \pm 1.9
	<i>G. pacificus</i> / CCMP1650	0.226 \pm 5.07%	0.950	13	53	0.74 \pm 0.7%	12.1 \pm 0.0	20.1 \pm 5.9
	<i>G. pacificus</i> / CCMP1650 (*)	0.244 \pm 2.59%	0.991	15	73	0.70 \pm 2.2%	13.5 \pm 0.4	20.3 \pm 1.6
	<i>G. scabrosus</i> / KW070922_1	0.140 \pm 1.84%	0.996	15	77	1.04 \pm 0.4%	27.9 \pm 3.8	1.5 \pm 0.2
<i>Gambierdiscus</i> sp. / Viet Nam	0.124 \pm 2.42%	0.992	15	74	1.21 \pm 4.0%	40.8 \pm 19.6	70.0 \pm 44.8	

(*) = cultured at the CCFHR laboratory (Beaufort, NC, USA) (section 2.2.1). The other strains were cultured at the IFREMER laboratory (Nantes, France) (section 2.2.2).

days_{tot} = the total duration of the culture throughout the study expressed in days.

DSF = dichloromethane soluble fraction

MSF = aqueous methanol soluble fraction

ESV = Estimated Spherical Volume (MultisizerTM 3 Coulter Counter®).

ND = not determined.

RSD = Relative Standard Deviation.

SD = Standard Deviation.

Supplementary info

Figure S1. Sigmoidal dose-response curve of CTX3C and CTX1B standards on the neuro-2a (N2a) assay plotted using GraphPad Prism 6.0 (Hardison et al., 2016). Error bars represent the standard deviation (SD, n=12 for CTX3C, n=14 for CTX1B). The CTX standard used as reference in this study was CTX3C only.




Figure S2. Sigmoidal dose-response curve of MTX standard on the human erythrocyte lysis assay (ELA) plotted using GraphPad Prism 6.0. Error bars represent the standard deviation (SD) of four replicates.

