

---

## Setting of a procedure for experimental fertilisation of Pacific oyster (*Crassostrea gigas*) oocytes

Y.P. Song<sup>a</sup>, M. Suquet<sup>a, \*</sup>, I. Quéau<sup>a</sup> and L. Lebrun<sup>a</sup>

<sup>a</sup> Ifremer, Station Expérimentale d'Argenton, Département PFOM, 11 presqu'île du vivier, 29840 Argenton, France

\*: Corresponding author : M. Suquet, Tel. : +33 2 9889 5755; fax: +33 2 9889 5777, email address : [Marc.Suquet@ifremer.fr](mailto:Marc.Suquet@ifremer.fr)

---

### Abstract:

The quality of oyster spermatozoa has become an issue for modern aquaculture. However, reliable protocols used to assess gamete quality are lacking. The aim of this work is to define a standardised experimental fertilisation protocol for Pacific oyster oocytes.

Six experiments have been carried out in this study. The optimal conditions for Pacific oyster experimental fertilisation have been defined: (1) oocytes can be conserved in seawater for at least 4 h before fertilisation, (2) oocyte concentration: 100–1000 ml<sup>-1</sup> seawater, (3) fertilisation volume: 10–100 ml, (4) spermatozoa:oocyte ratio: 400 to maintain discriminating conditions, (5) gamete contact time: longer than 10 min and (6) fertilisation and incubation temperatures: both 19 °C. The effect of individual male significantly influenced the fertilisation success. The results of this study will be exploited for spermatozoal quality analysis in combination with other techniques.

**Keywords:** Fertilisation; Experimental; Pacific oyster; *Crassostrea gigas*

## 1. Introduction

---

It is widely admitted that reproductive success of *C. gigas* is highly variable and that this variability is partially due to inconsistent gamete quality (Boudry et al., 2002). Gamete quality can be characterized by fertilisation rates and by embryo development yields. It results from multiple cellular and subcellular parameters of the gametes. The quality of gametes has been extensively studied in many freshwater and seawater fishes (Billard et al., 1995) but it is little known about molluscs and especially bivalves. Shellfish aquaculture requires a strict control of the reproduction process including a better assessment of gamete quality. In other domains, like ecotoxicology, the gametes have been used as a bio-indicator of modification and perturbation of environment (Kime and Nash, 1999). For these purposes, an objective and reproducible experimental fertilisation procedure is indispensable.

From a series of laboratory experiments, it appears that fertilisation success in several sessile and sedentary mollusc broadcast spawners depends on gamete concentration, gamete age, contact time between gametes, distance between spawning individuals and hydrodynamic conditions (Levitan, 1991; Hodgson et al., 2007; Adams et al., 2008). The experimental techniques can greatly influence the fertilisation success. An experimental incubation for Pacific oyster embryos has been established in previous work (Suquet et al., 2007). The aim of this work is to establish an experimental fertilisation protocol for Pacific oyster oocytes.

## 2. Materials and methods

---

### 2.1. Oyster conditioning and gamete stripping

Pacific oysters were collected in Charentes (France) and transferred to the experimental hatchery of Ifremer (Argenton, France). They were fed two microalgae (*Isochrysis galbana* clone Tahitian (T-Iso) and *Chaetoceros gracilis*) at a ratio of dry mass of algae to oyster of 2 : 100 and maintained in 200L tanks with running seawater at 19°C in order to obtain gonad maturation (Chavez-Villalba et al., 2002).

Gametes were collected from the oyster by the dry stripping method (Allen and Bushek, 1992). Briefly, gonads were dissected out and placed in a small glass bowl with 5ml salt stock solution "Store Gigas" (Brizard et al., 2001) for sperms or with 5ml filtered at 1µm seawater (FSW) for oocytes. The gonads were cut up into small chunks and the gonad material was agitated in the solution to release the gametes. The sperm suspension was filtered at 20 µm to remove the large chunks of gonad material. The oocyte suspension was successively filtered at 110 µm and 60 µm to remove the large and small chunks of gonads. After dilution to 1/1000 or 1/10000 in seawater, sperm concentration was determined by Coulter Counter in all the experiments. The concentration of oocytes was determined by microscopic count (3\*50µl). After oyster stripping, female gametes were conserved in a 2L container with FSW and at 19°C, while male gametes were stored in "Store Gigas" solution at 4°C prior fertilisation. Fertilisation occurred in a 1.8L beaker.

As all experimental conditions of fertilisation could not be tested at once, six separate experiments have been conducted. Thus, the results of previous experiment could be considered in optimising the following one. All the experiments were carried out with triplicate trials. When not stated, fertilisation standard procedure was as follows: after 30 minutes of contact between spermatozoa and oocytes, beakers were filled up to 1.8L with FSW at 19°C for incubation. The D-larval yield (48 hours) was used to estimate the fertilisation success (number of D-larvae.total number of embryos<sup>-1</sup>). Non viable larvae (translucent or non D shaped) were not included.

## **2.2. Experimental design**

### **2.2.1. Effects of oocyte conservation duration in FSW**

After 1, 2, 3, 4 hours of maintenance in FSW prior to fertilisation, 50000 oocytes from three females were transferred into a beaker to obtain a spermatozoa : oocyte ratio of 150 : 1 in an adjusted volume of 50ml. Spermatozoa were pooled from three males.

### **2.2.2. Effect of oocyte concentration**

Five oocyte concentrations were tested ( $100.\text{ml}^{-1}$ ,  $500.\text{ml}^{-1}$ ,  $1000.\text{ml}^{-1}$ ,  $5000.\text{ml}^{-1}$  and  $10000.\text{ml}^{-1}$ ) using three individual females. The fertilisation was conducted in 40 ml FSW at a sperm : oocyte ratio of 200 : 1. Spermatozoa were pooled from three males.

### **2.2.3. Effect of fertilisation volume**

Three fertilisation volumes (10ml, 50ml and 100ml) were tested. Oocytes collected from three females were fertilised using a sperm pool from three males. Oocyte concentration was adjusted at  $500 \text{ ml}^{-1}$  and spermatozoa : oocyte ratio was fixed at 200 : 1.

### **2.2.4. Effect of spermatozoa : oocyte ratio**

Four sperm : oocyte ratios were tested 10 : 1, 100 :1, 500 :1, 2000 :1. Oocytes were pooled from three oysters and the concentration was adjusted to  $500 \text{ ml}^{-1}$  for fertilisation. Three males were individually tested in a fertilisation volume of 50ml.

### **2.2.5. Effect of gamete contact time**

Six contact times between spermatozoa and oocytes were tested :10 seconds, 10, 20, 30, 60 and 90 minutes. The oocytes from three females were pooled and the concentration of oocytes used for fertilisation was  $500.\text{ml}^{-1}$ . Three males were individually tested. The fertilisation was performed in 50 ml FSW at a sperm : oocyte ratio of 400. At the indicated contact time, the samples were respectively filtered with a  $20 \mu\text{m}$  mesh in order to remove spermatozoa.

### **2.2.6. Effect of temperature**

Four combinations of temperature were tested for fertilisation and incubation respectively: (1)  $19^{\circ}\text{C} - 19^{\circ}\text{C}$ , (2)  $15^{\circ}\text{C} - 19^{\circ}\text{C}$ , (3)  $19^{\circ}\text{C} - 15^{\circ}\text{C}$  and (4)  $15^{\circ}\text{C} - 15^{\circ}\text{C}$ . The fertilisation occurred in 50 ml FSW; spermatozoa : oocyte ratio was fixed at 400; the oocyte concentration was adjusted to  $500.\text{ml}^{-1}$ ; Oocytes were pooled from three females and spermatozoa were collected from three males. After 20 minutes fertilisation, the beaker was fully filled.

## **2.3. Data analysis**

The percentage data were arcsin square-root transformed to achieve homogeneity of variance prior to two way analysis of variance (ANOVA). When the D-larval yield was significantly different, Tukey's multiple rank comparison was used.

## **3. Results and discussion**

---

### **3.1. Effects of oocyte conservation duration in FSW**

During the first four hours, the maintenance of oocytes in seawater had no significant effect on D-larval yield (Fig. 1A). As a consequence, fertilisation experiment may last a minimum of

4 hours. In other molluscs, fertilisation success for oocyte from *Patella ulyssiponensis* and *Patella vulgata* remains higher than 40% for up to 12 hours old and 6 hours old in seawater, respectively (Hodgson et al., 2007). This reflects an adaptive response to sedentary life and to external fertilisation of these species. The ecological significance of gamete longevity of free spawning invertebrates has been debated by a number of authors. Some have argued that gamete longevity is not important because hydrodynamic processes will rapidly dilute gametes (Levitan, 1991; André and Lindegarth, 1995). Others have suggested that increased gamete longevity could enhance fertilisation success when sperm availability varies (Williams and Bentley, 2002; Yund and Meidel, 2003).

### **3.2. Effect of oocyte concentration**

When the oocyte concentration increased from 1000 ml<sup>-1</sup> to 5000 ml<sup>-1</sup>, there was a significant decrease of D-larval yield ( $P < 0.05$ ). A second decrease was observed between 5000 ml<sup>-1</sup> to 10000 ml<sup>-1</sup> (Fig. 1B). It has been suggested that the deleterious effect of a high oocyte density in *Crassostrea rhizophorae* is caused by a water-soluble substance, originating from the zygote rather than oxygen depletion nor the physical crowding of eggs (Rampersad et al., 1994).

### **3.3. Effect of fertilisation volume**

Between 10 and 100 ml, fertilisation volume had no significant effect on D-larval yield (Fig. 2A). However, a significant effect of females was observed ( $P < 0.05$ ).

### **3.4. Effect of spermatozoa : oocyte ratio**

A significant increase of D-larval yield was recorded when the spermatozoa : oocyte ratio increased from 10 to 100 and from 100 to 500 ( $P < 0.01$ ). Then, a plateau was observed (Fig. 2B). The effect of male on fertilisation success was significant. This experiment confirms that gamete ratio is a crucial factor for the production of larvae. Santos and Nascimento (1985) reported that fertilisation in *Crassostrea rhizophorae* is optimum with a sperm : oocyte ratio ranging from 100:1 to 5000: 1, resulting thus in more than 50% larvae. Maximum fertilisation rates were observed for ratios ranging from 1000:1 to 10000:1 in thawed Pacific oyster sperm (Adams et al., 2008). Excessive amounts of sperm (50000:1) during fertilisation in Pacific oyster were shown to decrease the percentage of viable larvae, probably because of polyspermy (Stephano and Gould., 1988). For this reason, the spermatozoa to oocyte ratio was fixed at 400 in order to maintain discriminating conditions.

### **3.5. Effect of gamete contact time**

A significant increase of D-larval yield was observed between 10 seconds and 10 minutes contact duration between spermatozoa and oocytes ( $P < 0.001$ ). There was no significant difference between 10 to 90 minutes of contact between gametes (Fig. 3A). The effect of male was significant on fertilisation success. In this experiment fertilisation occurred within the first 10 minutes of gamete contact time for the Pacific oyster. Similarly, most fertilisation occurred within 5 minutes in sea urchin (*Diadema antillarum*) and asteroid (*Haliotis laevigata*) eggs (Levitan, 1991; Babcock and Keesing, 1999).

### **3.6. Effect of temperature**

Compared to other temperature conditions, a significant decreased D-larval yield was observed for the condition: 15°C for fertilisation and 15°C for incubation (Fig. 3B;  $P < 0.05$ ). The effect in male was significant on D-larval yield. It has been shown that the D-larval yield significantly decreased as temperature decreased in the pearl oyster (*Pinctada imbricata*; O'Connor and Lawlar, 2004).

## 4. Conclusions and perspectives

---

An experimental fertilisation procedure of Pacific oyster oocytes has been established in this study for which the main results are presented in Table 1. This fertilisation procedure will be exploited for spermatozoal quality analysis of Pacific oyster in combination with other techniques. The effect of male on fertilisation success was significant, which confirms that future study of spermatozoal quality is necessary to support Pacific oyster aquaculture.

## Acknowledgements

---

The authors would like to thank C. Mingant (Ifremer) for his advice for rearing oysters.

## References

---

- Adams, S.L., Smith, J.F., Roberts, R.D., Janke, A.R., King, N.G., Tervit, H.R., Webb, S.C, 2008. Application of sperm cryopreservation in selective breeding of the Pacific oyster, *Crassostrea gigas* (Thunberg). *Aquac. Res.*, 39, 1434-1442.
- Allen, S.K, Bushek, D., 1992. Large scale production of triploid oysters, *Crassostrea virginica* (Gmelin) using “stripped” gametes. *Aquaculture* 103, 241-251.
- André, C., Lindegarth, M., 1995. Fertilization efficiency and gamete viability of a sessile, free-spawning bivalve, *Cerastoderma edule*. *Ophelia* 43, 215-227.
- Babcock, R., Keesing, J., 1999. Fertilization biology of the abalone *Haliotis laevigata*: laboratory and field studies. *Fish. Aquat. Sci.* 56, 1668-1678.
- Billard, R., Cosson, J., Crim, L.W., Suquet, M., 1995. Sperm physiology and quality. In: Bromage, N. R., Roberts, R. (Eds.) *Broodstock management and egg and larval quality*. Cambridge University Press, Cambridge, 25-52.
- Boudry, P., Collet, B., Cornette, F., Hervouet, V., Bonhomme, F., 2002. High variance in reproductive success of the Pacific oyster (*Crassostrea gigas*, Thunberg) revealed by microsatellite - based parentage analysis of multifactorial crosses. *Aquaculture* 204, 283-296.
- Brizard, R, Bernardi, M., Boudry, P., Haffray, P., Labbé, C., Maise, G., Maurouard, E., Robert, R., Roger, J.L., 2001. Bilan du projet Cryoyster, Ifremer, 36pp
- Chavez-Villalba, J., Pommier, J., Andriamiseza, J., Pouvereau, S., Barret, J., Cochard, J.C, Le Penne, M., 2002. Broodstock conditioning of the oyster *Crassostrea gigas*: origin and temperature effect. *Aquaculture* 214, 115-130.
- Hodgson, A.N., Le Quesne, W.J.F, Hawkins, S.J., Bishop, J.D.D., 2007. Factors affecting fertilisation success in two species of patellid limpet (Mollusca: Gastropoda) and development of fertilisation kinetics models. *Mar. Biol.* 150, 415-426.
- Kime, D.E., Nash, J.P., 1999. A strategy for assessing the effects of xenobiotics on fish reproduction. *Sci. Total Environ.* 225, 3-11.

Levitan, D.R., 1991. Influence of body size and population density on fertilisation success and reproductive output in a free-spawning invertebrate. Biol. Bull. 181, 261-268.

O'Connor, W.A., Lawlar, N.F., 2004. Salinity and temperature tolerance of embryos and juveniles of the pearl oyster, *Pinctada imbricata* Röding. Aquaculture 229, 493-506.

Rampersad, J.N., Agard, J.B.R., Ammons D., 1994. Effects of gamete concentration on the in vitro fertilisation of manually extracted gametes of the oyster (*Crassostrea rhizophorae*). Aquaculture 123, 153-162.

Santos, A.E., Nascimento, I.A., 1985. Influence of gamete density, salinity and temperature on the normal embryonic development of the mangrove oyster *Crassostrea rhizophorae*. Aquaculture 47, 335-352.

Stephano, J.L., Gould, M., 1988. Avoiding polyspermy in the oyster (*Crassostrea gigas*). Aquaculture 73, 295-307.

Suquet, M., Amourda, C., Mingant, C., Quéau, I., Lebrun, L., Brizard, R., 2007. Setting of a procedure for experimental incubation of Pacific oyster (*Crassostrea gigas*) embryos. Aquaculture 273, 503-508

William, M.E., Bentley, M.G., 2002. Fertilization success in marine invertebrates: the influence of gamete age. Biol. Bull. 202, 34-42.

Yund, P.O., Meidel, S.K., 2003. Sea urchin spawning in benthic boundary layers: are eggs fertilised before advirting away from females? Limnol. Oceanogr. 48, 795-801.

## Tables

---

Table 1. Recommended experimental fertilisation conditions for Pacific oysters.

Oocyte conservation time in FSW	1-4 hours
Oocyte concentration	100-1000 ml <sup>-1</sup>
Fertilisation volume	10-100 ml
Spermatozoa : oocyte ratio	400
Gamete contact time	>10 minutes
Fertilisation and incubation temperatures	19°C-19°C

## Figures

Figure 1. A: Effect of oocyte conservation duration in FSW on D-larval yield, B: Effect of oocyte concentration on D-larval yield (mean  $\pm$  SD, n=3; different letters refer to significantly different results).





Figure 2. A: Effect of fertilisation volume on D-larval yield, B: Effect of sperm : oocyte ratio on D-larval yield (mean  $\pm$  SD, n=3, different letters refer to significantly different results).






Figure 3. A: Effect of gamete contact time on D-larval yield, B: Effect of temperature on D-larval yield (mean  $\pm$  SD, n=3, different letters refer to significantly different results).

