

Simulation of the combined effects of artisanal and recreational fisheries on a Mediterranean MPA ecosystem using a trophic model

Camille Albouy^{1,*}, David Mouillot¹, Delphine Rocklin^{1,2}, Jean M. Culioli³, François Le Loc'h⁴

¹Université Montpellier 2 CC 093, UMR 5119 CNRS-UM2-IFREMER-IRD ECOLAG, 34095 Montpellier, France

²Laboratoire de Biologie Halieutique—Département Sciences et Technologies Halieutiques, IFREMER Brest, BP 70, Technopôle Brest Iroise, 29280 Plouzané, France

³Office de l'Environnement de la Corse, Reserve Naturelle des Bouches de Bonifacio, 20250 Corte, France

⁴IRD, UMR 212 EME, CRHMT, avenue Jean Monnet 34203 Sète cedex, BP 171, France

*Email: albouycamille@gmail.com

Marine Ecology Progress Series: 412: 207–221 (2010)

Supplement 1. Additional data

Supplement 1A. Input data and references by biotic group for the Bonifacio Straits Natural Reserve (BSNR) model. *B*: biomass; *P/B*: production per unit of biomass; *Q/B*: consumption rate per unit of biomass; *EE*: ecotrophic efficiency; *Z*: total mortality; *F*: fishing mortality rate; *M*: non-predation natural mortality rate

Group	<i>B</i>	<i>P/B</i>	<i>Q/B</i>	<i>EE</i>	Diets
1 <i>Tursiops truncatus</i>	Life Linda 2000 to 2001 data from visual survey units of individuals has been transformed to t km ⁻² with the mean body weight per species	Data from Coll et al. (2006) corrected to consider differences of temperature with the equation of Opitz (Opitz 1996)			Blanco et al. (2001)
2 <i>Sphyraena sphyraena</i>	Unpublished BSNR data \ estimates from fishing data	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)		See Supplement 1C
3 <i>Scyliorhinus canicula</i>	Unpublished BSNR data \ estimates from fishing data	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)		See Supplement 1C

4	Piscivorous fish	Unpublished BSNR data \ estimates from fishing data and visual census	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	See Supplement 1C
5	Small pelagic feeders	Unpublished BSNR data \ estimates from fishing data	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	See Supplement 1C
6	<i>Epinephelus marginatus</i>	Unpublished BSNR data \ visual census data	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	Renones et al. (2002)
7	Opportunist piscivorous fish	Unpublished BSNR data \ estimates from fishing data and visual census	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	See Supplement 1C
8	Cephalopods	Unpublished BSNR data \ estimates from fishing data	Data from Opitz (1996) corrected to consider differences of temperature between areas with the Opitz equation		Ambrose & Nelson (1983)
9	Birds	Unpublished data from visual survey units of individuals has been transformed to $t \text{ km}^{-2}$ with the mean body weight per species	Data from Pinnegar & Polunin (2004) corrected to consider differences of temperature between areas with the Opitz equation		Hickling (1983)
10	Benthic invertebrate feeders	Unpublished BSNR data \ estimates from fishing data and visual census	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	See Supplement 1C
11	Zooplanktivorous fish	Data from the Pinnegar & Polunin (2004) model \ unpublished fishing data	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	See Supplement 1C
12	Mollusc feeders	Unpublished BSNR data \ estimates from fishing data and visual census	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	See Supplement 1C

13	Benthic invertebrate feeders 2	Unpublished BSNR data \ estimates from fishing data and visual census	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)		See Supplement 1C
14	Shrimp		Data from Coll et al. (2006), corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)		Fixed to current values	Opitz 1996
15	Macroplanktivorous fish	Unpublished BSNR data \ estimates from fishing data and visual census	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)		See Supplement 1C
16	Decapods	Data from Sala (1997)	Data from Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)	Data from Arias-Gonzales (1994) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)		Pinnegar & Polunin (2004)
17	Lobster & spiny lobster	Unpublished BSNR data \ estimates from fishing and sampling data	Data from Marin (1987) and Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)			Opitz (1996)
18	Gastropods		Data from Opitz (1996) corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)		Fixed to current values	Opitz (1996), Pearson & Gage (1984)
19	Zooplankton		Data from Pinnegar & Polunin (2004) corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)		Fixed to current values	Kleppel (1993), Kleppel et al. (1996)
20	Polychaetes	Data from Sala (1997)	Data from Pinnegar & Polunin (2004) corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)			Opitz (1996), Fauchald & Jumars (1979)

21	Suspension feeders		Data from Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)	Fixed to current values	Chintiroglou & Koukouras (1992), Arias-Gonzales (1994), Coma et al. (1995), Ribes et al. (1999)
22	Echinoderms	Data from the Pinnegar & Polunin (2004) model	Data from Opitz (1996) and Pauly et al. (1993) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)		Arias-Gonzales (1994), Frantzis et al. (1988), Pearson & Gage (1984), Verlaque & Nedelec (1983)
23	Protozoan plankton		Data from Pinnegar & Polunin (2004) corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)	Data from Sanchez & Olaso (2004) corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)	Fixed to current values Olivieri et al. (1993)
24	<i>Sarpa salpa</i>	Unpublished BSNR data \ estimates from fishing data	$Z = F + M$; $M =$ empirical equation from Pauly (1980)	Empirical relationship from Pauly et al. (1990)	Havelange et al. (1997)
25	Other crustaceans	Data from the Pinnegar & Polunin (2004) model	Data from Pinnegar & Polunin (2004) corrected to consider differences of temperature between areas with the Opitz equation (Opitz 1996)		Arias-Gonzales (1994), Caine (1977), Vannier et al. (1998), Viejo & Arrontes (1996)
26	Amphipods		Data from Vetter (1996) and Schwinghamer et al. (1986) and corrected using the Opitz equation (Opitz 1996)	Data from Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)	Fixed to current values Greze (1968)
27	Bivalves		Data from Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)	Fixed to current values	Opitz (1996)

28	Macroplankton		Data from Pinnegar & Polunin (2004) modified to consider area's temperature (Opitz 1996)	Fixed to current values	Dauby (1980), Hobson & Chess (1976), Kinsey & Hopkins (1994), Purcell (1981), Sullivan et al. (1994)
29	Phytoplankton		Data from Pinnegar & Polunin (2004) and modified to consider area's temperature (Opitz 1996)	Fixed to current values	
30	Macro-algae		Data from Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)	Fixed to current values	
31	<i>Posidonia oceanica</i>	Data from Gobert (2002) modified to include the meadow surface	Data from Opitz (1996) corrected to consider differences in temperature between areas with the Opitz equation (Opitz 1996)		
32	Detritus	Data from the Pinnegar & Polunin (2004) model			

Supplement 1B. Final diet matrix of the Bonifacio Straits Natural Reserve (BSNR) model. BIF: benthic invertebrate feeders

Prey/predator	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
1 <i>Tursiops truncatus</i>																												
2 <i>Sphyræna sphyræna</i>																												
3 <i>Scyliorhinus canicula</i>																												
4 Piscivorous fish	0.200	0.025		0.070		0.022		0.01																				
5 Small pelagic feeders	0.528		0.460	0.022	0.050		0.008																					
6 <i>Epinephelus marginatus</i>																												
7 Opportunist piscivorous fish		0.715		0.109	0.263	0.103	0.020		0.106																			
8 Cephalopods	0.032		0.090	0.099	0.010	0.213	0.125	0.030	0.005	0.010																		
9 Birds																												
10 BIF	0.034	0.101		0.090	0.108		0.027			0.000																		
11 Zooplanktivorous fish	0.202		0.090	0.114	0.220	0.011	0.042	0.001	0.366	0.001	0.009																	
12 Mollusc feeders		0.125		0.073			0.155		0.275	0.005																		
13 BIF2		0.034		0.011	0.257	0.101	0.090	0.150	0.246	0.017			0.000															
14 Shrimp				0.006	0.009	0.013	0.153	0.035		0.057		0.163		0.002	0.113													
15 Macrocarnivorous fish				0.020			0.022																					
16 Decapods				0.365		0.418	0.276	0.636		0.489		0.127	0.179		0.036	0.083		0.008										
17 Superior crustaceans	0.004		0.160	0.004		0.057	0.001			0.024																		
18 Gastropods						0.020	0.007	0.041		0.004		0.178	0.037	0.070	0.001	0.030	0.218	0.042		0.005		0.009						
19 Zooplankton					0.007								0.773	0.028	0.113	0.001	0.063		0.006	0.042		0.001		0.009	0.027		0.009	0.027
20 Polychaetes			0.005				0.012			0.109	0.084	0.034	0.406	0.050	0.001	0.012		0.125	0.089		0.001							
21 Suspensivores			0.005				0.003			0.001	0.009					0.010	0.375	0.090	0.027		0.073							
22 Echinoderms							0.013			0.098		0.008	0.012	0.016		0.001	0.032	0.001	0.001									
23 Protozoan plankton																0.050			0.394	0.050	0.063		0.247		0.009	0.050	0.100	0.010
24 <i>Sarpa salpa</i>						0.016																						
25 Other crustaceans			0.160	0.004		0.026	0.029	0.011		0.063	0.022	0.067	0.019	0.055		0.254		0.066		0.056		0.003						
26 Amphipods			0.040	0.001	0.009		0.007	0.049		0.114	0.074	0.067	0.214	0.087	0.150	0.262		0.006		0.048								
27 Bivalves							0.003	0.038					0.272	0.060	0.004	0.009												
28 Macroplankton					0.065		0.007				0.014		0.210	0.692	0.020					0.059	0.313	0.011		0.110				0.280
29 Phytoplankton										0.006									0.400	0.074	0.250			0.000	0.018		0.150	
30 Macro-algae			0.002							0.009	0.069	0.011	0.097	0.001	0.070	0.225	0.441		0.160		0.781	0.050		0.444	0.561		0.050	
31 <i>Posidonia oceanica</i>												0.091									0.078		0.890					
32 Detritus													0.237		0.136	0.150	0.221	0.200	0.390	0.375	0.043	0.303		0.021	0.362	0.750		
33 Import																						0.400		0.499			0.660	

Supplement 1C. Composition of fish groups and their associated references for diets. Asterisks show groups targeted by artisanal fishing, boat and shore fishing, or spearfishing

Group	Species	Artisanal fishing	Boat and shore fishing	Spear fishing	Source
Piscivorous	<i>Conger conger</i>	*		*	Cau & Manconi (1984)
	<i>Dentex dentex</i>	*	*	*	Morales-Nin & Moranta (1997)
	<i>Lophius piscatorius</i>	*			Macpherson (1981)
	<i>Muraena helena</i>	*		*	Cau & Manconi (1984)
	<i>Phycis phycis</i>	*		*	Papaconstantinou & Caragitsou (1989)
	<i>Synodus saurus</i>	*			Golani (1993)
Opportunist piscivorous fish	<i>Apogon imberbis</i>				Pinnegar & Polunin (2000)
	<i>Labrus viridis</i>	*		*	Bell & Harmelin-Vivien (1983)
	<i>Pagellus acarne</i>	*	*	*	Morato et al. (2001)
	<i>Pagrus pagrus</i>	*	*	*	Papaconstantinou & Caragitsou (1989)
	<i>Scomber</i> sp.	*		*	Cabral & Murta (2002)
	<i>Scorpaena porcus</i>	*		*	Arculeo et al. (1993)
	<i>Scorpaena scrofa</i>	*		*	Bell & Harmelin-Vivien (1983)
	<i>Serranus scriba</i>	*		*	Bell & Harmelin-Vivien (1983)
	<i>Uranoscopus scaber</i>	*			Sanz (1985)
Benthic invertebrate feeders	<i>Labrus merula</i>	*		*	Bell & Harmelin-Vivien (1983)
	<i>Pagellus erythrinus</i>	*	*	*	Rosecchi (1983)
	<i>Sciaena umbra</i>	*		*	Derbal & Kara (2007)
	<i>Scorpaena notata</i>	*		*	Bell & Harmelin-Vivien (1983)
	<i>Thalassoma pavo</i>				Bell & Harmelin-Vivien (1983)
Macroplanktivorous fish	<i>Serranus cabrilla</i>	*	*	*	Bell & Harmelin-Vivien (1983)
	<i>Spondyliosoma cantharus</i>	*	*		Bell & Harmelin-Vivien (1983)
	<i>Trachinus draco/araneus</i>	*			Morte et al. (1999)
	<i>Trisopterus minutus</i>	*			Le Loc'h (2004)


Small pelagic feeders	<i>Merluccius merluccius</i>	*		Le Loc'h (2004)
	<i>Sarda sarda</i>	*		* Campo et al. (2006)
	<i>Seriola dumerilii</i>	*	*	* Mazzola et al. (2000)
	<i>Trachurus mediterraneus</i>	*		Ben Salem (1988)
	<i>Zeus faber</i>	*		Bell & Harmelin-Vivien (1983)
Mollusc feeders	<i>Coris julis</i>		*	Bell & Harmelin-Vivien (1983)
	<i>Diplodus puntazzo</i>	*		* Sala & Ballesteros (1997)
	<i>Diplodus sargus</i>	*	*	Sala & Ballesteros (1997)
	<i>Diplodus vulgaris</i>	*	*	* Sala & Ballesteros (1997)
	<i>Symphodus</i> sp.	*		Bell & Harmelin-Vivien (1983)
	<i>Trigla lucerna</i>	*		Macpherson (1981)
Zooplanktivorous fish	<i>Atherina boyeri</i>			Pinnegar & Polunin (2000)
	<i>Boops boops</i>			Bell & Harmelin-Vivien (1983)
	<i>Chromis chromis</i>			Bell & Harmelin-Vivien (1983)
	<i>Engraulis encrasicolus</i>			Tudela & Palomera (1997)
	<i>Mugilidés</i>			Blaber (1976)
	<i>Oblada melanura</i>	*	*	Moreno & Castro (1995)
	<i>Spicara maena</i>	*		Khoury (1987)
Benthic invertebrate feeders 2	<i>Diplodus annularis</i>	*	*	Rossechi & Nouaze (1989)
	<i>Gobius</i> sp.			Dierter Zander & Berg (1984)
	<i>Lipophrys pavo</i>			Dierter Zander & Berg (1984)
	<i>Mullus surmuletus</i>	*		* Pinnegar & Polunin (2000)
	<i>Parablennius rouxi</i>			Pinnegar & Polunin (2000)
	<i>Solea</i> sp.	*		* Bell & Harmelin-Vivien (1983)
	<i>Symphodus tinca</i>	*		* Khoury (1987)
	<i>Tripterygion</i> sp.			Bell & Harmelin-Vivien (1983)
Single species groups	<i>Epinephelis marginatus</i>	*		Renones et al. (2002)
	<i>Sarpa salpa</i>			Havelange et al. (1997)
	<i>Sphyraena sphyraena</i>	*	*	* Barreiros et al. (2002)

Supplement 1D. Temperature assessment

An average temperature for the modelled area is primordial for assessing the consumption biomass ratio for fish (Eq. 6), and production per unit of biomass (P/B) and consumption rate per unit of biomass (Q/B) (for other groups). Values for other groups were taken from the worldwide literature and adapted to our system using the Opitz equation (Opitz 1996).

To estimate an annual average sea surface temperature (SST) of the Bonifacio Straits Natural Reserve (BSNR) we collected a grid (resolution of 1° of weekly SST values from the National Climatic Data Center (NCDC) National Operational Model Archive and Distribution System Meteorological Data Server (NOMADS) of the US National Oceanic and Atmospheric Administration (NOAA) Satellite and Information Service (www.osdpd.noaa.gov/PSB/EPS/SST/al_climo_mon.html). Then, maps were interpolated at the 0.1° resolution needed for our analyses via ordinary kriging. Weekly SST values were averaged over the 2000 to 2001 period to provide 14 SST variables for each 0.1° grid cell. Twelve monthly means were used to calculate the mean annual SST then an average SST was extracted. The mean annual temperature for the BSNR was 18.6°C for the 2000 to 2001 period.

Supplement 1E. Ecosim output, with the variation of relative biomass for phytoplankton (red line), the variation of relative biomass for piscivorous fish in the absence of fishing effort ($F = 0$) (light blue line), the variation of relative biomass for piscivorous fish and actual fishing effort ($F = 1$) (dark blue line), and the variation of relative biomass for piscivorous fish and a 4-fold increase in fishing effort ($F = 4$) (green line)



LITERATURE CITED

- Ambrose R, Nelson B (1983) Predation by *Octopus vulgaris* in the Mediterranean. PSZN I: Mar Ecol 4:251–261
- Arculeo M, Froggia C, Riggio S (1993) Food partitioning between *Serranus scriba* and *Scorpaena porcus* (Perciformes) on the infralittoral ground of the south Tyrrhenian Sea. Cybium 17:251–258
- Arias-Gonzalez J (1994) Fonctionnement trophique d'un écosystème récifal: secteur de Tiahura, île de Moorea, Polynésie française. Thèse doctorale, Ecole pratique des hautes études, Paris
- Barreiros JP, Santos RS, de Borba AE (2002) Food habits, schooling and predatory behaviour of the yellowmouth barracuda, *Sphyraena viridensis* (Perciformes: Sphyraenidae) in the Azores. Cybium 26:83–88
- Bell J, Harmelin-Vivien M (1983) Fish fauna of French Mediterranean *Posidonia oceanica* seagrass meadows. 2. Feeding habits. Tethys 11:1–14
- Ben Salem M (1988) Régime alimentaire de *Trachurus trachurus* (Linnaeus, 1758) et de *T. mediterraneus* (Steindacher, 1868), (poissons, Téléostéens, Carangidae) de la province Atlantico Méditerranéenne. Cybium 12:247–253
- Blaber SJM (1976) Food and feeding ecology of Mugilidae in the St. Lucia lake system. Biol J Linn Soc 8:267–277
- Blanco C, Salomon O, Raga J (2001) Diet of the bottlenose dolphin (*Tursiops truncatus*) in the western Mediterranean Sea. J Mar Biol Assoc UK 81:1053–1058

- Cabral HN, Murta AG (2002) The diet of blue whiting, hake, horse mackerel and mackerel off Portugal. *J Appl Ichthyol* 18:14–23
- Caine E (1977) Feeding mechanisms and possible resources partitioning of the Caprellidae from Puget Sound, USA. *Mar Biol* 42:331–336
- Campo D, Mostarda E, Castriota L, Scarabello MP, Andaloro F (2006) Feeding habits of the Atlantic bonito, *Sarda sarda* (Bloch, 1793) in the southern Tyrrhenian Sea. *Fish Res* 81:169–175
- Cau A, Manconi P (1984) Relationship of feeding, reproductive-cycle and bathymetric distribution in *Conger conger*. *Mar Biol* 81:147–151
- Chintiroglou C, Koukouras A (1992) The feeding habits of three Mediterranean Sea anemone species, *Anemonia viridis*, *Anemonia equina* and *Cereus pedunculatus*. *Helgol Meeresunters* 46:53–68
- Coll M, Palomera I, Tudela S, Sarda F (2006) Trophic flows, ecosystem structure and fishing impacts in the South Catalan Sea, Northwestern Mediterranean. *J Mar Syst* 59:63–96
- Coma R, Gili J, Zabala M (1995) Trophic ecology of a benthic marine hydroid, *Campanularia everta*. *Mar Ecol Prog Ser* 119:211–220
- Dauby P (1980) Cycle annuel du zooplancton de surface dans la baie de Calvi (Corse). Biomasse totale et plancton copépodien. *Oceanol Acta* 3:403–407
- Derbal F, Kara H (2007) Régime alimentaire du corb *Sciaena umbra* (Scianidae) des côtes de l'est algérien. *Cybium* 31:189–197
- Dierter Zander C, Berg J (1984) Feeding ecology of littoral gobiid and blennioid fishes of the Banyuls area (Mediterranean Sea). II. Prey selection and size preference. *Vie Milieu* 34:149–157
- Fauchald K, Jumars P (1979) The diet of worms: a study of polychaete feeding guilds. *Oceanogr Mar Biol Annu Rev* 17:193–284
- Frantzis A, Berthon J, Maggiore F (1988) Relations trophiques entre les oursins *Arbacia lixula* et *Paracentrotus lividus* et le phytobenthos infralittoral superficiel dans la baie de Port-Cros. *Sci Rep Port-Cros Natl Park, France* 14:81–140
- Gobert S (2002) Variations spatiale et temporelle de l'herbier à posidonie *oceanica*. (I.) Delile (baie de la revellata-calvi-corse). Thèse doctorale Université de Liège, Liège
- Golani D (1993) The biology of the red-sea migrant, *Saurida undosquamis* in the Mediterranean and comparison with the indigenous confamilial *Synodus saurus* (Teleostei, Synodontidae). *Hydrobiologia* 271:109–117
- Greze I (1968) Feeding habits and food requirements of some amphipods in the Black Sea. *Mar Biol* 1:316–332
- Havelange S, Lepoint G, Dauby P, Bouquegneau JM (1997) Feeding of the sparid fish *Sarpa salpa* in a seagrass ecosystem: diet and carbon flux. *PSZN I Mar Ecol*:18:289–297
- Hickling RAO (1983) *Enjoying ornithology: a celebration of fifty years of British trust for ornithology, 1933–1983*. Calton (T. and A. D. Poyser), London
- Hobson E, Chess J (1976) Trophic interactions among fishes and zooplankters near shore at Santa Catalina Island, California. *Fish Bull* 74:567–598
- Khoury C (1987) Ichtyofaune des herbiers de posidonies du parc national de Port-Cros: composition, éthologie alimentaire et rôle dans le réseau trophique. Thèse doctorale Université Aix-Marseille II, Marseille

- Kinsey S, Hopkins T (1994) Trophic strategies of euphausiids in a low latitude ecosystem. *Mar Biol* 118:651–661
- Kleppel GS (1993) On the diets of calanoid copepods. *Mar Ecol Prog Ser* 99:183–195
- Kleppel GS, Burkart CA, Carter K, Tomas C (1996) Diets of calanoid copepods on the West Florida continental shelf: relationships between food concentration, food composition and feeding activity. *Mar Biol* 127:209–217
- Le Loc'h F (2004) Structure, fonctionnement et évolution des communautés benthiques des fonds meubles exploités du plateau continental nord Gascogne. Thèse de Doctorat Thesis, Université de Bretagne Occidentale, Brest
- Macpherson E (1981) Resource partitioning in a Mediterranean demersal fish community. *Mar Ecol Prog Ser* 4:183–193
- Marin J (1987) Exploitation, biologie et dynamique du stock de langouste rouge de corse *Palinurus elepha fabricius*. Thèse de Doctorat Université d'Aix Marseille II, Marseille
- Mazzola A, Favalaro E, Sara G (2000) Cultivation of the Mediterranean amberjack, *Seriola dumerili* (Risso, 1810), in submerged cages in the western Mediterranean sea. *Aquaculture* 181:257–268
- Morales-Nin B, Moranta J (1997) Life history and fishery of the common dentex (*Dentex dentex*) in Mallorca (Balearic Islands, western Mediterranean). *Fish Res* 30:67–76
- Morato T, Sola E, Gros MP, Menezes G (2001) Feeding habits of two congener species of seabreams, *Pagellus bogaraveo* and *Pagellus acarne*, off the Azores (northeastern Atlantic) during spring of 1996 and 1997. *Bull Mar Sci* 69:1073–1087
- Moreno T, Castro JJ (1995) Community structure of the juvenile of coastal pelagic fish species in the Canary Islands waters. *Sci Mar* 59:405–413
- Morte S, Redon M, Sanz-Brau A (1999) Feeding habits of *Trachinus darco* off the eastern coast of Spain (western Mediterranean). *Vie Milieu* 49:287–291
- Olivieri RA, Cohen A, Chavez FP (1993) An ecosystem model of Monterey Bay, California. In: Christensen V, Pauly D (eds) *Trophic models of aquatic ecosystems*. ICLARM, Manila
- Opitz S (1996) Trophic interactions in Caribbean coral reefs. ICLARM, Manila
- Papaconstantinou C, Caragitsou E (1989) Feeding interaction between two sympatric species *Pagrus pagrus* and *Phycis phycis* around Kastellorizo Island (Dodecanese, Greece). *Fish Res* 7:329–342
- Pauly D (1980) On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *J Cons Int Explor Mer* 39:175–192
- Pauly D, Christensen V, Sambilay V (1990) Some features of fish food consumption estimates used by ecosystem modellers. *ICES CM* 17:1–8
- Pauly D, Sambilay J, Opitz S (1993). Trophic models of aquatic ecosystems; estimates of relative food consumption by fish and invertebrate populations, required for modelling the Bolino Reef ecosystem, Philippines. ICLARM, Manila
- Pearson M, Gage JD (1984) Diets of deep sea brittle stars. *Mar Biol* 82:247–258
- Pinnegar JK, Polunin NVC (2000) Contributions of stable-isotope data to elucidating food webs of Mediterranean rocky littoral fishes. *Oecologia* 122:399–409
- Pinnegar J, Polunin N (2004) Predicting indirect effects of fishing in Mediterranean rocky littoral communities using a dynamic simulation model. *Ecol Model* 172:249–267

- Purcell J (1981) Dietary composition and diel feeding patterns of epipelagic siphonophores. *Mar Biol* 65:83–90
- Renones OA, Polunin NVC, Goni RJ (2002) Size related dietary shifts of *Epinephelus marginatus* in a western Mediterranean littoral ecosystem: an isotope and stomach content analysis. *J Fish Biol* 61:122–137
- Ribes M, Coma R, Gili J (1999) Natural diet and grazing rate of the temperate sponge *Dysidea avra* (Demospongiae, Dendroceratida) throughout an annual cycle. *Mar Ecol Prog Ser* 176:179–190
- Rosecchi E (1983) Regime alimentaire du pageot, *Pagellus erythrinus*, Linne 1758 (Pisces, Sparidae) dans le Golfe du Lion. *Cybium* 7:17–29
- Sala E (1997) The role of fishes in the organization of a Mediterranean sublittoral community: II. Epifaunal communities. *J Exp Mar Biol Ecol* 212:45–60
- Sala E, Ballesteros E (1997) Partitioning of space and food resources by three fish of the genus *Diplodus* (Sparidae) in a Mediterranean rocky infralittoral ecosystem. *Mar Ecol Prog Ser* 152:273–283
- Sanchez F, Olaso I (2004) Effects of fisheries on the Cantabrian Sea shelf ecosystem. *Ecol Model* 172:151–174
- Sanz A (1985) Contribucion al estudio de la biologia de *Uranoscopus scaber* Linnaeus, 1758 (Osteichthyes, Uranoscopidae) del Mediterraneo Occidental. *J Investig Pesquera* 49:35–46
- Schwinghamer P, Hargrave B, Peer D, Hawkins CM (1986) Partitioning of production and respiration among size groups of organisms in an intertidal benthic community. *Mar Ecol Prog Ser* 31:131–142
- Sullivan B, Garcia J, Klein-MacPhee G (1994) Prey selection by the scyphomedusan predator *Aurelia aurata*. *J Mar Biol* 121:335–341
- Tudela S, Palomera I (1997) Trophic ecology of the European anchovy *Engraulis encrasicolus* in the Catalan Sea (northwest Mediterranean). *Mar Ecol Prog Ser* 160:121–134
- Vannier J, Abe K, Ikuta K (1998) Feeding in myodocopid ostracods: a functional morphology and laboratory observations from videos. *Mar Biol* 132:391–408
- Verlaque M, Nedelec H (1983) Biologie de *Paracentrotus lividus* sur substrat rocheux en corse (Méditerranée France): alimentation des adultes. *Vie Milieu* 33:191–201
- Vetter EW (1996) Secondary production of a southern Californie *Nebalia* (Crustacea: Leptostraca). *Mar Ecol Prog Ser* 137:95–101
- Viejo RM, Arrontes J (1996) Interactions between mesograzers inhabiting *Fucus vesiculosus* in northern Spain. *J Exp Mar Biol Ecol* 162:97–111