
Transmission of viruses through shellfish: when specific ligands come into play

Françoise S Le Guyader^{1,*}, Robert L Atmar², Jacques Le Pendu³

¹ Laboratoire de Microbiologie, IFREMER, BP 21105, 44311 Nantes Cedex 03, France

² Department of Medicine, Baylor College of Medicine, 1 Baylor Plaza, MS BCM 280, Houston, TX 77030, USA

³ INSERM, U892, Université de Nantes, Institut de Recherche Thérapeutique, 8 quai Moncoussu BP 70721, 44007 Nante Cedex 1, France

*: Corresponding author : Françoise S. Le Guyader, email address : sleguyade@ifremer.fr ;
Soizick.Le.Guyader@ifremer.fr

Abstract :

Shellfish are known as vectors for human pathogens and despite regulation based on enteric bacteria they are still implicated in viral outbreaks. Among shellfish, oysters are the most common vector of contamination, and the pathogens most frequently involved in these outbreaks are noroviruses, responsible for acute gastroenteritis in humans. Analysis of shellfish-related outbreak data worldwide show an unexpected high proportion of NoV GI strains. Recent studies performed *in vitro*, *in vivo* and in the environment indicate that oysters are not just passive filters, but can selectively accumulate norovirus strains based on viral carbohydrate ligands shared with humans. These observations contribute to explain the GI bias observed in shellfish-related outbreaks compared to other outbreaks.

Highlights :

► Shellfish are known to transmit NoV to humans. ► High proportion of NoV GI is implicated in oyster-related outbreaks. ► Oysters accumulate selectively NoVs based on glycan ligands. ► First demonstration that a food may play a role in viral selection.

Although first described ~100 years ago [1], it has only recently become very clear that food plays an important role in virus transmission. In 2007, the CDC identified viruses as the causative agent of 46% of illnesses due to food consumption in outbreaks with an identifiable etiologic agent. Noroviruses (NoVs) were the most common cause, being responsible for 193 outbreaks, while *Salmonella*, the second leading cause, was responsible for 136 outbreaks [2]. Recent estimates from the CDC are that there are 9.4 million episodes of foodborne illness caused annually by 31 major pathogens in the United States, and NoVs are responsible for 58% of these illnesses. Beside NoVs, foodborne transmission has been documented for at least 10 viral families, but repeated implication has been demonstrated only for few families (Table 1) [3]. If viral zoonotic transmission (e.g. hepatitis E) is not considered, the two primary routes for food contamination are infected food-handlers and the production process (such as contact of the food with sewage-contaminated waters) [4, 5]. Several factors influence the transmission process, including the manner of contamination, binding or attachment of the virus to the food, survival and persistence of the virus on the food, the manner of food preparation (raw, cooked, peeled), and the susceptibility of the person eating the food to the contaminating virus [6]. The food itself also has an important role. For example, lettuce maintains a higher quantity of viable hepatitis A virus and for a longer period of time compared to fennel and carrots [7]. Recognition of foodborne illness also is influenced by public sensitivity and awareness of such illness, which can bias the tendency to report an illness. All but 3 of the 36 outbreak notifications involving viruses reported during an 11-year period (2000-2010) in the European Food Alert System for Food and Feed (RASFF) were due to NoVs. The other three were recent reports of HAV linked to dried tomatoes. Among the NoV foodborne outbreaks, 11 were associated with berries and 22 with oysters [4]. Although reporting bias may play a role in the predominance of outbreaks associated with berries and oysters, as they are known to be high-risk foods, these data also highlight the association between shellfish and viral gastroenteritis.

Norovirus

NoVs belong to the *Caliciviridae* family, a group of nonenveloped, icosahedral viruses with a single-stranded, positive sense RNA genome [8]. These viruses are highly diverse and are currently divided into 5 genogroups [9]. Genogroups I, II and IV contain human strains. Each genogroup is further subdivided into genotypes based upon analyses of the amino acid sequence of the major capsid protein, VP1. Other genotyping systems based upon shorter sequences [10] or analysis of the polymerase gene [11] have also been described. New strains and genogroups infecting animals also have been described [12]. NoV infection causes gastroenteritis that is characterized by vomiting and diarrhoea [13]. The prevalence of vomiting along with the short incubation period (1-2 days) and short clinical illness (1-3 days) have been used epidemiologically to identify probable outbreaks of NoV-associated gastroenteritis [14, 15]. The infectious dose 50% has been estimated to be as low as fewer than 20 virions [16]. NoVs bind to Histo-blood group antigens (HBGAs), phylogenetically highly conserved complex glycans present on many different cell types and proposed as an attachment factor necessary to initiate infection in people [17-19].

NoVs are the major cause of epidemic non bacterial gastroenteritis worldwide and have been identified as the cause of 73% to more than 95% of outbreaks [8]. These outbreaks involve all age groups in a wide variety of settings, with a large dominance of GII strains that can constitute up to 90% of clinical strains [5, 13]. Over the past 10 years, NoV sequence analyses of outbreak strains collected from around the world show that GII.4 viruses have accounted for ~70% of all human cases [20].

Shellfish-related NoV outbreaks

Shellfish are known to be a high-risk food for viral outbreaks but clear strain identification in shellfish is still often difficult. One of the first reports providing the sequence of a NoV strain described an outbreak in the US. A GI.4 strain was found in oyster samples, but the sequence was not identical to those detected in patients' stools [21]. At the same time in Japan, a mixture of GI and GII NoVs was detected both in stool and the related oyster samples but no sequencing was performed [22]. Since then, improvements in detection methods and the development and harmonization of molecular typing strategies have simplified data comparisons, allowing a compilation of outbreak reports that used comparable methods (Table 2).

One characteristic of shellfish-related outbreaks is their frequent association with multiple virus strains observed both in infected patients and in the involved shellfish. When a number of different virus strains are detected in patients, association of the infection with shellfish consumption can be difficult if only a few stools from an outbreak are collected. Thus, it is essential to collect as many stool samples as possible from affected individuals so that all strains that may be present can be identified. It is also important to rapidly identify the outbreak in order to trace the oyster production and to quickly collect the samples related to the outbreak. These data can be used with collected epidemiological data to fully understand the role played by shellfish in the outbreak.

Primers and probe sets specific for each NoV genogroup have been developed for detection by real time RT-PCR [23, 24]. However, genotyping remains a challenge, especially in shellfish where low viral concentrations are observed and in stools containing several different strains. In addition, a cocktail of primers is often required to detect the various NoV strains because of the diversity of these viruses [11, 25, 26].

Most outbreaks of shellfish-associated NoV disease are linked to oyster consumption, presumably because oysters are the most commonly consumed shellfish and they are usually consumed raw (although some outbreaks have been linked to cooked oysters) [27]. Overall, contamination by multiple NoV strains has been reported in 65% of reported outbreaks, with GI and GII NoVs detected, respectively, in 71% and 88% of stool samples and in 75% and 92% of shellfish samples. The frequency of each genogroup detected in shellfish-related outbreaks is clearly distinct from that of other NoV outbreaks. GI strains are more frequently encountered in shellfish-related outbreaks, and the GII.4 genotype is not as dominant (Figure 1). Among GI NoVs, the most frequently reported genotype is GI.1, followed by GI.4 and GI.2 (Figure 1). Among GII NoVs, the GII.4 genotype is the most frequently reported from both stool and shellfish samples. The GII.b variant was reported four times in patient's stool from oyster-related outbreaks, but confirmed in shellfish only once. Its frequent involvement in human to human outbreaks raises the possibility of another source of infection for these individuals involved in the alleged shellfish-related outbreaks [28, 29].

Some reports provide only stool analyses without shellfish data, such as the description of GI.1 and GII.3 strains implicated in an oyster-related outbreak reported from the UK [30]. In Japan GI NoVs alone were detected in four out of 11 outbreaks related to oyster consumption, with the remaining 7 outbreaks being associated with a mixture of GI and GII NoVs. In that study, GI.1 strain was detected in 3 of the 11 eleven outbreaks [31]. A previous study, also from Japan, reported the presence of a mixture of GI and GII NoVs in stools from 19 out of 21 oyster-outbreaks. In contrast, of 45 outbreaks not linked to shellfish consumption, all but 3 were due to GII NoVs, with both GI and GII strains being found in the remaining three [10].

Screening of shellfish not involved in outbreaks for the presence of NoVs has also been performed in several countries. Highly variable frequencies of contamination have been

reported. These studies have also observed a relatively higher frequency of GI NoV contamination than seen in community outbreaks (Table 3). Both studies that reported sequencing results identified GI.1 strains in the contaminated shellfish.

Norovirus bioaccumulation and persistence in oysters

On numerous occasions viral contamination in shellfish has persisted following measures, such as depuration or relaying, that have been used successfully to remove bacterial pathogens [32]. For example, in a laboratory-based study there was only a 7% decrease in the levels of bioaccumulated Norwalk virus compared to a 95% reduction in bacterial levels following 48 hours of depuration [33]. In another study, a GII.6 NoV persisted for at least 10 days under depuration conditions while a feline calicivirus was promptly eliminated [34]. A third study reported that, after a contaminating event in a French production area, the percentage of samples positive for GI and GII NoVs, respectively, were 59% and 70%. The prevalence decreased to 41% and 17%, respectively, after 4 weeks, suggesting a greater persistence in oyster tissues of GI NoVs compared to GII strains [35].

These observations led to the hypothesis that NoVs may bind specifically to oyster tissues through carbohydrates, as observed in humans, and that this binding may facilitate bioaccumulation and increase persistence in shellfish. Using immunohistochemistry, we demonstrated that NoV VLPs specifically bind to glycans of *Crassostrea gigas* oyster tissues, and that strain-specific variation in binding occurs. GI.1 NoVs bind to the midgut and digestive diverticula but not to gills or mantle, whereas GII.3 and GII.4 NoVs bind to all of these tissues. Human saliva from type A and O secretors, but not of type B secretors, inhibited binding of the GI.1 Norwalk VLPs, in accordance with the strain HBGA binding specificity. In addition, introduction of a mutation in the virus like particles (VLPs) glycan-binding site that abrogates glycan binding was sufficient to eliminate binding to oyster tissues, demonstrating specificity of the binding [36]. Binding was also inhibited by a lectin and anti-blood group A antibodies, indicating that the GI.1 NoV binds to *Crassostrea gigas* as well as *Crassostrea virginica* oyster tissues through an A-like antigen [37]. The A-like antigen is also implicated in the binding of GII.3 and GII.4 strains to oyster digestive tissues. Binding of these GII strains to the oyster's gills and mantle occurs through a sialic acid residue [38].

The influence of ligand expression on NoV binding to oyster tissues was first demonstrated using VLPs. GI.1 VLPs were very efficiently bioaccumulated by *Crassostrea gigas* oysters and were detected by immunohistochemistry even at a low level of exposure, whereas a mutant VLP that was unable to recognize the A-like antigen was only detected in oyster tissues at a thousand fold higher concentration [36]. These results were confirmed using a GI.1-positive stool that bioaccumulated very efficiently in a dose-dependant manner. When these experiments were performed at different times of the year, there was a clear seasonal impact on bioaccumulation efficiency that paralleled expression of the HBGA ligand in oyster digestive tissue [38]. The quantitative approach also showed that the GI.1 NoV directly accumulates in digestive tissues with negligible concentration in other tissues. Performing bioaccumulation using two GII NoV positive stools (one stool positive with a GII.4 and one with a GII.3 strain) led to very different results. These two strains bound to digestive tissues, gills and mantle with a similar pattern [39]. The GII.4 strain, as well as GII.4 VLPs, were bioaccumulated at very low levels, although they were found in a number of tissues as also reported by others [40, 41]. In contrast, the GII.3 strain was efficiently bioaccumulated, although less well than the GI.1 strain, with a transient retention in the gills likely due to binding to sialic acid [39]. In contrast to the findings with the GI.1 strain, no seasonal impact was observed in the bioaccumulation of the two GII NoVs or of the sialic acid containing ligand present in all tissues. Our interpretation of these data is that the GI.1 strain is efficiently accumulated and retained through an HBGA A-like ligand present in the gut. GII

strains are less well accumulated because of a sialic acid containing ligand expressed in all tissues that contributes to their retention in the gills and leads to their destruction (or elimination) by an unknown mechanism. The latter process would be more efficient in the case of a GII.4 than of a GII.3 strain.

Shellfish species may also impact bioaccumulation as demonstrated comparing two oysters species (*Crassostrea ariakensis* and *C. virginica*). The GI.1 strain was more efficiently concentrated by *C. ariakensis* and persisted for a longer time compared to *C. virginica* [42]. It will be interesting to compare the glycan ligand expression between these species.

Since many environmental conditions may interfere with oyster's filter capacity and consequently with contamination, a field study was conducted to determine if the above observations performed in laboratory conditions are valid in the environment. Thus, concentrations of GI and GII NoVs in waters collected during a year were compared to concentrations in oyster digestive tissues. As expected, much higher concentrations of GII NoVs than of GI were detected in waters. GI NoVs were concentrated to a greater degree than GII strains, with GI viruses requiring 30 viral RNA copies/L water to bioaccumulate 1 viral RNA copy/g oyster tissue compared to GII viruses that required ~1200 viral copies per liter of water to observe 1 viral copy per gram of oyster tissue. These data provide additional evidence for the specific selection and persistence of GI NoVs in oysters. This field study was conducted in an area with a large amount of cattle breeding. Bovine NoVs (GIII) were detected in 14% of water samples at high levels, but only one shellfish sample contained a GIII NoV strain [43]. The α Gal HBGA epitope, identified as the virus-specific glycan ligand in bovine tissues [44], was absent from oyster tissues, potentially explaining the poor bioaccumulation efficiency observed for GIII NoV strains.

Conclusion

These data suggest a selective transmission of NoV strains via oysters through specific binding to carbohydrate ligands. Ligands that facilitate bioaccumulation (the A-like antigen) or that contribute to the elimination of the virus (the sialic acid-containing ligand) may both influence NoV accumulation and survival in oysters (Figure 2). For a long time, oysters were believed to act as filters or ionic traps, passively concentrating particles. However, this is clearly not the case for NoVs, especially for NoV GI.1 that is more actively and efficiently concentrated than GII strains. The differential accumulation efficiency provides a possible explanation for the unexpectedly high proportion of GI strains associated with shellfish-related outbreaks.

This new concept demonstrating a special relationship between oysters and NoV should be explored for other enteric viruses, including Aichi virus and oysters, sapovirus and clams, and other foods such as NoV and berries or hepatitis A virus and tomatoes. Food trade may contribute to dispersal of a virus strain, as virus-contaminated imported shellfish have been responsible for outbreaks (Table 1) [66]. A better understanding of virus-food interactions may provide strategies to prevent contamination, to increase viral elimination, and thus to increase consumer safety.

Acknowledgments.

This work was supported in part by a grant (CIMATH) from the Region des Pays de la Loire, by DGAI (Direction Générale de l'Alimentation) and by a grant from the NIH (PO1 AI057788).

References

1. Jubb G, Glass MD, Oxon DPH: **Third outbreak of epidemic poliomyelitis at West Kirby.** *The Lancet* 1915,**9**: 67.
2. Boore A, Herman KM, Perez AS, Chen CC, Cole D J, Mahon BE, Griffin PM, Williams IT, Hall AJ: **Surveillance of foodborne disease outbreaks-United States 2007** *MMWR report* 2010, **59**: 973-979.
3. Duizer E, Koopmans M: **Emerging foodborne viral diseases.** in *Food-borne viruss: progress and challenges*. Edited by Koopmans MPG, Cliver DO, Bosch A. ASM Press; 2008: 117-145.
4. EFSA Panel on Biological Hazards: **Scientific opinion on An update on the present knowledge on the occurrence and control of foodborne viruses.** *EFSA Journal* 2011, **9**:1-96.
5. Koopmans M, **Progress in understanding norovirus epidemiology.** *Curr Op Inf Dis* 2008, **21**: 544-552.
6. Le Guyader FS, Atmar R L: **Binding and inactivation of viruses on and in food, with a focus on the role of the matrix.** in *Food-borne viruss: progress and challenges*. Edited by Koopmans MPG, Cliver DO, Bosch A. ASM Press; 2009: 189-208.
7. Croci L, De Medici D, Scalfarro C, Fiore A, Toti L: **The survival of hepatitis A virus in fresh produce.** *Int J Food Microbiol* 2002, **73**: 29-34.
8. Atmar RL: **Noroviruses: state of the art.** *Food Environ Virol* 2010, **2**: 117-126.
9. Zheng DP, Ando T, Fankhauser RL, Beard RS, Glass R, Monroe SS: **Norovirus classification and proposed strain nomenclature.** *Virology* 2006, **346**: 312-323.
10. Kageyama T, Shinohara M, Uchida K, Fukushi S, Hoshino FB, Kojima S, Takai R, Oka T, Takeda N, Katayama K: **Coexistence of multiple genotypes, including newly identified genotypes, in outbreaks of gastroenteritis due to Norovirus in Japan.** *J Clin Microbiol* 2004, **42**: 2988-2995.
- 11.* Kroneman A, Vennema H, Deforche K, Avoort HV D, Penaranda S: **An automated genotyping tool for enteroviruses and noroviruses.** *J Clin Virol* 2011,**51**: 121-125.
This paper describes a typing tool for NoVs that provides genotype information from sequence data in the polymerase or major capsid gene
12. Mesquita JR, Barclay L, Nascimento MSJ, Vinje J: **Novel norovirus in dogs with diarrhea.** *Emerg Inf Dis* 2010. **16**: 980-982.
- 13.** Glass RI, Parashar UD, Estes MK: **Norovirus gastroenteritis.** *N Engl J Med* 2009, **361**: 1776-1785.
An excellent review of noroviruses.
14. Turcios RN, Widdowson M-A, Sulka AC, Mead PS, Glass RI: **Reevaluation of epidemiological criteria for identifying outbreaks of acute gastroenteritis due to norovirus: United States, 1998-2000.** *Clin Inf Dis* 2006, **42**: 964-969.
15. Kaplan JE, Feldman R, Campbell DS, Lookabaugh C, Gary GW: **The frequency of a Norwalk-like pattern of illness in outbreaks of acute gastroenteritis.** *Am J Pub Health* 1982, **72**: 1329-1332.
16. Teunis PFM, Moe CL, Liu P, Miller SE, Lindesmith L, Baric RS, Le Pendu J, Calderon RL: **Norwalk virus: how infectious is it?** *J Med Virol* 2008, **80**: 1468-1476.
- 17.** Tan M, Jiang X: **Norovirus-host interaction: multi-selections by human histo-blood group antigens.** *Trends Microbiol* 2011, **19**: 382-388.
An excellent review of carbohydrate-NoVs interactions and of their importance in NoVs biology and evolution.
18. Le Pendu J, Ruvoen-Clouet N, Kindberg E, Svensson L: **Mendelian resistance to human norovirus infections.** *Sem Immunol* 2006, **18**: 375-386.
19. Donaldson EF, Lindesmith LC, Iobue AD, Baric RS: **Viral shape-shifting: norovirus evasion of the human immune system.** *Nature reviews* 2010, **8**: 231-241.

- 20.* Siebenga JJ, Lemey P, Kosakovsky-Pond SL, Rambaut A, Vennema H, Koopmans M: **Phylogenetic reconstruction reveals norovirus GII.4 epidemic expansions and their molecular determinants.** *PLoS Pathogens* 2010, **6**: e1000884.

A good description of the evolution of GII.4 NoVs.

21. Le Guyader F, Neill FH, Estes MK, Monroe SS, Ando T, Atmar RL: **Detection and analysis of a Small Round-Structured virus strain in oysters implicated in an outbreak of acute gastroenteritis.** *Appl Environ Microbiol* 1996, **62**: 4268-4272.
22. Sugieda M, Nakajima K, Nakajima S: **Outbreaks of Norwalk like virus associated gastroenteritis traced to shellfish : coexistence of two genotypes in one specimen.** *Epidemiol Infect* 1996, **116**: 339-346.
23. Trujillo AA, McCaustland KA, Zheng D-P, Hadley LA, Vaughn G, Adams SM, Ando T, Glass RI, Monroe SS: **Use of Taq-Man real-time reverse transcription PCR for rapid detection quantification and typing of norovirus.** *J Clin Microbiol* 2006, **44**: 1405-1412.
24. Kageyama T, Kojima S, Shinohara M, Uchida K, Fukushi S, Hoshino FB, Takeda N, Katayama K: **Broadly reactive and highly sensitive assay for Norwalk-like viruses based on real-time quantitative reverse transcription-PCR.** *J Clin Microbiol* 2003, **41**: 154-157.
25. Vinje J, Hamidjaja RA, Sobsey MD: **Development and application of a capsid VP1 (regionD) based reverse transcription PCR assay for genotyping of genogroup I and II noroviruses.** *J Virol Meth* 2004, **116**: 109-117.
26. Mattison K, Grudeski E, Auk B, Charest H, Drews SJ, Fritzingler A, Gregorious N, Hayward S, Houde A, Lee BE, Pang XL, Wong J, Booth TF, Vinje J: **Multicenter comparison of two norovirus ORF2-based genotyping protocols.** *J Clin Microbiol* 2009, **47**: 3927-3932.
27. Alfano-Sobsey E, Sweat D, Hall A, Breedlove F, Rodriguez R, Greene S, Pierce A, Sobsey M, Davies M, Ledford SL: **Norovirus outbreak associated with undercooked oysters and secondary household transmission.** *Epidemiol Infect* 2011, doi:10.1017/S0950268811000665.
- 28.* Phillips G, Tam CC, Rodrigues L C, Lopman B: **Risk factor for symptomatic and asymptomatic norovirus infection in the community.** *Epidemiol Infect* 2010, doi 10.1017/S0950268810002839.

This study identified risk factors for the development of NoV-associated illness, including contact with another case, foreign travel and consumption of shellfish.

29. Greig JD, Ravel A: **Analysis of foodborne outbreak data reported internationally for source attribution.** *Int J Food Microbiol* 2009, **130**: 77-87.
30. Gallimore C, Cheesbrough JS, Lamden K, Bingham C, Gray J: **Multiple norovirus genotypes characterised from an oyster-associated outbreak of gastroenteritis.** *Int J Food Microbiol* 2005, **103**: 323-330.
- 31.** Nakagawa-Okamoto R, Arita-Nishida T, Toda S, Kato H, Iwata H, Akiyama M, Nishio O, Kimura H, Noda M, Takeda N, Oka T: **Detection of multiple sapovirus genotypes and genogroups in oyster-associated outbreaks.** *Jpn J Inf Dis* 2009, **62**: 63-66.

This study demonstrated that persons involved in oyster-related outbreaks were infected with multiple enteric viruses, including NoVs and sapoviruses.

32. Richards GP, McLeod C, Le Guyader FS: **Processing strategies to inactivate enteric viruses in shellfish.** *Food Environ Virol* 2010, **2**: 183-193.
33. Schwab KJ, Neill FH, Estes MK, Metcalf TG, Atmar RL: **Distribution of Norwalk virus within shellfish following bioaccumulation and subsequent depuration by detection using RT-PCR.** *J Food Prot* 1998, **61**: 1674-1680.
34. Ueki Y, Shoji M, Suto A, Tanabe T, Okimura Y, Kikuchi Y, Saito N, Sano D, Omura T: **Persistence of caliciviruses in artificially contaminated oysters during depuration.** *Appl Environ Microbiol* 2007, **73**: 5698-5701.
35. Le Guyader FS, Le Saux J-C, Ambert-Balay K, Krol J, Serais O, Parnaudeau S, Giraudon H, Delmas G, Pommepuy M, Pothier P, Atmar RL: **Aichi virus, norovirus,**

astrovirus, enterovirus and rotavirus involved in clinical cases from a French oyster-related gastroenteritis outbreak. *J Clin Microbiol* 2008, **46: 4011-4017.**

36. Le Guyader FS, Loisy F, Atmar R L, Hutson AM, Estes MK, Ruvoen-Clouet N, Pommepuy M, Le Pendu J: **Norwalk virus specific binding to oyster digestive tissues.** *Emerg Inf Dis* 2006, **12**: 931-936.
37. Tian P, Bates A H, Jensen HM, Mandrell RE: **Norovirus binds to blood group A-like antigens in oyster gastrointestinal cells.** *Let Appl Microbiol* 2006, **43**: 645-651.
- 38.* Maalouf H, Zakhour M, Le Pendu J, Le Saux J-C, Atmar RL, Le Guyader FS: **Norovirus genogroup I and II ligands in oysters: tissue distribution and seasonal variations.** *Appl Environ Microbiol* 2010, **76**: 5621-5630.

This study showed that the glycan ligands found in shellfish led to differential binding of NoVs to oyster tissues, and the binding varied with the expression of the ligands throughout the year.

- 39.* Maalouf H, Schaeffer J, Parnaudeau S, Le Pendu J, Atmar R L, Crawford SE, Le Guyader F S: **Strain-dependent norovirus bioaccumulation in oysters.** *Appl Environ Microbiol* 2011, **77**: 3189-3196.

This study showed that binding of NoVs to shellfish tissues varied by virus genotype.

40. McLeod C, Hay B, Grant C, Greening G, Day D: **Localization of norovirus and poliovirus in Pacific oysters.** *J Appl Microbiol* 2009, **106**: 1220-1230.
41. Wang D, Wu Q, Yao L, Wei M, Kou X, Zhang J: **New target tissue for food-borne virus detection in oysters.** *Let Appl Microbiol* 2008, **47**: 405-409.
- 42.** Nappier SP, Graczyk TK, Schwab KJ: **Bioaccumulation, retention, and depuration of enteric viruses by *Crassostrea virginica* and *Crassostrea ariakensis* oysters.** *Appl Environ Microbiol* 2008, **74**: 6825-6831.

This paper compares the abilities of two oyster species to bioaccumulate, retain and depurate different enteric viruses in various salinities conditions. The study provides indirect evidence for pathogen selection by oysters.

43. Zakhour M, Maalouf H, diBartolo I, Haugarreau L, Le Guyader FS, Ruvoen-Clouet N, Le Saux J-C, Ruggeri FM, Pommepuy M, Le Pendu J: **Bovine norovirus ligand, environmental contamination and potential cross-species transmission via oyster.** *Appl Environ Microbiol* 2010 **76**: 6404-6411.
44. Zakhour M, Ruvoen-Clouet N, Charpilienne A, Langpap B, Poncet D, Peters T, Bovin N, Le Pendu J: **The gal epitope of the histo-blood group antigen family is a ligand for bovine norovirus newbury 2 expected to prevent cross-species transmission.** *PlosPathgens* 2009, **5**: e1000504.
45. Shieh YSC, Monroe SS, Fankhauser RL, Langlois GW, Burkhardt W, Baris RS: **Detection of Norwalk-like virus in shellfish implicated in illness.** *J Infect Dis* 2000, **181**: 360-366.
46. Le Guyader FS, Neill FH, Dubois E, Bon F, Loisy F, Kohli E, Pommepuy M, Atmar RL: **A semi-quantitative approach to estimate Norwalk-like virus contamination of oysters implicated in an outbreak.** *Int J Food Microbiol* 2003, **87**: 107-112.
47. Boxman ILA, Tilburg JJHC, te Loeke NAJM, Vennema H, Jonker K, de Boer E, Koopmans M: **Detection of noroviruses in shellfish in the Netherlands.** *Int J Food Microbiol* 2006, **108**: 391-396.
48. Prato R, Lopalco PL, Chironna M, Barbuti G, Germinario C, Quarto M: **Norovirus gastroenteritis general outbreak associated with raw shellfish consumption in South Italy.** *BMC Inf Dis* 2004, **4**: 1-6.
49. Le Guyader FS, Bon F, DeMedici D, Parnaudeau S, Bertone A, Crudeli S, Doyle A, Zidane M, Suffredini E, Kholi E, Maddalo F, Morini M, Gallay A, Pommepuy M, Pothier P, Ruggeri FM: **Detection of multiple noroviruses associated with an international gastroenteritis outbreak linked to oyster consumption.** *J Clin Microbiol* 2006, **44**: 3878-3882.
50. Webby RJ, Carville KS, Kirk MD, Greening G, Ratcliff RM, Creara SK, Dempsey K, Sarna M, Stafford R, Patel M, Hall G: **Internationally distributed frozen oyster**

meat causing multiple outbreaks of norovirus infection in Australia. *Clin Infect Dis* 2007, **44**: 1026-1031.

51. David ST, McIntyre L, MacDougall L, Kelly D, Liem S, Schallie K, McNabb A, Houde A, Mueller P, Ward P, Trottier Y-L, Brassard J: **An outbreak of norovirus caused by consumption of oysters from geographically disperse harvest sites, British Columbia, Canada, 2004.** *Foodborne Path Dis* 2007, **4**: 349-358.
52. Sasaki Y, Kai A, Hayashi Y, Shinkai T, Nogushi Y, Hasegawa M, Sadamasu K, Mori K, Tabej Y, Nagashima M, Morozumi S, Yamamoto T: **Multiple viral infections and genomic divergence among noroviruses during an outbreak of acute gastroenteritis.** *J Clin Microbiol* 2006, **44**: 790-797.
53. Simmons G, Garbutt C, Hewitt J, Greening G: **A New Zealand outbreak of norovirus gastroenteritis linked to the consumption of imported raw Korean oysters.** *New Zealand Med J* 2007, **120**: 1-7.
54. Nenonen NP, Hannoun C, Olsson M B, Bergstrom T: **Molecular analysis of an oyster-related norovirus outbreak.** *J Clin Virol* 2009, **45**: 105-108.
55. Le Guyader FS, Kroll J, Ambert-Balay K, Ruvoen-Clouet N, Desaubliaux B, Parnaudeau S, Le Saux J-C, Ponge A, Pothier P, Atmar RL, Le Pendu J: **Comprehensive analysis of a norovirus-associated gastroenteritis outbreak, from the environment to the consumer.** *J Clin Microbiol* 2010, **48**: 915-920.
56. Ilzuka S, Oka T, Tabara K, Omura T, Katayama K, Takeda N, Noda M: **Detection of sapoviruses and noroviruses in an outbreak of gastroenteritis linked genetically to shellfish.** *J Med Virol* 2010, **82**: 1247-1254.
- 57.* Nishida T, Nishio O, Kato M, Chuma T, Kato H, Iwata H, Kimura H: **Genotyping and quantitation of norovirus in oysters from two distinct sea areas in Japan.** *Microbiol Immunol* 2007, **51**: 177-184.

This study describes NoV quantification in oysters and molecular typing based on partial sequences of the capsid gene. These data are valuable as sequencing from shellfish samples is difficult and thus such reports are still uncommon.

58. Lowther JA, Henshilwood K, Lees DL: **Determination of norovirus contamination in oysters from two commercial harvesting areas over an extended period, using semiquantitative real-time reverse transcription PCR.** *J Food Prot* 2008, **71**: 1427-1433.
59. Lowther JA, Avant JM, Gizysztof K, Rangdale EE, Lees DN: **Comparison between quantitative real-time reverse transcription PCR results for norovirus in oysters and shelf-reported gastroenteritis illness in restaurant customers.** *J Food Prot* 2010, **73**: 305-311.
60. Gentry J, Vinje J, Guadagnoli D, Lipp E K: **Norovirus distribution within a estuarine environment.** *Appl Environ Microbiol* 2009, **75**: 5474-5480.
61. Le Guyader FS, Parnaudeau S, Schaeffer J, Bosch A, Loisy F, Pommepuy M, Atmar RL: **Detection and quantification of noroviruses in shellfish.** *Appl Environ Microbiol* 2009, **74**: 618-624.
62. Woods JW, Burkhardt III W: **Occurrence of norovirus and hepatitis A virus in US oysters.** *Food Environ Virol* 2010, **2**: 176-182.
63. Vilarino ML, Le Guyader FS, Polo D, Schaeffer J, Krol J, Romalde JL: **Assessment of human enteric viruses in cultured and wild bivalve molluscs.** *Inter Microbiol* 2009, **12**: 145-151.
64. Nenonen NP, Hannoun C, Horl P, Hernroth B, Bergstrom T: **Tracing of norovirus outbreak strains in mussels collected near sewage effluents.** *Appl Environ Microbiol* 2008, **74**: 2544-2549.
65. Suffredini E, Pepe T, Ventrone I, Croci L: **Norovirus detection in shellfish using two real-time RT-PCR methods.** *New Microbiol* 2011, **34**: 9-16.
- 66.** Polo D, Vilarino ML, Manso CF, Romalde JL: **Imported mollusks and dissemination of human enteric viruses.** *Emerg Inf Dis* 2010, **16**: 1036-1037.

This letter illustrates the risk of introduction of specific strains following the importation of shellfish.

Table 1: Viruses transmitted by food

Family	Genus (Name)	Capsid	Genome	Illness & incubation	Food transmission
<i>Adenoviridae</i>	<i>Adenovirus</i> (type 40-41)	Icosahedral, 65-80 nm	DNA, 35 kb	Gastroenteritis (moderate)	Rare
<i>Astroviridae</i>	<i>Astrovirus</i>	Icosahedral, 28-30 nm	ssRNA, 6.8 kb	Gastroenteritis (moderate)	Rare
<i>Caliciviridae</i>	<i>Norovirus</i> <i>Sapovirus</i>	Icosahedral, 27-32 nm	ssRNA, 7.6 kb	Gastroenteritis, 1-3 days	Frequent: shellfish, berries, food handler
<i>Coronaviridae</i>	<i>Coronavirus</i> (SARS)	Enveloped, 170 nm	ssRNA, 27-32 kb	Common cold, pneumonia, enteric disease	Suspected zoonotic, food handler
<i>Flaviviridae</i>	<i>Flavivirus</i> , Tick borne encephalitis virus (TBEV)	Enveloped, 40-60 nm	ssRNA, 11 kb	Fever, vomiting, fatigue, pain in the neck, back, encephalitis, 7-14 days	Rare: cow sheep goat milk
<i>Hepeviridae</i>	<i>Hepevirus</i> (Hepatitis E virus)	Icosahedral, 32-34 nm	ssRNA, 7.2 kb	Hepatitis, 3-8 weeks	Rare: pig meat, oyster
<i>Orthomyxviridae</i>	<i>Influenza A</i> (H5N1 virus)	Enveloped, 120-300 nm	Segmented ssRNA, 13.6 kb	Flu (fever, muscle pain),	Rare: bird meat (chicken, duck, geese..)
<i>Paramyxoviridae</i>	<i>Henipavirus</i> (Nipah virus)	Enveloped, 150-350 nm	ssRNA, 15 kb	Influenza -like illness, febrile encephalitis	Rare, food suspected in two outbreaks
<i>Picornaviridae</i>	<i>Kobuvirus</i> (Aichi virus)	Icosahedral, 27-32 nm	ssRNA, 8.2 kb	Gastroenteritis, 1-2 days	Uncommon: shellfish
	<i>Enterovirus</i>	Icosahedral, 20-30 nm	ssRNA, 7.2 kb	Diverse clinical syndromes, 3 to 10 days	Rare
	<i>Hepatovirus</i> (Hepatitis A virus)	Icosahedral, 27-32nm	ssRNA, 7.4 kb	Hepatitis, 2 to 6 weeks	Frequent: shellfish, vegetables, food handler
<i>Reoviridae</i>	<i>Rotavirus</i>	icosahedral, 3 layers, 70nm	dsRNA, 11 genes 3.3- 0.6 bp	Gastroenteritis, 1-3 days	Rare

Grey shading: viruses frequently transmitted via food.

Table 2: Norovirus genotypes reported from shellfish-related outbreaks.

Date	Country	Stool			Shellfish				Ref.
		# pos/ # analyzed	NoV GI genotype	NoV GII genotype	species	# pos/ # analyzed	NoV GI genotype	NoV GII genotype	
May 1998	US	1/2	nd	4	oyster	2/3	nd	4	[45]
March 2000	France	4/4	1, 2, 3	nd	oyster	2/2	1	nd	[46]
February 2001	Netherlands	8/9	1, 4	b, 7	oysters (France)	5/5	4	7	[47]
Dec 1998- Feb 2002	Japan	84/108 ^a	1-5, 7-9, 11-14	1, 3-12, 14, 16	oyster - no sample				[10]
March- Apr 2002	Italy	24	4	8, b	mussels	5/11	4	II.b	[48]
December 2002	France	29/53	4, 6	4, 8, b	oysters	3/3	4	4, 8	[49]
Nov 2003-Jan 2004	Australia	8/?	2, 4	5, 6, 7, 9, 12	oysters (Japan)	1/1	nd	4	[50]
January 2004	UK	10/11	1, 2	3, 4	oyster - no sample				[30]
Jan /March 2004	Canada	26/50	1, 2	3, 4, 5,	oysters	12/19	1	12	[51]
October 2005	Japan	18/37	nd	1, 4, 5, 6	oyster - no sample				[52]
June 2006	New Zealand	4/4	nd	3, 6, 12	oysters (Korea)	4/6	3	3, 6, 8, 12	[53]
February 2006	France	12/12	1, 2, 4	2, 4, 7, 17, b	oysters	9	1, 2, 4	4, 17	[35]
Jan 2002- March 2007	Japan	71 ^b	1-5, 8, 10, 13-15	3-6, 8, 12	oyster - no sample				[31]
January 2007	Sweden	1/1	1	nd	oysters	1/1	1	3	[54]
February 2008	France	4/5	nd	4	oysters	4/4	nd	4	[55]
June 2008	Japan	11/24	1	4, 8	clams	3	1	8	[56]
December 2009	US	3/6	nd	12	oysters- no sample				[27]

Nd: not detected,

Two manuscripts report data from 21 (a) and 11 (b) individual outbreaks

Table 3: Frequency of NoV GI and GII in shellfish contamination in non-outbreak samples from different countries.

Shellfish	Country	# samples ¹	# NoV positive	NoV GI		NoV GII		Ref.
				# positive	%	# positive	%	
Oysters	Japan	1512	75	26 ²	35	49	65	[57]
	UK	237	139	116	83	112	80	[58]
	UK	66	55	21	38	19	34	[59]
	US	10	5	5	100	0	-	[60]
	France	100	45	19	42	36	80	[61]
	US	381	15	4	27	11	73	[62]
Clams	Spain	41	14	1	7	13	93	[63]
Mussels	Sweden	40	23	19	83	4	17	[64]
	Italy	90	31	10	32	31	100	[65]
Mollusks	Spain ²	50	16	12	75	4	25	[66]

1: individual samples consisted of pools of 4-36 individual shellfish except for the study [64] in which individual mussels were assayed.

2: mollusks (clams, oysters or cockles) were imported from Morocco, Peru, Vietnam and South Korea.

Figure 1: Genotype frequency in stool and shellfish samples. NoV GI (red) and GII (blue) genotype detected in stool (plain bar) and shellfish (striped bar) samples.




Figure 2. Influence of oyster in the selection of NoV transmission. 1: Shedding in the environment of large amounts of GII NoVs (blue) and much lower amounts of GI strains (red) due to the overwhelming predominance of NoV GII in human outbreaks. Shedding of NoV GIII (green) in cattle is also shown. 2: Viruses present in seawater are ingested by oysters. GI NoVs particles are very rapidly directed to the gut, whereas GII particles are retained in mantle or gills possibly via a sialic acid containing ligand. GIII NoVs are probably randomly distributed. 3: NoV GI and GII are accumulated in the gut via an HBGA A-like ligand, most GII and GIII particles outside the gut are presumably destroyed. 4: Upon consumption of a NoV-contaminated oyster, infection caused by GI and GII strains occur with similar frequency because of the selective accumulation and retention of GI viral particles. GIII NoV transmission is unlikely to happen as few particles persist in oysters and humans do not express the glycan ligand.

