

Easterly denitrification signal and nitrogen fixation feedback documented in the western Pacific sediments

Guodong Jia^{1,2} and Zhiyang Li²

Received 14 October 2011; revised 14 November 2011; accepted 16 November 2011; published 24 December 2011.

[1] A sedimentary $\delta^{15}\text{N}$ record in the equatorial western Pacific (WP) shows glacial-interglacial variability from 6.2 to 11.2‰ during the last two climatic cycles, similar to the denitrification record in the eastern tropical Pacific (ETP). Contrastively, a record in the South China Sea (SCS) exhibits less changes from 4.4 to 6.4‰ and is quite alike previously published results in marginal seas in the WP. By ruling out several possible causes for the $\delta^{15}\text{N}$ variability, the $\delta^{15}\text{N}$ record in the equatorial WP is interpreted as the source nitrate $\delta^{15}\text{N}$ signals advected from the ETP. Comparison of several $\delta^{15}\text{N}$ records for the last 25 ka distributed in the WP brings out a pattern of northward decrease in $\delta^{15}\text{N}$ values and variability from the equator to off Mindano and then to marginal seas, supposed to be caused by the northward increase of local N_2 fixation. Therefore, the less glacial-interglacial changes in some $\delta^{15}\text{N}$ records in the WP could imply that the glacial decrease in subsurface $\delta^{15}\text{N}$ due to less denitrification in source waters from the ETP would have been isotopically compensated by a synchronous decrease in local N_2 fixation. **Citation:** Jia, G., and Z. Li (2011), Easterly denitrification signal and nitrogen fixation feedback documented in the western Pacific sediments, *Geophys. Res. Lett.*, 38, L24605, doi:10.1029/2011GL050021.

1. Introduction

[2] The balance between surface N_2 fixation and water column denitrification (WCD) can modulate the $\delta^{15}\text{N}$ signal of the upper ocean. N_2 fixation introduces atmospheric N_2 ($\delta^{15}\text{N} \sim 0$) into the oceanic fixed-N pool with little isotopic fractionation; whereas WCD strongly discriminates against ^{15}N , leaving higher $\delta^{15}\text{N}$ in seawater [Altabet, 2006]. Photic zone nitrate $\delta^{15}\text{N}$ signals can be documented in sedimentary organic matter (OM) produced in surface waters where N is completely utilized [Altabet, 2006]. $\delta^{15}\text{N}$ records from WCD zones in the eastern tropical Pacific (ETP) have provided a pattern of enhanced WCD during interglacials and reduced WCD during glacials [e.g., Ganeshram et al., 2000]. Moreover, water column studies reveal that the nitrate isotope signals of WCD can be transported out of the WCD zones, e.g., in the tropical central and western Pacific (WP), by ocean circulations [Kienast et al., 2008; Sigman et al., 2009]. And sedimentary $\delta^{15}\text{N}$ records similar to those from WCD zones have been observed in higher latitudes outside the WCD zones [Kienast et al., 2002; Martinez et al., 2006;

Galbraith et al., 2008], suggesting a wide impact of WCD variability during the late Quaternary. However, there is still a lack of firm sedimentary record in WP showing signals from the WCD zones. For example, most $\delta^{15}\text{N}$ records in the WP, e.g., the South China Sea (SCS), the Sulu Sea (SS), and the Okinawa Trough (OT), exhibit much less or even unchanged temporal variability (<2‰) [Kienast, 2000; Horikawa et al., 2006; Kao et al., 2008]. The invariant $\delta^{15}\text{N}$ records (~ 5 ‰) in the SCS have been even interpreted as representing a constant mean oceanic $\delta^{15}\text{N}$ during the late Quaternary [Kienast, 2000], and hence a minor impact of WCD zones.

[3] The WP is an oligotrophic region where N_2 fixation supplies significant N to support primary production. Were local N_2 fixation temporally coupled with easterly advected WCD signals, N_2 fixation could have produced the isotopically light N to offset the isotopic signals from WCD zones [Galbraith et al., 2004]. We expect that this presumption can reconcile those less variant $\delta^{15}\text{N}$ records in the WP. Additionally, because N_2 fixation in the WP is minimum near the equator relative to high-latitude sites [Hashihama et al., 2009; Shiozaki et al., 2010; Somes et al., 2010], the western equatorial Pacific (WEP) is the most likely region to document the easterly WCD signals from ETP. In this paper, $\delta^{15}\text{N}$ records from two sediment cores, one in the WEP and the other in the southern SCS, are reported. As will be shown, the two records exhibit quite different glacial-interglacial patterns, which could give us better understanding of the spatial and temporal changes of oceanic $\delta^{15}\text{N}$ in the WP.

2. Oceanographic Settings

[4] The WEP is a crossroad where several major ocean currents terminate and originate [Fine et al., 1994] (Figure 1). The westward North and South Equatorial Currents (NEC and SEC, respectively) bifurcate at $\sim 15^\circ$ latitudes, and the equatorward branches are the Mindanao Current (MC) in the north, the seasonal reverse New Guinea Coastal Current (NGCC) and the permanent New Guinea Coastal Undercurrent (NGCUC) in the south. Most part of NGCUC crosses the equator east of 135°E and feeds the eastward Equatorial Undercurrent (EUC) [Zenk et al., 2005]. Some part of MC feeds the North Equatorial Counter Current (NECC) and the other part is lost to the Indonesian throughflow (ITF). The cyclonic circulation composed of the NEC, MC, and NECC produces the cyclonic Mindanao Dome (MDD) centering at 7°N , 130°E . MDD is intensified during the Asian winter monsoon [Masumoto and Yamagata, 1991].

[5] The northern branch of NEC eventually forms the main part of the Kuroshio Current (KC), which is the main nutrient source to the marginal seas in the WP, e.g., the East China Sea, the SCS, and the SS. The SCS is connected to the WP via the Luzon Strait (sill depth ~ 2000 m) in the northeast,

¹State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, China.

²CAS Key Laboratory of Marginal Sea Geology, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou, China.




Figure 1. Study area and sediment cores. Red circles are study cores in this work, and stars are previously studied ones. Light gray areas are emerging lands during the Last Glacial Maximum. Arrows sketch the major currents presented in the text. MDN: Mindanao.

through which inflow of North Pacific Intermediate Water and KC brings nutrients from the eastern Pacific to the SCS and SS.

3. Samples and Experiments

[6] Core MD01-2386 (1°7.8'N, 129°47.6'E, 2816 water depth) was retrieved in 2001 at the northern shelf margin of the Halmahera Sea; and core MD01-2392 (9°51.1'N, 110°12.6'E, 1966 water depth) in the southwest SCS (Figure 1). The age model of MD86 is based on five radiocarbon dates over the past 18 ka [Jiang *et al.*, 2004] and $\delta^{18}\text{O}$ curve of planktonic foraminiferal *Globigerinoides ruber*. The age model of MD92 is based on aligning *G. ruber* $\delta^{18}\text{O}$ record with that in a nearby core MD97-2151, independently corroborated by ^{14}C dates and a few biostratigraphic, geomagnetic and tephrochronologic markers [Xie *et al.*, 2007] (Figure 2).

[7] Sedimentary $\delta^{15}\text{N}$ was analyzed on freeze-dried, homogenized bulk samples. Total organic carbon (TOC), total nitrogen (TN) and $\delta^{13}\text{C}_{\text{TOC}}$ were analyzed on decarbonate samples treated with 1M HCl. The measurements were performed on a CE EA1112 C/N/S analyzer coupled to a DELTA^{plus}XL mass spectrometer. The analytical precision is $\pm 0.3\text{‰}$ for $\delta^{15}\text{N}$ and $\pm 0.2\text{‰}$ for $\delta^{13}\text{C}$. Biogenic silica (BSi) in MD86 was extracted, according to Mortlock and Froelich [1989], with 2 M Na_2CO_3 at 85°C for 5 h and analyzed by molybdate blue spectrometry.

4. Results and Discussion

[8] Downcore $\delta^{15}\text{N}$ record in MD86 shows significant glacial-interglacial variability from 6.2 to 11.2‰ during the last two climate cycles. The lower values occur during cold periods and higher values during warm periods (Figure 2d). By comparison, $\delta^{15}\text{N}$ in MD92 exhibits less changed lower values between 4.4 and 6.4‰ (Figure 2b). TN ranges from 0.05 to 0.14‰ in MD86 and from 0.05 to 0.11‰ in MD92. TN correlates linearly with TOC in both cores ($r^2 = 0.91$ in

MD86 and 0.70 in MD92), with positive intercepts of TN of 0.016 and 0.003, respectively, suggesting minute but fixed inorganic N backgrounds in bulk N. Therefore, downcore $\delta^{15}\text{N}$ variations are predominantly controlled by organic N.

[9] Values of $\delta^{13}\text{C}_{\text{TOC}}$ (−22.1 to −19.8‰) and TOC/TN ratio (6.0 to 10.0) in both cores lie in the typical ranges for marine OM, and exhibit no trends or variability associated with climatic cycles. These values suggest minimum influence of terrigenous OM that can confound the bulk $\delta^{15}\text{N}$ signature as an indicator of phytoplankton-utilized $\delta^{15}\text{N}$. Thus, sedimentary $\delta^{15}\text{N}$ in this study is, indeed, reflective of the $\delta^{15}\text{N}$ of the nitrate-fueling phytoplankton production.

4.1. Possible Causes for the Downcore $\delta^{15}\text{N}$ Variability at Site MD86

[10] Previously published $\delta^{15}\text{N}$ records at sites 3bPC, 5bPC, and SC-8 in the WEP (Figure 1) have been interpreted as the changes in size and location of MDD, assuming incomplete nitrate utilization (INU) of the upwelled nitrate and progressive uptake of the nitrate pool along flow path [Nakatsuka *et al.*, 1995]. However, in the MDD the upwelled waters are present up to 75 m water depth, not reaching the surface [Udarbe-Walker and Villanoy, 2001]; and surface nitrate is undetectable even close to the dome center [Kienast *et al.*, 2008]. Low particulate $\delta^{15}\text{N}$ observed in the westernmost WEP (< 4‰) is related with isotopically light N input [Yoshikawa *et al.*, 2005], i.e., N_2 fixation [Kienast *et al.*, 2008], rather than with INU. Therefore, changes in MDD and INU are unlikely responsible for the $\delta^{15}\text{N}$ variability observed in MD86.

[11] Nowadays the well oxygenated water at site MD86 ($\text{DO} > 3 \text{ mg L}^{-1}$ through the water column) [Bassinot and Baltzer, 2002] prevents local denitrification. This scenario could have held true in the past because its oceanographic




Figure 2. Downcore data of foram $\delta^{18}\text{O}$ in (a) MD01-2392 and (c) MD01-2386, and $\delta^{15}\text{N}$ in (b) MD92 and (d) MD86, and (e) TOC and (f) BSi in MD86. OIS: Oxygen Isotope Stage.




Figure 3. Comparison of $\delta^{15}\text{N}$ records in this study with those reported for the last 25 ka. ODP893a is located in the Santa Barbara Basin. Other cores are illustrated in Figure 1.

setting is open with a sharp bathymetric feature (Figure 1), suggesting minimum influence of sea level changes on local circulations. The absence of local denitrification, together with the much reduced WCD zones during the glacials that could have emanated to the WP discussed below, would have diminished the importance of N_2 fixation for the glacial lower $\delta^{15}\text{N}$ values, because glacial $\delta^{15}\text{N}$ in MD86 is not only much higher than that from N_2 fixation, but also slightly higher than that in the WCD zones.

[12] A decrease in the vertical supply of ^{15}N -enriched nitrate to the photic zone due to deepening of thermocline has been proposed as a cause for the late Holocene $\delta^{15}\text{N}$ decline observed off Mindanao [Kienast *et al.*, 2008]. In MD86, tracers of paleoproductivity (TOC, BSi) exhibit higher values during the glacials (Figures 2e and 2f), implying shallower thermocline, which is also indicated by the larger foram $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ differences between the subsurface *Pulleniatina obliquiloculata* and surface *G. ruber* at the site [Jiang *et al.*, 2004]. This occurrence could have been caused by the intensified MDD and vertical mixing related with the enhanced East Asian winter monsoon during the glacials. The coincident lower $\delta^{15}\text{N}$ values and shallower thermocline in MD86 during the glacials, however, contradict the application of thermocline shift as the cause of $\delta^{15}\text{N}$ variations.

[13] Our $\delta^{15}\text{N}$ record in MD86 is grossly similar to those from WCD zones, as shown by $\delta^{15}\text{N}$ curve at ODP 893a in the Santa Barbara Basin [Emmer and Thunell, 2000] (Figure 3), implying a link between them. Presently, subsurface water in the WP is characterized by low N^* ($\text{N}^* = \text{N} - 16 \times \text{P} + 2.9 \text{ mmol}$) and higher than the oceanic “mean” values for nitrate $\delta^{15}\text{N}$ (6–8‰), indicating the long distance advection of nitrate from WCD zones [Kienast *et al.*, 2008]. Core-top $\delta^{15}\text{N}$ of 7.9‰ in MD86 is well within the range of subsurface $\delta^{15}\text{N}$, suggesting the fidelity of sedimentary $\delta^{15}\text{N}$ at the site to document subsurface $\delta^{15}\text{N}$. Therefore, it is conceivable that $\delta^{15}\text{N}$ signals of WCD zones during climatic cycles could have been transported to the WP via subsurface and documented faithfully in MD86. The higher values and

larger amplitude than WCD signals at the site likely result from progressive uptake of the nitrate pool, the production of sinking OM, and its transportation and regeneration back to nitrate along the flow path [Sigman *et al.*, 2009]. Together with the WCD signals found in the Pacific high latitudes [Kienast *et al.*, 2002; Martinez *et al.*, 2006; Galbraith *et al.*, 2008], the results in MD86 suggest basin-wide subsurface $\delta^{15}\text{N}$ changes in concert with the variations of WCD zones during climatic cycles, implying a great impact of changes in WCD zones [Ganeshram *et al.*, 2000].

4.2. Spatial Pattern of $\delta^{15}\text{N}$ Record in the WP

[14] Contrary to the results in MD86, the less changed $\delta^{15}\text{N}$ in MD92 repeats the published records distributed throughout the SCS [Kienast, 2000], except one in a drift deposit receiving allochthonous OM input [Kienast *et al.*, 2005]. In addition, sedimentary $\delta^{15}\text{N}$ from nearby seas, e.g., in cores MD04 in the OT and SUP8 in the SS (Figure 1), also show low values and small amplitudes (3.2–6.2‰ range) [Horikawa *et al.*, 2006; Kao *et al.*, 2008] (Figure 3).

[15] However, in the open WEP the few reported $\delta^{15}\text{N}$ records display heavy values and large amplitudes. For example, the records close to the MDD center (MD67 in Figure 3) show elevated values (6–7‰) during 10–7 ka, followed by a $>3\%$ decrease during 7–3 ka [Kienast *et al.*, 2008]; and the records at sites 3bPC, 5bPC, and SC-8 exhibit glacial-interglacial variability between 9–12‰ or 11–15‰ [Nakatsuka *et al.*, 1995]. The systematically heavy values at the latter three sites are attributed to the zonal transport of surface nitrate from ETP [Nakatsuka *et al.*, 1995]. Comparatively, our record at site MD86 shows lower values than those of Nakatsuka *et al.* [1995], which might be due to the westernmost location of the site, where the influence of lateral advection of surface nitrate from ETP is minimal [Yoshikawa *et al.*, 2006].

[16] We notice a spatial pattern of $\delta^{15}\text{N}$ records showing northward decrease in absolute values and variability for the last 25 ka from the equator to off Mindanao and then to the OT, the SCS and the SS (Figure 3). Interestingly, this pattern co-occurs with the spatial pattern of the northward increase in N_2 fixation in the WP, with a minimum near the equator and “hot spots” in the KC and marginal seas due to the northward increase of dust deposition from the Asian continent [Wong *et al.*, 2002; Hashihama *et al.*, 2009; Someset *et al.*, 2010; Shiozaki *et al.*, 2010]. In the absence of N_2 fixation, sedimentary $\delta^{15}\text{N}$ records in the WP would have paralleled those in WCD zones [Galbraith *et al.*, 2004], as shown in MD86 where N_2 fixation is minimum. Therefore, we speculate that the WCD signals would have been greatly masked by local N_2 fixation in off Mindanao, SCS, SS, and OT because of the opposing $\delta^{15}\text{N}$ signals of N_2 fixation and WCD. Thus the less changed $\delta^{15}\text{N}$ during climatic cycles in these WP records would imply that the glacial decrease in subsurface nitrate $\delta^{15}\text{N}$ due to less WCD zones would have been isotopically compensated by a decrease in local N_2 fixation. This occurrence is quite similar to what has been proposed in the Caribbean Sea [Ren *et al.*, 2009]. Nevertheless, more records associated with N_2 fixation in the WP are necessary to confirm this supposition.

[17] **Acknowledgments.** This work is granted by Chinese NSF (40876028) and the National Basic Research Program (2009CB421206). Zhimin Jian at Tongji Univ. provides foram $\delta^{18}\text{O}$ data in MD01-2386.

R. S. Robinson and an anonymous reviewer are appreciated for their valuable comments. This is contribution IS-1408 from GIGCAS.

[18] The Editor thanks Rebecca S. Robinson and an anonymous reviewer for their assistance in evaluating this paper.

References

- Altabet, M. A. (2006), Isotopic tracers of the marine nitrogen cycle: Present and past, in *Marine Organic Matter: Biomarkers, Isotopes and DNA*, edited by J. K. Volkman, pp. 251–293, Springer, Berlin, doi:10.1007/978-2-008.
- Bassinot, F., and A. Baltzer (2002), Scientific report of the WEPAMA cruise, MD122/IMAGES VII, 453 pp., Inst. Fr. pour la Rech. et la Technol. Polaires, Plouzane, France.
- Emmer, E., and R. C. Thunell (2000), Nitrogen isotope variations in Santa Barbara Basin sediments: Implications for denitrification in the eastern tropical North Pacific during the last 50,000 years, *Paleoceanography*, *15*, 377–387, doi:10.1029/1999PA000417.
- Fine, R. A., R. Lukas, F. M. Bingham, M. J. Warner, and R. H. Gammon (1994), The western equatorial Pacific: A water mass crossroads, *J. Geophys. Res.*, *99*, 25,063–25,080, doi:10.1029/94JC02277.
- Galbraith, E. D., M. Kienast, T. F. Pedersen, and S. E. Calvert (2004), Glacial-interglacial modulation of the marine nitrogen cycle by high-latitude O₂ supply to the global thermocline, *Paleoceanography*, *19*, PA4007, doi:10.1029/2003PA001000.
- Galbraith, E. D., M. Kienast, S. L. Jaccard, T. F. Pedersen, B. G. Brunelle, D. M. Sigman, and T. Kiefer (2008), Consistent relationship between global climate and surface nitrate utilization in the western subarctic Pacific throughout the last 500 ka, *Paleoceanography*, *23*, PA2212, doi:10.1029/2007PA001518.
- Ganeshram, R. S., T. F. Pedersen, S. E. Calvert, G. W. McNeill, and M. R. Fontugne (2000), Glacial-interglacial variability in denitrification in the world's oceans: Causes and consequences, *Paleoceanography*, *15*, 361–376, doi:10.1029/1999PA000422.
- Hashihama, F., K. Furuya, S. Kitajima, S. Takeda, T. Takemura, and J. Kanda (2009), Macro-scale exhaustion of surface phosphate by dinitrogen fixation in the western North Pacific, *Geophys. Res. Lett.*, *36*, L03610, doi:10.1029/2008GL036866.
- Horikawa, K., M. Minagawa, Y. Kato, M. Murayama, and S. Nagao (2006), N₂ fixation variability in the oligotrophic Sulu Sea, western equatorial Pacific region over the past 83 kyr, *J. Oceanogr.*, *62*, 427–439, doi:10.1007/s10872-006-0066-2.
- Jiang, L. B., Z. M. Jian, and X. R. Cheng (2004), Oxygen and carbon stable isotopic records of planktonic foraminifers from the western equatorial Pacific since the Last Glacial Maximum [in Chinese with English abstract], *Mar. Geol. Quat. Geol.*, *24*(2), 67–71.
- Kao, S. J., K. K. Liu, S. C. Hsu, Y. P. Chang, and M. H. Dai (2008), North Pacific-wide spreading of isotopically heavy nitrogen during the last deglaciation: Evidence from the western Pacific, *Biogeosciences*, *5*, 1641–1650, doi:10.5194/bg-5-1641-2008.
- Kienast, M. (2000), Unchanged nitrogen isotopic composition of organic matter in the South China Sea during the last climatic cycle: Global implications, *Paleoceanography*, *15*, 244–253, doi:10.1029/1999PA000407.
- Kienast, M., M. J. Higginson, G. Mollenhauer, T. I. Eglinton, M.-T. Chen, and S. E. Calvert (2005), On the sedimentological origin of down-core variations of bulk sedimentary nitrogen isotope ratios, *Paleoceanography*, *20*, PA2009, doi:10.1029/2004PA001081.
- Kienast, M., M. F. Lehmann, A. Timmermann, E. Galbraith, T. Bolliet, A. Holbourn, C. Normandeau, and C. Laj (2008), A mid-Holocene transition in the nitrogen dynamics of the western equatorial Pacific: Evidence of a deepening thermocline?, *Geophys. Res. Lett.*, *35*, L23610, doi:10.1029/2008GL035464.
- Kienast, S. S., S. E. Calvert, and T. F. Pedersen (2002), Nitrogen isotope and productivity variations along the northeast Pacific margin over the last 120 kyr: Surface and subsurface paleoceanography, *Paleoceanography*, *17*(4), 1055, doi:10.1029/2001PA000650.
- Martinez, P., F. Lamy, R. R. Robinson, L. Pichevin, and I. Billy (2006), Atypical $\delta^{15}\text{N}$ variations at the southern boundary of the east Pacific oxygen minimum zone over the last 50 ka, *Quat. Sci. Rev.*, *25*, 3017–3028, doi:10.1016/j.quascirev.2006.04.009.
- Masumoto, Y., and T. Yamagata (1991), Response of the western tropical Pacific to the Asian winter monsoon: The generation of the Mindanao Dome, *J. Phys. Oceanogr.*, *21*, 1386–1398, doi:10.1175/1520-0485(1991)021<1386:ROTWTP>2.0.CO;2.
- Mortlock, R. A., and P. N. Froelich (1989), A simple method for the rapid determination of biogenic opal in pelagic marine sediments, *Deep Sea Res., Part A*, *36*, 1415–1426, doi:10.1016/0198-0149(89)90092-7.
- Nakatsuka, T., N. Harada, E. Matsumoto, N. Handa, T. Oba, M. Ikehara, H. Matsuoka, and K. Kimoto (1995), Glacial-interglacial migration of an upwelling field in the western equatorial Pacific recorded by sediment $^{15}\text{N}/^{14}\text{N}$, *Geophys. Res. Lett.*, *22*, 2525–2528, doi:10.1029/95GL02544.
- Ren, H., D. M. Sigman, A. N. Meckler, B. Plessen, R. S. Robinson, Y. Rosenthal, and G. H. Haug (2009), Foraminiferal isotope evidence of reduced nitrogen fixation in the Ice Age Atlantic Ocean, *Science*, *323*, 244–248, doi:10.1126/science.1165787.
- Shiozaki, T., K. Furuya, T. Kodama, S. Kitajima, S. Takeda, T. Takemura, and J. Kanda (2010), New estimation of N₂ fixation in the western and central Pacific Ocean and its marginal seas, *Global Biogeochem. Cycles*, *24*, GB1015, doi:10.1029/2009GB003620.
- Sigman, D. M., P. J. DiFiore, M. P. Hain, C. Deutsch, and D. M. Karl (2009), Sinking organic matter spreads the nitrogen isotope signal of pelagic denitrification in the North Pacific, *Geophys. Res. Lett.*, *36*, L08605, doi:10.1029/2008GL035784.
- Somes, C. J., A. Schmittner, and M. A. Altabet (2010), Nitrogen isotope simulations show the importance of atmospheric iron deposition for nitrogen fixation across the Pacific Ocean, *Geophys. Res. Lett.*, *37*, L23605, doi:10.1029/2010GL044537.
- Udarbe-Walker, M. J. B., and C. L. Villanoy (2001), Structure of potential upwelling areas in the Philippines, *Deep Sea Res., Part I*, *48*, 1499–1518, doi:10.1016/S0967-0637(00)00100-X.
- Wong, G. T. F., S.-W. Chung, F.-K. Shiah, C.-C. Chen, L.-S. Wen, and K.-K. Liu (2002), Nitrate anomaly in the upper nutricline in the northern South China Sea: Evidence for nitrogen fixation, *Geophys. Res. Lett.*, *29*(23), 2097, doi:10.1029/2002GL015796.
- Xie, H. Q., G. D. Jia, P. A. Peng, and L. Shao (2007), Sea surface temperature variations in the southern South China Sea over the past 160 kyr, *Acta Oceanol. Sin.*, *46*, 49–55.
- Yoshikawa, C., T. Nakatsuka, and H. Kawahata (2005), Transition of low-salinity water in the Western Pacific Warm Pool recorded in the nitrogen isotopic ratios of settling particles, *Geophys. Res. Lett.*, *32*, L14615, doi:10.1029/2005GL023103.
- Yoshikawa, C., Y. Yamanaka, and T. Nakatsuka (2006), Nitrate-nitrogen isotopic patterns in surface waters of the western and central equatorial Pacific, *J. Oceanogr.*, *62*, 511–525, doi:10.1007/s10872-006-0072-4.
- Zenk, W., G. Siedler, A. Ishida, J. Holfört, Y. Kashino, Y. Kuroda, T. Miyama, and T. J. Müller (2005), Pathways and variability of the Antarctic Intermediate Water in the western equatorial Pacific Ocean, *Prog. Oceanogr.*, *67*, 245–281, doi:10.1016/j.pocean.2005.05.003.

G. Jia and Z. Li, CAS Key Laboratory of Marginal Sea Geology, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, 511 Kehua St., Guangzhou 510640, China. (jiagd@gig.ac.cn)