
Characterizing catches taken by different gears as a step towards evaluating fishing pressure on fish communities

Fauconnet Laurence ^{1,*}, Trenkel Verena ¹, Morandeau Gilles ², Caill-Milly Nathalie ²,
Rochet Marie-Joelle ¹

¹ IFREMER, Département Écologie et Modèles pour l'Halieutique, B.P. 21105, 44311 Nantes Cedex 03, France

² IFREMER, Laboratoire Ressources Halieutiques Aquitaine, UFR Côte Basque, 1 allée du Parc Montauray, 64600 Anglet, France

* Corresponding author : Laurence Fauconnet, email address : laurence.fauconnet@ifremer.fr

Abstract :

To implement an ecosystem approach to fisheries management, there is a need to characterize the total pressure exerted by fisheries at the community level. French onboard observer data were used to derive catch metrics and compare fishing distribution across community components between two sites in the Southern Bay of Biscay. Sample-based rarefaction curves were used to standardize metrics across different active and passive gears, and correct for sample size differences. Six metrics for species, length and functional catch composition were tested. Length and functional metrics were found the most relevant metrics to highlight differences in catches between gears, sites, and gear-site interactions. Significant differences were found between gears, mainly in mean length and proportion of piscivores. None of the gears had the most diverse catch across all metrics. Small differences were found between sites, mainly in length range and species richness.

Keywords : Catch diversity, Gear comparison, Multivariate analyses, Southern Bay of Biscay

1. Introduction

The ecosystem approach to fisheries (EAF) aims at maintaining ecosystem productivity for present and future generations by balancing multiple societal objectives (Garcia et al., 2003). One goal of fisheries management under an EAF is to keep fishing impacts on the ecosystem within acceptable limits, where the ecosystem structure and functioning is not threatened. The causal relationships formalized under the Driver - Pressure - State - Impact - Response (DPSIR) framework can help management. In particular, pressure can be adjusted by managers to keep the state of marine communities within, or move it towards, acceptable limits (Piet et al., 2006).

While methods exist and are commonly used to characterize fishing pressure on target populations, the limited knowledge on the biology and ecology and lack of fisheries data for most species imply that fishing pressures can not be characterized by fishing mortality or harvest rate at the community level (Piet et al., 2006). It has been hypothesized that both the total amount of fishing, and the way fishing pressure is distributed among ecosystem components determine fishing impacts on the community level (Garcia et al., 2012). Therefore, to develop an EAF, there is a need to characterize fishing pressure at the community level, i.e. the mortality caused by all fishing gears deployed in a given fishing ground on commercial and non-commercial species. Indicators are necessary tools to support this task as they provide information on the range and intensity of effort and mortality (Jennings, 2005; Piet et al., 2006). Two aspects of fishing pressure can be considered at the community level : fishing intensity and distribution across community components. In this study, we focus on how pressure is distributed across community components.

Pressures exerted on marine communities have long been considered only through the landings as declared by fishers and recorded on markets. However, landings represent only part of what is caught by fishers. Discards can make up a significant part of the catch, depending on the gear, area, season and species (Cornou et al., 2013; Hall et al., 2000), including for passive

gears (Morandeau et al., 2014). Most individuals when discarded are dead, and even if few studies have been undertaken on the survival of species that are released alive, a high level of mortality is assumed (Hall et al., 2000; Revill, 2012). Onboard observer programmes were developed to address the need
35 to identify and quantify the whole catch, distinguished between landings and discards (Alfaro-Shigueto et al., 2010; Attwood et al., 2011). By providing information on the amount, diversity and body size of the catch, onboard observer data are a valuable source to describe fisheries catches at the community level in its multiple dimensions such as species, length, and functional composition.

40 Onboard observer programmes further provide data on the characteristics and conditions of the fishing operations and on the main fishing metiers. Fishing mortality is likely to differ between gears (Piet et al., 2006). Therefore fishing pressure should be characterized by gear. Given the large diversity of gear characteristics, a gear can be defined at different levels of precision. The fishing
45 method or gear group as defined by the European Union (EU) Data Collection Framework (DCF; European Union, 2008), e.g. bottom trawls or mid-water trawls, subsequently called 'gear' was chosen in this study.

The catch composition reflects both the selective properties of the gear and how it is operated, and the available fish community. In order to study the effect
50 of the gear on the catch composition, we selected two sites in the Southern Bay of Biscay that are structurally and ecologically broadly similar, but differ in their exploitation though they are partly exploited by similar gears (see Section 2.1). Demersal and pelagic fisheries operate in both sites. In the most Southern site, the coastal area in ICES rectangle 16E8, the area located within 3 miles
55 from the coast and part of the 3-6 miles band, is prohibited to bottom and pelagic trawlers (figure 1 ; Sanchez et al., 2013). This site is consequently mostly harvested by passive fishing gears (figure 1). In the second site located further North, the coastal area in ICES rectangle 19E8, trawling is allowed due to exemptions limiting the application of the trawling ban inside the 3 miles limit
60 (Le Tixerant, 2006). This site is mostly exploited by active gears (figure 1). These study sites are well suited to test the relevance of metrics and highlight

differences between gears and sites.

Data from the French onboard observer programme were used to compare the catch for all species between gears and sites. However, the onboard ob-
65 server sampling plan was not established for this purpose, but for estimating discarded amounts per fishing métier. Therefore, the sample size was heterogeneous between gears and sites. Sample size is known to affect catch composition, especially its diversity (Magurran & McGill, 2011). Besides, different gears use different capture processes, mainly based on fish behaviour (Huse et al., 1999).
70 A fishing operation from a given gear is not directly comparable with a fishing operation from another gear, especially when comparing passive and active gears. Therefore, metrics needed to be standardized before they could be compared.

The objectives of this study were : i) to propose a method to standardize
75 and compare the distribution of catches across community components between passive and active fishing gears based on different sample size, and ii) to propose relevant metrics to characterize the catches that can highlight differences between gears.

2. Materials and methods

80 2.1. Study sites

The structure and sediments of the continental shelf in the Southern Bay of Biscay are homogeneous all along the coast of Aquitaine (Le Suavé et al., 2000). Sediments are mostly sandy, except in the deep environment of the Capbreton canyon, which is composed by a mix of rocks, coarse sediments
85 and mudflats. This geological formation favours the presence of species and life stages which live in deeper areas, such as mature hake (*Merluccius merluccius* ; Sanchez & Gil, 2000). The Southern Bay of Biscay is important for migratory species like meagre (*Argyrosomus regius*) in particular for feeding (Sourget & Biaï, 2009). The Southern Bay of Biscay is also the geographic
90 Northern limit of some species belonging to the Sparidae family (Quéro & Vayne,




Figure 1: Map of sampled fishing operations observed by gear onboard fishing vessels (2003-2012) in the Southern site and in the Northern site in the Southern Bay of Biscay (inset).

2005). Habitats and associated communities of the two sites are influenced by the plume of major rivers: Adour River for the Southern site, Gironde for the Northern site (figure 1). River plumes provide habitat for spawning and feeding for many species such as hake, monkfish (*Lophius piscatorius* and *L. budegassa*), sea bass (*Dicentrarchus labrax*), common sole (*Solea solea* ; Le Pape et al., 2003), turbot (*Scophthalmus maximus*), mackerels (*Scomber scombrus* and *S. colias* ; Borja et al., 2002), anchovy (*Engraulis encrasicolus* ; Borja et al., 1998). For those reasons, the two sites, situated 100 km apart, are considered ecologically broadly similar.

100 A major difference between the sites lies in the fact that, because of differences in access conditions for trawlers, they are harvested by different combinations of fishing gears. The Southern site is exploited by pelagic (purse seiners, baitboaters and pelagic trawlers) and demersal (gillnetters, longliners and pots) fisheries, most of which use passive gears. Pelagic species constitute the most abundant fish in the catch with mackerels, pilchard (*Sardina pilchardus*), horse mackerel (*Trachurus trachurus*), anchovy and tunas (*Thunnus alalunga* and *T.*

105

thynnus). Pelagic species are caught by a few boats on a small number of trips. The main demersal target species are hake, monkfishes, sea bass, common sole, turbot and Sparidae. About 70% of all boats operating in this area are smaller than 12 m length and perform a large number of short fishing trips (Leblond et al., 2010). The Northern site, where trawling is allowed, is characterized by pelagic and demersal fisheries targeting the same species along with cephalopods (*Loligo spp*, *Sepia officinalis*), which deploy mostly active gears. Pelagic species are mainly exploited by pelagic trawlers. Demersal species are exploited by bottom trawlers and gillnetters, the latter are the most important metiers in this area (92% of the activity in number of months ; Leblond et al., 2010). Eighty percent of the boats that fished at least once in this area in 2008 were longer than 12 m.

2.2. Onboard observer programme

Data from the French onboard observer programme contribute to the characterization of fishing pressure at the community level by providing information about the catch composition, as well as the characteristics and conditions of the fishing operation.

According to the sampling plan of the national programme, observers randomly select professional fishing boats to embark on, and once aboard randomly sample fishing operations (FOs). A FO includes all actions from the shooting to the hauling of the gear. The geographical positions, target species, gear and mesh size used, fishing time, and other information on the fisher's strategy and conditions of the FOs are recorded.

On sampled FOs, the whole catch is also recorded for both the landed and the discarded parts. All species of fish and commercial invertebrates are identified to the most precise level possible, ideally to the species level, counted, weighed (weight is sometimes calculated using the length/weight relationship) and measured.

The level of species identification can vary according to the observer's experience and/or the species. To circumvent this issue, 32 taxa which are difficult

to identify were grouped here at the family or genus level (*Allotheutis*, *Alosa*,
Argentina, *Arnoglossus*, *Callionymus*, *Hippocampus*, *Labrus*, *Loligo*, *Lophius*,
Microchirus, *Mullus*, *Mustelus*, *Octopus*, *Pagellus*, *Scomberomorus*, *Scorpaena*,
140 *Scyliorhinus*, *Sepia*, *Seriola*, *Serranus*, *Solea* except *S. solea*, *Sparus*, *Syng-*
nathus, *Torpedo*, *Trachurus*, *Trisopterus*, and families Carangidae, Gobiidae,
Mugilidae, Palinuridae, Rajidae except *Raja clavata* and *Leucoraja naevus*, and
Triglidae).

All observations from 2003 to 2012 were analysed together because of the
145 limited quantity of data available on study sites. Additional observations carried
out between July 2011 and December 2012 on coastal netters and longliners
deployed around the Capbreton canyon in the context of the regional programme
LOUPE (Observation of the habitat and associated communities in the context
of the fisheries of the Capbreton Canyon) were also used in this study. They
150 followed the same protocol.

The geographical distribution of observations collected on each site of our
case study is shown by gear in figure 1. Sampling effort was measured as the
number of observed fishing operations and vessels. The main group of target
species as declared beforehand by fishers, the mesh size mode and range informed
155 about the fishers' strategy per gear and site. For each gear-site combination,
the total number of individuals caught was calculated per FO and the mean
value over FOs was used as a proxy for the pressure intensity of an average
gear deployment. To determine which components were extracted from the
community by each gear, we characterized the catch in species, grouped by
160 combined taxon and main habitat (see Appendix), and in length.

2.3. Catch metrics

Since marine communities have different dimensions, three kinds of catch
metrics were calculated (table 1): (i) species-based metrics to provide informa-
tion on the number of species under fishing pressure and their relative abun-
165 dance, (ii) length-based metrics to provide information on the length of the
catch and its range, and (iii) functional metrics to provide information on the

170 trophic composition of the catch. Mean length and median length provided similar results, so just mean length is reported below. For the functional metrics, each species was classified as piscivore or non-piscivore, based on the main diet of adult individuals (see Appendix).

Table 1: List of catch metrics (n_i = number of individuals for i th species ; N = total number of individuals ; L_j = j th length class in cm ; N_j = number at j th length class ; $L^{0.95}$ = quantile 95% of length distribution ; $L^{0.05}$ = quantile 5% of length distribution ; W = weight ; $pisc$ = piscivores ; C = whole catch across all species = landings + discards) and method used for their estimation (RC = asymptote of rarefaction curve ; \tilde{x} = median of all resamples from sample sizes larger than 60% of the maximum sample size - see Section 2.3 and figure 2 ; \bar{x}_{logit} = mean on all logit-transformed replicates from sample sizes larger than 60% of the maximum sample size - see Section 2.3 and figure 2).

Type	Name	Code	Calculation	Description	Method
Species	Richness	S		Number of species	RC
	Evenness	$E_{1/D}$	$\frac{1/D}{S}$ with $D = \sum_i (\frac{n_i}{N})^2$	Variability in species abundance (Simpson)	RC
Length	Mean length	\bar{L}	$\frac{\sum L_j \times N_j}{\sum N_j}$	Length of individuals in the catch in cm	\tilde{x}
	Length range	ΔL	$L^{0.95} - L^{0.05}$	Interpercentile range 5-95% of length distribution in cm	\tilde{x}
Functional	Piscivore weight proportion	PWR	$\frac{\sum W_{pisc}}{\sum W_C}$	Proportion of piscivores in catch	\bar{x}_{logit}
	Piscivore number proportion	PNR	$\frac{\sum N_{pisc}}{\sum N_C}$	In comparison with PWR, says if piscivores are larger than other functional groups	\bar{x}_{logit}

The FOs of passive and active gears cannot be directly compared because they use different capture processes based on different fish behaviours (Huse et al., 1999). Also for a given gear, the number of observed FOs per site differed (ta-

ble 2). Sample size is known to affect the estimates of most selected metrics,
175 especially species richness (Gotelli & Colwell, 2001; Magurran & McGill, 2011).
Sample-based, i.e. FO-based, rarefaction curves were used to circumvent these
problems. Individuals from the same FO can not be considered as independant
entities. Indeed, they are likely to reflect spatial aggregation because the tar-
geting undertaken by professional fishers focuses on places with high resource
180 concentrations (Huse et al., 2000). FO-based, instead of individual-based, rar-
efaction curves were thus selected because they preserve the spatial structure
of the data (Gotelli & Colwell, 2001; Magurran & McGill, 2011). Rarefaction
curves tend to favor the species with the highest occurrences, contrary to meth-
ods that use the frequency of the rarest species to estimate the frequencies
185 of undetected species, that provide more accurate estimates of the total species
richness of a community (Chao et al., 2005). Since the purpose in this study is to
characterize not the whole community, but the part of the community, including
the number of species, available to each gear, rarefaction curves were preferred.
Because they are based on a re-sampling method, they could be used to estimate
190 uncertainty in metric estimates ascribable to sample size (Efron & Tibshirani,
1994). Sample-based rarefaction curves were used to standardize metric esti-
mates across gears to allow comparison between them (Gotelli & Colwell, 2001).
For this, a bootstrap was carried out by randomly sampling with replacement
FOs per combination of gear-site. The size of the re-sample varied from 1 to
195 20 FOs, or 1 to 50 FOs for gear-site combinations with more than 100 sampled
FOs. A thousand replicates were drawn for each re-sample size. Metrics were
calculated for each replicate.

With increasing sample size, the metric value converged toward an asymp-
totic value. The speed and shape of the convergence differed between metrics
200 types, so different methods were used for their estimation (summarized in table
1; figure 2). Species metrics kept varying with increasing sample size, though
they started to level off. They were estimated by determining the asymptote
value and its associated standard deviation after fitting a Michaelis-Menten
function: $S(n) = \frac{Smax \times n}{B+n}$, with S the value of the metric, $Smax$ the asymp-

205 totic value of the metric to be estimated, n the sample size and B the sample
size to get half the asymptotic value - 'half life' value (figure 2a). Length and
functional metrics converged rapidly with increasing sample size. Length met-
rics were estimated by calculating the median and standard deviation based on
all replicates from the largest sample sizes. A threshold of 60% of the maxi-
210 mum sample size was chosen. Therefore, estimates of length metrics were cal-
culated as the median of all replicates from 12 to 20 or from 30 to 50 FOs
(figure 2b). Metrics of functional composition, bounded between 0 and 100,
were first logit-transformed ($y = \log(\frac{x}{1-x})$) to make the values symmetrical and
unbounded (Jørgensen & Pedersen, 1998). They were calculated as the mean
215 value of all the logit-transformed replicates from 12 to 20 or 30 to 50 FOs (fig-
ure 2c), then converted back to the original scale. The standard deviation was
calculated on the original scale as $\tilde{\sigma}_X = \tilde{X}(1 - \tilde{X})\hat{\sigma}_Y$ with $\tilde{X} = \frac{\exp(\hat{Y})}{1+\exp(\hat{Y})}$, \hat{Y}
the logit-transformed values and $\hat{\sigma}_Y$ the standard deviation calculated for the
logit-transformed values (Jørgensen & Pedersen, 1998).




Figure 2: Methods for estimating metrics devoid of sample size effect and for standardizing catch characteristics between different passive and active gears, illustrated for gillnets in the Southern site : (a) rarefaction curve and asymptotic value of a fitted Michaelis-Menten model for species richness, (b) median of replicates resampled with sample size larger than 60% of the maximum sample size, for mean length, and (c) mean of logit-transformed replicates resampled with sample size larger than 60% of the maximum sample size, for piscivore weight proportion.

220 *2.4. Metric comparison among gears and sites*

Rarefaction curves were used to estimate catch metrics standardized for sampling effort. One estimate was obtained for each gear-site combination. However, the estimates were associated with an uncertainty due to the variability between FOs. In order to account for this variability, a parametric resampling
225 was carried out by randomly drawing 500 samples from a gaussian distribution $\mathcal{N}(Smax, \sigma^2)$ with $Smax$ the estimated value and σ its associated standard deviation, for each gear-site combination. They are referred to as 'resamples'.

A two-way analysis of variance (ANOVA) was undertaken for all resamples separately for each metric to highlight significant effects of gear and site. For this
230 ANOVA (A1) linear models with metric as response variable, and gear and site as factors, were fitted and the effects of the factors on each metric were tested by a Fisher's test. All gears were included. The percentage of variance explained by each factor was calculated as the sum of squares for each factor divided by the total sum of squares. Since the sample size was artificially increased by the
235 parametric resampling, only the relative importance of factors, and not the full results, was displayed and discussed.

Multivariate analyses were also undertaken using the resamples from the different gears and sites as individuals and catch metrics as variables. A principal component analysis (PCA1) was carried out to examine the relationships
240 between species, length and functional metrics. Groups of resampled replicates based on their center of gravity were visualised through starplots by site (North and South) and by gear.

For the comparison of gears deployed on both sites, a second ANOVA (A2) was performed including a gear-site interaction effect. The gear-site interaction
245 informs on whether the differences between gears varied between sites. Such differences neither ascribable to the gear nor to the site (i.e. the environment of the FOs) would likely to be due to the way the gear was rigged and/or deployed in the site and thus resulting from different fishers' strategies. Bottom and pelagic trawls as well as longlines were removed from this analysis as they were
250 only present on one of the sites.

A second PCA (PCA2), similar to PCA1 but using only on resamples from the gears deployed in both sites was performed to highlight differences between sites by a given gear.

3. Results

255 3.1. Description of observer data for study sites

The number of observed fishing operations differed markedly between gears and sites ranging from 14 to 175 (table 2). All gears were mainly targeting bottom-dwelling species, the main target species were flatfish, gadoids and other demersal fish (table 2). The main mesh size used in both sites for all gears, 260 except for bottom trawls, was 100 mm. However, a wide range of mesh sizes were used by each gear, with a wider range in the South than in the North.

Table 2: Number of sampled fishing operations (FO) and characteristics from each site and gear in number of sampled vessels, main group of target species (FLF = flatfish, DEF = demersal fish except gadoids, GAF = gadoid fish, MXF = diverse or unspecified fish) and mesh size mode with mesh size range in mm.

Site	Gear code	Gear name	No of FO	No of vessels	Targets	Mesh size [min-max] (mm)
South	TN	Trammel nets	112	11	FLF	100 [85-270]
	GN	Gillnets	175	14	DEF, GAF, MXF	100 [50-190]
	LL	Longlines	12	5	GAF, DEF	-
North	TN	Trammel nets	168	15	FLF	100 [70-100]
	GN	Gillnets	36	5	DEF	100 [84-110]
	PT	Pelagic trawls	14	7	DEF, GAF, MXF	100 [16-100]
	BT	Bottom trawls	62	16	FLF, MXF	70 [40-80]

The mean number of individuals per FO differed between fishing gears (figure 3 a). Bottom and pelagic trawls caught more individuals per FO than nets and

longlines, with trammel nets catching an intermediate number of individuals.
 265 In the South, the mean number of individuals caught was smaller than in the
 North, even for the same gear. The variability between FOs was large, especially
 for pelagic trawls.

Demersal fishes represented an important part of the catch for all gears
 (figure 3 b). Benthic species also constituted an important part of the catch,
 270 especially benthic fish and crustaceans for bottom trawls, trammel nets and,
 to a lesser extent, gillnets in the Southern site. Pelagic species were caught in
 smaller proportions, even by pelagic trawls for which pelagic fishes represented
 less than 20% of the total catch weight.

Differences in catch lengths were observed, with bottom trawls catching the
 275 smallest individuals and longlines catching the largest ones (figure 3 c). All
 gears caught individuals of lengths comprised between 20 and 40 cm.




Figure 3: Characterization of catches in the Southern Bay of Biscay (2003-2012): (a) fishing intensity in mean number of individuals caught by each gear and site with vertical bars = 95% confidence intervals, and (b-c) fishing distribution across community components : (b) composition of the catch in combined taxonomic group (CRU = crustacean; FIS = fish; MOL = mollusc) and main habitat (BEN = benthic; DEF = demersal; PEL = pelagic), and (c) length composition of the catch, for each gear (BT = bottom trawls ; GN = gillnets ; LL = longlines ; PT = pelagic trawls ; TN = trammel nets) in the Southern site (black) and the Northern site (grey).

3.2. Catch metrics

Species richness varied widely between gears (figure 4 a). Gillnets, trammel
nets and bottom trawls caught a high number of species, while pelagic trawls
280 caught a lower number of species and longlines caught less than 20 species. For
gears being used in both sites (trammel nets and gillnets), richness was generally
higher in the South than in the North, but this difference was not significant
due to a high inter-FO variability.

There was no significant difference in evenness among gears (figure 4 b). For
285 the nets, the catch was generally more even in the South than in the North,
though this difference was not significant.

Mean length varied between gears with longlines and gillnets catching the
largest individuals, while bottom trawls caught smaller ones (figure 4 c). Both
pelagic trawls and trammel nets caught intermediate lengths, with mean length
290 around 30 cm. Differences between the North and South differed among gears,
mean length was similar for trammel nets, but smaller in the South for gillnets.

The length range was similar between gillnets, trammel nets and pelagic
trawls, while it was larger for longlines and smaller for bottom trawls (figure 4
d). This means bottom trawls caught a narrower length range than other gears.
295 The range of lengths caught was wider in the South than in the North for both
gillnets and trammel nets.

The proportion of piscivores greatly differed between gears, with the propor-
tion in weight being consistently larger than the proportion in numbers (figure 4
e, f). Longlines, gillnets and pelagic trawls had a catch dominated by piscivores,
300 while piscivores were inferior to 40% in the catch of trammel nets and bottom
trawls (figure 4 e, f). The proportion of piscivores was similar between both
sites for trammel nets but it was significantly lower for gillnets in the South
than in the North.




Figure 4: Estimated catch metrics in the Southern (black) and Northern (grey) sites : (a) richness, (b) Simpson's evenness, (c) mean length in cm, (d) length range in cm, (e) piscivore weight proportion, and (f) piscivore number proportion, by gear (BT = bottom trawls ; GN = gillnets ; LL = longlines ; PT = pelagic trawls ; TN = trammel nets). Vertical bars represent two standard deviations.

3.3. Comparison of catch between study sites harvested by different combinations
of gears

305

Results from the ANOVA undertaken on all gears (A1) showed significant effects of both site and gear on each metric ($p(F) < 0.05$). Gear explained a higher variance proportion for all metrics (table 3), compared to site. The unexplained part remained high ($> 70\%$) for evenness. By contrast, for mean
310 length and piscivore proportions, gears explained the majority of variance ($> 85\%$), suggesting that those metrics are the most relevant to detect differences among gears.

Table 3: Percentage of variance explained by gear and site for each catch metric from the analysis of variance undertaken on all gears deployed in each site (A1).

Metric	Gear	Site	% residuals
Richness	56.6	12.3	31.1
Evenness	26.0	3.8	70.1
Mean length	89.0	2.9	8.1
Length range	69.7	18.0	12.3
Piscivore weight proportion	85.3	4.9	9.8
Piscivore number proportion	86.8	5.3	7.9

The first two axes of the principal component analysis on all gears (PCA1) explained 79% of the total variance, with the first axis explaining 61%. Along
315 this axis, a strong positive correlation was found between both piscivore proportions, mean length and length range (figure 5 a). The higher the proportion of piscivores, the larger the individuals in the catch and the wider the length range. Richness was negatively correlated with length and functional metrics. The richer the catch, the smaller the proportion of piscivores, the smaller the
320 individuals, but also the narrower the length range. Evenness was nearly independent from length and functional metrics but was negatively correlated with richness. This means that catches made of a small number of species were more

even.




Figure 5: Principal component analysis of catch metrics for all gears deployed in each site (PCA1) with 61% of the variance explained by the 1st axis, 18% explained by the 2nd axis. (a) Catch metrics loadings (Nb = number ; Wt = weight). Groups of resamples (b) by gear (BT = bottom trawls ; GN = gillnets ; LL = longlines ; PT = pelagic trawls ; TN = trammel nets) and (c) by site.

The grouping of resamples by gear clearly differentiated the gears, mainly
325 along the length and functional metrics (figure 5 b). Bottom trawls and trammel
nets caught smaller individuals, a narrower length range and a smaller propor-
tion of piscivore while longlines caught the largest individuals with a wide length
range and a large proportion of piscivores (figure 5 b). Pelagic trawls and gill-
nets were mainly differentiated by species metrics with gillnets having a richer
330 catch (figure 5 b).

Small differences between the Southern and the Northern sites were observed, the length of the individuals caught in the Southern site were slightly larger and more dominated by piscivores than the catch in the Northern site (figure 5 c).

3.4. Comparison of catch by similar gears between study sites

335 Results from the ANOVA undertaken only on gears deployed in both sites, trammel nets and gillnets (A2) showed a significant effect of both gear and site on all metrics, as well as an effect of the gear-site interaction on all metrics except evenness (table 4).

340 While gear explained most of the variance for mean length and piscivore proportions (> 66%), site explained a greater variance part for length range and richness (table 4). The gear-site interaction explained a small percentage of the variance for all metrics, except for mean length and piscivore weight proportion (> 16%). The unexplained part remained high (> 50%) for evenness and richness.

Table 4: Percentage of variance explained by gear, site and gear-site interaction for each catch metric from the analysis of variance undertaken only on common gears deployed in both sites, trammel nets and gillnets (A2).

Metric	Gear	Site	Gear-site interaction	% residuals
Richness	11.7	30.6	5.9	51.9
Evenness	16.1	8.7	0.0	75.1
Mean length	69.2	9.4	16.6	4.8
Length range	9.6	60.5	7.4	22.5
Piscivore weight proportion	66.9	11.9	17.2	4.1
Piscivore number proportion	73.9	12.5	7.8	5.9

345 Results from PCA2, undertaken only on gears deployed in both sites, clearly separates gear from site effects, which were mixed in PCA1 due to differences in gear deployment between the two sites. Grouping resamples by gear clearly

differentiated gillnets and trammel nets, mainly along the first PCA axis, with gillnets catching larger individuals and larger proportions of piscivores than trammel nets (figure 6 b). Differences between the Southern and the Northern sites were observed mainly along the second PCA axis (figure 6 c). The catch was of wider length range, richer and more even in the Southern site than in the Northern site.

With PCA2, we can distinguish the metrics that discriminate gears from sites, while the differences between gears and sites were confused in PCA1 with differences in gear deployment. Similarly to results from PCA1, a strong positive correlation between mean length and both piscivore ratios and a negative correlation with richness (figure 6 a) was found in PCA2. But contrary to PCA1, results from PCA2 showed no correlation between length range and mean length, and the correlation between richness and evenness was positive, meaning that the richer the catch, the more even.

4. Discussion

The first objective of this study was to propose a method to standardize and compare the distribution of catches across community components between passive and active fishing gears based on different sample sizes. Sample-based rarefaction curves were used to address this objective. This method, even if limited by the small number of observations for some gear-site combination, was found appropriate for this purpose because metrics converged for all combinations of gear-site.

The second objective of this study was to propose suitable metrics to characterize the distribution of catches across community components in species, length and function, that can highlight differences between gears. Length and functional metrics were found to be the most relevant metrics, in contrast to species metrics, for which a large part of the variance remained unexplained by gear, site, or gear-site interaction effect. Significant differences were found between gears, mainly in mean length and piscivore proportions. Smaller differ-




Figure 6: Principal component analysis of catch metrics for the gears deployed in both sites (PCA2) with 56% of the variance explained by the 1st axis, 24% explained by the 2nd axis. (a) Catch metrics loadings (Nb = number ; Wt = weight). Groups of resamples (b) by gear (GN = gillnets ; TN = trammel nets) and (c) by site.

ences were found between sites, mainly in length range and species richness. The effect of the gear-site interaction was also significant when comparing catches from the gears deployed on both sites only. This means that taking account of the fishers' strategy, in addition to the gear, would provide a better understanding of the catch composition.

4.1. Estimation method

To characterize total fishing pressure at the community level, we need to take account of the pressures exerted by all gears deployed in a given area as well as the state of the community. None of the gears is likely to sample the whole community (Fraser et al., 2007; Huse et al., 2000). Instead, each gear will provide a restricted view of the marine community. Therefore, the state of the community remains unknown and pressure can only be characterized indirectly through catch. Characterizing the catch taken from the community implies to simultaneously take account of both active and passive gears. For passive gears standardization of sampling effort is difficult. In this study, the number of sampled fishing operations differed between gears and sites. This is known to affect diversity, especially species richness (Magurran & McGill, 2011). Besides, fishing operations from different gears are not directly comparable, since gears have different capture processes which are based on different species behaviours. To overcome these problems, rarefaction curves were used to characterize the part of the community available to each gear. Since convergence with increasing sample size was observed on all gears and sites, we assume that rarefaction curves accordingly reached this objective and are appropriate to compare different passive and active gears.

The potential bias introduced by sample size when estimating biodiversity is widely recognized. However, it has not been as much examined for other metrics, such as individual length or functional composition of the catch. To avoid bias, we widened the use of sample-based rarefaction curves to length and functional metrics. Interestingly, convergence differed greatly between metrics. In contrast to species metrics which kept varying with increasing sample size, the length

and functional metrics had a fast convergence with increasing sample size. It means that, despite a wide range of mesh sizes used for each gear, selection in length and functional composition from the community by the different gears was rather homogeneous. However, this fast convergence was favored because
410 these metrics quantify an average property from a sample, while species metrics quantify unique observations, what partly explains why they converged more slowly.

However, a drawback of rarefaction curve is that it can be biased if sample
415 size is too small (Gotelli & Colwell, 2001). The limited number of observations for pelagic trawlers in the North and longliners in the South ($n < 20$) might therefore bias our results on the differences by gear and site, especially for species metrics. The small number of observations further limited this study, by compelling to analyze all years and quarters together, which might mask
420 seasonnal or annual effects. This data limitation has three main reasons. First, the number of trips to sample, as calculated for the sampling plan, is limited by technical and financial constraints. Second, the sampling plan is not fully realized owing to weather conditions and low acceptance of the programme by some professional fishers. Third, we had to focus on a small spatial scale, con-
425 strained by the need to find two sites broadly ecologically similar, differing in the way they were harvested, but partly exploited by similar gears. The sampling plan was not established for the purpose of this study but for estimating discarded amounts per fishing métier on all the French maritime areas. This study suggests that the amount of data necessary to characterize fishing pres-
430 sure on marine communities, especially on species, is larger than to estimate discards. If onboard observer programmes were redesigned so as to be used to characterize fishing pressure on marine communities, the sampling plan should include all gears deployed in all areas, with a minimum sample size for each.

4.2. *Catch metrics*

435 A wide variety of metrics could be used to characterize catches at the community level. The ones selected for this study were intended to be simple and easy

to interpret while describing the distribution of catches across different dimensions of the marine community : species, length, and function. Those metrics have been widely calculated from survey data as 'state' indicators to characterize fishing impacts on communities, including for the whole Bay of Biscay (e.g. 440 Rochet & Trenkel, 2005; Rochet et al., 2010; Shin et al., 2010). But we know of few studies estimating these metrics from catch data to characterize fishing pressure except Stergiou et al. (2002); Viana et al. (2013). However, because gears apply pressure on different components of the community (Piet et al., 2006), 445 the use of such metrics to characterize pressure per gear appears necessary.

Length metrics, particularly mean length, and functional diversity in both number and weight were the most relevant to detect differences between gears and sites. The gears catching the largest fish were also catching the most piscivores in both number and weight. These patterns tend to confirm that piscivores 450 were larger than other parts of the catch. Therefore, the classification of species into functional groups based on their diet as adults did not bias our results. The mean lengths of the catch found in this study, between 19 and 46 cm, were larger than estimates calculated with the EVHOE bottom trawl survey data from the same sites, between 11 and 17 cm (Fauconnet, unpublished data). The 455 same was true for piscivore proportions. Our estimates varied from 20 to 94% in weight and from 7 to 80% in number, while estimates from survey were much smaller : <4% in weight, and <1% in number (Fauconnet, unpublished data). This provides empirical evidence that commercial gears in general, but some more than others, select towards larger individuals and more piscivores than 460 survey gear.

Species metrics were the metrics for which the smallest part of the variance was explained by the gears and sites. Small differences in evenness across gears can be due to fisheries targeting and catching mainly benthic-demersal species, many of which are solitary species. A catch made of gregarious species 465 would have resulted less even than this. The small difference in evenness between fishing gears and sites can also result from the patchy sampling due to fishers targeting places where the resource is highly concentrated. Payne et al.

(2005) found that Simpson's measure of evenness gives precise and unbiased estimates regardless of the underlying patchiness of the distribution, but it does
470 not perform as well if the sampling is biased towards sites with greater number
of organisms, as it is the case here. Evenness turned out not to be appropriate
to detect differences between commercial gears. The grouping of species at
genus or family level undertaken to get a consistent level of species identification
across the whole dataset might have masked differences in richness and evenness
475 between gears or sites. This highlights the importance of the quality of species
identification in observer programmes, and that particular care should be taken,
especially if the data are to be used for studies at the community level.

4.3. Comparison between gears and sites

Our results suggest that catch composition in species, length and function
480 differ among gears. However, none of the gears was found to have the most, or
least, diverse catch across all metrics. Notably, our results are not consistent
with the prejudice that trawls have a more diverse catch than passive gears. In-
deed, we found that the gears catching the largest individuals were also catching
the widest length range. Bottom trawls caught smaller individuals than nets
485 and longlines as expected, but with a narrower length range. This restricted
length range can be explained by large fish being more efficient at escaping
the trawl than smaller individuals (Huse et al., 2000). With respect to species,
pelagic trawls had a slightly more even catch than gillnets but caught fewer
species. These findings highlight that the criteria passive - active gears is not
490 relevant to compare fishing pressures on marine communities. It further sug-
gests that the aspect of diversity to be considered needs to be clearly explicited
for management purpose, if fishing pressures are to be managed.

By comparing catches between two sites that are broadly similar ecologically,
and differ in the way they are harvested but are partly exploited by similar gears,
495 we were able to distinguish the effect of the gear from the effect of the site on the
catch composition. Considering all gears, small differences were found between
sites, with the catch from the South slightly larger-sized and with more piscivore

than the catch from the North. However, the wider length range, and greater richness when only comparing gears deployed in both sites, characterizing the Southern site also proved that the catch was more heterogeneous in length and richer in this site compared to the Northern site. Those differences can be due to a difference of harvesting between sites (i.e. pressure), in particular to the trawling ban in the Southern site, or to the sites themselves (i.e. state). Even if the sites were selected to be as similar as possible, some structural and biological differences may exist between the two sites. For instance, the presence of the Capbreton canyon in the Southern site is known to attract mature hake (Sanchez & Gil, 2000), while the Adour river plume plays an important role as a nursery for many species. The importance of both biological functions on this site might explain the wider length range observed. Studies on the annual variations would help in determining whether catches differed between sites because the underlying communities were different to start with (i.e. differences in state), or because they were harvested by different gear combinations (i.e. differences in pressure). This would further complete the knowledge of fishing pressure by taking account, besides their absolute values, of metric trajectories (Jennings, 2005). The Southern Bay of Biscay has been harvested for over a century, mostly by trawlers but also netters and longliners (Quero & Cendrero, 1996). The catch composition has considerably changed since the beginning of the harvest. Back then, some large demersal piscivores were highly frequent in the catch, while now they are no longer found (Quero & Cendrero, 1996). This historical record is qualitative though, and quantitative data have not been available for long enough to study in details the effects of fishing history on marine communities.

The gear-site interaction was tested to determine whether differences between sites varied between gears, which would likely result from differences in the way the gears were rigged and/or deployed in each site, i.e. from different fishers' strategies. The gear-site interaction which could only be tested for trammel nets and gillnets was significant for all metrics, particularly mean length and piscivore weight ratio, but not for evenness. The diversity of targets and

strategies for those gears implies that the gear level may be too general, con-
530 trary to the study by Stergiou et al. (2002) which found that the gear level,
independant of mesh size and season, was informative to characterize pressure
on species composition and diversity. Taking account of more information, such
as target species and/or mesh size, could enable to more accurately characterize
535 fishing pressure (Cornou et al., 2013; Dubé et al., 2012). Based on the catch
composition, two main groups of gears related to the type of target species can
be distinguished. The gears targeting benthic species, bottom trawls and tram-
mel nets, tended to catch more species with a smaller mean length, a narrower
length range and a smaller proportion of piscivores than the gears used to target
gadoids or other demersal fish, i.e. gillnets, longlines and pelagic trawls. The
540 diversity of target species can also play an important role. While longlines of-
ten target a small number of species, bottom trawls are usually less specialized
(Sanchez et al., 2013). Better understanding how targeting affects the catch
composition might help assessing fishing pressures on marine communities.

4.4. Conclusions

545 This study aimed at characterizing fishing pressure at the community level
based on onboard observer data. Total pressure on marine communities was
estimated here as total catch, i.e. including discards. The latter can be a
significant part of the catch and has been shown by Viana et al. (2013) to be
important to estimate fisheries' ecological footprint. This study highlights the
550 importance in observer programmes to consistently and exhaustively sample the
total catch, i.e. both landings and discards of all species. However, even total
catch is likely to underestimate pressure exerted by fisheries on marine commu-
nities. Indeed, it does not account for the mortality on individuals that were not
brought onboard, for example those that escaped the gear with potential severe
555 injuries (Ingólfsson & Jørgensen, 2006). Pressure on habitats and on benthic
organisms could not be examined with the available data either. Indeed, for
practical reasons, the protocol cannot include data collection on more species,
and 'community' had to be restricted in this study to fish and commercial in-

vertebrates. Some community components are likely to undergo more pressure
560 than others, since targets and many catch components are similar between the
different gears deployed in both sites. How the different gears compete with
each other remains to be studied to highlight community components that are
under higher pressure. In this study, we focused on how catches were distributed
565 across community components. However pressure exerted on highly abundant
or productive species will likely have different impacts than pressure exerted
on rare or vulnerable species. A complementary approach would be to quan-
tify the intensity of pressure undergone by each component, and their ability
to sustain it, for example with methods such as the Productivity Susceptibility
Analysis (Smith et al., 2007). This study suggests that fishing pressure indeed
570 varies among gears and among sites exploited by different gear combinations.
The impacts of this contrasted pressure on marine communities remain to be
studied.

Acknowledgements

We thank two anonymous referees for their comments and suggestions. Lau-
575 rence Fauconnet received a PhD fellowship from the Région Pays de la Loire.
Support provided by the French Department for Ecology, Sustainable develop-
ment and Energy (Direction des Pêches Maritime et de l'Aquaculture), and the
Pew Charitable Trusts. The opinions expressed are those of the authors and
do not necessarily reflect the views of the funding organizations. The authors
580 would like to thank Olivier Gaudou and Joël Diméet for their valuable contribu-
tion to the French observer programme and for their help in preparing the data
for the analyses. We thank Jocelyne Martin and Pascal Laffargue for help in
grouping the species. We also thank all fishers and observers who participated
in the French observer programme.

585 **References**

- Alfaro-Shigueto, J., Mangel, J. C., Pajuelo, M., Dutton, P. H., Seminoff, J. A., & Godley, B. J. (2010). Where small can have a large impact: Structure and characterization of small-scale fisheries in Peru. *Fish. Res.*, *106*, 8–17. doi:10.1016/j.fishres.2010.06.004.
- 590 Attwood, C. G., Petersen, S. L., & Kerwath, S. E. (2011). Bycatch in South Africa's inshore trawl fishery as determined from observer records. *ICES J. Mar. Sci.*, *68*, 2163–2174. doi:10.1093/icesjms/fsr162.
- Borja, A., Uriarte, A., & Egaña, J. (2002). Environmental factors and recruitment of mackerel, *Scomber scombrus* L. 1758 , along the north-east Atlantic
595 coasts of Europe. *Fish. Oceanogr.*, *11*, 116–127.
- Borja, A., Uriarte, A., Egaña, J., Motos, L., & Valencia, V. (1998). Relationships between anchovy (*Engraulis encrasicolus*) recruitment and environment in the Bay of Biscay (1967-1996). *Fish. Oceanogr.*, *7*, 375–380.
- Chao, A., Chazdon, R. L., Colwell, R. K., & Shen, T.-J. (2005). A new statistical
600 approach for assessing similarity of species composition with incidence and abundance data: A new statistical approach for assessing similarity. *Ecol. Lett.*, *8*, 148–159. doi:10.1111/j.1461-0248.2004.00707.x.
- Cornou, A.-S., Diméet, J., Tétard, A., Gaudou, O., Dubé, B., Fauconnet, L., & Rochet, M.-J. (2013). *Observations à bord des navires de pêche professionnelle - Bilan de l'échantillonnage 2012*. Technical Report Ifremer. URL:
605 <http://archimer.ifremer.fr/doc/00167/27787/25978.pdf>.
- Dubé, B., Diméet, J., Rochet, M.-J., Tétard, A., Gaudou, O., Messannot, C., Fauconnet, L., Morizur, Y., Biseau, A., & Salaun, M. (2012). *Observations à bord des navires de pêche professionnelle - Bilan de l'échantillonnage 2011*. RBE/STH/LTBH 2012-008 Ifremer. URL:
610 <http://archimer.ifremer.fr/doc/00109/21976/19586.pdf>.

- Efron, B., & Tibshirani, R. (1994). *An Introduction to the Bootstrap*. Number 57 in Monographs on Statistics & Applied Probability (Chapman and Hall/CRC ed.).
- 615 European Union (2008). Commission Decision of 6 November 2008 adopting a multiannual Community programme pursuant to Council Regulation (EC) No 199/2008 establishing a Community framework for the collection, management and use of data in the fisheries sector and support for scientific advice regarding the common fisheries policy.
- 620 Fraser, H. M., Greenstreet, S. P., & Piet, G. J. (2007). Taking account of catchability in groundfish survey trawls: implications for estimating demersal fish biomass. *ICES J. Mar. Sci.*, *64*, 1800–1819.
- Garcia, S. M., Kolding, J., Rice, J., Rochet, M.-J., Zhou, S., Arimoto, T., Beyer, J. E., Borges, L., Bundy, A., Dunn, D., Fulton, E. A., Hall, M., Heino, M., Law, R., Makino, M., Rijnsdorp, A. D., Simard, F., & Smith, A. D. M. 625 (2012). Reconsidering the consequences of selective fisheries. *Science*, *335*, 1045–1047. doi:10.1126/science.1214594.
- Garcia, S. M., Zerbi, A., Aliaume, C., Do Chi, T., & Lasserre, G. (2003). *The ecosystem approach to fisheries: issues, terminology, principles, institutional foundations, implementation and outlook*. Number 443 in FAO Fisheries Technical Paper. Rome: Food & Agriculture Organisation.
- 630 Gotelli, N. J., & Colwell, R. K. (2001). Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecol. Lett.*, *4*, 379–391.
- 635 Hall, M. A., Alverson, D. L., & Metuzals, K. I. (2000). By-catch: Problems and solutions. *Mar. Pollut. Bull.*, *41*, 204–219.
- Huse, I., C Gundersen, A., & H Nedreaas, K. (1999). Relative selectivity of Greenland halibut (*Reinhardtius hippoglossoides*, Walbaum) by trawls, longlines and gillnets. *Fish. Res.*, *44*, 75–93.

- 640 Huse, I., Løkkeborg, S., & Soldal, A. V. (2000). Relative selectivity in trawl,
longline and gillnet fisheries for cod and haddock. *ICES J. Mar. Sci.*, *57*,
1271–1282.
- Ingólfsson, O. A., & Jørgensen, T. (2006). Escapement of gadoid fish beneath
a commercial bottom trawl: relevance to the overall trawl selectivity. *Fish.*
645 *Res.*, *79*, 303–312.
- Jennings, S. (2005). Indicators to support an ecosystem approach to fisheries.
Fish Fish., *6*, 212–232.
- Jørgensen, E., & Pedersen, A. R. (1998). *How to obtain those nasty standard er-*
rors from transformed data - and why they should not be used: Internal report
650 *volume 7*. Aarhus Universitet, Det Jordbrugsvidenskabelige Fakultet. URL:
<http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.47.9023&rep=rep1&type=pdf>.
- Le Pape, O., Chauvet, F., Désaunay, Y., & Guérault, D. (2003). Rela-
tionship between interannual variations of the river plume and the ex-
tent of nursery grounds for the common sole (*Solea solea*, L.) in Vi-
655 laine Bay. Effects on recruitment variability. *J. Sea Res.*, *50*, 177–185.
doi:10.1016/S1385-1101(03)00061-3.
- Le Suavé, R., Bourillet, J.-F., & Coutelle, A. (2000). *La marge nord*
du golfe de Gascogne: Connaissances générales et apport des nouvelles
synthèses de données multifaisceaux. Technical Report Ifremer. URL:
660 <http://archimer.ifremer.fr/doc/00071/18199/>.
- Le Tixerant, M. (2006). *Cartographie de la réglementation des*
pêches professionnelles-Régions Aquitaine et Poitou Charente.
Technical Report Comités Régionaux des Pêches et des Élevages
Marins de la région Aquitaine et de la région Poitou-Charentes,
665 Association du Grand Littoral Atlantique (AGLIA). URL:
<http://www.aglia.org/etudes.asp?Id=70&Page=14#Bottom>.

Leblond, E., Daures, F., Berthou, P., Merrien, C., Pitel-Roudaut, M., Brigaudeau, C., Demaneche, S., Jezequel, M., L'Equipe D'Exploitation Du Systeme Harmonie Du Sih, Bodere, E., & Le Blond, S. (2010). Synthèse des flottilles de pêche 2008 - Flotte Mer du Nord - Manche - Atlantique - Méditerranée. URL: <http://archimer.ifremer.fr/doc/00003/11456/8013.pdf>.

Magurran, A. E., & McGill, B. J. (2011). *Biological Diversity - Frontiers in Measurement and Assessment*. New York: Oxford University Press.

Morandeau, G., Macher, C., Sanchez, F., Bru, N., Fauconnet, L., & Caill-Milly, N. (2014). Why do fishermen discard? Distribution and quantification of the causes of discards in the Southern Bay of Biscay passive gear fisheries. *Mar. Policy*, *48*, 30–38. doi:10.1016/j.marpol.2014.02.022.

Payne, L. X., Schindler, D. E., Parrish, J. K., & Temple, S. A. (2005). Quantifying spatial pattern with evenness indices. *Ecol. Appl.*, *15*, 507–520.

Piet, G., Quirijns, J., Robinson, L., & Greenstreet, S. (2006). Potential pressure indicators for fishing, and their data requirements. *ICES J. Mar. Sci.*, *64*, 110 – 121.

Quero, J.-C., & Cendrero, O. (1996). Incidence de la pêche sur la biodiversité ichthyologique marine: le bassin d'Arcachon et le plateau continental sud Gascogne. *Cybium*, *20*, 323–356.

Quéro, J.-C., & Vayne, J.-J. (2005). *Les Poissons de mer des pêches françaises*. Delachaux Et Niestlé.

Revill, A. (2012). *Survival of discarded fish - A rapid review of studies on discard survival rates*. Request for services commitment no. S12.615631. European Commission, Directorate - General for maritime affairs and fisheries, Policy development and coordination, Brussels, MAREA2. URL: <http://nsrc.org/wp-content/uploads/2012/08/EU-discard-survival-short-study-version-001.p>

- 695 Rochet, M.-J., & Trenkel, V. M. (2005). Factors for the variability of dis-
cards: assumptions and field evidence. *Can. J. Fish. Aquat. Sci.*, *62*, 224–235.
doi:10.1139/f04-185.
- Rochet, M.-J., Trenkel, V. M., Carpentier, A., Coppin, F., De Sola, L. G.,
Léauté, J.-P., Mahé, J.-C., Maiorano, P., Mannini, A., Murenu, M., Piet,
G., Politou, C.-Y., Reale, B., Spedicato, M.-T., Tserpes, G., & Bertrand,
700 J. A. (2010). Do changes in environmental and fishing pressures impact ma-
rine communities? An empirical assessment. *J. Appl. Ecol.*, *47*, 741–750.
doi:10.1111/j.1365-2664.2010.01841.x.
- Sanchez, F., & Gil, G. (2000). Hydrographic mesoscale structures and Poleward
Current as a determinant of hake (*Merluccius merluccius*) recruitment in
705 southern Bay of Biscay. *ICES J. Mar. Sci.*, *57*, 152–170.
- Sanchez, F., Morandeau, G., Bru, N., & Lissardy, M. (2013). A re-
stricted fishing area as a tool for fisheries management: Example of the
Capbreton canyon, southern Bay of Biscay. *Mar. Policy*, *42*, 180–189.
doi:10.1016/j.marpol.2013.02.009.
- 710 Shin, Y.-J., Bundy, A., Shannon, L. J., Simier, M., Coll, M., Fulton, E. A.,
Link, J. S., Jouffre, D., Ojaveer, H., & Mackinson, S. (2010). Can simple be
useful and reliable? Using ecological indicators to represent and compare the
states of marine ecosystems. *ICES J. Mar. Sci.*, *67*, 717–731.
- Smith, A. D. M., Fulton, E. J., Hobday, A. J., Smith, D. C., & Shoulder, P.
715 (2007). Scientific tools to support the practical implementation of ecosystem-
based fisheries management. *ICES J. Mar. Sci.*, *64*, 633–639.
- Sourget, Q., & Biais, G. (2009). *Ecologie, biologie et ex-
ploitation du maigre du golfe de Gascogne*. Rapport fi-
nal convention SMIDDEST - IFREMER - CNRS. URL:
720 http://oai.eau-adour-garonne.fr/oai-documents/58983/GED_00000000.pdf.

Stergiou, K. I., Moutopoulos, D. K., & Erzini, K. (2002). Gill net and long-lines fisheries in Cyclades waters (Aegean Sea): species composition and gear competition. *Fish. Res.*, *57*, 25–37.

725 Viana, M., McNally, L., Graham, N., Reid, D. G., & Jackson, A. L. (2013). Ignoring discards biases the assessment of fisheries' ecological fingerprint. *Biol. Lett.*, *9*, 20130812–20130812. doi:10.1098/rsbl.2013.0812.

APPENDIX

Table 1: Classification of species, genus or family by main habitat, taxonomic and functional groups (nei = not elsewhere identified).

Taxon	Main habitat	Taxonomic group	Functional group
<i>Alloteuthis</i>	demersal	mollusc	non-piscivore
<i>Alopias vulpinus</i>	pelagic	fish	piscivore
<i>Alosa</i>	pelagic	fish	non-piscivore
<i>Argentina</i>	pelagic	fish	non-piscivore
<i>Argyrosomus regius</i>	demersal	fish	piscivore
<i>Arnoglossus</i>	benthic	fish	non-piscivore
<i>Auxis rochei rochei</i>	pelagic	fish	piscivore
<i>Balistes capriscus</i>	demersal	fish	non-piscivore
<i>Belone belone</i>	pelagic	fish	non-piscivore
<i>Beryx decadactylus</i>	demersal	fish	piscivore
<i>Boops boops</i>	pelagic	fish	non-piscivore
<i>Brama brama</i>	demersal	fish	piscivore
<i>Buccinum undatum</i>	benthic	mollusc	non-piscivore
<i>Buglossidium luteum</i>	benthic	fish	non-piscivore
<i>Callionymus</i>	benthic	fish	non-piscivore
<i>Cancer pagurus</i>	benthic	crustacean	non-piscivore
<i>Carangidae</i>	pelagic	fish	piscivore
<i>Cepola macrophthalma</i>	demersal	fish	non-piscivore
<i>Ciliata mustela</i>	demersal	fish	non-piscivore
<i>Citharus linguatula</i>	benthic	fish	non-piscivore
<i>Clupea harengus</i>	pelagic	fish	non-piscivore
<i>Conger conger</i>	benthic	fish	piscivore
<i>Crangon crangon</i>	demersal	crustacean	non-piscivore
<i>Dalatias licha</i>	demersal	fish	piscivore
<i>Dasyatis pastinaca</i>	benthic	fish	piscivore
<i>Delphinus delphis</i>	pelagic	mammal	piscivore
<i>Dicentrarchus labrax</i>	demersal	fish	piscivore
<i>Dicentrarchus punctatus</i>	demersal	fish	piscivore
<i>Dicologlossa cuneata</i>	benthic	fish	non-piscivore
<i>Diplodus cervinus</i>	demersal	fish	non-piscivore
<i>Diplodus puntazzo</i>	demersal	fish	non-piscivore

Taxon	Main habitat	Taxonomic group	Functional group
<i>Diplodus sargus</i>	demersal	fish	non-piscivore
<i>Diplodus vulgaris</i>	demersal	fish	non-piscivore
<i>Echiichthys vipera</i>	benthic	fish	non-piscivore
<i>Engraulis encrasicolus</i>	pelagic	fish	non-piscivore
<i>Euthynnus alletteratus</i>	pelagic	fish	piscivore
<i>Gadus morhua</i>	demersal	fish	piscivore
<i>Galeorhinus galeus</i>	demersal	fish	piscivore
<i>Galeus melastomus</i>	demersal	fish	piscivore
<i>Gobiidae</i>	benthic	fish	non-piscivore
<i>Helicolenus dactylopterus</i>	demersal	fish	non-piscivore
<i>Heptanchias perlo</i>	demersal	fish	piscivore
<i>Hexanchus griseus</i>	demersal	fish	piscivore
<i>Hippocampus</i>	benthic	fish	non-piscivore
<i>Homarus gammarus</i>	benthic	crustacean	non-piscivore
<i>Illex coindetii</i>	pelagic	mollusc	non-piscivore
<i>Labrus</i>	demersal	fish	non-piscivore
<i>Lepidorhombus whiffiagonis</i>	benthic	fish	piscivore
<i>Leucoraja naevus</i>	benthic	fish	piscivore
<i>Limanda limanda</i>	benthic	fish	non-piscivore
<i>Lithognathus mormyrus</i>	demersal	fish	non-piscivore
<i>Loligo</i>	pelagic	mollusc	non-piscivore
<i>Lophius</i>	benthic	fish	piscivore
<i>Maja brachydactyla</i>	benthic	crustacean	non-piscivore
<i>Melanogrammus aeglefinus</i>	demersal	fish	non-piscivore
<i>Merlangius merlangus</i>	demersal	fish	piscivore
<i>Merluccius merluccius</i>	demersal	fish	piscivore
<i>Microchirus</i>	benthic	fish	non-piscivore
<i>Micromesistius poutassou</i>	pelagic	fish	non-piscivore
<i>Mola mola</i>	pelagic	fish	piscivore
<i>Molva molva</i>	demersal	fish	piscivore
<i>Mugilidae</i>	demersal	fish	non-piscivore
<i>Mullus</i>	benthic	fish	non-piscivore
<i>Mustelus</i>	demersal	fish	piscivore

Taxon	Main habitat	Taxonomic group	Functional group
<i>Myliobatis aquila</i>	benthic	fish	piscivore
<i>Necora puber</i>	benthic	crustacean	non-piscivore
<i>Nephrops norvegicus</i>	benthic	crustacean	non-piscivore
<i>Octopus</i>	benthic	mollusc	non-piscivore
<i>Pagellus</i>	demersal	fish	piscivore
<i>Pagrus pagrus</i>	demersal	fish	piscivore
<i>Palinuridae</i>	benthic	crustacean	non-piscivore
<i>Pecten maximus</i>	benthic	mollusc	non-piscivore
<i>Pegusa lascaris</i>	benthic	fish	non-piscivore
<i>Petromyzon marinus</i>	demersal	fish	non-piscivore
<i>Phocoena phocoena</i>	pelagic	mammal	piscivore
<i>Phrynorhombus norvegicus</i>	benthic	fish	non-piscivore
<i>Platichthys flesus</i>	benthic	fish	piscivore
<i>Pleuronectes platessa</i>	benthic	fish	piscivore
<i>Pollachius pollachius</i>	demersal	fish	piscivore
<i>Pollachius virens</i>	demersal	fish	piscivore
<i>Polyprion americanus</i>	demersal	fish	piscivore
<i>Prionace glauca</i>	pelagic	fish	piscivore
<i>Phycis blennoides</i>	demersal	fish	non-piscivore
<i>Phycis phycis</i>	demersal	fish	non-piscivore
<i>Raja clavata</i>	benthic	fish	piscivore
<i>Rajidae nei</i>	benthic	fish	piscivore
<i>Salmo salar</i>	demersal	fish	piscivore
<i>Salmo trutta trutta</i>	demersal	fish	piscivore
<i>Sarda sarda</i>	pelagic	fish	piscivore
<i>Sardina pilchardus</i>	pelagic	fish	non-piscivore
<i>Sarpa salpa</i>	demersal	fish	non-piscivore
<i>Scomber colias</i>	pelagic	fish	non-piscivore
<i>Scomber scombrus</i>	pelagic	fish	non-piscivore
<i>Scomberomorus</i>	pelagic	fish	piscivore
<i>Scophthalmus maximus</i>	benthic	fish	piscivore
<i>Scophthalmus rhombus</i>	benthic	fish	piscivore
<i>Scorpaena</i>	benthic	fish	piscivore

Taxon	Main habitat	Taxonomic group	Functional group
<i>Scyliorhinus</i>	demersal	fish	piscivore
<i>Sepia</i>	demersal	mollusc	non-piscivore
<i>Seriola</i>	pelagic	fish	piscivore
<i>Serranus</i>	demersal	fish	piscivore
<i>Solea nei</i>	benthic	fish	non-piscivore
<i>Solea solea</i>	benthic	fish	non-piscivore
<i>Sparus</i>	demersal	fish	piscivore
<i>Spondylisoma cantharus</i>	demersal	fish	non-piscivore
<i>Sprattus sprattus</i>	pelagic	fish	non-piscivore
<i>Squalus acanthias</i>	demersal	fish	piscivore
<i>Stenella coeruleoalba</i>	pelagic	mammal	piscivore
<i>Syngnathus</i>	benthic	fish	non-piscivore
<i>Torpedo</i>	benthic	fish	piscivore
<i>Trachinus draco</i>	benthic	fish	non-piscivore
<i>Trachurus</i>	pelagic	fish	non-piscivore
<i>Triglidae</i>	benthic	fish	non-piscivore
<i>Trisopterus</i>	demersal	fish	non-piscivore
<i>Umbrina canariensis</i>	demersal	fish	non-piscivore
<i>Umbrina cirrosa</i>	demersal	fish	non-piscivore
<i>Zeus faber</i>	demersal	fish	piscivore