

Dinophysis spp. cells concentrated from nature for experimental purposes, using size fractionation and reverse migration

Serge Y. Maestrini^{1,*}, Brigitte R. Berland², Daniel Grzebyk², Anna-Maria Spanò¹

¹Centre de Recherche en Ecologie Marine et Aquaculture de L'Houmeau (CNRS-IFREMER), BP 5,
F-17137 L'Houmeau, France

²Centre d'Océanologie de Marseille (URA CNRS 41), Station marine d'Endoume, Chemin de la Batterie des Lions,
F-13007 Marseille, France

ABSTRACT: A method based on size fractionation plus reverse phototactic migration was used to collect large *Dinophysis* spp.-dominated populations. Cells were active and survived handling and transportation. Experiments involving various treatments for establishing permanent cultures are described; none have yet been successful.

KEY WORDS: *Dinophysis* · Dense-cell suspension · Attempted culture

INTRODUCTION

The recent increase, both in space and time, of damage related to the presence of dinoflagellates *Dinophysis* (Anderson 1989, Smayda 1990, Hallegraeff 1993) has considerably stimulated research on their life cycles, reproductive strategies, nutrition, toxin production and taxonomy, using newly developed methods. Failure to culture any of the *Dinophysis* species, however, has so far prevented the usual set of ecophysiological experiments, seriously limiting current knowledge. Notwithstanding these difficulties, various working hypotheses have been tested using material isolated, with considerable effort, from seawater.

In offshore areas, the thermocline has been reported as the layer where *Dinophysis* spp. accumulate (Delmas et al. 1992, and references therein); a research vessel is needed there to allow CTD and 'Niskin'-type sampling bottles to be operated. A few locations further inshore, however, represent more convenient sampling sites. Granéli et al. (1993), for instance, sampled natural populations of *Dinophysis* spp. a few tens of meters from the pier of Kristineberg Marine Station,

Sweden, and Subba Rao & Pan (1993) sampled these organisms near the Bedford Institute of Oceanography, Canada.

In France, the port of Antifer, near Le Havre, provides exceptionally good conditions for collecting *Dinophysis* spp. cells, since high densities, up to 160 000 cells l⁻¹, have occurred in summer nearly every year since 1987 (Lassus et al. 1993). The dominant species is similar to *D. acuminata*, though it is likely to be a different, undescribed species (hereafter referred as *D. cf. acuminata*; Lassus & Bardouil 1991). *D. acuta* and *D. sacculus* are usually also present.

Here we report (1) useful information gained during attempts to cultivate these dinoflagellates, and (2) a method we have developed to provide a substitute material for experiments, i.e. a dense cell suspension, obtained from the natural population.

MATERIAL AND METHODS

Sampling. From 1989 to 1992, from July to September, as soon as the cell density reached 5000 l⁻¹ (Fig. 1), samples of up to 2000 l of surface water were collected during flood tide, using a bucket, from the pier of Antifer harbor (Fig. 2). Concentration (see below) was

*E-mail: smaestri@ifremer.fr




Fig. 1. Temperature and cell density of *Dinophysis cf. acuminata* in Antifer harbor, France, summer 1992

done immediately after sampling in uncontrolled-temperature conditions. Cultivation attempts were made in laboratory facilities, at least 1 d after sampling.

Cultivation. First attempts were made with 3 typical algal culture media free from silicate: medium f/2 of Guillard & Ryther (1962), Antia & Cheng's (1970) medium, Chan's (1978) medium; standard glassware and handling methods were used. Later, the following improvements were made: polycarbonate or Teflon bottles, cleaned according to the ultraclean protocol (Guillard & Keller 1984), were used in order to prevent leaching from the vessel walls of toxic substances such as heavy metals; offshore deep water, purified by passage on Florisil resin to remove potential organic inhibiting substances (Gentien & Arzul 1990), was used to make up media; ultrapure chemical compounds were employed; and sterilisation was by filtration or by autoclavation in Teflon bottles.

As supplements, various materials were tried according to results reported in the literature, including soil extract, humic acids (Carlsson & Granéli 1993) and various cocktails of either inorganic compounds such as those of Fe (Okaichi et al. 1989, Wells et al. 1991), Mn, Mo or Se (Lindström & Rodhe 1978, Keller et al. 1987, Harrison et al. 1988), or organic substances: dextrans at various concentrations (15 to 500×10^3 M or 5 to 500 mg l⁻¹) (Tranvik et al. 1993), and organic nitrogenous substances such as urea, glutamic acid and hypoxanthin (Antia et al. 1975). Growth substances such as gibberellic acid, indol acetic acid and kinetin (Bentley-Mowat & Reid 1969, Paster & Abott 1970) were tried with or without a mixture of 10 vitamins. Polyamines (cadaverin, dimethyl amine, putrescine, spermine, spermidine) at 10^{-6} to 10^{-9} M, lectins of *Phaseolus* at 1 to 100 mg l⁻¹, and porcine blood platelets at 10 to 500 µg l⁻¹ were also tested.

Several global concentration sets were used, from low (i.e. 5 µM nitrogen) to high concentration (i.e. 500 µM nitrogen).

During these trials, temperature was kept constant (18 to 20°C), and light intensity dim (200 µmol photons m⁻² s⁻¹) in view of reports that such conditions give better activity (Keller & Guillard 1985), while stirring was nil or reduced, since turbulence usually inhibits growth in dinoflagellates (White 1976, Thomas et al. 1991, Berdalet 1992).

The inoculum used was either a single or several *Dinophysis* cells isolated under a microscope with a micropipette, or a sample of a natural population.

On the basis of results obtained and those of Hansen's (1991), we ultimately tried to grow *Dinophysis cf. acuminata*, *D. acuta* and *D. sacculus* by providing varied food: bacteria, picoplankton, nanoplankton (*Cryptomonas salina*; according to Larsen 1988) and yeast. We also performed experiments with a double chamber well, with *Dinophysis* spp. cells in one chamber and 'companion' species in the other, so as to investigate any possible stimulating effect of external metabolites. *In situ* incubations in dialysis bags were also carried out in the field at the point where the cells were sampled.

RESULTS

Cultivation

All attempts to provide a culture failed. The best result we obtained was to keep several cells alive for 5 mo, and to obtain 16 cells from a single one in 4 mo (both in 400 µl wells, 20 µm filtered seawater, dim light, 12/12 h light-dark, 18°C). Generally cells became progressively depigmented and then died.

Nutrition

We never observed any external phagotrophy with a peduncle or a velum, like we did several times with a *Protoperdinium* spp. which was frequently a companion species in samples. On the other hand, we observed several *Dinophysis cf. acuminata* cells which stopped swimming, sank to the bottom, attached the epitheca and the flagellar pore to the wall of the culture flask, and later resumed swimming.

Cell behavior

All 3 species of *Dinophysis* appeared not to be fundamentally fragile: (1) they were collected several

times in very turbulent water and swam quite well soon afterwards in a laboratory vessel, whereas other motile species did not swim but sank; (2) they did not die when mailed in a 100 ml polyethylene bottle in a regular parcel taking 48 h, nor when transported by car for 8 h without temperature control in a 25 l polycarbonate bottle covered by a dark cloth. In contrast, they suffered badly from concentrated decayed material.

DISCUSSION

Our results on cultivation are consistent with previous reports. Barker (1935), who first tried to cultivate *Dinophysis* species on the medium 'Erd-Schreiber' of Foyn (1934) prepared with aged seawater, maintained several cells alive for 1 mo. Later, Maranda & Shimizu (1987) tried, in addition to the medium f/50 of Guillard & Ryther (1962), several specific media prepared with



Fig. 2. Sampling of surface water from the pier of Antifer harbor




Fig. 3. Accumulation strips (white arrows) of *Dinophysis* spp. in the upper part of the polycarbonate bottle




Fig. 4. Extremely concentrated population of *Dinophysis* cf. *acuminata* (not usual)

artificial or natural seawater to grow some dinoflagellates; namely, the media MLH/4 and MLH/40 that Tuttle & Loeblich (1975) devised to grow *Cryptothecodinium cohnii*, a heterotrophic species, and the medium OX 7/20 of Droop (1959) supplemented with terpenoid quinones and steroids (Droop & Pennock 1971), originally used to grow the phagotrophic species *Oxhyrrhis marina*. In experiments by Durand Clément et al. (1988), *Dinophysis* spp. cells survived for 4 to 5 wk and some divided, in seawater enriched according to Provasoli et al. (1957) or simply with inorganic N and P compounds and soil extract. Sampayo (1993) reported better results by using the natural-seawater medium of Miquel (1890–93) and the artificial-seawater medium ASP-7 of Provasoli (1963); she could maintain alive several cells of *D. acuminata* and *D. acuta* for up to 5 mo, with some cellular divisions occurring.

Ishimaru et al. (1988) reported having obtained a culture of *Dinophysis fortii* and *D. acuminata* fed with the cryptomonad *Plagioselmis* sp. They obtained 22 cells from a single one in 3 wk, which is not sufficient to be regarded as a culture in the full sense. Their results are nevertheless important because they support the possibility of phagotrophy which was already suspected by many authors, including us, on the basis that *Dinophysis* spp. cells lose their pigments and divide best in the presence of small living particles. Furthermore, Granéli et al. (1995) reported they had observed *Dinophysis* cells which appeared to have captured a small *Thalassiosira* cell.

That *Dinophysis* cf. *acuminata* could be mixotrophic would not be surprising, since other dinoflagellates of different genera frequently are (Schnepf & Elbrächter 1992, Bockstahler & Coats 1993) and some *Dinophysis* species are heterotrophic (Hallegraeff & Lucas 1988) and even predatory (Hansen 1991). On this basis, further research should focus on both the photosynthetic and the heterotrophic capabilities of *Dinophysis* spp., and protistological approaches should be used as well,




Fig. 5. Diagram of protocol used to concentrate *Dinophysis* spp. cells

Table 1. Typical cell concentrations (10^3 cells l^{-1}) and relative cell density (%; number within parentheses) of algal components in the natural population, the 20–77 µm fractionated fraction and the surface layer of the nonturbulent bottle.

Experiment of 31 August 1992; sample taken at Antifer

Alga	Natural seawater	20–77 µm fraction	Surface layer
<i>Dinophysis</i> spp.	2.5 (11)	12.9 (14)	317.4 (69)
Other dinoflagellates	8.8 (39)	17.5 (19)	137.0 (29)
Diatoms	11.3 (50)	61.8 (67)	7.7 (2)

since results of Jacobson & Andersen (1994) suggest the prey could be in the >77 µm size fraction.

By taking advantage of their resistance and their motility, it is possible to concentrate *Dinophysis* spp. populations, with markedly reduced diatom components. The following protocol (Figs. 3, 4 & 5), based on that used by Vernet et al. (1989) for *Prorocentrum micans*, has proved efficient: (1) Seawater, preferably harvested with a bucket, or a peristaltic pump when necessary, is successively filtered through 77 µm and 20 µm meshes (Fig. 5), so as to concentrate the cells by a factor of 5 to 10. (2) The concentrated 20–77 µm fraction is gently siphoned into 10 l polycarbonate bottles and left to stand for at least 6 h; most of the dead cells and diatoms sink and collect at the bottom, while *Dinophysis* spp. and some other motile cells swim near the surface. (3) The surface layer is gently siphoned off; Table 1 shows a typical result. (4) In a few favorable cases, *Dinophysis* spp. concentrated in 5 to 6 mm strips on the side away from the light when the bottle was laterally illuminated at low intensity with 1 or 2 fluorescent tubes (Fig. 3); siphoning with a curved glass tube provided very dense *Dinophysis*, yet never totally free from co-occurring companion species (Fig. 4).

Acknowledgements. This study was supported by the 'Programme National Efflorescences Algales Toxiques'. We warmly thank Captain Cirot, M. Ferme and their teams for help during sampling at Antifer; Mrs F. Mornet for phytoplankton counting; Dr Ian Jenkinson (ACRO, La Roche Canillac) for improving the English version; and one anonymous referee for very helpful comments.

LITERATURE CITED

- Anderson DM (1989) Toxic algal blooms and red tides: a global perspective. In: Okaichi T, Anderson DM, Nemoto T (eds) Red tides: biology, environmental science, and toxicology. Proc 1st Int Conf on Red Tides, 10–14 November 1987, Takamatsu, Japan. Elsevier Sci Pub, New York, p 11–16
- Antia NJ, Berland BR, Bonin DJ, Maestrini SY (1975) Comparative evaluation of certain organic and inorganic sources of nitrogen for phototrophic growth of marine microalgae. J mar biol Ass UK 55:519–539

- Antia NJ, Cheng JY (1970) The survival of axenic cultures of marine planktonic algae from prolonged exposure to darkness at 20°C. *Phycologia* 9:179–84
- Barker HA (1935) The culture and physiology of the marine dinoflagellates. *Arch Mikrobiol* 6:157–181
- Bentley-Mowat JA, Reid SM (1969) Effect of gibberellins, kinetin and other factors on the growth of unicellular marine algae in culture. *Bot mar* 12:185–199
- Berdalet E (1992) Effects of turbulence on the marine dinoflagellate *Gymnodinium nelsonii*. *J Phycol* 28(3):267–272
- Bockstahler KR, Coats DW (1993) Grazing of the mixotrophic dinoflagellate *Gymnodinium sanguineum* on ciliate populations of Chesapeake Bay. *Mar Biol* 116(3):477–487
- Carlsson P, Granéli E (1993) Availability of humic bound nitrogen for coastal phytoplankton. *Estuar coast Shelf Sci* 36:433–447
- Chan TA (1978) Comparative physiological studies of marine diatoms and dinoflagellates in relation to irradiance and cell size. I. Growth under continuous light. *J Phycol* 14:396–402
- Delmas D, Herbland A, Maestrini SY (1992) Environmental conditions which lead to increase in cell density of the toxic dinoflagellates *Dinophysis* spp. in nutrient-rich and nutrient-poor waters of the French Atlantic coast. *Mar Ecol Prog Ser* 89:53–61
- Droop MR (1959) Water-soluble factors in the nutrition of *Oxhyrrhis marina*. *J mar biol Ass UK* 38:605–620
- Droop MR, Pennock JF (1971) Terpenoid quinones and steroids in the nutrition of *Oxhyrrhis marina*. *J mar biol Ass UK* 51:455–470
- Durand Clément M, Clément JC, Moreau A, Jeanne N, Puisieux-Dao S (1988) New ecological and ultrastructural data on the dinoflagellate *Dinophysis* sp. from the French coast. *Mar Biol* 97:37–44
- Foyn B, (1934) Lebenszyklus, Cytologie und Sexualität der Chlorophyceen *Cladophora suhriana* Kützinger. *Archiv Protistenkd* 83:1–56
- Gentien P, Arzul G (1990) Exotoxin production by *Gyrodinium* cf. *aureolum* (Dinophyceae). *J mar biol Ass UK* 70:571–587
- Granéli E, Anderson DM, Carlsson P, Finenko G, Maestrini SY, Smayda TJ, Sampayo MA de M (1995) Nutrition, growth rate and sensibility to grazing for the dinoflagellates *Dinophysis acuminata*, *D. acuta* and *D. norvegica*. *La Mer* 33(3): in press
- Granéli E, Anderson DM, Maestrini SY, Paasche E (1993) Light and dark carbon fixation by the marine dinoflagellate genera *Dinophysis* and *Ceratium*. In: Li WKW, Maestrini SY (eds) Measurement of primary production from the molecular to the global scale. *Proc Int Symp, La Rochelle, France, 21–24 April 1992. ICES mar Sci Symp* 197:274
- Guillard RRL, Keller MD (1984) Culturing dinoflagellates. In: Spector DL (ed) *Dinoflagellates*. Academic Press, New York, p 391–442
- Guillard RRL, Ryther JH (1962) Studies on marine planktonic diatoms. I. *Cyclotella nana* Hustedt and *Detonula confervacae* (Cleve) Gran. *Can J Microbiol* 8:229–39
- Hallegraeff GM (1993) A review of harmful algal blooms and their apparent global increase. *Phycologia* 32(2):79–99
- Hallegraeff GM, Lucas IAN (1988) The marine dinoflagellate genus *Dinophysis* (Dinophyceae): photosynthetic, neritic and non-photosynthetic, oceanic species. *Phycologia* 27(1):25–42
- Hansen PJ (1991) *Dinophysis* — a planktonic dinoflagellate genus which can act both as a prey and a predator of a ciliate. *Mar Ecol Prog Ser* 69:201–204
- Harrison PJ, Yu PW, Thompson PA, Price NM, Phillips DJ, (1988) Survey of selenium requirements in marine phytoplankton. *Mar Ecol Prog Ser* 47:89–96
- Ishimaru T, Inoue H, Fukuyo Y, Ogata T, Kodama M (1988) Cultures of *Dinophysis fortii* and *D. acuminata* with the cryptomonad, *Plagioselmis* sp. In: Aibara K et al. (eds) *Mycotoxins and phycotoxins*. Jap Ass Microtoxicol, Fac Pharm Sci, Univ Tokyo, p 19–20
- Jacobson DM, Andersen RA (1994) The discovery of mixotrophy in photosynthetic species of *Dinophysis* (Dinophyceae): light and electron microscopical observations of food vacuoles in *Dinophysis acuminata*, *D. norvegica* and two heterotrophic dinophysoid dinoflagellates. *Phycologia* 33(2):97–110
- Keller MD, Guillard RRL (1985) Factors significant to marine dinoflagellate culture. In: Anderson DM, White AW, Baden DG (ed) *Toxic dinoflagellates*. Elsevier Sci Pub, New York, p 113–116
- Keller MD, Selvin RC, Claus W, Guillard RRL (1987) Media for the culture of oceanic ultraphytoplankton. *J Phycol* 23:633–638
- Larsen J (1988) An ultrastructural study of *Amphidinium poecilochroum* (Dinophyceae), a phagotrophic dinoflagellate feeding on small species of cryptophytes. *Phycologia* 27(3):366–377
- Lassus P, Bardouil M (1991) Le complexe '*Dinophysis acuminata*': identification des espèces le long des côtes françaises. *Cryptogamie Algol* 12(1):1–9
- Lassus P, Proniewski F, Maggi P, Truquet P, Bardouil M (1993) Wind-induced toxic blooms of *Dinophysis* cf. *acuminata* in the Antifer area (France). In: Smayda TJ, Shimizu Y (eds) *Toxic phytoplankton blooms in the sea. Proc 5th Int Conf on Toxic Marine Phytoplankton, 28 Oct–1 Nov 1991, Newport, USA*. Elsevier Sci Pub, New York, p 519–523
- Lindström K, Rodhe W (1978) Selenium as a micronutrient for the dinoflagellate *Peridinium cinctum* fa. *westii*. *Mitt Int Verein Limnol* 21:168–173
- Maranda L, Shimizu Y (1987) Diarrhetic shellfish poisoning in Narragansett Bay. *Estuaries* 10(4):298–302
- Miquel P (1890–93) De la culture artificielle des diatomées. *Le diatomiste* 9–10, 93–99 and 121–128
- Okaichi T, Montani S, Hiragi J, Hasui A (1989) The role of iron in the outbreaks of *Chattonella* red tide. In: Okaichi T, Anderson DM, Nemoto T (eds) *Red tides: biology, environmental science, and toxicology. Proc 1st Int Conf on Red Tides, 10–14 November 1987, Takamatsu, Japan*. Elsevier Sci Pub, New York, p 357–362
- Paster Z, Abbott BC (1970) Gibberelic acid: a growth factor in the unicellular alga *Gymnodinium breve*. *Science* 169:601–606
- Provasoli L (1963) Growing marine seaweeds. In: *Proceedings of the 4th International Seaweed Symposium*. Pergamon Press, Oxford, p 9–17
- Provasoli L, McLaughlin JJA, Droop MR (1957) The development of artificial media for marine algae. *Arch Mikrobiol* 25:392–428
- Sampayo MA de M (1993) Trying to cultivate *Dinophysis* spp. In: Smayda TJ, Shimizu Y (eds) *Toxic phytoplankton blooms in the sea. Proc 5th Int Conf Toxic Marine Phytoplankton, 28 Oct–1 Nov 1991, Newport, USA*. Elsevier Sci Pub, New York, p 807–810
- Schnepf E, Elbrächter M (1992) Nutritional strategies in dinoflagellates. A review with emphasis on cell biological aspects. *Eur J Protistol* 28:3–24
- Smayda TJ (1990) Novel and nuisance phytoplankton blooms in the sea: evidence for a global epidemic. In: Granéli E, Sundström B, Edler L, Anderson DM (eds) *Toxic marine*

- phytoplankton. Proc 4th Int Conf Toxic Mar Phytoplankton, 26-30 June 1989, Lund, Sweden. Elsevier Sci Pub, New York, p 29-40
- Subba Rao DV, Pan Y (1993) Photosynthetic characteristics of *Dinophysis norvegica* Claparede & Lachmann, a red-tide dinoflagellate. J Plankton Res 15(8):965-976
- Thomas WH, Gibson CH, Vernet M (1991) Inhibition of *Gonyaulax polyedra* Stein by quantified small-scale turbulence. J Phycol, Suppl 27(3):72
- Tranvik LJ, Sherr EB, Sherr BF (1993) Uptake and utilization of 'colloidal DOM' by heterotrophic flagellates in seawater. Mar Ecol Prog Ser 92:301-309
- Tuttle RC, Loeblich AR III (1975) An optimal growth medium for the dinoflagellate *Cryptothecodinium cohnii*. Phycologia 14(1):1-8
- Vernet M, Neori A, Haxo FT (1989) Spectral properties and photosynthetic action in red-tide populations of *Prorocentrum micans* and *Gonyaulax polyedra*. Mar Biol 103: 365-371
- Wells ML, Mayer LM, Guillard RRL (1991) Evaluation of iron as a triggering factor for red tide blooms. Mar Ecol Prog Ser 69:93-102
- White AW (1976) Growth inhibition caused by turbulence in the toxic marine dinoflagellate *Gonyaulax excavata*. J Fish Res Bd Can 33:2598-2602