



European
Commission

JRC SCIENTIFIC AND POLICY REPORTS

Scientific, Technical and Economic Committee for Fisheries (STECF)

Multiannual management plans SWW and NWW (STECF-15-08)

Edited by Ernesto Jardim & Iago Mosqueira

This report was reviewed by the STECF during its 49th plenary meeting
held from 6 to 10 July 2015 in Varese, Italy

Report EUR XXXXX EN

European Commission
Joint Research Centre (JRC)
Institute for the Protection and Security of the Citizen (IPSC)

Contact information
STECF secretariat
Address: Maritime Affairs Unit, Via Enrico Fermi 2749, 21027 Ispra VA, Italy
E-mail: stecf-secretariat@jrc.ec.europa.eu
Tel.: 0039 0332 789343
Fax: 0039 0332 789658

JRC Science Hub
<https://ec.europa.eu/jrc>

Legal Notice

This publication is a Science and Policy Report by the Joint Research Centre, the European Commission's in-house science service. It aims to provide evidence-based scientific support to the European policy-making process. The scientific output expressed does not imply a policy position of the European Commission. Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use which might be made of this publication.

JRC XXXX
EUR XXXX EN
ISBN XXXXX
ISSN 1831-9424
doi:XXXXXXXX

Luxembourg: Publications Office of the European Union, 2015
© European Union, 2015
Reproduction is authorised provided the source is acknowledged

How to cite this report:

Scientific, Technical and Economic Committee for Fisheries (STECF) – Multiannual management plans SWW and NWW (STECF-15-08). 2015. Publications Office of the European Union, Luxembourg, EUR XXXX EN, JRC XXXX, XXX pp.

Abstract

The STECF was tasked with an analysis of the likely effects of proposed management plans for the Southwestern (Bay of Biscay and Iberia) and Northwestern (Celtic sea) waters. Quantitative analyses were carried out to compare the likely effect of those management plans and of the direct application of the CFP on both stocks and fleets involved in these fisheries.

TABLE OF CONTENTS

Multiannual management plans SWW and NWW (STECF-15-08)	6
Request to the STECF	6
STECF considerations	6
Conclusions of the STECF	7
Expert Working Group EWG-15-04 & 09 report	9
1 Executive summary	10
2 Introduction	10
2.1 Background	10
2.2 Terms of reference	10
2.3 Detailed terms of reference	11
2.3.1 Basic data and assumptions	11
2.3.2 Indicators	11
2.3.2.1 Biological:	11
2.3.2.2 Economic (by fleet segment):	11
2.3.3 Governance	11
2.3.4 Detailed scenarios	11
2.3.5 Number and scope of MAPs.	12
2.3.6 Fishery approach	12
2.3.7 List of stocks considered (provided by DGMARE).....	13
2.3.8 List of fisheries with their target species (provided by DGMARE)	15
3 Description of the fisheries	17
3.1 North Western Waters	17
3.1.1 Celtic Sea proper (VIIe-k including Western Channel)	18
3.1.2 Irish Sea (VIIa).....	19
3.1.3 West of Scotland (ICES Subarea VIa)	19
3.2 South Western Waters	20
3.2.1 Bay of Biscay	20
3.2.2 Iberian waters	22
4 Methods and data	23
4.1 Addressing the ToRs	23
4.1.1 Evaluating scenarios using quantitative methods.....	23
4.1.2 Employment and fleet dependency	24
4.1.3 Number and scope of MAPs	24

4.1.4	Management of by-catch (fishery approach)	24
4.2	Provisional Fmsy ranges	24
4.3	Multi-model approach	25
4.4	Scenarios	26
4.4.1	Management scenarios	26
4.4.2	Fleet scenarios	28
4.4.3	Scenario summary	28
4.5	Data	29
5	ToR 3.1-3.3 - Evolution of EU fisheries under different scenarios	29
5.1	North Western Waters	30
5.1.1	State of the fisheries in 2017	30
5.1.2	State of the fisheries in 2021	32
5.1.3	State of the fisheries in 2025	34
5.2	South Western Waters - Bay of Biscay	36
5.2.1	State of the fisheries in 2017	36
5.2.2	State of the fisheries in 2021	39
5.2.3	State of the fisheries in 2025	43
5.3	South Western Waters - Iberian Waters	47
5.3.1	State of the fisheries in 2017	47
5.3.2	State of the fisheries in 2021	49
5.3.3	State of the fisheries in 2025	51
5.4	Employment and Dependency in the NWW	52
5.5	Employment and Dependency in the SWW	54
5.6	Reconciling TACs by using FMSY ranges	57
6	ToR 3.4 – Number and scope of MAPs	57
7	ToR 3.5 – Fishery approach NWW	58
8	ToR 3.5 – Fishery approach SWW	60
9	ToR 3.5 – Multi-species TACs for by-catch stocks	64
10	Conclusions	66
10.1	ToR 3.1-3.3	66
10.2	ToR 3.4	67
10.3	ToR 3.5	67
11	CONTACT DETAILS OF STECF MEMBERS AND EWG-15-04 & EWG-015-09 PARTICIPANTS	68
12	List of Background Documents	74
13	List of electronic annexes	74

14	ANNEX I – CODES AND ACRONYMS	75
14.1	COUNTRIES CODES	75
14.2	SPECIES CODES USED.....	75
14.3	DCF CODES	75
14.4	ACRONYMS	77

SCIENTIFIC, TECHNICAL AND ECONOMIC COMMITTEE FOR FISHERIES (STECF)

Multiannual management plans SWW and NWW (STECF-15-08)

THIS REPORT WAS REVIEWED DURING THE PLENARY MEETING HELD IN Varese (Italy), 6-10 June 2015

Request to the STECF

STECF is requested to review the report of the STECF Expert Working Group meeting, evaluate the findings and make any appropriate comments and recommendations.

Given the generic approach undertaken for the evaluation of Multi-annual plans associated with the North Western Waters and the South Western Waters Region, the STECF evaluation of the relevant sections (NWW/SWW) of EWG 15-04 and EWG 15-09 are considered together in the following evaluation.

STECF considerations

STECF notes the considerable amount of work carried out by the EWG and concludes that the different methodologies used to address all the TORs follows the best practices in the field of simulation modelling for providing scientific policy advice.

STECF notes that TORs 3.1 to TOR 3.2 of the EWG 15-04 and EWG 15-09 have been addressed using simulation testing. Five different models have been used to conduct the simulations of the EWG:

- Iberian waters simulation model (FLBEIA).
- Iberian waters multi-fleet state-space model
- Bay of Biscay Spanish fleets simulation model (FLBEIA).
- Bay of Biscay French fleets simulation model (IAM).
- Celtic Sea (FLBEIA)

At the present time, models covering other areas in the NWW (e.g. Irish Sea, Western Channel and West of Scotland) are not available.

Using each of the above models, two management options were simulated. Option one (baseline) which included:

- Single species Fmsy objectives
- Achieving objectives in 2016
- Inter-species flexibility (LO)
- Inter-annual flexibility (LO)
- Existing management plans

and option two (named MAP) which when implemented will repeal the existing management plans, includes:

- FMSY ranges instead of single species Fmsy
- Achieving objectives in 2016
- Inter-species flexibility (LO)

- Inter-annual flexibility (LO)
- De minimis exemption (LO)
- Survivability exemption (LO)
- Biomass safeguards

The results provided in the EWG Report are expressed in relative terms in order to highlight the relative differences between the two management options.

For most of the stocks concerned, F_{MSY} ranges have not yet been provided by ICES and so were derived using a regression analysis approach based on North Sea and Baltic F_{MSY} estimates (ICES WKFMSYREF3).

The models used were unable to incorporate all fleets and stocks that exist in each of the management areas. However, for the stocks and fleets that could be included in the analysis, the simulations take account of the catches of all stocks and the fleet revenues obtained from them. Furthermore, for the Northern Hake stock, which is common to the two Bay of Biscay simulation models, the parametrization was made consistent and the results obtained from both models were similar.

The potential impact of the LO was not evaluated by the EWG due to time constraints and uncertainty associated with how it is likely to be implemented; namely which decisions will be taken by the MS regarding inter-annual and inter-species flexibilities, which may result in large changes in fishing mortality.

STECF notes that EWG 15-04 and EWG 15-09 used the same method used by EWG 15-04, to highlighted fleets with ‘high’ and ‘low’ employment together with their economic dependency on the species identified in the MAP (relative to the total landings’ value of each fleet). Such an analysis allows the identification of potential employment impacts created by the implementation of the MAP, as well as identifying the fleets most impacted.

All of the EWGs computed a number of economic indicators such as fixed costs, variable costs, revenue and GVA. STECF notes that the forecasts of economic indicators are largely based on the transformation of catch, effort and capacity, and do not reflect other potential economic dynamics due to the due to the difficulties in forecasting changes in prices of fish, costs of fuel, wages, etc. Indicators based on profits are considered to be uninformative and potentially misleading and were deliberately not computed for the reasons outlined in Section 4.1 of this report.

STECF notes that for TOR 3.4 no quantitative analysis was carried out, the EWGs’ findings are based on experts’ knowledge.

STECF notes that TOR 3.5 has been undertaken using correlations between species’ catches. The analyses indicate it is unlikely that setting TACs for the target/driver stocks will be sufficient to manage exploitation rates on by-catch/non-driver stocks.

Conclusions of the STECF

Based on the results of simulations of the provisions of the proposed management plan, STECF concludes that, setting fishing opportunities in line with single-species F_{MSY} ranges will provide managers with additional flexibility compared to the basic provisions of the 2013 CFP. Such flexibility is likely to help alleviate the problem of mismatches in quota availability in mixed-species fisheries thereby reducing the risk of early closure of some fisheries due to choke species. Adopting F_{MSY} ranges will therefore increase the likelihood that desired exploitation rates will be achieved and will reduce the risk that some fishing fleets will go out of business.

STECF considers that it is crucial that managers take note that persistent fishing at the upper limits of the F_{msy} ranges across all or most stocks simultaneously negates the flexibility introduced by the

FMSY ranges and greatly increases the risk of overfishing. Such an approach will also increase the risk that the objectives of the CFP will not be achieved..

STECF concludes that single species biomass safeguards for all stocks should be maintained to provide a basic level of protection.

STECF notes that for the fleets affected by the SWW MAP, those providing the highest employment are generally not dependent to a great extent on the species that will be regulated through the MAP proposals.

STECF notes that in the NWW there are some fleets which provide significant levels of employment and seem to be very dependent on the species that will be regulated through the MAP proposals. Nevertheless, there are a number of fleets in the NWW area that are not included in the employment analysis because of an absence of appropriate data. .Regarding the number and scope of MAPs as currently defined, STECF considers that a MAP covering a wider geographic area has advantages in terms of reducing management overheads and avoiding multiple regulations affecting the sector. A larger MAP area however, may have disadvantages associated with reducing the emphasis on local management measures and this may discourage the involvement of stakeholders, although this effect will depend on how the process of regionalization operates within the MAP.

To evaluate the question of whether management of the species that drive the fisheries adequately allows for the management of by-catch species, the EWG carried out an analysis of correlations between catches of driver species identified in the plan and a variety of by-catch species. The analysis suggested only limited correlation. In view of this, the STECF notes that it is unlikely that relying on the TAC of the driver species to manage other species will be effective, in accordance with CFP requirements. STECF however notes that when analysis was performed at the fleet level, there were more obvious correlations, suggesting some scope to use fleet related management measures for the driver species as a way of managing some of the bycatch species. STECF therefore concludes that management of exploitation rates of non-driver (or bycatch) species is unlikely to occur as an automatic consequence of the management of the main (driver) stocks by TAC considered in the MAP.

REPORT TO THE STECF

**EXPERT WORKING GROUPS ON
Multiannual management plans SWW & NWW (EWG-15-04 &
09)**

Vigo 25-29 May 2015 & Sète 15-19 June 2015

This report does not necessarily reflect the view of the STECF and the European Commission and in no way anticipates the Commission's future policy in this area

1 EXECUTIVE SUMMARY

2 INTRODUCTION

2.1 Background

According to the reformed CFP (Regulation (EU) No 1380/2013), the objective of sustainable exploitation of marine biological resources is more effectively achieved through a multiannual approach to fisheries management, and hence multiannual plans reflecting the specificities of different fisheries shall be adopted as a priority.

Multiannual plans should, where possible, cover multiple stocks where those stocks are jointly exploited. The multiannual plans should establish the framework for the sustainable exploitation of stocks and marine ecosystems concerned, defining clear time-frames and safeguard mechanisms for unforeseen developments. Multiannual plans should also be governed by clearly defined management objectives in order to contribute to the sustainable exploitation of the stocks and to the protection of the marine ecosystems concerned. Those plans should be adopted in consultation with Advisory Councils, operators in the fishing industry, scientists and other stakeholders having an interest in fisheries. Prior to including measures in a multiannual plan, account shall be taken of their likely environmental, economic and social impact.

2.2 Terms of reference

The purpose of the request to STECF is to obtain the scientific grounds for the assessment of the ecological, economic and social effects of a range of possible measures applicable in the context of multiannual plans applicable to demersal fisheries (excluding those for deep-water fish) in:

- 1) North-western EU waters: subareas VI, VII
- 2) South-western EU waters: subareas VIII, IX

These multiannual plans will be hereinafter referred to as NWW MAP and SWW MAP respectively.

STECF is requested to analyse the evolution of EU fisheries and to describe their likely situation in the short and medium term in each of the two areas mentioned under two main management options:

Option 1: There are no MAPs; fisheries continue to be managed under the existing rules of the CFP. This includes the existing multiannual plans¹, Regulation (EU) 1380/2013 (the Basic Regulation), the Technical Measures Regulation (Regulation (EC) No 850/98) and the Omnibus Regulation (in the process of finalisation at this point in time).

Option 2: In addition to the existing rules, two MAPs enter into force from 2017. The existing MAPs are repealed from 1 January 2017, except the Western Waters Regime². Under this scenario, STECF will be requested to analyse alternative measures that could be part of the plans.

For each of the scenarios, STECF is requested to run the appropriate forecast models in order to describe the likely situation of the fisheries as in 1 January 2017, 2021 and 2025 using the indicators given below.

2.3 Detailed terms of reference

2.3.1 Basic data and assumptions

Simulations are to be carried out on the basis of the most recent ICES analysis available and on data that exist or that can be collected through the data collection framework (Regulation (EC) No 199/2008). This includes information on population status and dynamics, and reference points, taken as point estimates and, where applicable, ranges of likely values for those reference points. Whenever the later are unavailable, STECF is requested to estimate approximate values just for the purpose of this evaluation, using a simplified methodology on the basis of the same principles as those of ICES (mainly to allow 5% variation in yield and constrain upper limits on the basis of Bpa).

2.3.2 Indicators

2.3.2.1 Biological:

- Abundance (SSB) and fishing mortality relative to Fmsy (F/Fmsy) of main stocks
- Abundance (total biomass) of the main predator stocks. Description of the significance of this indicator in terms of ecosystem status.
- Mean individual size of each of the main species and overall mean individual size of all the main species combined. Description of the significance of this indicator in terms of ecosystem status.

2.3.2.2 Economic (by fleet segment):

- GVA
- Gross cash flow
- Net profit
- Social (by fleet segment):
- Employment and, where possible, associated wages.

2.3.3 Governance

STECF is requested to call the attention to situations where there are difficulties to abide by the rules, leading to is a high probability of non-compliance with law (e.g. "choke" effects potentially leading to discarding or illegal landings). Where STECF believes that measures can be adopted to alleviate the difficulties for the industry (improved selectivity, quota swapping) these should be described.

2.3.4 Detailed scenarios

Scenario under option 1:

Setting of TACs: For all stocks with an analytical assessment and a catch forecast (ICES categories 1 and 2), TACs are proposed in accordance with Fmsy (point estimates) or proxies to it, but in reality it can be expected that TAC reductions beyond 15% will not be finally adopted and that in 30% of the cases Fmsy will be exceeded by a significant margin. For stocks without a full analytical assessment (ICES categories 3 and beyond) TACs will be rolled over.

The landings obligation (LO) applies to all demersal fish subject to quota regulations from 2018 on. This is a knife-edge approximation to a gradual phasing in of the LO from 2016 to 2019. In the absence of MAPs, the landing obligation applies strictly, without exceptions (survivability, *de minimis*, etc).

Only the existing technical measures (Reg. 850/98 and Omnibus) apply. No new rules can be put in place in any other EU legislation. Improvements in selective fishing, e.g. to minimise choke effects or to avoid catch of juvenile fish, might be voluntarily adopted by some fishermen, but then these measures should be taken as less effective (by 50%) than legally binding ones, given they are not expected to be adopted by all vessels. .

Scenario under option 2:

Setting of TACs: Until 2016, TACs will be set as in option 1. From 2017 on, flexibility will be introduced in the Fmsy estimates by the introduction of ranges of Fmsy values consistent with MSY. Decisions in Council are supposed to be on TACs that keep the fishing mortality within Fmsy ranges. Where a stock is or fall below safeguard levels, the strategy would be to rebuild it above such levels in 5 years. Should time permits, STECF is requested to explore the consequences of extending that period to 10 years.

The landing obligation will apply as for option 1, but now the exceptions for survivability and *de minimis* can be applied.

In addition to the existing technical measures, additional measures may be introduced by the regionalisation process in order to minimise choke effects or to avoid catch of juvenile fish. These are to be taken as adopted one year after the entry into force of the plans and are to be considered 100% effective in their intended goals.

2.3.5 Number and scope of MAPs.

While initially two MAPs are conceived for western EU waters (essentially, bounded by the 48°N parallel), STECF is requested to examine the possible advantages and consequences of the following alternatives:

- i. A single MAP covering all fisheries operating on the Western EU waters
- ii. Separate MAPs covering fisheries in well characterised regions for STECF to determine on the basis of preliminary work already carried out (STECF Report 12-14).
- iii. Separate MAPs for the main groups of fisheries, covering all most important fishing activities. Those fisheries would be characterised by a reduced number of target species and a set of by-catch species. The main fisheries, chosen on the basis of work being currently undertaken by Member States on discard plans for demersal species, are set out as an appendix to this document

The above-mentioned exercise is to be based of qualitative expert judgement rather than on mathematical simulations.

2.3.6 Fishery approach

Within the alternative sub-option iii) above, STECF is requested to examine whether setting MSY-compatible TACs uniquely for the target species is sufficient to grant conservation effects on the by-catch species. These conservation effects are to be evaluated against MSY reference points and precautionary stock levels (Bpa) of the by-catch species. Where possible, STECF should explore whether appropriate combinations of F values for target species can be found so the by-catch production is maximised within their Fmsy constraints.

Where managing the target species gives insufficient conservation guarantee to by-catch species, STECF is also requested to assess the possibility of improving the conservation of by-catch species by adopting multi-species by-catch quotas.

2.3.7 List of stocks considered (provided by DGMARE)

A. Stocks for which fishing opportunities are set as part of the NWW MAP

a. Stocks for which Fmsy ranges can be provided (Cat. 1 and 2):

- Blue ling (*Molva dypterygia*) in Subdivision Vb, and Subareas VI and VII
- Cod (*Gadus morhua*) in Subarea IV (North Sea), Division VIIId (Eastern Channel) and IIIa West (Skagerrak)
- Cod in Divisions VIIe-k (Celtic Sea cod)
- Cod in Division VIIa (Irish Sea)
- Cod in Division VIa (West of Scotland)
- European sea bass (*Dicentrarchus labrax*) in Divisions IVbc, VIIa, and VIIId to h (Irish Sea, Celtic Sea, English Channel and southern N,Sea)
- Greenland halibut (*Reinhardtius hippoglossoides*) in Subareas V, VI, XII and XIV
- Haddock (*Melanogrammus aeglefinus*) in Subarea IV and Divisions IIIa West and VIa
- Haddock in Division VIb (Rockall)
- Hake (*Merluccius merluccius*) in Division IIIa, Subareas IV, VI and VII and Divisions VIIa,b,d (Northern stock)
- Megrim (*Lepidorhombus* spp) in Divisions IVa and VIa
- Nephrops (*Nephrops norvegicus*) in Division VIa (North Minch, FU 11)
- Nephrops in Division VIa (South Minch, FU 12)
- Nephrops in Division VIa (Firth of Clyde + Sound of Jura, FU 13)
- Nephrops in Division VIIa (Irish Sea East, FU 14)
- Nephrops in Division VIIa (Irish Sea West, FU 15)
- Nephrops in Division VIIb,c,j,k (Porcupine Bank, FU 16)
- Nephrops in Division VIIb (Aran Grounds, FU 17)
- Nephrops in Division VIIa,g,j (South East and West of IRL, FU 19)
- Nephrops in the Smalls (FU 22)
- Plaice (*Pleuronectes platessa*) in Division VIIId (Eastern Channel)
- Plaice in Division VIIe (Western Channel)
- Sole in Divisions VIIf, g (Celtic Sea)

- Sole in Division VIId (Eastern Channel)
- Sole in Division VIIe (Western Channel)
- Sole in Division VIIa (Irish Sea)
- Whiting (*Merlangius merlangus*) Subarea IV (North Sea) and Division VIId (Eastern Channel)
- Whiting in Division VIIe-k
- Whiting in Division VIa (West of Scotland)
- Haddock in Division VIIa (Irish Sea)

b. Stocks for which only Fmsy proxies can be provided (Cat. 3 and 4):

- Anglerfish (*Lophius budegassa*) in Divisions VIIb-k and VIIa,b,d
- Anglerfish (*Lophius piscatorius* and *L. budegassa*) in Division IIIa and Subareas IV and VI
- Anglerfish (*Lophius piscatorius*) in Divisions VIIb-k and VIIa,b,d
- Greater silver smelt (*Argentina silus*) in Subareas I, II, IV, VI, VII, VIII, IX, X, XII, and XIV, and Divisions IIIa and Vb (other areas')
- Haddock in Division VIIa (Irish Sea)
- Ling (*Molva molva*) in Divisions IIIa and IVa, and in Subareas VI, VII, VIII, IX, XII, and XIV ('other areas')
- Megrim (*Lepidorhombus* spp.) in ICES Division VIb (Rockall')
- Megrim (*Lepidorhombus whiffjagonis*) in Divisions VIIb-k and VIIa,b,d
- Nephrops in the FU 20 (Labadie) and FU 21 (Jones and Cockburn)
- Haddock in Division Vb
- Haddock in Divisions VIIb,c,e-k
- Saithe (*Pollachius virens*) in Subarea IV (North Sea) Division IIIa West (Skagerrak) and Subarea VI (West of Scotland and Rockall)
- Plaice in Divisions VIIh-k (Southwest of Ireland)
- Plaice in Divisions VIIf,g (Celtic Sea)
- Plaice in Division VIIa (Irish Sea)
- Sole in Divisions VIIh-k (Southwest of Ireland)
- Whiting in Division VIIa (Irish Sea)
- Tusk (*Brosme brosme*) in Divisions IIIa, Vb, VIa, and XIIb, and Subareas IV, VII, VIII, and IX (other areas)
- Tusk in Division VIb (Rockall)

c. Stocks for which Fmsy values or proxies cannot be determined (Cat. 5 and 6)

- Pollack (*Pollachius pollachius*) in Subareas VI and VII (Celtic Sea and West of Scotland)
- Saithe (*Pollachius virens*) in Subarea VII

B. Stocks for which fishing opportunities are set as part of the SWW MAP

a. Stocks for which Fmsy ranges can be provided (Cat. 1 and 2):

- Black-bellied anglerfish (*Lophius budegassa*) in Divisions VIIIc and IXa
- White-bellied anglerfish (*Lophius piscatorius*) in Divisions VIIIc and IXa
- Hake in Division IIIa, Subareas IV, VI and VII and Divisions VIIIa,b,d (Northern stock)
- Hake in Division VIIIc and IXa (Southern stock)
- Four-spot megrim (*Lepidorhombus boscii*) in Divisions VIIIc and IXa
- Megrim (*Lepidorhombus whiffiagonis*) in Divisions VIIIc and IXa
- Sole in Divisions VIIIab

b. Stocks for which only Fmsy proxies can be provided (Cat. 3 and 4):

- Anglerfish (*Lophius budegassa*) in Divisions VIIb-k and VIIIa,b,d
- Megrim (*Lepidorhombus whiffiagonis*) in Divisions VIIb-k and VIIIa,b,d
- Nephrops in Divisions VIIIa,b (Bay of Biscay, FU 23, 241)
- Nephrops in North Galicia (FU 25')
- Nephrops in West Galicia and North Portugal ("FIT 7,6-77)
- Nephrops in South-West and South Portugal (FU 28-29)
- Nephrops in Gulf of Cadiz (FI J 30)
- Nephrops in the Cantabrian Sea (FU 31)

c. Stocks for which Fmsy values or proxies cannot be determined (Cat. 5 and 6)

- Pollack in Division VIIIab
- Pollack in Division VIIIc
- Pollack in Division IXa
- Sole in Divisions VIIIc and IXa

2.3.8 List of fisheries with their target species (provided by DGMARE)

Table 1. List of fisheries included in the SWW MAP

Area	Species defining the fishery	Fishing gear
VIIIabde	Common sole	OTB between 70-100 mm GTR larger or equal to 150 mm

		BT larger or equal to 70 mm
VIIIabde	Hake	PTB larger or equal to 100 mm OTB larger or equal to 100 mm (20% limit of hake catches)FR LLS GNS larger or equal to 80mm ES 1
VIIIabde	Nephrops	OTB larger or equal to 70 mm*
VIIIc & IXa	Hake	PTB larger or equal to 70 mm*1 OTB larger or equal to 70 mm*1 GNS between 80-99 1 LLS *2
VIIIc & IXa	Nephrops	OTB larger or equal to 70 mm*
IXa	Common sole and plaice	GTR larger or equal 100 mm

*Only applies inside functional units

*1 Only applies to fishing days under effort regime for the southern hake recovery plan FRA

*2 (hook size, conger) ES

Table 2. List of fisheries included in the evaluation of the NWW MAP.

Area	Species defining the fishery	Fishing gear
VIa	Cod, Haddock, Whiting and Saithe	OTB, SSC, OTT, PTB, SDN, SPR, TBN, TBS, TB, SX, SV, all mesh sizes
VIa	Nephrops	OTB, SSC, OTT, PTB, SDN, SPR, FPO, TBN, TB, TBS, SX, SV, FIX, all mesh sizes
VI, VII	Hake	OTB, SSC, OTT, PTB, SDN, SPR, TBN, TBS, TB, SX, SV, GNS, GN, GND, GNC, GTN, GTR, all mesh sizes
VI, VII	Hake	GNS, GN, GND, GNC, GTN, GTR, all mesh sizes
VI, VII	Hake	LL, LLS, LLD, LX, LTL, LHP, LHM, all mesh sizes
VII	Nephrops	OTB SSC, OTT, PTB, SDN, SPR, FPO, TBN, TB, TBS, SX, SV, FIX, all mesh sizes
VIIa	Cod, Haddock, Whiting and Saithe	OTB, SSC, OTT, PTB, SDN, SPR, TBN, TBS, TB, SX, SV, all mesh sizes
VIIId	Common Sole	TBB
VIIId	Common Sole	OTT, OTB, TBS, TBN, TB, PTB, smaller than 100mm
VIIId	Common Sole	GNS, GN, GND, GNC, GTN, GTR, all mesh sizes
VIIId	Cod, Haddock, Whiting and Saithe	OTB, SSC, OTT, PTB, SDN, SPR, TBN, TBS, TB, SX, SV, all mesh sizes
VIIe	Common Sole	TBB, all mesh sizes
VII excl. VIIa; VIIId and VIIe for	Common Sole	TBB, all mesh sizes

Common Sole		
VII excl. VIIa; VIIId and VIIe for Common Sole	Common Sole	GNS, GN, GND, GNC, GTN, GTR, all mesh sizes
VII excl. VIIa; VIIId and VIIe for Common Sole	Cod, Haddock, Whiting and Saithe	OTB, SSC, OTT, PTB, SDN, SPR, TBN, TBS, TB, SX, SV, all mesh sizes

3 DESCRIPTION OF THE FISHERIES

This section describes the major fleets and stocks of each region with the Western Waters of the European Union, as well as the major oceanographic and geologic characteristics of the area. For more detailed information, refer Annexes I-IV, where a thorough description of the fleets and stocks can be found.

3.1 North Western Waters

The Celtic Seas comprise the shelf area west of Scotland (ICES Subarea VIa), the Irish Sea (VIIa), west of Ireland (VIIb), as well as the Celtic Sea proper (VIIIf-k) and western Channel (VIIe).

The variety of habitats in the Celtic Sea accommodates a diverse range of fish, crustacean and cephalopod species that support a wide variety of fisheries targeting different species assemblages. The Celtic Sea groundfish community consists of over a hundred species and the most abundant 25 comprise 99% of the total estimated biomass and around 93% of total estimated numbers (Trenkel and Rochet, 2003). This ecoregion has important commercial fisheries for cod, haddock, whiting and a number of flatfish species. Hake (*Merluccius merluccius*) and anglerfish (*Lophius spp*) are also fished across the whole area. The shelf slope (500-1800 m) comprises a distinct species assemblage, including roundnose grenadier (*Coryphaenoides rupestris*), black scabbardfish (*Aphanopus carbo*), blue ling (*Molva macrophthalma*) and orange roughy (*Hoplostethus atlanticus*), as well as deep-sea squalidae (sharks) and macrouridae. The major commercial invertebrate species is the Norway lobster (*Nephrops norvegicus*), targeted by trawl fisheries throughout the Celtic Sea. Common cuttlefish (*Sepia officinalis*) are also exploited in the Celtic Sea, whilst there is dredging for scallops and smaller bivalves in the western English Channel, Irish Sea and west of Scotland. Pot fisheries take place for lobster (*Homarus gammarus*) and edible crab (*Cancer pagurus*) in coastal areas of this region. The most commonly used gear types in the Celtic Sea are otter trawls, beam trawls, netters, dredges and pots.

The following maps (Figure 1) illustrate the spatial distribution of the catches of main targets species described in the Annex 2 and the catches per gear in the Celtic Sea, based on STECF catch data. Each statistical rectangle is split depending on the proportion of each species/gear catches and their size are proportional to the total amount of catches.




Figure 1 Spatial distribution of the catches of main targets species and catches per gear in the Celtic Sea, based on STECF catch data.

3.1.1 Celtic Sea proper (VIIe-k including Western Channel)

This is the dominant trawl activity in the Celtic Sea OTB and OTT and two major mesh size ranges 100-119 and 70-99mm codend. Within the DCR Level 6 métier OTB&OTT_DEF_70–99 there are two distinct métiers targeting mainly gadoids and benthic species (mainly anglerfish). The former has been declining in importance in recent years whereas the latter has become more important. The fleet targeting *Nephrops* is OTB&OTT_CRU_70–99. Again there are two distinct métiers recognized by WGCSE one focused almost exclusively on large volumes of small *Nephrops* (i.e. where *Nephrops* accounts for >60% of the landed weight) and one with more mixed *Nephrops* and demersal fish catches. The former is focused on the Celtic Sea deep or “Smalls” mainly whereas the latter is more spread out throughout the Celtic Sea where there is suitable habitat for *Nephrops*.

Beam trawl (TBB_DEF_70–99) targeting flatfish, operated and monitored by respectively UK, Belgium and Ireland. The distribution of the activity covers certain grounds where sole, anglerfish, cuttlefish and megrim are abundant and the seabed is suitable for beam trawling. This DCR level 6 métiers GNS_DEF_120-219_0_0 includes set gillnets mainly targeting anglerfish (*Lophius* spp.) and those targeting gadoids.

Common cuttlefish (*Sepia officinalis*) are also exploited in the Celtic Sea, whilst there is dredging for scallops and smaller bivalves in the western English Channel, Irish Sea and west of Scotland. Pot fisheries take place for lobster (*Homarus gammarus*) and edible crab (*Cancer pagurus*) in coastal areas of this region

The main gill (GN1) and trammel (GT1) nets effort are from the French and English fisheries. The GN1 effort is widely spread in the Celtic sea, but most the effort is close to the English and French shore (Figure 2.1-6). Both fleets mainly target demersal species including hake and pollack (*Pollachius pollachius*). The French fleet also targets for crustacean species (Spider crab and common crab). Also a Spanish small fleet (only 2 vessels) target hake operated in Divisions VII j and VIIk. A pilot survey in 2006 showed a discard rate < 5%, so discards sampling programme was not focussed on gillnets. There is an important Irish gillnet fishery targeting cod in VIIe between January and March. Much of this fishery is operated by vessels under 12m. The trammel net effort is less wide spread than the

gillnet fishery and most of the effort is carried out close to the Brittany coast. The targets species for this fishery are sole, anglerfish and crustaceans (Spider crab and common crab).

3.1.2 *Irish Sea (VIIa)*

The main gear in the Irish Sea is demersal trawls. Several sub fleets exist within this fleet. The largest of these are the otter trawls, with a small proportion of demersal seines. The otter trawl vessels of this fleet primarily utilize 80 mm mesh codends. The majority of this fleet belong to targeted Nephrops fisheries. Two main Nephrops fisheries exist in the Irish Sea, one in the East (FU14) and one in the West (FU15). These fisheries are generally seasonal and confined to the summer months although the season has been extending in recent years. A number of other species are caught in relatively low levels by this fishery, including cod, haddock, plaice, anglerfish, and to a lesser extent sole. Although relative landings of cod within this fishery are low compared with the quantities of Nephrops landed, this fleet's contribution to the total cod landed within the Irish Sea is generally high. A small proportion of the demersal trawl fleet utilizes 100–119 mm meshes and targets the traditional whitefish fishery. This takes a mixture of species, specifically cod, haddock and whiting which used to be an important fishery within the Irish Sea, but has declined to low levels since 2003 following the adoption of larger meshed gear.

A beam trawl fleet operates within this area and the majority of vessels employ meshes in the range of 80–89 mm. This fleet primarily targets flatfish species, plaice and sole in particular. There is also a fishery for ray species. These fisheries have bycatches of anglerfish, and low catches of cod, haddock and whiting. Gillnetting also occurs in the Irish Sea. However, this is a very small fleet within the Irish Sea, accounting for around 1% of effort. Effort is focused to the south/southwestern area of the Irish Sea and is a subsection of a larger fleet operating within the Celtic Sea. In addition there is some gillnetting activity around the Isle of Man, however this is minimal. The primary target of those operating in the southern area is cod. Low landings of other species including haddock, saithe and anglerfish also occur. In relation to mesh size, although a number of different ranges are used, 150–219 mm has dominated in the last couple of years, moving away from 100–149 mm which used to be the primary mesh range used.

3.1.3 *West of Scotland (ICES Subarea VIa)*

The demersal fisheries in Subarea VI are predominantly conducted by otter trawlers fishing for prawns (Nephrops); cod, haddock, hake, saithe, and whiting (gadoids); anglerfish and megrim. Other species including lemon sole, plaice, witch, red mullet, halibut, turbot and pollack form a proportionally small but valuable part of the catch. Trawlers may target a particular species assemblage in particular areas, but invariably catch some mixture of species. Generally one can consider there to be:

An inshore fishery targeting prawns (with smaller catches of gadoids). The fishery mainly uses trawls with a mesh size of 80mm although there is also some creel fishing. There are separate fisheries in the Minch, the Firth of Clyde and the Sound of Jura. These fisheries mostly involve Scottish vessels;

- A shelf fishery for the gadoids. This mainly involves trawls with a mesh size of 120mm. Scottish vessels predominate, with smaller numbers of vessels from Ireland, Northern Ireland, England, France, Spain and Germany;
- A fishery close to the shelf edge targeting anglerfish and megrim. This is mainly a trawl fishery involving Scottish and Irish vessels. In addition, French vessels catching anglerfish may be targeting saithe and other demersal species or fishing in deep water for roundnose grenadier, blue ling or orange roughy. Spanish and UK gillnetters and longliners, work along the shelf

edge targeting anglerfish, hake and ling but occasionally moving into deeper water to fish for deep-water sharks;

- A fishery at Rockall targeting haddock on the bank (<200 m) and anglerfish on the slope (>200 m). This is mainly a trawl fishery involving Scottish and Irish vessels, with sporadic involvement of Russian vessels on the southwest part of the bank that falls within international waters.

In addition to these main demersal fisheries, some inshore vessels on the west coast of Scotland turn to scallop dredging when Nephrops catches or prices drop. A seasonal sprat fishery often develops in the south Minch in November and December, which is targeted by vessels of all sizes (including those that usually target Nephrops).

3.2 South Western Waters

3.2.1 Bay of Biscay

Bay of Biscay (Figure 2 **Error! Reference source not found.**) is a highly productive system. It creates the perfect conditions to multispecies fleets to make use of this productivity.




Figure 2. Case study area: Bay of Biscay

More than 200 species are caught in the Bay of Biscay with 20 species contributing to 80% of the landings. Main species in value are sole, Nephrops, hake, monkfish and seabass. Bay of Biscay concentrates important mixed demersal French and Spanish fisheries of trawlers, netters and longliners with a high degree of technical interactions between fleets through species.

The fisheries in the Bay of Biscay are mainly managed through conservation measures imposed by the Spanish and French administrations. Sole, Nephrops, Hake and Monkfish are thus submitted to TAC

and quotas system, minimum landing sizes, technical measures (mesh sizes limits and selectivity measures), (EC Reg. No. 850/98 and 1239/98). Effort reallocation for the different fleets may be restricted by constraints in terms of TAC and national quotas consumption.

The demersal Spanish fleets operating in this area are composed of bottom trawlers, longliners and netters. These fleets are managed through TAC and TAE, apart from some other technical and physical measures. These two regulations (TAC and TAE) come from different origins.

The TAC was first implemented when Spain joined the EU in 1986. Setting TACs involves the fixing of maximum quantities of fish that can be caught from a specific stock over a given period of time. This operation requires cooperation among the various parties enabling those involved to come to an agreement regarding TACs and an allocation key for sharing them. The EU went on to share fishing opportunities in the form of quotas among Member States. A formula was devised to divide TACs according to a number of factors, including countries' past catch record. This formula is still used today, on the basis of what is known as the principle of 'relative stability' which ensures Member States a fixed percentage share of fishing opportunities for commercial species. Even if the share has been maintained stable over time, the growing scarcity of the key stocks has eroded significantly the fishing opportunities for these fleets.

The TAE is previous to the TAC regulation. In 1981 it was decided to list all the Spanish vessels operating in Divisions VIIIa,b,d and Sub-areas VI and VII, in order to create the access rights to these fisheries (a single fishing right per vessel). The idea was to maintain fixed these rights even if the number of vessel decreased. When Spain joined the EU the number of vessels in that list was close to 300 and the so-called "300 list" was created. These fishing rights became transferable by area.

Finally, concerning technical measures, some mesh size limitations and minimum landing sizes for some stocks have been implemented. Further information on how this fishery is managed can be found in [Iriando et al. \(2013\)](#), [Prellezo et al. \(2009\)](#) and [Prellezo \(2010\)](#).

In 2013, 792 French vessels operated in the Bay of Biscay demersal fisheries. It represented around 25% of the total French vessels operating in the Atlantic and 49% of the French vessels operating in the Bay of Biscay. The bay of Biscay French demersal fisheries total gross revenue was calculated at around 249 million euros in 2013, and total direct employment amounted to 2256 fishermen.

Most important species caught by French vessels, in value, in the demersal fisheries in the Bay of Biscay are Common sole (17%), Nephrops (10%), European hake (10%), monkfishes (9%), Common cuttlefish (4%) and Sea Bass (4%) (percentages of the total gross revenue for those fleets).

Two main fleets of bottom trawlers and netters operate in these fisheries among which several strategies and specializations are observed. A fleet typology was developed together with stakeholders in the framework of the partnership bio-economic working group (PBEWG) and the European GEPETO project to provide a more detailed approach than DCF segmentation of fleets' situation, strategies and potential impact of management plans (Figure 1). 21 fleets were considered in the analysis (see table 1). These fleets are subsets of DCF fleet segments. Hereafter, Sole gillnetters, mixed gillnetters, Nephrops trawlers, Mixed demersal and Mixed demersal coastal trawlers, hake longliners and hake gillnetters are considered, each fleet being divided in vessel length (VL) categories. 3 fleets can be considered as small scale fleets (SSF) according to EC definition (Vessels <12m using passive gears exclusively). These SSF represent 38% of the vessels number, most of them being Sole gillnetters.

Main fleet segments in terms of vessels are mixed bottom trawlers (210 vessels) and mixed netters (263 vessels). Nephrops trawlers account for around 150 vessels and sole netters for around 130

vessels. Hake specialized fleets (longliners and gillnetters) only account for around 30 vessels but concentrates a large part of the French landings.

The main fleets in terms of gross revenue are the Nephrops trawlers (specialized) VL1224 (12% of the total gross revenue of French demersal fleets in the Bay of Biscay), the Hake gillnetters VL1840 (12%), the Mixed demersal trawlers North Bay Biscay_VL1824 (10%) and the Sole gillnetters_VL1218 (10%).

The Bay of Biscay demersal fisheries are complex mixed fisheries with high level of technical interactions between fleets through species caught by different fleets and joint productions. Joint productions occur at the trip and métier level for a given season and area. Estimation of production functions and joint production thus requires disaggregated data. At the year level, mixed production of fleets observed can result from practicing different métiers along the year and in different area.

As expected, data highlight that trawlers have more multi-species catches than netters or longliners with as a consequence less ability to reconcile catches.

Landings by fleet show that fleets operating on sole also catch hake but in different proportions according to fleets: For Sole gillnetters VL1824, hake represents 11% of their total landings (17% of their total value) while it represents 1% (4%) and 4% (5% in value) for Sole gillnetters VL1012 and VL1218 respectively. For Nephrops trawlers (specialized) VL1224, Nephrops is the first species (55% in value) but Hake or Sole appears to be significant level in the landings

For specialized fleets on hake (hake longliners and gillnetters), sole catches are not observed. Those fleets don't operate on the distribution area for sole which is more coastal.

Analyses of landings by fleet-métier enable to precise correlations between species and show in particular that sole landings by netters are due to trammelnet métier for sole while catches of hake by the same fleets are due to gillnet métier. There is thus ability for these fleets to reconcile both objectives while hake and sole (and other species) are caught by same bottom trawlers métiers. Proportion of species varies however according to main strategies of bottom trawling (demersal trawl cephalopods, Nephrops, sole or anglerfish) which also correspond to different spatio-temporal allocation of the effort.

3.2.2 *Iberian waters*

The Atlantic Iberian waters (ICES Divisions VIIIc and IXa) include three areas with different oceanographic characteristics: Gulf of Cadiz with Mediterranean influence, Atlantic front under a strong upwelling process, and Cantabrian Sea (southern area of Bay of Biscay). They include the transition between subtropical and sub-polar areas. Politically, the Atlantic Iberian waters are divided into the Spanish and Portuguese national waters. The current analysis of the Iberian waters only considers the Atlantic front and the Cantabrian Sea.

Vessels that operate in Atlantic Iberian waters belong to the national fleets of Spain and Portugal. Therefore, the vessels fishing Iberian stocks (ICES VIIIc and IXa) have to apply for a fishing licence to operate in the respective National waters. Both countries classify their national vessels in fleet categories depending of the gear type (trawl, purse seine, gillnet or longline), and both countries leave an independent group for the small-scale fleet.

These fleets operate on a narrow continental shelf where they exploit a variety of fishing resources by using different type of gears (trawl, gillnet, long lines...), forming a common demersal mixed-fisheries fleet. Although recent changes in fishing strategies and gears design have led some traditional demersal fleets to also exploit pelagic species, is not simple the combined management of demersal and pelagic stocks. On the one hand, most of the landings of pelagic stocks are made by fleets (purse seine, hand lines...) without any effect on demersal stocks. On the other hand, the populations of large

pelagic species usually inhabit wide oceanic areas, so their life cycle is developed beyond the geographical limits of the case study.

4 METHODS AND DATA

4.1 Addressing the ToRs

In order to address the questions asked by the request to STECF, the work was organized around three issues/subjects/questions:

- Which are the potential changes in the EU fisheries under the different scenarios set by the request (items 3.1-3.3 of the request).
- Which are the advantages and consequences of different configurations of the MAPs with relation to their spatial scope (item 3.4 of the request).
- Will management applied to the driver species be able to constraint the catches of the non-driver species (item 3.5 of the request). The second part of this item was lifted from the EWG1502 report (STECF, 2015).

Although the ToRs provided a list of species and fisheries, the time available to include them in FLBEIA and IAM was too short. It's important to note that adding species and fisheries to a simulation algorithm is not a trivial process, which may require several months of work.

4.1.1 *Evaluating scenarios using quantitative methods*

Following the best practices in the field of scientific policy advice, the evaluation of the regulation proposal was carried out using simulation testing. For the purpose of this report recent developments in the modelling tools for fisheries management were used, produced under the European projects GEPETO, SOCIOEC, MYFISH and DAMARA; as well as the national projects Bio-economic partnership working group project funded by the French Directorate of Sea Fisheries and Aquaculture since 2009, ANR ADHOC project (2010-2014) funded by the French National Research Agency, SIMLO (FEP 04-2014-00650) and projects funded by the Directorate of Fisheries and Aquaculture of the Basque Government.

The ToRs set a number of questions that were not possible to approach using a single comprehensive model. The settings are complex and the forecasts require strong assumptions to be made, in particular the effect that the Landings Obligation (LO) will have on the fleets behaviour is very uncertain. On the other hand the removal of HCRs from the MAP legislation, introduced an extra level of complexity to be simulated, which was new for the current model frameworks and techniques.

The new framework for MAPs requires a shift in the analysis concepts, from a situation where scientists were required to assist policy makers designing a MAP by studying the trade-offs of candidate HCRs, to a situation where scientists are required to evaluate the added value of implementing a MAP when compared with a baseline. To deal with this new framework a new approach had to be developed in a very short time frame.

The EWG used several models available and defined the scenarios in forms that were expected to provide the necessary information to support the advice. The time frame available was very limited, which conditioned the possibility to test different options to implement the scenarios in each model.

To depict the trade-offs between the MAP and the baseline scenario, the results were presented in relative terms to the baseline. As such the figures present a direct comparison between management options.

4.1.2 *Employment and fleet dependency*

For the purpose of estimating how employment may be impacted by a change in FMSY, an analysis was undertaken to highlight fleets with “high” and “low” employment. Employment numbers were then combined with the economic dependency of each fleet segment (landings values of the selected species compared to the total landings values in FAO Area 27) to identify fleets that are likely to be impacted by the MAP while being at the same time large employers.

The data submitted by the MS for the Annual Economic Report (STECF, 2013) for the year 2012 was used. The aggregation used for the social and economic data does not allow the analysis to be performed separately by sub-region, neither at the detailed level of fishing activities or métiers used in the MAP simulations. For instances, the DTS group includes several types of trawlers (e.g. demersal trawlers, pair trawlers, beam trawlers), different mesh sizes and target species. These aggregations limited the conclusions that can be drawn concerning the fleets dependency on the MAP driver species.

The first step involved taking relevant data from the AER database, for the fleets operating in the Celtic Sea, Bay of Biscay and Iberian Waters, including: employment (total number employed), landings value for the MAP main species (hake, Norway lobster, sole, megrims, anglerfishes and pollack) and the total landings value at the FAO Area 27 (Northeast Atlantic). The economical dependency on the fishing activity in the WW on those target species was calculated. The final evaluation included the total employment for fleets operating in the area, focusing on the value of landings from these species compared with each fleet overall total landing values, in order to estimate fleet dependency on these stocks.

4.1.3 *Number and scope of MAPs*

To address this question, the EWG discussed which may be the main elements of a MAP. Based on the discussion the EWG elaborated on the pros and cons of each option. This ToR was addressed through qualitative expert knowledge. There was no quantitative support to this ToR.

4.1.4 *Management of by-catch (fishery approach)*

To explore the potential impact of management measures applied to the “target” species into the “by-catch” species, the EWG used, as in EWG 15-04 report. The rationale is that if caught together, a management reducing or increasing the TAC (effort) on one of the main species might impact the other species which part of the catch assemblage.

4.2 **Provisional Fmsy ranges**

One of the most important elements of the new MAPs is the list of Fmsy ranges for each stock considered by the MAP. In the case of the European Western Waters these values should have been provided by ICES. However, for the stocks in this area the ICES advice is scheduled for late 2015. As such, to carry on with the evaluation of the MAP proposals, the EWG computed provisional Fmsy ranges which try to keep the fundamental concepts required by DGMARE, the fishing mortality ranges that produce 95% of the estimated catch when the stock is fished at MSY levels. Annex VI presents three working documents detailing the methodologies used. The values are presented in Table 3.

Table 3. Values of Fmsy, and their lower and upper range, as used in the analyses.

Stock	Fmsy	Lower limit	Upper limit	Method
Hake (south)	0.24	0.17	0.36	YPR (WD: Abad et.al)
Hake (north)	0.27	0.18	0.37	PLM (WD: Jardim)
Horse mackerel (south)	0.11	0.08	0.16	PLM (WD: Jardim)
Megrim (south)	0.17	0.08	0.19	YPR (WD: Abad et.al)

Sole (Bay of Biscay)	0.26	0.17	0.36	PLM (WD: Jardim)
Blue whiting	0.30	0.20	0.41	PLM (WD: Jardim)
Four spot megrim (south)	0.17	0.11	0.24	PLM (WD: Jardim)
Horse mackerel (western)	0.13	0.09	0.18	PLM (WD: Jardim)
White anglerfish (south)	0.19	0.13	0.26	PLM (WD: Jardim)
Haddock (VIIb-k)	0.40	0.26	0.60	EqSim (WD: Gerritsen and Lordan)
Cod (VIIe-k)	0.40	0.27	0.55	EqSim
Whiting (VIIe-k)	0.32	0.21	0.44	PLM (WD: Jardim)
Sole (VIIfg)	0.31	0.21	0.43	PLM (WD: Jardim)
Plaice (VIIfg)	0.3	0.21	0.43	PLM (WD: Jardim)

The current Fmsy value (0.26) set for the Bay of Biscay sole is based on Fmax, as estimated during WGHMM 2010 (ICES, 2010). The technical basis for this choice relies mainly on the fact that there is no clear stock-recruitment relationship for this stock. ICES notes that this value is ill defined (ICES Advice 2014, book 7) as the current Fmax (0.46 as estimated in 2014) is higher than the one that was calculated using the 2010 data. ICES considers that the basis for FMSY may need to be re-evaluated. Several attempts at estimating it have been made by the ICES WGBIE working group in 2014 and 2015 without success. ICES will again consider this issue during a workshop on Fmsy ranges for western waters stocks scheduled for the fall of 2015.

4.3 Multi-model approach

The scope of the MAPs was too wide to be addressed by any of the models currently available, which included all demersal fisheries and stocks in the South Western Waters (Bay of Biscay and Iberian waters) and the North Western Waters (West of Scotland, Celtic Sea, Irish Sea and Western Channel). As such existing models were further developed and calibrated to specific fisheries to analyse impacts of management measures at the regional level (Iberian waters, Celtic Sea and Bay of Biscay). The approach taken by the EWGs was to invite the scientists involved in modeling these areas to contribute to the evaluation. As a result three models were available; FLBEIA, IAM and a State-Space model.

These are all bio-economic models, although they're based in different modelling concepts. FLBEIA is based on an MSE algorithm with yearly time steps, where the allocation of effort across fishing strategies (metiers) is based on historical effort allocation or on the attempt to maximize profit. Furthermore, total effort is restricted by the TAC advices. Production functions are based on Cobb-Douglas for stocks explicitly modelled or on linear relationship with effort for other species. IAM uses stochastic forecast with quarterly or yearly time steps for biological dynamics and yearly time steps for fleets' behaviours. Production functions are based either on Baranov equations for stocks explicitly modelled or on linear relationship with effort for other species. The IAM adjusts and reconciles effort by fleet and metier to meet F or TAC objectives. Allocation of effort across fishing strategies (metiers) was assumed to be based on historical allocation resulting from fishermen behaviours. Both, IAM and FLBEIA are multi-species, multi-fleets and multi-metiers models. The state-space model is a bio-economic multi species equilibrium model.

For the NWW area the EWG used an FLBEIA application to the Celtic Sea. For the SWW area there were an IAM application for the Bay of Biscay, focus on the French fleets operating in the area, an FLBEIA application to the Bay of Biscay, focus on the Spanish fleets operating in the area, and an

FLBEIA application to Iberian Waters covering all the relevant fleets. A summary of the scope and main concepts of these models is presented in Table 4.

Annexes I-V contain detailed descriptions of each of the models, their use in the analyses and additional results that were considered interesting.

4.4 Scenarios

Two sets of scenarios were investigated: the management scenarios and the fleet scenarios. The first relates to decision making options that were simulated to evaluate the trade-offs across options and inform decision makers of the effects/impacts that their decisions may have. The fleet scenarios aimed to inform on the likely responses from the fleets to the decisions taken. Such scenarios are the most difficult to forecast, as the reactions of the sector can vary widely and unexpectedly. Hence, the fleet scenarios are inevitably based on strong assumptions about likely responses, which may or may not be entirely accurate.

4.4.1 Management scenarios

The management scenarios were designed to evaluate whether a MAP with the characteristics proposed by DGMARE (see background), would be more successful at achieving the objectives set by Art° 2 of the CFP, than implementing the basic CFP provisions (baseline scenario).

The basic CFP provisions were interpreted as a situation where the current MAPs would continue to be applied and the CFP provisions added on top of those. The CFP provisions in this context are the LO flexibilities and the technical measures.

Technical measures were not possible to simulate. These measures will have to be implemented through co-decision with regional bodies and currently it's unknown which and to what extent these will be implemented. Time constraints didn't allow the EWG to explore through simulation this aspect. It would require a large number of scenarios to be run, in particular in the absence of guidance about which measures are likely to be implemented.

With regards to the LO, the interpretation was that inter-species and inter-annual flexibilities should be part of the baseline, while the *de minimis* and survivability exemptions should be part of the MAPs. However, due to time constraints, these rules were not implemented. There was a significant effort allocated to code these effects, nevertheless the results obtained were not satisfactory and were not included in the report. The LO is introduced in 2018 through the limitation of discards, and the uplift of the TAC to cover the total removals and not just landings.

The new MAP framework does not include HCRs, meaning that the Council has the freedom to decide on how it wishes to fix fishing opportunities and achieve the objectives of the CFP. The EWG was therefore faced with the problem of how to evaluate the provisions of the MAP in the absence of an HCR to derive a target fishing mortality rate. The EWG decided that the best alternative would be to use an "envelope" approach. Such approach considered the potential consequences of fishing at the limits (upper and lower) of the F_{MSY} ranges, to simulate both high and low exploitation cases, and thereby inform managers on the range of potential outcomes of alternative tactical management decisions, without giving advice about the 'best' way to get to the target.

Note that in this approach each scenario has two management options that lead to two simulations:

- upp – TAC_{Y+1} is set as the catch that results from exploiting the stock at $FMSY^{upp}$
- low - TAC_{Y+1} is set as the catch that results from exploiting the stock at $FMSY^{low}$

Table 4. Overview of the models used for the WW management plans evaluation

Western Waters models for ex-ante evaluation	FLBEIA IW	FLBEIA BoB	IAM	FLBEIA CS
Fishery description				
Multispecies (M) / Single species (S)	M	M	M	M
Seasonal				
Vessels LoA group				
< 12 m (small scale fishery)			X	x
12-24 m			X	x
24-40			X	x
>40 (long distance fishery)				x
Type of gear used				
passive	X	X	X	x
active	X	X	X	x
polyvaent	X		X	
Fleets disaggregation Level				
Economic fleet segments			X	
Métier 4 (gear type)	X	X	X	x
Model characteristics				
Optimisation			X	
Simulation	X	X	X	x
MSE	X	X	X	x
MSE - full feedback loop with stock assessment model	At present just time lag but it would be possible			At present just time lag but it would be
MSE - implementation error	Derived from mixed fisheries dynamics			Derived from mixed fisheries
Time step	Annual	Annual	Year (quarterly SS3 dynamics)	Annual N
Spatial (Y/N) in case of Y resolution (...)				Celtic Sea (VIIbc,e-k)
Spatial coverage (North Sea, Skagerrak (Sk), Eastern Channel (EC))	Iberian Waters	BoB	BoB	
Population dynamics				
Biological structure				
age (A)	X	X	X	X
size (S)				
biomass (B)	X	X	X	X
Processes: dynamic recruitment (Drec), growth (Gr), Migration (Mig)	Drec	Drec	Drec	Drec
Simulate recruitment failure (Y/N)	X	X	X	X
Fleet dynamics				
based on F (F) / effort (E)	E	E	E	E
selectivity (model or fixed)	In the present conditioning it is fixed but it can be modeled		m/f	fixed
Economic dynamics				
Price elasticity	In the present conditioning it is fixed but it can be modeled		potentially	conditioning it is fixed but it can
Costs	In the present conditioning it is fixed but it can be modeled		X	In the present conditioning it is
Employment or FTE	Variable, function of number of vessels		X	Variable, function of
Fuel costs	In the present conditioning it is fixed but it can be modeled		X	conditioning it is fixed but it can
MANAGEMENT OPTIONS (Yes/No/Development)				
De minimis	Available but not used		X	
Interspecies quota flexibility	Available but not used			
Swaps	Available but not used			
Borrow and banking	Available but not used			
ICES data limited stocks	HCR for DLS			X
F target	X	X	X	X
TAC & quotas	X	X	X	X
Biomass safeguards	X	X	X	X
Combined TACs (multiple species in one TAC)				
Diferenciated management between driver and non-driver stocks				
Multidimensional Fmsy ranges				
Harvest control rules	X	X	X	X
Temporary closure of fishery				
Area closures				?
INDICATORS (Yes/No/Development)				
Impact on biodiversity				
Abundance of main stocks	X	X	X	X
Evolution of main predator and prey stock				
Profitability	X	X	X	X
Income	X	X	X	X
Supply	X	X	X	X
Fuel consumption			X	
Employment	X	X	X	X
Compliance				
Stocks				
	Hake (south)	Hake (north)	Hake (north)	Cod VIIbc,e-k
	Horse Mackerel (South)	Megrim	Sole	Haddock VIIb-k
	Megrim	Blue whiting	Nephrops	whiting VIIbc,e-k
	4 Spot Megrim	H.Mackerel West	Monkfish	ole VIIfg
	Monkfish	Mackerel	Megrim	plaice VIIfg
	Blue whiting		Sea bass	Nephrops FU22
	Horse Mackerel (Western)		Pilchard	Anglerfish7&8
	Mackerel		Anchovy	
	Others		Mackerel	
			Horse mackerel	
			Pollack	
			whiting	
			Blue whiting	
			Rays	
			Cephalopods	
			Red mullet	

where $FMSY^{upp}$ and $FMSY^{low}$ are the upper and lower limits of the FMSY range, respectively.

In both cases the biomass safeguards were set at the precautionary biomass (Bpa). In the absence of an HCR to define the tactics to recover the stock, recovery was simulated as a linear increase in SSB up to the safeguard. There were two recovery periods simulated, 5 and 10 years, as requested by the ToR.

4.4.2 Fleet scenarios

The likely responses of the fishing sector to any management decisions are of major importance when forecasting potential stock and fleet impacts. The range of potential responses is very wide, which makes it extremely difficult to forecast. Although a large effort was allocated to modelling fleet response to management, the results obtained were not satisfactory. In most cases there were large differences with what was observed in the past, and the EWGs were not able to find justifications for such differences. Consequently, only one fleet behaviour was simulated, in which the fleets distribute their fishing effort in the same way they've done in the past, reflecting a strong inertia to change in face of the new management options.

4.4.3 Scenario summary

In summary, 1 fleet scenario and 3 management scenarios were investigated. Implementation of the provisions of the MAP comprised 2 options to perform the envelope analysis. Table 5 summarizes each scenario and how they were used to address the different ToRs.

Table 5. Summary of scenarios analysed.

Management scenario			Fleet scenario
Name	Runs	Description	Historical inertia
Baseline	cfp	Target:	Fmsy
		Time to target:	2016
		Landings obligations:	2018
MAP fast recovery	map.low	Target:	lower limit of Fmsy range
		Time to target:	2016
		Landings obligations:	2018
		Safeguards:	Bpa
		Recovery period:	5 years
	map.upp	Target:	upper limit of Fmsy range
		Time to target:	2016
		Landings obligations:	2018
		Recovery period:	5 years
MAP slow recovery	map10y.low	Target:	lower limit of Fmsy range
		Time to target:	2016
		Landings obligations:	2018
		Recovery period:	10 years
	map10y.upp	Target:	upper limit of Fmsy range
		Time to target:	2016

	Landings obligations:	2018
	Safeguards:	Bpa
	Recovery period:	10 years

4.5 Data

A summary of the data and parameters used to tune and condition the models is presented in Table 6. For more details check the model annexes (Annexes I-V).

Table 6. Summary of data and parameters used.

	FLBEIA IB	FLBEIA BoB	FLBEIA CS	IAM	State-space
Population dynamics	ICES 2013	ICES 2014	ICES 2012 or ICES 2014 (where available, truncated to 2012)	ICES 2014	ICES 2013
Fleet exploitation	GEPETO project	ICES 2014	STECF 2013	Ifremer/Fisheries Information System/DPMA 2013	ICES 2013
Fleet economics	STECF AER 2013	STECF AER 2014 Prices from AZTI DB	STECF AER 2013	DCF – DPMA 2013	STECF AER 2013
Fleet interactions	GEPETO	DCF – IEO 2013	STECF FDI (2013)		ICES 2013
Fmsy	ICES 2014 or estimated in meeting				
Bpa	ICES 2014	ICES 2014	ICES 2014	ICES 2014	
Employment	STECF AER 2014				

Processing of model outputs for final analysis and visualization was conducted using the FLR packages (Kell et al, 2007; <http://flr-project.org>) for the R language (R Core Team, 2015) version 3.1. This toolset is also employed by the software implementing the FLBEIA method.

5 ToR 3.1-3.3 - EVOLUTION OF EU FISHERIES UNDER DIFFERENT SCENARIOS

To compute the effects of the MAPs proposal, a set of simulations were run, implementing the options described by the ToRs. The results were reported for the years 2017, 2021, and 2025, as required.

The results were presented as ratios between the MAP proposal scenarios and the baseline scenario, as such focusing on the differences between the two options, fishing under the CFP provisions or under a MAP framework, which is the simplest way of showing the effect of the MAP. Note that a value of 1 means that there aren't differences between the MAP scenario and the baseline, a value <1 means that there was a reduction of the variable when compared with the baseline, e.g. a value of 0.5 in F would mean that F in the MAP scenario was half of the baseline F, and vice-versa for values >1.

This section is split into North Western Waters, further broke down in Celtic Sea (CS), and South Western Waters, further broke down into Bay of Biscay (BoB) and Iberian Waters (IW).

5.1 North Western Waters

5.1.1 State of the fisheries in 2017




Figure 3. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the NWW MAP in 2017, and for seven stocks in the area.

In 2017 the differences between scenarios are small, with the exception of TAC and quota uptake. TACs are expected to be lower in the case of MAP-low. Monkfish and Nephrops have fixed dynamics. Quota uptake in the MAP-low is higher than the baseline. Note that catches don't change much across scenarios, reflecting the inertia in effort allocations.




Figure 4. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the NWW MAP in 2017, and for 14 fleets operating in the area.

From a fleet perspective the results show a variety of effect among fleets. Most of the fleets are not sensitive to the MAP scenarios. The Spanish and UK fleets of demersal trawlers and seiners over 10m, and the UK and Irish fleets of TBB over 10m, show larger catches and effort in the MAP-upp scenario in relation to the baseline. The opposite trend occurs in the case of the MAP-low scenario.

5.1.2 State of the fisheries in 2021




Figure 5. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the NWW MAP in 2021, and for seven stocks in the area.

The outlook for 2021 under the tested condition shows how the different species incorporating in the model are more or less sensitive to the targeted FMSY value. Gadoids show high differences in terms of biomass, SSB than flatfish. However, catches of sole can vary quite extensively.




Figure 6. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the NWW MAP in 2021, and for fourteen fleets operating in the area.

In 2021, fishing at the upper limit of the Fmsy ranges will produce larger catches for most fleets, ~30%. The Spanish and Irish fleets of demersal trawls and seines over 10m don't show the same increase in catches.

5.1.3 State of the fisheries in 2025




Figure 7. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the NWW MAP in 2025, and for seven stocks in the area.

Catch inter annual variability are more stable than in the previous periods as well as catches of whiting and plaice. Catches of sole are still quite distinct depending on the scenarios. The mixed fisheries interaction are well illustrated by the lack of consistency between, TACs and catches over the studied period.




Figure 8. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the NWW MAP in 2025, and for 14 fleets operating in the area.

Some fleets are still not sensitive to the different MAP scenario. As before demersal trawlers and seiners over 10m from Spain, Ireland and the UK don't increase their catches when fishing at the upper limits of the Fmsy range.

Catches by fleet in the scenario MAP-upp seem to increase less in relative terms than effort, which may be reflected in larger costs by catch per unit of effort.

5.2 South Western Waters - Bay of Biscay

5.2.1 State of the fisheries in 2017




Figure 9. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, FLBEIA) in 2017, and for 13 stocks in the area.

The outlook for 2017 under the conditions described above shows that in relation to the baseline the biomass, ssb and biological risks are the same. The TACs, fishing mortality and catch are expected to be higher in the MAP-upp scenario and lower in the MAP-low scenario. As a consequence quota uptake is higher for the MAP-upp scenario and inter-annual variability of catches lower. This scenario affects the short term effects of decreasing fishing mortality to reach the target fishing mortality.

Figure 10. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, IAM) in 2017, and for three stocks in the area.

The results obtained by the IAM model are consistent with FLBEIA, showing a similar pattern. Risk to Bpa and Blim and SSB and biomass are the same (or very close) in each of the scenarios for nephrops

hake and sole. Fishing mortality and thus catches of hake, nephrops and sole are expected to be higher in the scenario upp than in the scenario low.




Figure 11. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, FLBEIA) in 2017, and for three Spanish fleets operating in the area.

From a fleet perspective the results show similar effects for all fleets, with larger catches for the scenario MAP-upp (~30%), requiring larger effort to be deployed (~30%) with relation to the baseline. The MAP-low scenario shows ~ 30% less for both indicators. Fixed costs are the same, once that the number of fleets in the fishery doesn't change. The economic indicators are all very similar reflecting mainly a scaling of catches (revenue) and effort (variable costs).




Figure 12. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, IAM) in 2017, and for 21 french fleets operating in the area.

With relation to the French demersal fleets operating in the Bay of Biscay, results show two groups of fleets:

- fleets specialized on hake (larger netters and longliners) that are not choked by sole and can increase or decrease F (and thus landings, effort, value of landings and costs) according to hake Fmsy ranges;
- fleets catching hake and sole for which F is constrained by sole.

Decrease in effort, landings and revenue by fleet observed is the results of the decrease of effort by fleet and métier according to reconciliation and fleets joint productions. As a result, we observe in the simulations that fleets specialized on hake (hake gillnetters and longliners) have lower decrease in effort than other fleets catching also for sole. Variability of impacts on hake fleets compared to baseline is lower than for other fleets. Economic impacts (positive or negative) will largely depend on fishing possibilities decided in the range of possible F.

Mixed bottom trawlers (12-18m in particular), characterized by mixed productions with low ability to adjust species correlations, would be the most impacted fleets, assuming no possible reallocation of effort and constant number of vessels. 2017 is the first year of transition phases and thus characterized by high negative impacts of scenarios compared to initial situation before recovery.

5.2.2 State of the fisheries in 2021




Figure 13. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, FLBEIA) in 2021, and for 13 stocks in the area.

Biomass is expected to be lower in the MAP-upp scenario and higher in the MAP-low scenario with relation to the baseline. The catches of hake and megrim are expected to be lower for both scenarios. In the case of hake there will be more catches in the MAP-low scenario than in MAP-upp. In the case of TACs, it is expected to be lower in both scenarios for hake and higher for megrim in the MAP-low scenario.




Figure 14. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, IAM) in 2021, and for three stocks in the area.

The results obtained by the IAM model in 2021 show higher SSB and biomass of sole, nephrops and hake the MAP-lowscenario as a result of decrease in F. Risk to Blim and Bpa are however not increased. Results show that F objectives for hake and sole are reached (or almost) for all the scenarios. It highlights that reconciliation of objectives for sole and hake is possible. Modeled fleets only account however for a part of the fishing mortality on hake and that choke effects for other non-explicitly modeled fleet is not taken into account. In the case of nephrops or sole, modeled fleets account for more than 90% of the total mortality on those stocks.

Decrease in F observed for nephrops is to be linked with management objectives for sole and hake, however it should be underlined that correlations between nephrops and sole are modeled in this application at the fleet-métier level and that spatio-temporal allocation of effort can modify correlation between species. Distribution of sole and nephrops landings thus show that landings of both species have low overlap and that there is possibility for fishermen to catch both species almost separately. Nephrops and hake are more joint by the spatial distribution of both species.




Figure 15. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, FLBEIA) in 2021, and for three fleets operating in the area.

The number of vessels is expected to be lower in the MAP-low scenario for Spanish longliners.

Revenues for the three fleets will be lower for both scenarios, reflecting the decrease in catches.




Figure 16. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, IAM) in 2021, and for 21 fleets operating in the area.

In 2021, the previous differences found between Flow and the baseline tends to decrease, as yields of Flow scenarios tend to increase due to the biomass increase. Improvement of fleets' performance compared to 2017 is observed for the Nephrops fleet, due to the biomass recovery (and assumption of no TAC constraints).

5.2.3 State of the fisheries in 2025




Figure 17. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, FLBEIA) in 2025, and for 13 stocks in the area.

The TAC for megrim is expected to be 60% higher for MAP-low scenario comparing with the baseline. As in 2021, the biomass is expected to be higher for the MAP-low scenario and lower for MAP-upp scenario, always above Blim and Bpa.




Figure 18. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, IAM) in 2025, and for three stocks in the area.

Trends observed in 2025 are similar as observed in 2021. Differences in Biomass, SSB and Catches between Fupp and Flow scenario compared to baseline are increasing due to fishing mortality applied to stocks.




Figure 19. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, FLBEIA) in 2025, and for three fleets operating in the area.

In 2025 the differences between scenarios in economic indicators at fleet level are very similar to those observed in 2021.




Figure 20. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Bay of Biscay, IAM) in 2025, and for 21 fleets operating in the area.

The trend observed in 2021 is still observed in 2025.

The results highlight the high ranges of possible economic impacts according to the scenario adopted and TAC decided within the envelope of possibilities. Assumptions of the models need to be kept in mind when analyzing the results. The constant number of vessels and absence of reallocation of effort limit the results obtained.

Differences in impacted fleets according to scenarios, depend mainly on:

- métiers by fleet, reconciliation process and choke effects assuming no possible reallocation of effort
- dependence and contribution to the different species managed and ability to benefit from stocks recoveries.

5.3 South Western Waters - Iberian Waters

5.3.1 State of the fisheries in 2017




Figure 21. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Iberian Waters, FLBEIA) in 2017, and for nine stocks in the area.

The outlook for 2017 under the conditions described above shows that in relation to the baseline the biomass, ssb and biological risks are the same. As expected the TACs, fishing mortality and catch are higher in the MAP-upp scenario and lower in the MAP-low scenario. The quota uptake depends on the stocks, it is higher under MAP-low scenario for the stocks with dynamic (HKE, HOM, LDB, MEG and MON) and under MAP-upp scenario for the rest.




Figure 22. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Iberian Waters, FLBEIA) in 2017, and for five fleets operating in the area.

From a fleet perspective the results show fleet dependent effects. The effort and catch for Portuguese trawlers are similar in the three scenarios, but for the rest of the fleets the effort and total catch is up to 30% higher in MAP-upp scenario and around 25% lower in MAP-low scenario. The differences in landings are lower than in catch for all the fleets except for polyvalent gear fleet for which the differences are similar. Fixed costs are the same, once that the number of fleets in the fishery doesn't change. The economic indicators are in general higher in MAP-upp scenario but there are cases where the indicator is higher in MAP-lo scenario. For Spanish vessels using Hooks and Lines (HOK_SP) and Spanish fixed nets (DFN_SP), most economic indicators are higher in MAP-low scenario.

5.3.2 State of the fisheries in 2021




Figure 23. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Iberian Waters, FLBEIA) in 2021, and for nine stocks in the area.

In 2021 the differences in biological indicators between scenarios are higher. As expected biomass is always higher in MAP-low than in baseline scenario (>5-15%), and lower in MAP-up scenario (up to 10%). Similar trends are observed for SSB. On the contrary fishing mortality is lower in MAP-low scenario (< 10-20%) and higher in MAP-up scenario (~40% higher) than in baseline scenario for all the stocks except for Horse Mackerel (HOM)). The risk of SSB falling below Blim is null for all the stocks and the risk of falling below BPA is positive only for Horse Mackerel. For this stock the probability in MAP-up scenario is double than in the baseline scenario and in MAP-low scenario is 75% lower. Total catch is up to 30% higher in MAP-up scenario and in MAP-low scenario catch is only lower than in baseline scenario for Horse Mackerel (~15% lower). Inter-annual variability is always higher in MAP-low scenario. For stocks without dynamic the variability comes exclusively from the effort exerted by the fleets annually and for the rest is a product of the TAC advice. For the stocks with annual advice the variability is lower than for the rest of the stocks, 25% higher than in baseline in MAP-low scenario and 10% lower in MAP-up scenario. Quota uptake depends greatly on the stock and scenario. The TAC is always higher in MAP-up scenario and lower than in the baseline in MAP-low scenario for all the stocks except for Hake.




Figure 24. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Iberian Waters, FLBEIA) in 2021, and for five fleets operating in the area.

In 2021 all the indicators except fixed cost and number of vessels follow similar trends. They are all around 30% and 40% higher in the MAP-upp scenario than in the baseline. Regarding the MAP-low scenario for Spanish fleets (DFN_SP, DTS_SP and HOK_SP) the differences with the baseline are minimal. For the Portuguese fleets the differences are between 20% and 25%. In Spanish fixed nets (DFN_SP) and Portuguese polyvalent gears (PGP_PT) fleets the number of vessels is the same in all scenarios and so are the fixed costs. For Spanish Drift and Fixed Nets (DFN_SP) the number of vessels in the MAP_upp scenario does not change while for Spanish trawlers and Hook & liners (DTS_SP and HOK_SP) the number of vessels is 3% higher. In the MAP_low scenario the number of vessels decreases up to 5% in Spanish trawlers fleet (DTS_SP).

5.3.3 State of the fisheries in 2025




Figure 25. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Iberian Waters, FLBEIA) in 2025, and for nine stocks in the area.

In 2025 the differences between scenarios in Fishing mortality and biomass related indicators are similar to the differences in 2021. In the indicators related with catch the differences are in general higher. The catch in the MAP-low scenario is 10% to 20% lower than in the baseline and in the MAP-upp scenario from 10% to 30% higher. The differences in inter-annual variability are much higher than in 2021, trends also differ depending on the stock and scenario. For example, for Hake the inter-annual variability is 5 times higher in the MAP-upp scenario and for Megrim 6 times higher in the MAP-low scenario. The quota uptake is in the same range as in 2021. In 2025 the quota uptake in both MAP scenarios is similar and slightly higher than in the baseline. The differences in TAC are similar to those observed in 2021 except for hake. In 2025 the TAC for hake is 10% lower in the MAP_low scenario and 10% higher in the MAP_upp scenario than in the baseline.




Figure 26. Ratios of various indicators for the upper and lower MSY ranges against the baseline (CFP) scenario, for the SWW MAP (Iberian Waters, FLBEIA) in 2017, and for five fleets operating in the area.

In 2021 the differences between scenarios in economic indicators at fleet level are very similar to those observed in 2021. The higher differences between these two years are observed in number of vessels and fixed costs. In 2025 the number of vessels in Spanish fixed nets fleet (DFN_SP) is slightly higher than in the baseline in MAP_upp scenario and slightly lower in MAP_low scenario. In the case of Portuguese trawlers (DTS_PT) the difference between MAP_upp and baseline increases in 2025 up to 2.5%. For the rest of the fleets the differences still the same as in 2021.

5.4 Employment and Dependency in the NWW

A dependency index of the fleets fishing in the North Western Atlantic waters (areas 27.6 and 27.7) for the main demersal stocks was calculated for 2012. The index was estimated by country and fleet segment (main fishing technique + vessel length). Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), European hake (*Merluccius merluccius*), Norway lobster (*Nephrops norvegicus*), European plaice (*Pleuronectes platessa*), common sole (*Solea solea*), whiting (*Merlangius merlangus*), megrims (*Lepidorhombus whiffiagonis* and *Lepidorhombus spp.*), and monkfishes (*Lophius piscatorius*, *Lophiidae*, and *Lophius spp.*) were previously identified as main demersal target species for the fleets fishing in the area of study.

The dependency index identifies the importance of a species from an economic point of view for a fleet. The index is built by dividing a species value of landings from a fleet segment by the fleet segment's total value of landings.

We extracted the 2012 landings weight and value for these target species and totals by fleet segment from the 2014 Annual Economic Report (STECF, 2014) to estimate the indicator.

Table 7 presents the fleet segments with the largest number of fishers employed in the Northeast Atlantic waters (area 27). The highest employment in Northeast Atlantic waters can be found with the <10m fleets, which would be expected, given the nature of these fleets; 7 of the top 10 fleets are <10m. Note that there are a number of fleets that didn't have all the information required and as such was not possible to include them in this analysis.

Table 7, Top 10 higher employment fleet segments in area 27

country	gear	Lenght	Total employed fishers
ESP	PGP	VL0010	4223
ESP	DRB	VL0010	4013
PRT	PMP	VL0010	2852
GBR	FPO	VL0010	2846
PRT	PGP	VL0010	2415
FIN	PG	VL0010	1834
ESP	DTS	VL2440	1632
ESP	HOK	VL2440	1595
EST	PG	VL0010	1538
ESP	PS	VL2440	1123
Total in the area			74297

However, only 3 of these top 10 fleets catch the target species in areas 27.6 and 27.7: UK's vessels using pots and/or traps of less than 10 meters, and Spanish demersal trawlers and long-liners between 24 and 40 meters.

Table 8 presents the fleet segments with the largest number of fishers employed that catch the target species in the areas 27.6 and 27.7. It is also reported on the table the degree of dependency the fleets have on the catch of these target species in the area.

Table 8. Top 10 higher employment fleet segments in area 27.6 and 27.7, and dependency indicator.

country	gear	length	Fishers employed	Dependency target spp.
GBR	FPO	VL0010	2846	15.6
ESP	DTS	VL2440	1632	23.1
ESP	HOK	VL2440	1595	32.2
GBR	DTS	VL1824	1080	30.7
GBR	DFN	VL0010	1011	33.8
GBR	DTS	VL1218	971	59.7
GBR	HOK	VL0010	860	2.8

GBR	DTS	VL2440	798	23.0
FRA	DTS	VL1824	783	38.3
NLD	TBB	VL40XX	734	0.0
Total general			74297	12.9

This table shows that most of the fleet segments with the largest number of fishers employed that catch the target species in the areas 27.6 and 27.7 have a high degree of dependency. Consequently, these target demersal species constitute a key source for their revenues. Significant decreases on these target demersal species landings, and so revenues would hamper the economic performance of these fleets and so their capacity to keep the current levels of employment.

Dependency and employment indicators for all fleet segments that catch the target demersal species are presented in the Annex VII. In the Annex, it is also reported the share each fleet catches the main demersal target species in area 27.6 and 27.7 in comparison to all area 27 catches.

5.5 Employment and Dependency in the SWW

4.4.1 High Employment Fleets

The highest employment can be found with the <10m fleets, which would be expected, given the nature of these fleets; 4 of the top 10 fleets are <10m, employing 13503 individuals. However, the dependency measure mostly indicates low dependency on the MAP target species in the SWW, except for the Spanish trawl fleet 24-40m and hook fleet 12-18m, which show moderate dependency, ~15%.

Table 9. Top 10 higher employment fleet segments with the number of employed people and dependency degree.

Fleet segment	Employment (No of employees)	Dependency (%)
ESP AREA27 PGP VL0010	4,223	2
ESP AREA27 DRB VL0010	4,013	0
PRT AREA27 PMP VL0010	2,852	0
PRT AREA27 PGP VL0010	2,415	5
ESP AREA27 DTS VL2440	1,632	14
ESP AREA27 HOK VL2440	1,595	8
ESP AREA27 PS VL2440	1,123	0
ESP AREA27 HOK VL1218	1,040	16
PRT AREA27 PS VL1824	1,002	0
ESP AREA27 PS VL1824	998	0

The demersal trawl fleets (DTS and TBB) also have high employment, with the top 10 fleets employing 6088 individuals (Table 4.4.3.2). Concerning dependency on the species in the MAP, the FRA trawl fleet 12-18m shows the highest dependency (40%), followed by the PRT fleets 18-24m and 12-18m (25% and 21%, respectively).

Table 10. Employment and dependency degree for the demersal trawl fleets

Fleet segment	Employment (No of employees)	Dependency (%)
---------------	---------------------------------	----------------

ESP AREA27 DTS VL2440	1,632	14
FRA AREA27 DTS VL1824	783	5
FRA AREA27 DTS VL1218	619	40
PRT AREA27 DTS VL2440	567	10
IRL AREA27 DTS VL1824	441	0
ESP AREA27 DTS VL1824	425	9
FRA AREA27 DTS VL2440	423	1
ESP AREA27 DTS VL40XX	404	1
PRT AREA27 DTS VL40XX	403	0
FRA AREA27 DTS VL1012	391	19
ESP AREA27 DTS VL1218	336	4
IRL AREA27 DTS VL2440	288	0
GBR AREA27 DTS VL40XX	203	0
PRT AREA27 DTS VL0010	180	6
IRL AREA27 DTS VL1218	179	0
BEL AREA27 TBB VL2440	166	7
FRA AREA27 DTS VL0010	133	14
PRT AREA27 DTS VL1218	66	21
PRT AREA27 DTS VL1824	53	25
PRT AREA27 DTS VL1012	36	5

4.4.4 Low Employment Fleets

23 fleets employ less than 100 individuals. The majority of these fleets belong to France (10) and Portugal (8).

The dependency on landings of the MAP target species varies significantly among gear type, boat length, and MS. The French fleet using other active gears (FRA MGO VL1012) presents the highest dependency (44%) and is one with the lowest employment. 7 fleets show dependency on the MAP target species between 10 and 25%. These include fleets from all length groups, using all type of gears, belonging to France (3), Portugal (3) and Spain (1).

Six fleets have no dependency on the MAP target species. They are 2 fleets with dredges (PRT and ESP), 2 with drift or fixed nets (GBR and DEU), 1 using pots or traps (FRA) and 1 with pelagic trawl (DNK).

Table 11. Fleet segments with the number of employees < 100 and dependency degree.

Fleet segment	Employment (No of employees)	Dependency (%)
FRA AREA27 MGO VL1012	7	44
PRT AREA27 DTS VL1824	53	25
PRT AREA27 DTS VL1218	66	21
FRA AREA27 PGP VL1012	15	16
FRA AREA27 PMP VL1218	32	13
ESP AREA27 HOK VL0010	18	13
FRA AREA27 PGP VL0010	87	12
PRT AREA27 FPO VL1824	56	10

PRT AREA27 HOK VL1012	54	8
FRA AREA27 TM VL1218	58	8
FRA AREA27 TM VL1012	15	7
PRT AREA27 PGP VL1012	32	6
PRT AREA27 DTS VL1012	36	5
FRA AREA27 MGP VL0010	21	4
FRA AREA27 MGP VL1012	70	2
FRA AREA27 MGP VL1218	74	2
PRT AREA27 DRB VL1012	58	1
PRT AREA27 DRB VL1218	54	0
GBR AREA27 DFN VL1218	77	0
FRA AREA27 FPO VL1824	85	0
ESP AREA27 DRB VL1012	36	0
DNK AREA27 TM VL40XX	94	0
DEU AREA27 DFN VL2440	77	0

4.4.5 Highest Dependency Fleets

Table 4.4.5.1 presents the 13 fleets showing the highest degree of dependency on the MAP target species ($\geq 20\%$). The total number of people employed by these fleets is 3 780. These fleets include 9 using drift and fixed nets (FRA – 4, ESP – 3, PRT – 2), 3 trawl fleets (PRT – 2, FRA – 1) and 1 using other active gears (FRA). All length segments are involved.

Table 12. Fleets with highest dependency ($\geq 20\%$)

Fleet segment	Employment (No of employees)	Dependency (%)
ESP AREA27 DFN VL2440	117	51
FRA AREA27 DFN VL1824	278	48
FRA AREA27 MGO VL1012	7	44
FRA AREA27 DFN VL1218	330	41
FRA AREA27 DTS VL1218	619	40
PRT AREA27 DFN VL1824	351	35
ESP AREA27 DFN VL1824	342	32
FRA AREA27 DFN VL2440	327	31
PRT AREA27 DTS VL1824	53	25
ESP AREA27 DFN VL1218	588	23
PRT AREA27 DTS VL1218	66	21
FRA AREA27 DFN VL1012	579	20
PRT AREA27 DFN VL1012	124	20

STECF, 2013. The 2013 Annual Economic Report on the EU Fishing Fleet (STECF-13-15). Publications Office of the European Union, Luxembourg, EUR 26158 EN, JRC 84745, 302 pp.

5.6 Reconciling TACs by using FMSY ranges

A state-space model was designed to compute fishing mortality as the “endogenous” result of the fleets’ responses to the management measures. In the Annex V we show how fishing mortality interactions can be modelled as simple “state dependents” variables in a multi-fleet age-structured model. The model can be used to simulate the size of the fishing mortalities fluctuations around reference (equilibrium) targets.

The state-space multi-fleet model was calibrated to match 2012 $Fbar$ levels of the Southern Stock of hake for four species: hake, megrim, four-spot-megrim and monkfish. A vector autoregressive (VAR) model was used for the analysis of recruitments time series.

Two scenarios were simulated. In the first scenario, fleets’ behaviour was projected assuming that the response of the fleets will not be affected by management regulations. In the second scenario, the size of fluctuations is reduced to minimize (simultaneously) the distance of the four $Fbar$ values to the target.




Figure 27. Endogenous $Fbar$ fluctuations around the target. The y-axis and x-axis plot the deviation of $Fbar$ from the target and the time, respectively. The horizontal lines represent the $Fbar$ Ranges of each species.

Error! Reference source not found. shows the endogenous $Fbar$ fluctuations generated by the model. The y-axis plots the deviation of $Fbar$ from the target, and the x-axis, time. Note that differences in the recruitment variability (measured by the variance covariance matrix of the VAR process) and the existence of technical fishing interactions (captured in the state-space model) generate differences in the “natural” fluctuations of each species.

The figure clearly shows that a management procedure that tries to reconcile the several F targets simultaneously can be more successful achieving the single species targets defined in the CFP. Additionally it shows that the $Fmsy$ ranges will accommodate most of the natural fluctuation introduced by recruitment.

6 TOR 3.4 – NUMBER AND SCOPE OF MAPS

The MAP, as a strategic tool of the CFP, sets tactical objectives to achieve the CFP goals. The contents of the MAP, as defined in Art^o 10.1, can be grouped into measures that relate to the stocks (a-e,g) and measures that relate to the fleets (f).

The first set includes objectives regarding the exploitation of the stocks and risk-avoiding actions. The agreement between the Council, the Parliament and the Commission, translated those into Fmsy ranges, biomass safeguards and recovery periods for each stock (ref).

The second set is related with the implementation of landing obligations (Art^o 15) and should operate at a local level, adjusted to the fleet(s) dimension. These are technical measures, quota allocation and others measures to reduce unwanted catches.

From the point of view of the stock measures, having MAPs with a wider scope would limit both the number of stocks that will have to be split across regulations and the potential inconsistencies that may arise from having to make several regulations coherent. Nevertheless, it still remains largely a policy decision if the implementation of MAPs is better regulated by one, two or more regulations. As long as the objectives are still followed and the biomass safeguards applied, the outcomes of MAPs designed under the current framework, should not be impaired by their scope.

When considering fleet measures, the spatial scope is largely dependent on the fleet composition and the technical characteristics of the vessels. In such cases, having MAPs that focus on more homogenous regions, like the Bay of Biscay or Iberian waters, may encourage buy-in by Member States and regional/local bodies and establish a more homogeneous playing field.

7 TOR 3.5 – FISHERY APPROACH NWW

Landings discard rates data from the FDI database (STECF, 2013) were used to calculate the correlation between catches of species in the ICES divisions 7bcefg hjk. The correlations between target and by catch species are calculated on year 2003 to 2012. A threshold of ten tons of catches over the time period was applied to remove insignificant species. The data provides a detailed image of the catches and catch composition of the different gears operating in these areas.

Catches of cod, haddock and hake appear positively correlated with megrim (LEZ), anglerfish (ANF), surmullet (MUR), witch flounder (WIT), common mora (RIB), tope shark (GAG), plaice (PLE) horse mackerel (JAX), scallop (SCE), turbot (TUR) and herring (HER) (i.e. those with correlation superior to 0.5) and negatively correlated with smooth hound (SDV), tusk (USK), horse mackerel (HOM), Raja rays (SKA), gunards (GUX), catsharks (SCL), brill (BLL), John dory (JOD), rays (SRX), blackbanded trevally (RNJ), surmullets (MUX), Red gurnard (GUR), megrim (MEG), lobster (LBE), Porbeagle (POR), edible crab (CRE), common squid (SQC), spotted reay (RJM) and common cuttlefish (CTC) (i.e., those with correlation inferior to -0.55). Sole and saithe are weakly correlated to the other species catches (correlation coefficient rarely higher than 0.5 and 0.8 respectively).

Using the STECF database it was also possible to assess the yearly catch assemblage of the different gears and test for correlation between the level of catches of the target species and catches of the other species in the database. All targeted species and by catches do not appears for each gear type, which brings information on the species caught by each gear.

Table 13. Correlation matrix at the specie and area level. Target species are in column and by catch species in row. This is a subset of the matrix showing correlation coefficient higher that |0.5|.

	COD	HAD	HKE	NEP	POK	SOL	WHG
LEZ	0.91	0.78	0.86	-0.55	0.44	0.11	-0.27
ANF	0.84	0.66	0.81	-0.27	0.44	0.14	0.07
MUR	0.83	0.62	0.8	-0.46	0.63	0.35	-0.06
WIT	0.76	0.66	0.73	-0.45	0.88	0.59	0.09
RIB	0.72	0.88	0.71	-0.34	0.59	0.2	-0.22

GAG	0.68	0.69	0.78	-0.37	0.32	0.35	-0.15
PLE	0.68	0.62	0.62	-0.37	0.4	0.06	-0.24
JAX	0.61	0.75	0.51	-0.74	0.32	-0.16	-0.37
SCE	0.61	0.55	0.68	-0.35	0.34	0.63	-0.02
TUR	0.59	0.78	0.54	-0.45	0.82	0.35	-0.11
HER	0.5	0.29	0.48	-0.28	0.75	0.73	0.17
CTC	-0.55	-0.53	-0.47	0.52	0.12	0.35	0.55
RJM	-0.55	-0.54	-0.5	0.51	-0.03	0.26	0.35
SQC	-0.55	-0.55	-0.48	0.5	0.09	0.37	0.46
CRE	-0.55	-0.65	-0.43	0.72	-0.11	0.24	0.53
POR	-0.58	-0.47	-0.48	0.56	0.13	0.15	0.47
LBE	-0.58	-0.71	-0.48	0.68	-0.12	0.28	0.48
MEG	-0.59	-0.64	-0.5	0.62	-0.06	0.31	0.47
GUR	-0.59	-0.67	-0.45	0.66	-0.06	0.4	0.54
MUX	-0.6	-0.6	-0.43	0.68	0.03	0.33	0.61
RJN	-0.6	-0.6	-0.5	0.6	-0.05	0.28	0.38
SRX	-0.6	-0.62	-0.37	0.79	-0.04	0.14	0.54
JOD	-0.62	-0.66	-0.51	0.66	-0.07	0.29	0.42
BLL	-0.63	-0.68	-0.5	0.73	-0.12	0.24	0.48
SCL	-0.63	-0.7	-0.5	0.73	-0.13	0.25	0.49
GUX	-0.63	-0.74	-0.49	0.76	-0.2	0.25	0.46
SKA	-0.63	-0.77	-0.51	0.75	-0.24	0.21	0.59
HOM	-0.64	-0.63	-0.49	0.71	-0.17	0.12	0.29
LSK	-0.65	-0.77	-0.47	0.83	-0.33	0.14	0.33
SDV	-0.66	-0.75	-0.48	0.85	-0.27	0.12	0.42

Looking at a more detailed aggregation shows that the relationships between the target-bycatch dynamics are stronger at the fleet level than the stock or métier level. For example, at the species/TAC level the expected correlation between sole and plaice is very weak whereas it appears quite clearly for several gear (Table 14).

Table 14. Sole and Plaice correlation for different gears

	BT2	OTTER	TR1
PLE-SOL	0.3	0.72	0.7

Detailed analysis of this table illustrate the difficulty in management such a multi species and multi fleet fisheries. For example, management of the target species should positively impact anglerfish catches for trawlers (TR1, TR2, Otter and BT2) but the gillnets fleet it is the reverse Table 15.

As discussed in the EWG NSMAP 15-04 report, It should be noted that in the effort database, catches are aggregated over years, métiers and areas. However, fleets and species move during the year (changing fishing ground, spawning migrations, etc) which means that observed correlations might not reflect real technical interaction. Correlations between levels of catches of the main species and the “other species” presented here should be taken as indicative of the potential impact of management on species caught by the different gears.

Table 15. Correlation between catches of anglerfish with the main target species for different gears.

GT1

	HKE	WHG	COD	POK	SOL		
ANF	-0.43	-0.51	-0.25	-0.02	-0.11		
GN1							
	HKE	SOL	COD	POK	HAD	WHG	
ANF	-0.4	0.52	-0.14	-0.27	-0.28	-0.12	
TR3							
	HAD	WHG					
ANF	0.59	0.24					
TR2							
	COD	SOL	NEP	WHG	HAD	HKE	POK
ANF	-0.15	-0.21	0.38	0.57	0.42	0.68	0.61
TR1							
	HAD	COD	WHG	SOL	HKE	POK	NEP
ANF	0.89	0.9	0.72	0.47	0.82	0.58	-0.23
OTTER							
	POK	COD	HAD	WHG	HKE	SOL	
ANF	0.59	0.64	0.86	0.64	0.71	0.57	
BT2							
	HAD	COD	WHG	SOL	HKE		
ANF	0.77	0.39	-0.31	-0.42	-0.59		

8 TOR 3.5 – FISHERY APPROACH SWW

The data for the analysis was extracted from the STECF effort database (EWG 14-13: Fishing effort Part 2 [<http://stecf.jrc.ec.europa.eu/data-reports/>]). The availability and aggregation level of data have determined the selection of species and years for the correlation exercise. For the same reason, an adaptation of the list of fisheries for the SWW MAP detailed in the request to STECF has been done. It was not possible to analyze Pair and Otter-trawl fisheries separately because in the database they are aggregated under the Trawl category. It was also not possible to identify the fishery of plaice in IXa operated by trammel netters because no landing records for this species were found in the database.

In Iberian waters (VIIIc & IXa), at the stock level, there are very few correlations between the level of landings of hake and Nephrops and the rest of the species. Hake is only correlated with mackerel, the main contributor to landings (Table 16) and Nephrops is positive correlated with anglerfish (Table 17). Furthermore Nephrops landings present strong negative correlations with forkbeard, rays, squids and conger.

Table 16. Correlation between Hake landings and other species landings in Iberian waters.

MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
HKE	MAC	0.83	9455	41547
	JAX	0.02		33057
	WHB	0.19		20516
	ANF	0.20		2618
	SQC	-0.20		1930
	LEZ	0.40		779
	RAJ	0.32		422
	NEP	-0.03		299
	COE	0.02		280
	SOL	0.38		221
	FOX	0.30		191

Table 17. Correlation between Nephrops landings and other species landings in Iberian waters.

MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
NEP	MAC	0.16	299	41547
	JAX	0.38		33057
	WHB	0.23		20516
	HKE	-0.03		9455
	ANF	0.61		2618
	SQC	-0.64		1930
	LEZ	0.49		779
	RAJ	-0.68		422
	COE	-0.60		280
	SOL	0.28		221
	FOX	-0.69		191

The same analysis was carried out by fishery. In this analysis the aggregation of data at the fishery level possibly masks the potential correlation between target and by-catch species. The assemblages at this level do not correspond with métiers assemblages where fleets actually operate. Additionally the data is aggregated by year, which blurs the effects of seasonality. Only the results of some representative cases are showed in the report.

When looking at the correlations between hake and the other species in the trawl fishery, hake has weak positive correlations with mackerel and megrims (Table 18). It should be noted that hake and mackerel are targeted by two different métiers of the trawl fleet.

Table 18. Correlation between Hake landings and other species landings in Trawl fishery in Iberian waters.

Fishery	MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
Trawlers	HKE	WHB	0.25	5752	20224
		JAX	-0.09		16972
		MAC	0.56		15278
		SQI	-0.40		1627
		ANF	0.22		1495
		LEZ	0.52		764
		NEP	0.13		288
		RAJ	0.06		197

In the Bay of Biscay area (VIIIabde), at stock level, some correlations between hake, Nephrops and sole and the others were found. Hake correlates positively with seabass, megrims and blue whiting and negatively with cuttle fish, nephrops and red mullet (Table 19). Nephrops correlations are the opposite to hake ones, being positive correlated with red mullet, cuttlefish and squids (Table 20). Sole landings are correlated to sea bass and pollack, as it can be seen in Table 21.

Table 19. Correlation between Hake landings and other species landings in Bay of Biscay.

MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
HKE	JAX	-0.18	6705	9970
	MAC	-0.58		8753
	ANF	-0.11		4446
	CTC	-0.90		3493
	SOL	0.38		3115
	NEP	-0.83		2525
	BSS	0.61		1142
	MUL	-0.86		1078
	RAJ	-0.81		1056
	SQC	-0.80		905
	WHG	0.66		885
	POL	-0.32		757
	LEZ	0.85		547
	WHB	0.74		176

Table 20. Correlation between Nephrops landings and other species landings in Bay of Biscay.

MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
NEP	JAX	0.48	2525	9970
	MAC	0.59		8753
	HKE	-0.83		6705
	ANF	0.33		4446
	CTC	0.84		3493
	SOL	0.10		3115
	BSS	-0.22		1142
	MUL	0.97		1078
	RAJ	0.17		1056
	SQC	0.83		905
	WHG	-0.38		885
	POL	0.59		757
	LEZ	-0.60		547
	WHB	-0.73		176

Table 21. Correlation between Sole landings and other species landings in Bay of Biscay.

MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
SOL	JAX	0.37		9970
	MAC	-0.34		8753
	HKE	0.38		6705
	ANF	0.34		4446
	CTC	-0.12		3493
	NEP	0.10		2525
	BSS	0.72		1142
	MUL	0.13		1078
	RAJ	0.01		1056
	SQC	0.05		905
	WHG	0.51		885
	POL	0.65		757
	LEZ	0.51		547
	WHB	0.18		176

As in the case of Iberian waters, at the fishery level the correlations cannot identify the real interactions between species because the aggregation level of data does not match with fleets' activities. As an example, the trawl fishery with Nephrops as main species shows a strong negative correlation with hake (Table 22). In fact it is known there are several métiers of trawl targeting different demersal species.

Table 22. Correlation between Nephrops landings and other species landings in Trawl fishery in Bay of Biscay.

Gears	MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
Trawlers	NEP	ANF	0.46	2507	3415
		CTC	0.04		3108
		HKE	-0.72		2157
		SQU	0.25		1090
		RAJ	0.89		950
		MAC	-0.69		869
		SOL	0.47		812
		JAX	0.53		622
		MUX	0.62		607
		LEZ	-0.59		531
		PIL	-0.09		521
		WHG	0.03		463
		BSS	-0.23		352
		POL	0.89		208

Conversely, the analysis for the trammel net fishery in Bay of Biscay shows good correlations between sole and almost all of the by-catch species (Table 23).

Table 23. Correlation between Sole landings and other species landings in Trammel Net fishery in Bay of Biscay.

Fishery	MainSpecies	Bycatch	CorrCoef	LandMainSpecies	LandBycatch
TrammelNet	SOL	ANF	0.51	1749	391
		CTC	0.62		320
		BSS	0.81		260
		HKE	0.49		143
		POL	0.53		95
		RAJ	0.71		79
		WHG	0.93		66
		MUX	0.80		37
		MAC	0.21		12

In both areas, these results suggest that setting TACs individually for target species does not ensure conservation of the others species when global data by stock is analyzed. However, at the fishery level, different situations have been observed. A more detailed study per métier would allow detecting other important correlations.

Spatio-temporal allocation of effort by fleet and métier can modify correlations between species and tend to reconcile TAC.

9 TOR 3.5 – MULTI-SPECIES TACS FOR BY-CATCH STOCKS

A similar ToR was requested to the EWG 15 02, that evaluated the proposal of a MAP for the North Sea, and constituted the basis for the advice given by STECF (STECF, 2015). The EWG considered that the discussion and conclusions are still valid, and as such the text below is based on the work done by STECF (2015) with small edits.

In practice, grouping stocks already occurs in other areas. For example, in the North Sea there are grouped TACs for turbot and brill, for flounder and dab, and for lemon sole and witch flounder. Likewise, skates and rays are currently managed under a grouped TAC. The status for these stocks is generally estimated separately for the individual stocks, using one of the Data Limited Stock methodologies in ICES. Often, this means the stock status is assessed using survey trends.

In theory, the considerations on the sustainability of combined TACs are similar if several species are combined, or if several stocks of the same species are combined. In the North Sea, several stocks of *Nephrops* are combined into a single TAC. Examples of grouping TACs can also be seen in other areas. In the Northeast Atlantic for example, there are grouped species TACs for monkfish and megrim: the two species of monkfish sharing a single TAC, and two species of megrim sharing a single TAC.

One of the problems with addressing this ToR is the use of the term “by-catch”, without specifying exactly what it entails. There are many different definitions of “bycatch”. In the description of advantages and disadvantages of grouping quota that is given below, “bycatch” is defined as catches that are caught unintentionally while catching target species and target sizes. Bycatch can either be of a different species, or the undersized or juvenile individuals of the target species. However, what is a target species and what is a bycatch species depends on the fishery, and different vessels within a fleet may have different target species and bycatches. If combined TACs for so-called bycatch species are introduced, there will be a need to precisely define which species constitute the bycatch and this may need to be specified separately for different fisheries.

One of the **advantages** of combined TACs is that it provides increased flexibility for fishers to deal with the variability in bycatches. Hence catches within a quota can be substituted, so the species that potentially choke a fishery can be substituted by other species thereby allowing fishing on the target species to continue. Such increased flexibility could also improve the reporting of catches taken under the bycatch quota, because there would be less of an incentive to under- or mis-report the by catch species.

Furthermore, setting individual quotas for species that have until now been largely discarded is surrounded with a high level of uncertainty. Combining stocks may alleviate the problems with setting quota for such species individually, and create a buffer against uncertainty in the assessment and management of such stocks.

One of the **disadvantages**, by definition, is that combined TACs do not necessarily constrain the catches of individual species, because substitution between species subject to the combined TAC may take place. This could lead to overexploitation of some species, especially when combining vulnerable and invulnerable species.

The amount of substitution depends on several factors:

- the species composition and relative weight of those species in the bycatch: a large difference in the catch weights allows for easy substitution of a relatively large part of a small catch with a relatively small part of a large catch.
- the differences in net economic benefit (depending on price, and costs of exploitation) of the different bycatch species: a large difference in net economic benefit will generate an incentive to substitute lower value species with higher value species.

While one of the potential benefits of combined quotas is a reduction in the underreporting of catches, in the long run there is a risk of mislabelling of catches for pooled species that have a similar appearance and market price. This has previously been observed with anglerfish, skates and rays.

As mentioned above, to introduce combined TACs for bycatches, the terms “bycatch” and “target” need to be clearly defined, perhaps on a fishery or fleet basis. If vulnerability to overfishing of the by-

catch species that comprise the combined TAC is considered a flexible system in which the grouping is regularly evaluated. The costs of monitoring and managing such a system are likely to be high.

In order to mitigate the above disadvantages, the species composition of mixed-species TACs would need to be tracked to monitor the changes in the catchability and the vulnerability of the bycatch species to overfishing.

Combining species of different vulnerabilities that have large differences in price, and large differences in catch volumes should be avoided. There are a range of sources available for this information. For example, information on vulnerability indices by species (from Cheung et al. 2005, based on life history parameters) can be extracted from FishBase; prices can be found in the STECF Annual Economic report database; data on stock and catch status can be extracted from the STECF Consolidated Review of Advice and from ICES.

Finally, under a precautionary approach the combined-species TACs could be set lower than the sum of the individual species TACs to account for the increased risk of overexploitation of the individual species, due to the uncertainty associated with the conservation of the species grouped in a single TAC.

10 CONCLUSIONS

10.1 ToR 3.1-3.3

- Simultaneously managing a number of stocks at single species F_{MSY} levels is likely to fail and create inconsistencies between targets for different stocks.
- In the context of mixed fisheries, fishing opportunities can more easily be reconcile and made consistent with achieving the objectives of the CFP, using the flexibility provided by the F_{MSY} ranges.
- Adopting F_{MSY} ranges may increase the risk of overfishing if a decision is taken to persistently fish at the upper limits of the ranges. Taking into account the mixed fisheries constrains on matching the single species targets simultaneously, the benefits in terms of flexibility and adaptability would be lost, the probability of some stocks falling below B_{pa}/B_{lim} reference points may increase and the economic performance could be impacted negatively.
- Fishing at lower limits of the F_{msy} ranges generate larger SSB, lower catches and require less effort by fleet when compared with the baseline. The opposite pattern is observed when fishing at the upper limit of the F_{msy} ranges.
- The fleets that are responsible for most of the employment in this area don't seem to be very dependent on the species that will be regulated through this MAP.
- Biomass safeguards for all stocks should still be maintained and should provide a basic level of protection.
- Inter-annual catch constraints should be kept to stabilize inter-annual fishing opportunities.
- The scientific advisory process will have to be more focus on mixed fisheries.
- Sole safeguard and F reference point is likely to be reviewed by ICES.
- Horse mackerel is not achieving the F target and the safeguard is operating ~10% of the times, which may be an indication of inconsistencies in reference points.
- The evaluation of the Management Plan proposal provides a general comparison of the expected outcomes of managing the stocks using a MAP when compared with the basic provisions of the CFP.

- Knowledge about the mixed fisheries system is still partial and does not allow a full evaluation of the risks associated with all management options.
- Due to time constraints the models were not updated to incorporate all fleets and stocks that exist on the SWW.
- Not having an HCR introduces an extra level of uncertainty on future decisions. For the EWG work it represented a limitation on the capacity to simulate and evaluate the plan.
- The impact of the LO cannot be evaluated at this time due to the limited time available and the uncertainty associated with the implementation of the measures.

10.2 ToR 3.4

- The number and scope of MAPs is largely a policy decision. If the implementation is correct then the number and scope of the MAPs shouldn't impaired the achievement of the CFP objectives.
- Having larger MAPs may “promote” more coherent regulations in terms of objectives and safeguards for each stock and avoid over-regulating the sector.
- Having smaller MAPs increases the potential of over-regulating the sector but may promote more localized management measures and contribute to the involvement of stakeholders.

10.3 ToR 3.5

- Catch control measures over the species that drive the fisheries are not likely to drive the exploitation of non-driver species and as such will not guarantee the levels of conservation required by the CFP. Dynamics regarding the target species seem to occur at the fleet level.
- Grouping a number of single species TACs could introduce additional flexibility in the management of this system. However, the trade-off is that the potential to overexploit some stocks appears to increase. A set of mitigation principles were identified which should be considered if grouping of single species TACs is finally included in a management plan. Intense and strict monitoring will be essential to ensure that non-target species, or those less easily identified, are not overfished. The inclusion of fishing effort controls should also be considered in this case.

11 CONTACT DETAILS OF STECF MEMBERS AND EWG-15-04 & EWG-015-09 PARTICIPANTS

Information on STECF members and invited experts' affiliations is displayed for information only. In some instances the details given below for STECF members may differ from that provided in Commission COMMISSION DECISION of 27 October 2010 on the appointment of members of the STECF (2010/C 292/04) as some members' employment details may have changed or have been subject to organisational changes in their main place of employment. In any case, as outlined in Article 13 of the Commission Decision (2005/629/EU and 2010/74/EU) on STECF, Members of the STECF, invited experts, and JRC experts shall act independently of Member States or stakeholders. In the context of the STECF work, the committee members and other experts do not represent the institutions/bodies they are affiliated to in their daily jobs. STECF members and invited experts make declarations of commitment (yearly for STECF members) to act independently in the public interest of the European Union. STECF members and experts also declare at each meeting of the STECF and of its Expert Working Groups any specific interest which might be considered prejudicial to their independence in relation to specific items on the agenda. These declarations are displayed on the public meeting's website if experts explicitly authorized the JRC to do so in accordance with EU legislation on the protection of personnel data. For more information: <http://stecf.jrc.ec.europa.eu/adm-declarations>

STECF members:

Name	Address ¹	Tel.	Email
STECF members			
Abella, J. Alvaro (vice-chair)	ARPAT – AREA MARE Agenzia Regionale per la Protezione Ambientale della Toscana Articolazione Funzionale RIBM Risorse Ittiche e Biodiversità Marina Via Marradi 114, 57126 Livorno – Italia	Tel. 0039-0555- 3206956	alvarojuan.abella@arpat.toscana.it
Andersen, Jesper Levring (vice- chair)	Department of Food and Resource Economics (IFRO) Section for Environment and Natural Resources University of Copenhagen Rolighedsvej 25 1958 Frederiksberg Denmark	Tel.dir.: +45 35 28 68 92	jla@ifro.ku.dk
Bailey, Nicholas	Fisheries Research Services Marine Laboratory, P.O Box 101 375 Victoria Road, Torry Aberdeen AB11 9DB UK	Tel: +44 (0)1224 876544 Direct: +44 (0)1224 295398 Fax: +44 (0)1224 295511	baileyn@marlab.ac.uk n.bailey@marlab.ac.uk
Bertignac, Michel	Laboratoire de Biologie Halieutique IFREMER Centre de Brest BP 70 - 29280 Plouzane, France	tel : +33 (0)2 98 22 45 25 - fax : +33 (0)2 98 22 46 53	michel.bertignac@ifremer.fr
Cardinale, Massimiliano	Föreningsgatan 45, 330 Lysekil, Sweden	Tel: +46 523 18750	massimiliano.cardinale@slu.se
Curtis, Hazel	Sea Fish Industry Authority 18 Logie Mill Logie Green Road Edinburgh EH7 4HS	Tel: +44 (0)131 558 3331 Fax: +44 (0)131 558 1442	H.Curtis@seafish.co.uk

Name	Address ¹	Tel.	Email
STECF members			
Delaney, Alyne	Innovative Fisheries Management, -an Aalborg University Research Centre, Postboks 104, 9850 Hirtshals, Denmark	Tel.: +45 9940 3694	ad@ifm.aau.dk
Daskalov, Georgi	Laboratory of Marine Ecology, Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences	Tel.: +359 52 646892	gmdaskalov@yahoo.co.uk
Döring, Ralf	Thünen Bundesforschungsinstitut, für Ländliche Räume, Wald und Fischerei, Institut für Seefischerei - AG Fischereiökonomie, Palmaille 9, D-22767 Hamburg, Germany	Tel.: 040 38905-185 Fax.: 040 38905-263	ralf.doering@ti.bund.de
Gascuel, Didier	AGROCAMPUS OUEST 65 Route de Saint Briec, bat.4 CS 84215, F-35042 RENNES Cedex France	Tel:+33(0)2.23.48.55.3 4 Fax: +33(0)2.23.48.55.35	Didier.Gascuel@agrocampus-ouest.fr
Graham, Norman (chair)	Marine Institute, Fisheries Science Services (FSS), Rinville, Oranmore, Co. Galway, Ireland	Tel: + 353(0) 91 87200	norman.graham@marine.ie
Garcia Rodriguez, Mariano	Instituto Español de Oceanografía, Servicios Centrales, Corazón de María 8, 28002, Madrid, Spain		Mariano.Garcia@md.ieo.es
Gustavsson, Tore Karl-Erik	Independent Consultant, Göteborg, Sweden		tore.gustavsson@hotmail.com
Jennings, Simon	CEFAS Lowestoft Laboratory, Pakefield Road, Lowestoft Suffolk, UK NR33 0HT	Tel.: +44 1502562244 Fax: +44 1502513865	simon.jennings@cefas.co.uk
Kenny, Andrew	CEFAS Lowestoft Laboratory, Pakefield Road, Lowestoft Suffolk, UK NR33 0HT	Tel.: +44 1502562244 Fax: +44 1502513865	andrew.kenny@cefas.co.uk
Kraak, Sarah	Thünen-Institut für Ostseefischerei, Alter Hafen Süd 2, 18069 Rostock Germany	Tel. +49 3818116113	sarah.kraak@ti.bund.de
Kuikka, Sakari	University of Helsinki, Department of Environmental Sciences, P.O. Box 65 (Viikinkaari 1), FI-00014 University of Helsinki, FINLAND	Tel.: +358 50 3309233 Fax. +358-9-191 58754	skuikka@mappi.helsinki.fi
Martin, Paloma	CSIC Instituto de Ciencias del Mar Passeig Marítim, 37-49 08003 Barcelona Spain	Tel: 34.93.2309500 direct line : 34.93.2309552 Fax: 34.93.2309555	paloma@icm.csic.es
Malvarosa, Loretta	NISEA S.c.a.r.l.		malvarosa@nisea.eu

Name	Address ¹	Tel.	Email
STECF members			
Murua, Hilario	AZTI - Tecnalia / Unidad de Investigación Marina, Herrera kaia portualdea z/g 20110 Pasaia (Gipuzkoa), Spain	Tel: 0034 667174433 Fax: 94 6572555	hmurua@azti.es
Nord, Jenny	The Swedish Agency of Marine and Water Management (SwAM)	Tel. 0046 76 140 140 3	Jenny.nord@havochvatten.se
Nowakowski, Piotr	Maritime University of Szczecin. – Faculty of Food Science and Fisheries, Department of Fishing Technique, Szczecin		npfgd@poczta.onet.pl
Prelezzo, Raul	AZTI - Tecnalia / Unidad de Investigación Marina Txatxarramendi Ugarteia z/g 48395 Sukarrieta (Bizkaia), Spain	Tel: 94 6029400 Ext: 406- Fax: 94 6870006	rprelezo@suk.azti.es
Sala, Antonello	Fishing Technology Unit National Research Council (CNR) Institute of Marine Sciences (ISMAR) - Fisheries Section Largo Fiera della Pesca, 1 60125 Ancona - Italy	Tel: +39 071 2078841 Fax: +39 071 55313	a.sala@ismar.cnr.it
Scarcella, Giuseppe	Environmental Management Unit National Research Council (CNR) Institute of Marine Sciences (ISMAR) - Fisheries Section Largo Fiera della Pesca, 1 60125 Ancona - Italy	Tel: +39 071 2078846 Fax: +39 071 55313	g.scarcella@ismar.cnr.it
Somarakis, Stylianos	Department of Biology University of Crete VassilikaVouton P.O. Box 2208 71409 Heraklion Crete Greece	Tel.: +30 2610 394065, +30 6936566764	somarak@biology.uoc.gr
Stransky, Christoph	Thünen Institute [TI-SF] Federal Research Institute for Rural Areas, Forestry and Fisheries, Institute of Sea Fisheries, Palmaille 9, D-22767 Hamburg, Germany	Tel. +49 40 38905-228 Fax: +49 40 38905-263	christoph.stransky@ti.bund.de
Theret, Francois	Scapêche 17 Bd Abbé Le Cam 56100 Lorient France		ftheret@comata.com
Ulrich, Clara	DTU Aqua, National Institute of Aquatic Resources, Technical University of Denmark, Charlottenlund Slot, JægersborgAllé 1, 2920 Charlottenlund, Denmark		cu@aqu.dtu.dk
Vanhee, Willy	ILVO - Institute for Agricultural and Fisheries Research Unit Animal Sciences - Fisheries Ankerstraat 1, B-8400 Oostende, Belgium	Tel 00-32-59-34-22-55 Fax 00-32-59-33-06-29	willy.vanhee@ilvo.vlaanderen.be
van Oostenbrugge, Hans	Landbouweconomish Instituut-LEI, Fisheries Section, Burg. Patijnlaan 19 P.O.Box 29703 2502 LS The Hague The Netherlands	Tel:+31 (0)70 3358239 Fax: +31 (0)70 3615624	Hans.vanOostenbrugge@wur.nl

EWG-15-04 participants

STECF members			
Name	Address ¹	Telephone no.	Email
Raul PRELLEZO	AZTI - Tecnalia / Unidad de Investigación Marina Txatxarramendi Ugarteaz/g 48395 Sukarrieta (Bizkaia), Spain	+34 94 6029400	rprellezo@suk.azti.es

Invited experts			
Name	Address	Telephone no.	Email
Esther ABAD	Instituto Español de Oceanografía, Cabo Estai - Canido s/n, Vigo, Spain	+34 986492111	esther.abad@vi.ieo.es
Itsaso CARMONA	AZTI - Tecnalia / Unidad de Investigación Marina Txatxarramendi Ugarteaz/g 48395 Sukarrieta (Bizkaia), Spain	+34 94 6029400	icarmona@azti.es
Santiago CERVIÑO	Instituto Español de Oceanografía, Cabo Estai - Canido s/n, Vigo, Spain	+34 986492111	santiago.cervino@vi.ieo.es
José María DA ROCHA	Universidade de Vigo, Campus Lagoas Marcosende, Vigo, Spain	+34986812400	jmrocha@uvigo.es
Dorleta GARCIA	AZTI - Tecnalia / Unidad de Investigación Marina Txatxarramendi Ugarteaz/g 48395 Sukarrieta (Bizkaia), Spain	+34946029400	dgarcia@azti.es
Claire MACHER	IFREMER Département d'Economie Maritime BP 70 29280 Plouzané France	+33298224480	claire.macher@ifremer.fr
Mathieu MERZÉREAUD	IFREMER Département d'Economie Maritime BP 70 29280 Plouzané France		mathieu.merzereaud@ifremer.fr
María Paz SAMPEDRO	Instituto Español de Oceanografía, Paseo Marítimo Alcalde Francisco Vazquez, 10. A Coruña, Spain		paz.sampedro@co.ieo.es
Cristina SILVA	IPMA Av. de Brasília 1449-006 Lisboa Portugal	Tel.: +351 213 027 039	csilva@ipma.pt

JRC experts			
Name	Address	Telephone no.	Email
Ernesto JARDIM	Joint Research Centre (IPSC) Maritime Affairs Unit Via E. Fermi, 2749 21027 Ispra (Varese), Italy	+390332785311	ernesto.jardim@jrc.ec.europa.eu
Iago MOSQUEIRA	Joint Research Centre (IPSC) Maritime Affairs Unit Via E. Fermi, 2749 21027 Ispra (Varese), Italy	+39033285413	iago.mosqueira@jrc.ec.europa.eu

European Commission			
Name	Address	Telephone no.	Email
Angel Andrés CALVO SANTOS	DG MARE European Commission Rue de la Loi, 99 Bruxelles, Belgium	+32 2 2993630	angel-andres.calvo-santos@ec.europa.eu
Armando ASTUDILLO	DG MARE European Commission Rue de la Loi, 99 Bruxelles, Belgium	+32 2 2961191	armando.astudillo@ec.europa.eu

Observers			
Name	Address	Telephone no.	Email
Javier LÓPEZ	Southern Western Waters Regional Advisory Council 6 rue Alphonse Rio 56100 Lorient – FRANCE	+33 297 83 11 69	jlopez@oceana.org
Bjorn STOCKHAUSEN	Southern Western Waters Regional Advisory Council 6 rue Alphonse Rio 56100 Lorient – FRANCE	+33 297 83 11 69	bstockhausen@seas-at-risk.org
Alexandre MOUSSEIGNE	Southern Western Waters Regional Advisory Council 6 rue Alphonse Rio 56100 Lorient – FRANCE	+33 297 83 11 69	amousseigne@ccr-s.eu

EWG-15-09 participants

STECF members			
Name	Address ¹	Telephone no.	Email
Mariano GARCIA RODRIGUEZ	Instituto Español de Oceanografía, Servicios Centrales, Corazón de María 8, 28002, Madrid, Spain		Mariano.Garcia@md.ieo.es
Paloma MARTIN	Institut de Ciències del Mar CSIC Psg Marítim de la Barceloneta 37-49, 8003, Barcelona, Spain	+ 34932309500	paloma@icm.csic.es

Invited experts			
Name	Address	Telephone no.	Email

Paul DOLDER	CEFAS Lowestoft Laboratory, Pakefield Road, Lowestoft Suffolk, UK NR33 0HT	+441502 524259	paul.dolder@cefas.co.uk
Mariano GARCIA RODRIGUEZ	Instituto Español de Oceanografía, Servicios Centrales, Corazón de María 8, 28002, Madrid, Spain		Mariano.Garcia@md.ieo.es
Isabel GONZALEZ HERRAIZ	Instituto Español de Oceanografía, Paseo Marítimo Alcalde Francisco Vazquez- 10. A Coruña, Spain	+34981205362	isabel.herraiz@co.ieo.es
Jordi GUILLEN GARCIA	Institut de Ciències del Mar CSIC Psg Marítim de la Barceloneta 37-49, 8003, Barcelona Spain	+ 34932309500	jordiguillen@hotmail.com
Jordi LLEONART	Institut de Ciències del Mar CSIC Psg Marítim de la Barceloneta 37-49, 8003, Barcelona Spain	+ 34932309500	lleonart@icm.csic.es
Marianne ROBERT	Laboratoire de Technologie et de Biologie Halieutique, Station de Lorient - 8, rue François Toullec - 56100 Lorient, France	+330297873823	marianne.robert@ifremer.fr
Paloma MARTIN	Institut de Ciències del Mar CSIC Psg Marítim de la Barceloneta 37-49, 8003, Barcelona Spain	+ 34932309500	paloma@icm.csic.es
Francesc MAYNOU	Institut de Ciències del Mar CSIC Psg Marítim de la Barceloneta 37-49, 8003, Barcelona Spain	+ 34932309500	maynouf@icm.csic.es
Coilin MINTO	Galway-Mayo Institute of Technology, Galway Campus. Dublin Road. Galway, Ireland	+353 91 753161	coilin.minto@gmit.ie
Laura RECASENS	Institut de Ciències del Mar CSIC Psg Marítim de la Barceloneta 37-49, 8003, Barcelona Spain	+ 34932309500	laura@icm.csic.es

JRC experts			
Name	Address	Telephone no.	Email
Ernesto JARDIM	Joint Research Centre (IPSC) Maritime Affairs Unit Via E. Fermi, 2749 21027 Ispra (Varese), Italy	+390332785311	ernesto.jardim@jrc.ec.europa.eu
Giacomo Chato OSIO	Joint Research Centre (IPSC) Maritime Affairs Unit Via E. Fermi, 2749 21027 Ispra (Varese), Italy	+390332785948	giacomo-chato.osio@jrc.ec.europa.eu

European Commission			
Name	Address	Telephone no.	Email
Angel Andrés CALVO SANTOS	DG MARE European Commission Rue de la Loi, 99 Bruxelles, Belgium	+32 2 2993630	angel-andres.calvo-santos@ec.europa.eu
Armando ASTUDILLO	DG MARE European Commission Rue de la Loi, 99 Bruxelles, Belgium	+32 2 2961191	armando.astudillo@ec.europa.eu

Observers			
Name	Address	Telephone no.	Email
Benoit ARCHAMBAULT	North Western Waters Advisory Council, c/o Bord Iascaigh Mhara Crofton Road Dun Laoghaire Co. Dublin, Ireland	+353 1 214 4100	benoit.archambault@agrocampus-ouest.fr

12 LIST OF BACKGROUND DOCUMENTS

Background documents are published on the meetings websites on:

<http://stecf.jrc.ec.europa.eu/web/stecf/ewg1504> and <http://stecf.jrc.ec.europa.eu/web/stecf/ewg1509>

List of background documents:

1. EWG-15-04 / 09 - Declarations of invited and JRC experts (see also section 11 of this report – List of participants)

13 LIST OF ELECTRONIC ANNEXES

Electronic annexes are published on the meetings' websites on:

<http://stecf.jrc.ec.europa.eu/web/stecf/ewg1504> and <http://stecf.jrc.ec.europa.eu/web/stecf/ewg1509>

List of electronic annexes:

2. Annex I - Bio-economic impact assessment of multiannual management plans (MAPs) for the Spanish demersal fishing fleets in the Bay of Biscay
3. Annex II - Bio-economic impact assessment of multiannual management plans (MAPs) for Iberian demersal mixed fisheries
4. Annex III - Bio-economic Impact Assessment of the multi-annual management plan (MAP) for the Celtic Sea (ICES divisions VII bc, e-k) fisheries
5. Annex IV - IAM Description
6. Annex V - F ranges and Fleet Behaviour
7. Annex VI - Documents on FMSY ranges
 - a. WD1: Review of proxies for Fmsy ranges for Iberian Peninsula stocks (Southern hake, megrims and white anglerfish)
 - b. MSY reference points for haddock in VIIbce-k Working document to WGCSE 2015
 - c. Proxies for FMSY ranges using predictive linear models
8. Annex VII - Dependency and employment for the most dependent fleets. Dependency of North Western Atlantic waters (area 27.6 & 27.7) fleets on target demersal species
9. Annex VIII – Employment and degree of dependency by fleet

14 ANNEX I – CODES AND ACRONYMS

14.1 COUNTRIES CODES

<i>Alpha 3 code</i>	<i>Other codes used</i>	<i>Contry name</i>
BEL	Be	Belgium
FRA	Fr	France
GBR	Gb	United Kingdom
	En	<i>England</i>
	Sc	<i>Scotland</i>
NLD	Nl	Netherlands
ESP	Es	Spain
IRL	Ir	Ireland
PRT	Pt	Portugal

14.2 SPECIES CODES USED

Anf	Anglerfishes nei
Cod	Cod
Had	Haddock
Hal	Halibut
Her	Herring
Hke	Hake
Jax	Jack and horse mackerels nei
Mac	Mackerel
Nep	Nephrops
Nop	Nethrops
Ple	Plaice
Pok	Saithe(=Pollock)
San	Sandeels
Shr	Shrimps
Sol	Sole
Whg	Whiting

14.3 DCF CODES

FISHING_TECHNIQUE

DFN	Drift and/or fixed netters
DRB	Dredgers
DTS	Demersal trawlers and/or demersal seiners
FPO	Vessels using pots and/or traps
HOK	Vessels using hooks
MGO	Vessel using other active gears
MGP	Vessels using polyvalent active gears only
PG	Vessels using passive gears only for vessels < 12m
PGO	Vessels using other passive gears
PGP	Vessels using polyvalent passive gears only
PMP	Vessels using active and passive gears
PS	Purse seiners
TM	Pelagic trawlers
TBB	Beam trawlers

VESSEL_LENGTH classes

VL0010	Vessel between 0 meters and 10 meters in length.
VL1012	Vessel between 10 meters and 12 meters in length.
VL1218	Vessel between 12 meters and 18 meters in length.
VL1824	Vessel between 18 meters and 24 meters in length.
VL2440	Vessel between 24 meters and 40 meters in length.
VL40XX	Vessel greater than 40 meters in length.

FISHING GEAR

DRB	Boat dredges
DRH	Hand dredges
FPN	Stationary uncovered pound nets
FPO	Pots
FYK	Fyke nets
GNC	Encircling gillnets
GND	Driftnets
GNS	Set gillnets (anchored)

GTN	Combined gillnets-trammel nets
GTR	Trammel nets
HMD	Mechanised dredges including suction dredges
LA	Lampara nets
LHM	Handlines and pole-lines (mechanised)
LHP	Handlines and pole-lines (hand-operated)
LLD	Drifting longlines
LLS	Set longlines
LNB	Boat-operated lift nets
LNS	Shore-operated stationary lift nets
LTL	Troll lines
MIS	Miscellaneous Gear
NK	NOT KNOWN*
NO	NO GEAR
OTB	Bottom otter trawl
OTM	Midwater otter trawl
OTT	Otter twin trawl
PS	Purse seines
PTB	Bottom pair trawl
PTM	Pelagic pair trawl
SB	Beach seines
SDN	Danish seines
SPR	Pair seines
SSC	Scottish seines
SV	Beach and boat seines
TBB	Beam trawl

14.4 ACRONYMS

AER – Annual economic report

CFP - Common Fisheries Policy

ICES - International Council for the Exploration of the Sea

MSY – Maximum sustainable yield

CPUE – Catch per unit of effort

TAC – Total Allowable Catch

TAL – Total Allowable Landings

STECF - Scientific, Technical and Economic Committee for Fisheries

SG-MOS – Sub-group on management objectives and strategies

BoB – Bay of Biscay

NWW – North Western Waters

SWW – South Western Waters

HCR – Harvest Control Rules

MAP – Multi-annual plan

EwE - Ecopath with Ecosim model

LO - Landings obligation

FTE - Full Time Equivalent

FMSY – fishing mortality that provides maximum sustainable yield

SSB – Spawning stock biomass

HCR – Harvest Control Rule

Europe Direct is a service to help you find answers to your questions about the European Union
Freephone number (*): 00 800 6 7 8 9 10 11

(* Certain mobile telephone operators do not allow access to 00 800 numbers or these calls may be billed.

A great deal of additional information on the European Union is available on the Internet.
It can be accessed through the Europa server <http://europa.eu>.

How to obtain EU publications

Our publications are available from EU Bookshop (http://publications.europa.eu/howto/index_en.htm),
where you can place an order with the sales agent of your choice.

The Publications Office has a worldwide network of sales agents.
You can obtain their contact details by sending a fax to (352) 29 29-42758.

European Commission

EUR XXXXXX EN – Joint Research Centre – Institute for the Protection and Security of the Citizen
Title: Scientific, Technical and Economic Committee for Fisheries. Multiannual management plans SWW and NWW (STECF-15-04 & 09).

STECF members:

Graham, N., J., Abella, J. A., Andersen, J., Bailey, N., Bertignac, M., Cardinale, M., Curtis, H., Daskalov, G., Delaney, A., Döring, R., Garcia Rodriguez, M., Gascuel, D., Gustavsson, T., Jennings, S., Kenny, A., Kraak, S., Kuikka, S., Malvarosa, L., Martin, P., Murua, H., Nord, J., Nowakowski, P., Prellezo, R., Sala, A., Scarcella, G., Somarakis, S., Stransky, C., Theret, F., Ulrich, C., Vanhee, W. & Van Oostenbrugge, H.

JRC experts: Ernesto Jardim & Iago Mosqueira

EWG-15-04 members:

E. Abad, I. Carmona, S. Cervino, J.M. da Rocha, D. García, E. Jardim, C. Macher, M. Merzéréaud, I. Mosqueira, R. Prellezo, M.P. Sampedro, C. Silva.

EWG-15-09 members:

P. Dolder, M. Garcia Rodriguez, I. Gonzalez Herraiz, J. Guillen Garcia, E. Jardim, J. Lleonart, r. Marianne, P. Martin, F. Maynou, C. Minto, C. Osio, L. Recasens

Luxembourg: Publications Office of the European Union

2015 – 78 pp. – 21 x 29.7 cm

EUR – Scientific and Technical Research series – ISSN 1831-9424 (online), ISSN 1018-5593 (print)

ISBN XXXXXXXXXXXXX

doi:XXXXXXXXXXXXXXXXXXXX

STECF

The Scientific, Technical and Economic Committee for Fisheries (STECF) has been established by the European Commission. The STECF is being consulted at regular intervals on matters pertaining to the conservation and management of living aquatic resources, including biological, economic, environmental, social and technical considerations.

JRC Mission

As the Commission's in-house science service, the Joint Research Centre's mission is to provide EU policies with independent, evidence-based scientific and technical support throughout the whole policy cycle.

Working in close cooperation with policy Directorates-General, the JRC addresses key societal challenges while stimulating innovation through developing new methods, tools and standards, and sharing its know-how with the Member States, the scientific community and international partners.

Serving society

Stimulating innovation

Supporting legislation

doi:xx.xxxx/xxxxx

ISBN xxx-xx-xx-xxxxx-x

