

Auxiliary References

Ahmad, S. M., G. A. Babu, V. M. Padmakumari, and W. Raza (2008), Surface and deep water changes in the northeast Indian Ocean during the last 60 ka inferred from carbon and oxygen isotopes of planktonic and benthic foraminifera, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 262(3-4), 182-188, doi:10.1016/j.palaeo.2008.03.007.

Andreasen, D. H., M. Flower, M. Harvey, S. Chang, and A. C. Ravelo (2000), Late Pleistocene oxygen and carbon isotopic records from Sites 1011, 1012, and 1018, *Proc. Ocean Drill. Program Sci. Results*, 167, 141-144.

Arz, H. W., J. Pätzold, and G. Wefer (1999), The deglacial history of the western tropical Atlantic as inferred from high resolution stable isotope records off northeastern Brazil, *Earth Planet. Sci. Lett.*, 167, 105-117, doi:10.1016/S0012-821X(99)00025-4.

Baas, J. H., J. Mienert, F. Abrantes, and M. A. Prins (1997), Late Quaternary sedimentation on the Portugese continental margin: climate-related processes and products, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 130, 1-23.

Bard, E., et al. (1989), Sea-level estimates during the last deglaciation based on $\delta^{18}\text{O}$ and accelerator mass spectrometry ^{14}C ages measured in *Globigerina bulloides*, *Quat. Res.*, 31, 381-391.

Bard, E., et al. (1994), The North Atlantic atmosphere-sea surface ^{14}C gradient during the Younger Dryas climatic event, *Earth Planet. Sci. Lett.*, 126, 275-287.

Bard, E., F. Rostek, and G. Ménot-Combes (2004), Radiocarbon calibration beyond 20,000 ^{14}C yr B.P. by means of planktonic foraminifera of the Iberian Margin, *Quat. Res.*, 61, 204-214.

Bassinot, F. C., L. D. Labeyrie, E. Vincent, X. Quidelleur, N. J. Shackleton, and Y. Lancelot (1994), The astronomical theory of climate and the age of the Brunhes-Matuyama magnetic reversal, *Earth and Planetary Science Letters*, 126, 91-108.

Becquey, S., and R. Gersonde (2003), A 0.55-Ma paleotemperature record from the Subantarctic zone: Implications for Antarctic Circumpolar Current development, *Paleoceanography*, 18(1), 1014, doi:10.1029/2000PA000576.

Behling, H., H. W. Arz, J. Pätzold, and G. Wefer (2002), Late Quaternary vegetational and climate dynamics in southeastern Brazil, inferences from marine cores GeoB 3229-2 and GeoB 3202-1, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 179, 227-243.

Bickert, T., and G. Wefer (1996), Late Quaternary deep water circulation in the South Atlantic: Reconstruction from carbonate dissolution and benthic stable isotopes, in *The South Atlantic: Present and Past Circulation*, edited by G. Wefer et al., pp. 599-620, Springer, New York.

Bickert, T., W. H. Berger, S. Burke, H. Schmidt, and G. Wefer (1993), Late Quaternary stable

isotope record of benthic foraminifers: Site 805 and 806, Ontong Java Plateau, *Proc. Ocean Drill. Program Sci. Results*, 130, 411-420.

Bickert, T., W. B. Curry, and G. Wefer (1997), Late Pliocene to Holocene (2.6 – 0 Ma) western equatorial Atlantic deep-water circulation: inferences from benthic stable isotopes, *Proc. Ocean Drill. Program Sci. Results*, 154, 239-254.

Black, K. P., C. S. Nelson, and C. H. Hendy (1988), A spectral analysis procedure for dating Quaternary deep-sea cores and its application to a high-resolution Brunhes record from the Southwest Pacific, *Marine Geology*, 83, 21-30.

Bond, G., et al. (1997), A pervasive millennial-scale cycle in North Atlantic Holocene and Glacial climates, *Science*, 278, 1257-1266.

Boyle, E. A., and L. D. Keigwin (1985) Comparison of Atlantic and Pacific paleochemical records for the last 215,000 years: changes in deep ocean circulation and chemical inventories, *Earth Planet. Sci. Lett.*, 76, 135-150.

Broecker, W. S., et al. (1988), New evidence from the South China Sea for an abrupt termination of the last glacial period, *Nature*, 333, 156-158.

Came, R. E., D. W. Oppo, and W. B. Curry (2003), Atlantic Ocean circulation during the Younger Dryas: Insights from a new Cd/Ca record from the western subtropical South Atlantic, *Paleoceanography*, 18(4), 1086, doi:10.1029/2003PA000888.

Came, R. E., D. W. Oppo, and J. F. McManus (2007), Amplitude and timing of salinity and temperature variability in the high-latitude North Atlantic, *Geology*, 35, 315-318.

Channell, J. E. T., D. A. Hodell, and B. Lehman (1997), Relative geomagnetic paleointensity and $\delta^{18}\text{O}$ at ODP Site 983 (Gardar Drift, North Atlantic) since 350 ka, *Earth Planet. Sci. Lett.*, 153, 103-118.

Chapman, M. R., and N. J. Shackleton (1998), Millennial-scale fluctuations in North Atlantic heat flux during the last 150,000 years, *Earth Planet. Sci. Lett.*, 159, 57-70.

Chapman, M. R., and N. J. Shackleton (1999), Global ice-volume fluctuations, North Atlantic ice-rafting events, and deep-ocean circulation changes between 130 and 70 ka, *Geology*, 27, 795-798.

Chapman, M. R., N. J. Shackleton, and J.-C. Duplessy (2000), Sea surface temperature variability during the last glacial–interglacial cycle: assessing the magnitude and pattern of climate change in the North Atlantic, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 157(1), 1–25, doi:10.1016/S0031-0182(99)00168-6.

Charles, C. D., and R. G. Fairbanks (1992), Evidence from Southern Ocean sediments for the effect of North Atlantic deep-water flux on climate, *Nature*, 355, 416-419.

Charles, C. D., P. N. Froelich, M. A. Zibello, R. A. Mortlock, and J. J. Morley (1991), Biogenic opal in Southern Ocean sediments over the last 450,000 years: implications for surface water chemistry and circulation, *Paleoceanography*, *6*, 697-728, doi:10.1029/91PA02477.

Charles, C. D., J. Lynch-Stieglitz, U. S. Ninnemann, and R. G. Fairbanks (1996), Climate connections between the hemisphere revealed by deep sea sediment core/ice core correlations, *Earth Planet. Sci. Lett.*, *142*, 19-27.

Chen, M., and C. Huang (1998), Ice-volume forcing of winter monsoon climate in the South China Sea, *Paleoceanography*, *13*(6), 622-633.

Chen, J., J. W. Farrell, D. Murray, and W. L. Prell (1995), Timescale and paleoceanographic implications of a 3.6 m.y. oxygen isotope record from the northeast Indian Ocean (Ocean Drilling Program Site 758), *Paleoceanography*, *10*, 21-48, doi:10.1029/94PA02290.

Cheng, X., J. Tian, and P. Wang (2004), Stable isotopes from Site 1143, *Proc. Ocean Drill. Program Sci. Results*, *184*, 1-8.

Clemens, S. C., and W. L. Prell (2003), Oxygen and carbon isotopes from Site 1146, northern South China Sea, *Proc. Ocean Drill. Program Sci. Results*, *184*, 1-8, doi:10.2973/odp.proc.sr.184.214.2003.

Clemens, S. C., W. L. Prell, Y. Sun, Z. Liu, and G. Chen (2008), Southern Hemisphere forcing of Pliocene $\delta^{18}\text{O}$ and the evolution of Indo-Asian monsoons, *Paleoceanography*, *23*, PA4210, doi:10.1029PA001638.

Collins, J. A., et al. (2011), Interhemispheric symmetry of the tropical African rainbelt over the past 23,000 years, *Nat. Geosci.*, *4*, 42-45.

Cortijo, E. (1995), La variabilité climatique rapide dans l'Atlantique Nord depuis 128,000 ans: relations entre les calottes de glace et l'océan de surface, Ph. D. thesis, Univ. de Paris-Sud, Paris, France.

Cramer, B. S., J. R. Toggweiler, J. D. Wright, M. E. Katz, and K. G. Miller (2009), Ocean overturning since the Late Cretaceous: Inferences from a new benthic foraminiferal isotope compilation, *Paleoceanography*, *24*, PA4216, doi:10.1029/2008PA001683.

Curry, W. B. (1996), Late Quaternary deep circulation in the western equatorial Atlantic, in *The South Atlantic: Present and Past Circulation*, edited by G. Wefer et al., pp. 577-598, Springer, Berlin.

Curry, W. B., and G. P. Lohmann (1983), Reduced advection into Atlantic Ocean deep eastern basins during last glaciation maximum, *Nature*, *306*, 577-580, doi:10.1038/306577a0.

Curry, W. B., and D. W. Oppo (1997), Synchronous, high-frequency oscillations in tropical sea

surface temperatures and North Atlantic Deep Water productivity during the last glacial cycle, *Paleoceanography*, *12*, 1-14, doi:10.1029/96PA02413.

Curry, W. B., and D. W. Oppo (2005), Glacial water mass geometry and the distribution of $\delta^{13}\text{C}$ of ΣCO_2 in the western Atlantic Ocean, *Paleoceanography*, *20*, PA1017, doi:10.1029/2004/PA001021.

Curry, W. B., J.-C. Duplessy, L. D. Labeyrie, and N. J. Shackleton (1988), Changes in the distribution of $\delta^{13}\text{C}$ of deep water ΣCO_2 between the last glaciation and the Holocene, *Paleoceanography*, *3*, 317-341.

Curry, W. B., T. M. Marchitto, J. F. McManus, D. W. Oppo, and K. L. Laarkamp (1999), Millennial-scale changes in ventilation of the thermocline, intermediate, and deep waters of the glacial North Atlantic, *Geophys. Monogr. Ser.*, *112*, 59-76.

De Abreu, L., N. J. Shackleton, J. Schönfeld, M. Hall, and M. Chapman (2003), Millennial-scale oceanic climate variability off the Western Iberian margin during the last two glacial periods, *Marine Geology*, *196*, 1-20.

deMenocal, P., et al., (2000), Abrupt onset and termination of the African Humid Period: rapid climate responses to gradual insolation forcing, *Quat. Sci. Rev.*, *19*(1), 347-361, doi:10.1016/S0277-3791(99)00081-5.

Dickson, A. J., W. E. N. Austin, I. R. Hall, M. A. Maslin, and M. Kucera (2008), Centennial-scale evolution of Dansgaard-Oeschger events in the northeast Atlantic Ocean between 39.5 and 56.5 ka B.P., *Paleoceanography*, *23*, PA3206, doi:10.1029/2008PA001595.

Dickson, A. J., et al. (2009), Oceanic forcing of the Marine Isotope Stage 11 interglacial, *Nat. Geosci.*, *2*, 428-433.

Dudley, W. C., and C. S. Nelson (1994), The influence of non-equilibrium isotope fractionation on the Quaternary calcareous nanofossil stable isotope signal in the southwest Pacific Ocean, DSDP Site 594, *Mar. Micropaleontol.*, *24*, 3-27.

Duplessy, J.-C., C. Lalou, and A. C. Vinot (1970), Differential isotopic fractionation in benthic foraminifer and paleotemperatures reassessed, *Science*, *168*, 250-251.

Duplessy, J.-C. (1982), North Atlantic deep water circulation during the last climate cycle, *Bull. Inst. Geol. Bassin d'Aquitaine*, *31*, 379-391.

Duplessy, J.-C., N. J. Shackleton, R. K. Matthews, W. Prell, W. F. Ruddiman, M. Caralp, and C. H. Hendy (1984), ^{13}C record of benthic foraminifera in the last interglacial ocean: implication for the carbon cycle and the global deep water circulation, *Quaternary Research*, *21*, 225-243.

Duplessy, J.-C. (1996), Quaternary paleoceanography: unpublished stable isotope records. IGBP PAGES/World Data Center for Paleoclimatology Data Contribution Series #1996-035.,

NOAA/NGDC Paleoclimatology Program, Boulder, Colorado, USA.

Duplessy, J.-C., et al. (1988), Deepwater source variations during the last climatic cycle and their impact on the global deepwater circulation, *Paleoceanography*, 3(3), 343-360.

Duplessy, J.-C., et al. (1992), Changes in surface salinity of the North Atlantic Ocean during the last deglaciation, *Nature*, 358, 485-488.

Duplessy, J.-C., D. M. Roche, and M. Kageyama (2007), The deep ocean during the last interglacial period, *Science*, 316, 89-91.

Elliot, M., et al. (1998), Millennial-scale iceberg discharges in the Irminger Basin during the last glacial period: Relationship with the Heinrich events and environmental settings, *Paleoceanography*, 13(5), 433-446, doi:10.1029/98PA01792.

Elliot, M., L. Labeyrie, and J.-C. Duplessy (2002), Changes in North Atlantic deep-water formation associated with the Dansgaard-Oeschger temperature oscillations (60-10 ka), *Quat. Sci. Rev.*, 21(10), 1153-1165, doi:10.1016/S0277-3791(01)00137-8.

Eynaud, F., et al. (2007), Deglacial laminated facies on the NW European continental margin: The hydrographic significance of British-Irish Ice Sheet deglaciation and Fleuve Manche paleoriver discharges, *Geochem. Geophys. Geosystems*, 8(6), Q06019, doi:10.1029/2006GC001496.

Freudenthal, T., H. Meggers, J. Henderiks, H. Kuhlmann, A. Moreno, and G. Wefer (2002), Upwelling intensity and filament activity off Morocco during the last 250,000 years, *Deep Sea Research II*, 49, 3655-3674.

Ganeshram, R. S., and T. F. Pedersen (1998), Glacial-interglacial variability in upwelling and bioproductivity off NW Mexico: Implications for Quaternary paleoclimate, *Paleoceanography*, 13, 634-645.

Gardner, J. V., W. E. Dean, and P. Dartnell (1997), Biogenic sedimentation beneath the California Current system for the past 30 kyr and its paleoceanographic significance, *Paleoceanography*, 12(2), 207-225.

Gebhardt, H., et al. (2008), Paleonutrient and productivity records from the subarctic North Pacific for Pleistocene glacial terminations I to V, *Paleoceanography*, 23, PA4212, doi:10.1029/2007PA001513.

Gherardi, J. M., et al. (2005), Evidence from the Northeastern Atlantic basin for variability in the rate of the meridional overturning circulation through the last deglaciation, *Earth Planet. Sci. Lett.*, 240, 710-723.

Graham, D. W., B. H. Corliss, M. L. Bender, and L. D. Keigwin (1981), Carbon and oxygen isotopic disequilibria of recent deep-sea benthic foraminifera, *Marine Micropaleontology*, 6, 483-497.

Grousset, F. E., et al. (1993), Patterns of ice-rafted detritus in the glacial North Atlantic, *Paleoceanography*, 8, 175-192.

Hagen, S., and L. D. Keigwin (2002), Sea-surface temperature variability and deep water reorganization in the subtropical North Atlantic during Isotope Stage 2-4, *Marine Geology*, 189, 145-162.

Hall, I. R., I. N. McCave, N. J. Shackleton, G. P. Weedon, and S. E. Harris (2001), Intensified deep Pacific inflow and ventilation in Pleistocene glacial times, *Nature*, 412, 809-812.

Herbert, T. D., et al. (2001), Collapse of the California Current During Glacial Maxima Linked to Climate Change on Land, *Science*, 293, 71-76.

Hillenbrand, C. D., D. K. Fütterer, H. Grobe, and T. Frederichs (2002), No evidence for a Pleistocene collapse of the West Antarctic Ice Sheet from continental margin sediments recovered in the Amundsen Sea, *Geo-Mar Letters*, 22, 51-59.

Hodell, D. A., C. D. Charles, and U. S. Ninnemann (2000), Comparison of interglacial stages in the South Atlantic sector of the southern ocean for the past 450 kyr: implications for Marine Isotope Stage (MIS) 11, *Global Planet. Change*, 24, 7-26.

Hodell, D. A., et al. (2003), Oxygen isotope stratigraphy of ODP Leg 177 Sites 1088, 1089, 1090, 1093, and 1094, *Proc. Ocean Drill. Program Sci. Results*, 177, 1-26.

Hodell, D. A., J. E. T. Channell, J. H. Curtis, O. E. Romero, and U. Rohl (2008), Onset of "Hudson Strait" Heinrich events in the eastern North Atlantic at the end of the middle Pleistocene transition (~640 ka)?, *Paleoceanography*, 23, PA4218, doi:10.1029/2008PA001591.

Hoogakker, B. A. A., I. N. McCave, and M. J. Vautravers (2007), Antarctic link to deep flow speed variation during Marine Isotope Stage 3 in the western North Atlantic, *Earth Planet. Sci. Lett.*, 257, 463-473.

Hoogakker, B. A. A., et al. (2011), Dynamics of North Atlantic Deep Water masses during 1003 the Holocene, *Paleoceanography*, 26(4), PA4214, doi:10.1029/2011PA002155.

Hovan, S. A., D. K. Rea, and N.G. Pisias (1991), Late Pleistocene continental climate and oceanic variability recorded in Northwest Pacific sediments, *Paleoceanography*, 6(3), 349-370.

Hüls, M., and R. Zahn (2000), Millennial-scale sea surface temperature variability in the western tropical North Atlantic from planktonic foraminiferal census counts, *Paleoceanography*, 15, 659-678.

Imbrie, J. D., A. McIntyre, and A. C. Mix (1989), Oceanic response to orbital forcing in the late Quaternary: observational and experimental strategies, in *Climate and Geo-Sciences*, edited by A. Berger et al., pp. 121-164, Kluwer Academic, Boston, MA.

Isono, D., et al. (2009), The 1500-year climate oscillation in the midlatitude North Pacific during the Holocene, *Geology*, *37*, 591-594.

Jansen, E., and T. Veum (1990), Evidence for two-step deglaciation and its impact on North Atlantic deep water circulation, *Nature*, *343*, 612-616, doi:10.1038/343612a0.

Jullien, E., et al. (2006), Contrasting conditions preceding MIS3 and MIS2 Heinrich events, *Global Planet. Change*, *54*, 225-238.

Jung, S. J. A. (1996), Wassermassenaustausch zwischen NE-Atlantik und Nordmeer während der letzten 300000/80000 Jahre im Abbild stabiler O- und C- Isotope, *Berichte aus dem Sonderforschungsbereich 313*, Christian Albrechts Univ., Kiel.

Keigwin, L. D. (1995), Stable isotope stratigraphy and chronology of the upper Quaternary section at Site 883, Detroit Seamount, *Proc. Ocean Drill. Program Sci. Results*, *145*, 257-264.

Keigwin, L. D., and G. A. Jones (1989), Glacial-Holocene stratigraphy, chronology, and paleoceanographic observations on some North Atlantic sediment drifts, *Deep Sea Res. Part Ocean. Res. Pap.*, *36*(6), 845-867.

Keigwin, L. D., and G. A. Jones (1994), Western North Atlantic evidence for millennial-scale changes in ocean circulation and climate, *J. Geophys. Res.*, *99*(C6), 12397-12410.

Keigwin, L., G. Jones, S. Lehman, and E. Boyle (1991), Deglacial meltwater discharge, North Atlantic deep circulation, and abrupt climate change, *J. Geophys. Res.*, *96*(C9), 16811-16826, doi:10.1029/91JC01624.

Labeyrie, L. D., J.-C. Duplessy, and P. L. Blanc (1987), Variations in mode of formation and temperature of oceanic deep waters over the past 125,000 years, *Nature*, *327*, 477-482.

Labeyrie, L., et al. (1995), Surface and deep hydrology of the Northern Atlantic Ocean during the last 150000 years, *Philos. Trans. R. Soc. Lond. B. Biol. Sci.*, *348*(1324), 255-264.

Labeyrie, L., et al. (1996), Hydrographic changes of the Southern Ocean (southeast Indian sector) over the last 230 kyr, *Paleoceanography*, *11*(1), 57-76.

Labeyrie, L., et al. (1999), Temporal variability of the surface and deep waters of the North West Atlantic Ocean at orbital and millennial scales, *Geophys. Monogr. Ser.*, *112*, 77-98, doi:10.1029/GM112p0077.

Labracherie, M., et al. (1989), The last deglaciation in the Southern Ocean, *Paleoceanography*, *4*, 629-638.

Lackschewitz, K. S., et al. (1998), North Atlantic ice sheet fluctuations 10,000-70,000 yr ago as inferred from deposits on the Reykjanes Ridge, Southeast of Greenland, *Quat. Res.*, *49*, 171-182.

Lalicata, J. J., and D. W. Lea (2011), Pleistocene carbonate dissolution fluctuations in the eastern equatorial Pacific on glacial timescales: Evidence from ODP Hole 1241, *Marine Micropaleontology*, *79*, 41-51.

Lea, D. W., P. A. Martin, D. K. Pak, and H. J. Spero (2002), Reconstructing a 350 ky history of sea level using planktonic Mg/Ca and oxygen isotope records from a Cocos Ridge core, *Quat. Sci. Rev.*, *21*, 283-293.

Lea, D. W., et al. (2006), Paleoclimate history of Galápagos surface waters over the last 135,000 yr, *Quat. Sci. Rev.*, *25*, 1152-1167, doi:10.1016/j.quascirev.2005.11.010.

Lebreiro, S. M., et al. (2009), Sediment instability on the Portuguese continental margin under abrupt glacial climate changes (last 60 kyr), *Quat. Sci. Rev.*, *28*, 3211-3223, doi:10.1016/j.quascirev.2009.08.007.

Lee, M. Y., K. Y. Wei, and Y. G. Chen (1999), High resolution oxygen isotope stratigraphy for the last 150,000 years in the southern South China Sea: Core MD972151, *TAO*, *10*, 239-254.

Lisiecki, L. E., M. E. Raymo, and W. B. Curry (2008), Atlantic overturning responses to Late Pleistocene climate forcings, *Nature*, *456*, 85-88.

Little, M. G., et al. (1997), Rapid paleoceanographic changes in the Benguela Upwelling System for the last 160,000 years as indicated by abundances of planktonic foraminifera, *Palaeogeogr. Palaeoclim. Palaeoecol.*, *130*, 135-161.

Loubere, P., M. Richaud, Z. Liu, and F. Mekik (2003), Oceanic conditions in the eastern equatorial Pacific during the onset of ENSO in the Holocene, *Quat. Res.*, *60*, 142-148.

Lyle, M., et al. (1992), Paleoproductivity and carbon burial across the California Current: The multitracers transect, 42°N, *Paleoceanography*, *7*, 251-272.

Lyle, M., et al. (2000), Millennial-scale CaCO₃ and C_{org} events along the northern and central California margins: stratigraphy and origins, *Proc. Ocean Drill. Program Sci. Results*, *167*, 163-182, doi:10.2973/odp.proc.sr.167.214.2000.

Lyle, M., A. Mix, and N. Pisiias (2002), Patterns of CaCO₃ deposition in the eastern tropical Pacific Ocean for the last 150 kyr: Evidence for a southeast Pacific depositional spike during marine isotope stage (MIS) 2, *Paleoceanography*, *17*(2), 1013, doi:10.1029/2000PA000538.

Mackensen, A., and T. Bickert (1999), Stable carbon isotopes in benthic foraminifera: proxies for deep and bottom water circulation and new production, in *Use of proxies in paleoceanography – examples from the South Atlantic*, edited by G. Fischer and G. Wefer, pp.

229-254, Springer, Berlin.

Mackensen, A., H. Grobe, H. W. Hubberten, V. Spiess, and D. K. Fütterer (1989), Stable isotope stratigraphy from the Antarctic continental margin during the last one million years, *Marine Geology*, *87*, 315-321.

Mackensen, A., H. Grobe, H. W. Hubberten, and G. Kuhn (1994), Benthic foraminiferal assemblages and the $\delta^{13}\text{C}$ -signal in the Atlantic sector of the Southern Ocean: glacial-interglacial contrasts, in *Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change*, edited by R. Zahn et al., pp. 105-114, Springer, Berlin.

Mackensen, A., M. Rudolph, and G. Kuhn (2001), Late Pleistocene deep-water circulation in the subantarctic eastern Atlantic, *Global Planet. Change*, *30*, 197-229, doi:10.1016/S0921-8181(01)00102-3.

Marchitto, T. M., J. Lynch-Stieglitz, and S. R. Hemming (2005), Deep Pacific CaCO_3 compensation and glacial-interglacial atmospheric CO_2 , *Earth Planet. Sci. Lett.*, *231*, 317-336.

Martin, P. A., et al. (2002), Quaternary deep sea temperature histories derived from benthic foraminiferal Mg/Ca, *Earth Planet. Sci. Lett.*, *198*, 193-209.

Martinson, D. G., et al. (1987), Age dating and the orbital theory of the ice ages: development of a high-resolution 0 to 300,000-year chronostratigraphy, *Quat. Res.*, *27*, 1-29.

McIntyre, A., W. F. Ruddiman, K. Karlin, and A.C. Mix (1989), Surface water response of the Equatorial Atlantic Ocean to orbital forcing, *Paleoceanography*, *4*, 19-55, doi:10.1029/PA004i001p00019.

McManus, J. F., D. W. Oppo, and J. L. Cullen (1999), A 0.5 million year record of millennial-scale climate variability in the North Atlantic, *Science*, *283*, 971-975.

Millo, C., M. Sarnthein, A. Voelker, and H. Erlenkeuser (2006), Variability of the Denmark Strait Overflow during the Last Glacial Maximum, *Boreas*, *35*, 50-60.

Mix, A. C. (1986), Late Quaternary paleoceanography of the Atlantic Ocean: foraminiferal faunal and stable-isotopic evidence, Ph. D. thesis, Columbia Univ., New York, NY.

Mix, A. C., and R. G. Fairbanks (1985), North Atlantic surface-ocean control of Pleistocene deep-ocean circulation, *Earth Planet. Sci. Lett.*, *73*, 231-243.

Mix, A. C., et al. (1991), Carbon 13 in Pacific deep and intermediate waters, 0-370 ka: implications for ocean circulation and Pleistocene CO_2 , *Paleoceanography*, *6*, 205-226, doi:10.1029/90PA02303.

Mix, A. C., J. Le, and N. J. Shackleton (1995a), Benthic foraminiferal stable isotope stratigraphy of Site 846: 0-1.8 Ma, *Proc. Ocean Drill. Program Sci. Results*, *138*, 839-854.

- Mix, A. C., et al. (1995b), Benthic foraminifer stable isotope record from Site 849 (0-5 Ma): local and global climate changes, *Proc. Ocean Drill. Program Sci. Results*, 138, 371-412, doi:10.2973/odp.proc.sr.138.120.1995.
- Mohtadi, M., D. Hebbeln, S. N. Ricardo, and C. B. Lange (2006), El Niño-like pattern in the Pacific during marine isotope stages (MIS) 13 and 11?, *Paleoceanography*, 21, PA1015, doi:10.1029/2005PA001190.
- Mojtahid, M., et al. (2005), Palaeoclimatology and palaeohydrography of the glacial stages on Celtic and Armorican margins over the last 360000 yrs, *Mar. Geol.*, 224(1-4), 57–82, doi:10.1016/j.margeo.2005.07.007.
- Molyneux, E. G., I. R. Hall, R. Zahn, and P. Diz (2007), Deep water variability on the southern Agulhas Plateau: Interhemispheric links over the past 170 ka, *Paleoceanography*, 22, PA4209, doi:10.1029/2006PA001407.
- Mortyn, P. G., R. C. Thunell, D. M. Anderson, L. D. Stott, and J. Le (1996), Sea surface temperature changes in the Southern California Borderlands during the last glacial-interglacial cycle, *Paleoceanography*, 11(4), 415-430.
- Mulitza, S., et al. (2008), Sahel megadroughts triggered by glacial slowdown of Atlantic meridional overturning, *Paleoceanography*, 23, PA4206, doi:10.1029/2008PA001637.
- Murray, R. W., C. Knowlton, M. Leinen, A. C. Mix, and C. H. Polsky (2000), Export production and carbonate dissolution in the central equatorial Pacific Ocean over the past 1 Ma, *Paleoceanography*, 15, 570-592.
- Nelson, C. S., C. H. Hendy, A. M. Cuthbertson, and G. R. Jarrett (1986), Late Quaternary carbonate and isotope stratigraphy, subantarctic Site 594, Southwest Pacific, *Initial Reports DSDP*, 90, 1425-1436, doi:10.2973/dsdp.proc.90.144.1986.
- Nelson, C. S., P. J. Cooke, C. H. Hendy, and A. M. Cuthbertson (1993), Oceanographic and climatic changes over the past 160,000 years at Deep Sea Drilling Project Site 594 off southeastern New Zealand, southwest Pacific Ocean, *Paleoceanography*, 8, 435-458.
- Ninkovitch, D., and N. J. Shackleton (1975), Distribution, stratigraphic position and age of ash layer "L," in the Panama Basin region, *Earth Planet. Sci. Lett.*, 27, 20–34.
- Nürnberg, D., N. Brughmans, J. Schönfeld, U. Ninnemann, and C. Dullo (2004), Paleo-export productivity, terrigenous flux, and sea surface temperature around Tasmania - Implications for glacial/interglacial changes in the Subtropical Convergence Zone, *Geophys. Monogr. Ser.*, 151, 291-318.
- Oba, T., and M. Murayama (2004), Sea-surface temperature and salinity changes in the northwest Pacific since the Last Glacial Maximum, *Journal of Quaternary Science*, 19(4), 335-346.

Oba, T., et al. (2006), Paleoceanographic change off central Japan since the last 144,000 years based on high-resolution oxygen and carbon isotope records, *Global Planet. Change*, 53, 5-20.

Ohkushi, K., A. Suzuki, H. Kawahata, and L. P. Gupta (2003), Glacial-interglacial deep-water changes in the NW Pacific inferred from single foraminiferal $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$, *Marine Micropaleontology*, 48, 281-290.

Oppo, D. W., and R. G. Fairbanks (1987), Variability in the deep and intermediate water circulation of the Atlantic Ocean during the past 25,000 years: Northern Hemisphere modulation of the Southern Ocean, *Earth Planet. Sci. Lett.*, 86, 1-15, doi:10.1016/0012-821X(87)90183-X.

Oppo, D. W., and R. G. Fairbanks (1990), Atlantic Ocean thermohaline circulation of the last 150,000 years: relationship to climate and atmospheric CO_2 , *Paleoceanography*, 5, 277-288, doi:10.1029/PA005i003p00277.

Oppo, D. W., and M. Horowitz (2000), Glacial deep water geometry: South Atlantic benthic foraminiferal Cd/Ca and $\delta^{13}\text{C}$ evidence, *Paleoceanography*, 15(2), 147-160.

Oppo, D. W., and S. J. Lehman (1995), Suborbital timescale variability of North Atlantic Deep Water during the past 200,000 years, *Paleoceanography*, 10, 901-910.

Oppo, D. W., and Y. Sun (2005), Amplitude and timing of sea-surface temperature change in the northern South China Sea: Dynamic link to the East Asian monsoon, *Geology*, 33, 785-788.

Oppo, D. W., R. G. Fairbanks, A. L. Gordon, and N. J. Shackleton (1990), Late Pleistocene Southern Ocean $\delta^{13}\text{C}$ variability, *Paleoceanography*, 5, 43-54.

Ostermann, D. R., and W. B. Curry (2000), Calibration of stable isotopic data: an enriched $\delta^{18}\text{O}$ standard used for source gas mixing detection and correction, *Paleoceanography*, 15(3), 353-360.

Pahnke, K., and R. Zahn (2005), Southern Hemisphere Water Mass Conversion Linked with North Atlantic Climate Variability, *Science*, 307, 1741-1746, doi:10.1126/science.1102163.

Pahnke, K., R. Zahn, H. Elderfield, and M. Schulz (2003), 340,000-year centennial-scale marine record of Southern Hemisphere climatic oscillation, *Science*, 301, 948-952.

Pierre, C., J. F. Saliege, M. J. Urrutiaguer, and J. Giraudeau (2001), Stable isotope record of the last 500 k.y. at Site 1087 (Southern Cape Basin), *Proc. Ocean Drill. Program Sci. Results*, 175, 1-22, doi:10.2973/odp.proc.sr.175.230.2001.

Pichevin, L., P. Martinez, P. Bertrand, R. Schneider, and J. Giraudeau (2005), Nitrogen cycling on the Namibian shelf and slope over the last two climatic cycles: Local and global forcings, *Paleoceanography*, 20, PA2006, doi:10.1029/2004PA001001.

- Pichon, J. L., et al. (1992), Surface water temperature changes in the high latitudes of the southern hemisphere over the last glacial-interglacial cycle, *Paleoceanography*, 7, 289-318, doi:10.1029/92PA00709.
- Piotrowski, A. M., S. L. Goldstein, S. R. Hemming, and R. G. Fairbanks (2004), Intensification and variability of ocean thermohaline circulation through the last deglaciation, *Earth Planet. Sci. Lett.*, 225, 205-220.
- Pisias, N., and A. Mix (1997), Spatial and temporal oceanographic variability of the eastern equatorial Pacific during the late Pleistocene: Evidence from radiolaria microfossils, *Paleoceanography*, 12, 381-393.
- Praetorius, S. K., J. F. McManus, D. W. Oppo, and W. B. Curry (2008), Episodic reductions in bottom-water currents since the last ice age, *Nat. Geosci.*, 1, 449-452.
- Raymo, M. E., D. W. Oppo, and W. Curry (1997), The mid-Pleistocene climate transition: A deep sea carbon isotopic perspective, *Paleoceanography*, 12(4), 546-559.
- Raymo, M. E., et al. (2004), Stability of North Atlantic water masses in face of pronounced climate variability during the Pleistocene, *Paleoceanography*, 19, PA2008, doi:10.1029/2003PA000921.
- Richter, T. (1998), Sedimentary fluxes at the Mid-Atlantic Ridge - Sediment sources, accumulation rates, and geochemical characterisation, *GEOMAR Report 73*, GEOMAR Research Center for Marine Geosciences, Christian Albrechts University, Kiel.
- Ruddiman, W. F., and A. McIntyre (1981), Oceanic mechanism for amplification of the 23,000 year ice-volume cycle, *Science*, 212, 617-627.
- Rühlemann, C., S. Mulitza, P. M. Muller, G. Wefer, and R. Zahn (1999), Warming of the tropical Atlantic Ocean and slowdown of thermohaline circulation during the last deglaciation, *Nature*, 402, 511-514.
- Rühlemann, C., et al. (2004), Intermediate depth warming in the tropical Atlantic related to weakened thermohaline circulation: Combining paleoclimate data and modeling results for the last deglaciation, *Paleoceanography*, 19, PA1025, doi:10.1029/2003PA000948.
- Russon, T., et al. (2009), Middle-late Pleistocene deep water circulation in the southwest subtropical Pacific, *Paleoceanography*, 24, PA4205, doi:10.1029/2009PA001755.
- Saikku, R., L. Stott, and R. Thunell (2009), A bi-polar signal recorded in the western tropical Pacific: Northern and Southern Hemisphere climate records from the Pacific warm pool during the last Ice Age, *Quat. Sci. Rev.*, 28, 2374-2385, doi:10.1016/j.quascirev.2009.05.007.
- Salgueiro, E., et al. (2010), Temperature and productivity changes off the western Iberian margin during the last 150 kyr, *Quat. Sci. Rev.*, 29, 680-695.

Samson, C. R., E. L. Sikes, and W. R. Howard (2005), Deglacial paleoceanographic history of the Bay of Plenty, New Zealand, *Paleoceanography*, 20, PA4017, doi:10.1029/2004PA001088.

Sarnthein, M., et al. (1994), Changes in east Atlantic deepwater circulation over the last 30,000 years: Eight time slice reconstructions, *Paleoceanography*, 9, 209-267, doi:10.1029/93PA03301.

Sarnthein, M., et al. (2004), Mid-Holocene origin of the sea surface salinity low in the subarctic North Pacific, *Quat. Sci. Rev.*, 23, 2089-2099.

Sarnthein, M., et al. (2005), 95-Ky Cycles of ocean circulation in the far northwestern Pacific and South China Sea during the Brunhes Chron, in *Milutin Milankovitch Anniversary Symposium: Paleoclimate and the Earth Climate System*, Serbian Academy of Sciences and Arts, 110, edited by A. Berger et al., pp. 135-140, Serbian Acad. of Sci. and Arts, Belgrade.

Sarnthein, M., P. M. Grootes, J. P. Kennett, and M.-J. Nadeau (2007), ^{14}C reservoir ages show deglacial changes in ocean currents and carbon cycle, *Geophys. Monogr. Ser.*, 173, 175-196, doi:10.1029/173GM13.

Schlünz, B., R. Schneider, P. J. Müller, and G. Wefer (2000), Late Quaternary organic carbon accumulation south of Barbados: influence of the Orinoco and Amazon rivers?, *Deep Sea Research I*, 47, 1101-1124.

Schmiedl, G., and A. Mackensen (1997), Late Quaternary paleoproductivity and deep water circulation in the eastern South Atlantic Ocean: Evidence from benthic foraminifera, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 130, 43-80.

Schönfeld, J., R. Zahn, and L. de Abreu (2003), Surface and deep water response to rapid climate changes at the Western Iberian Margin, *Global Planet. Change*, 36, 237-264.

Shackleton, N. J. (1977), Carbon-13 in *Uvigerina*: Tropical rain forest history and the equatorial Pacific carbonate dissolution cycle, in *The Fate of Fossil Fuel in the Oceans*, edited by N. R. Andersen and A. Malahoff, pp. 401-427, Plenum Press, New York.

Shackleton, N. J., and M. A. Hall (1982), Oxygen isotope study of continuous scrape samples from site 480, *Initial Rep. Deep Sea Drill. Proj.*, 64, 1251-1254.

Shackleton, N. J., and M. A. Hall (1984), Oxygen and carbon isotope stratigraphy of Deep Sea Drilling Project Hole 552A: Plio-Pleistocene glacial history, *Initial Reports DSDP*, 81, 599-609.

Shackleton, N. J., and M. A. Hall (1989), Stable isotope history of the Pleistocene at ODP Site 677, *Proc. Ocean Drill. Program Sci. Results*, 111, 295-316.

Shackleton, N. J., and N. D. Opdyke (1973), Oxygen isotope and paleomagnetic stratigraphy of equatorial Pacific core V28-238: oxygen isotope temperatures and ice volumes on a 10^5 and 10^6 year scale, *Quaternary Research*, 3, 39-55.

Shackleton, N. J., J. Imbrie, and M. A. Hall (1983), Oxygen and carbon isotope record of East Pacific core V19-30: implications for the formation of deep water in the late Pleistocene North Atlantic, *Earth Planet. Sci. Lett.*, *65*, 233-244.

Shackleton, N. J., A. Berger, and W. R. Peltier (1990), An alternative astronomical calibration of the lower Pleistocene timescale based on ODP Site 677, *Transactions of the Royal Society of Edinburgh: Earth Sciences*, *81*, 251-261.

Shackleton, N. J., M. A. Hall, and E. Vincent (2000), Phase relationships between millennial-scale events 64,000-24,000 years ago, *Paleoceanography*, *15*, 565-569.

Shackleton, N. J., R. G. Fairbanks, T. Chiu, and F. Parrenin (2004), Absolute calibration of the Greenland time scale: implications for Antarctic time scales and for $\Delta^{14}\text{C}$, *Quat. Sci. Rev.*, *23*, 1513-1522, doi:10.1016/j.quascirev.2004.03.006.

Sikes, E. L., et al. (2009), Southern Ocean seasonal temperature and Subtropical Front movement on the South Tasman Rise in the late Quaternary, *Paleoceanography*, *24*, PA2201, doi:10.1029/2008PA001659.

Sirocko F. (1994), Abrupt change in monsoonal climate: evidence from the geochemical composition of Arabian Sea sediments, Ph. D. thesis, Christian Albrechts Univ., Kiel.

Sirocko, F., et al. (1993), Century-scale events in monsoonal climate over the past 24,000 years, *Nature*, *364*, 322-324.

Sirocko, F., D. Garbe-Schonberg, and C. Devey (2000), Processes controlling trace element geochemistry of Arabian Sea sediments during the last 25,000 years, *Global Planet. Change*, *26*, 217-303.

Skinner, L. C., and N. J. Shackleton (2004), Rapid transient changes in northeast Atlantic deep water ventilation age across Termination I, *Paleoceanography*, *19*(2), PA2005, doi:10.1029/2003PA000983.

Skinner, L. C., and N. J. Shackleton (2005), An Atlantic lead over Pacific deep-water change across Termination I: implications for the application of the marine isotope stage stratigraphy. *Quat. Sci. Rev.*, *24*, 571-580, doi:10.1016/j.quascirev.2004.11.008.

Skinner, L. C., N. J. Shackleton, and H. Elderfield (2003), Millennial-scale variability of deep water temperature and $\delta^{18}\text{O}_{\text{dw}}$ indicating deep-water source variations in the Northeast Atlantic, 0-34 cal. ka BP, *Geochem. Geophys. Geosystems*, *4*(12), 1098, doi:10.1029/2003GC000585.

Skinner, L. C., S. Fallon, C. Waelbroeck, E. Michel, and S. Barker (2010), Ventilation of the Deep Southern Ocean and Deglacial CO_2 Rise, *Science*, *328*, 1147-1151.

Slowey, N. C., and W. B. Curry (1995), Glacial-interglacial differences in circulation and carbon

cycling within the upper western North Atlantic, *Paleoceanography*, 10, 715-732.

Smart, C. W., C. Waelbroeck, E. Michel, and A. Mazaud (2010), Benthic foraminiferal abundance and stable isotopic changes in the Indian Ocean sector of the Southern Ocean during the last 20 kyr: Paleooceanographic implications, *Palaeogeogr. Palaeoclim. Palaeoecol.*, 297, 537-548.

Sowers, T., et al. (1993), A 135,000-year Vostok-SPECMAP common temporal framework, *Paleoceanography*, 8, 737-766, doi:10.1029/93PA02328.

Stott, L. D. (2007), Comment on “Anomalous radiocarbon ages for foraminifera shells” by W. Broecker et al.: A correction to the western tropical Pacific MD9821-81 record, *Paleoceanography*, 22, PA1211, doi:10.1029/2006PA001379.

Stott, L. D., M. Nuemann, and D. Hammond (2000), Intermediate water ventilation on the northeastern Pacific margin during the late Pleistocene inferred from benthic foraminiferal $\delta^{13}\text{C}$, *Paleoceanography*, 15, 161-169.

Stott, L. D., A. Timmermann, and R. Thunell (2007), Southern Hemisphere and deep-sea warming led deglacial atmospheric CO_2 rise and tropical warming, *Science*, 318, 435-438.

Thomson, J., et al. (1999), Implications for sedimentation changes on the Iberian margin over the last two glacial/interglacial transitions from $(^{230}\text{Th}_{\text{excess}})_0$ systematics, *Earth Planet. Sci. Lett.*, 165, 255-270.

Thomson, J., et al. (2000), Enhanced productivity on the Iberian margin during glacial/interglacial transitions revealed by barium and diatoms, *Journal of the Geological Society (London)*, 157, 667-677.

Thornalley, D. J. R., H. Elderfield, and I. N. McCave (2010), Intermediate and deep water paleoceanography of the northern North Atlantic over the past 21,000 years, *Paleoceanography*, 25, PA1211, doi:10.1029/2009PA001833.

Tian, J., P. Wang, X. Cheng, and Q. Li (2002), Astronomically tuned Plio- Pleistocene benthic $\delta^{18}\text{O}$ record from South China Sea and Atlantic-Pacific comparison, *Earth Planet. Sci. Lett.*, 203, 1015-1029, doi:10.1016/S0012-821X(02)00923-8.

Tiedemann, R., M. Sarnthein, and N. J. Shackleton (1994), Astronomic timescale for the Pliocene Atlantic $\delta^{18}\text{O}$ and dust flux records of Ocean Drilling Program site 659, *Paleoceanography*, 9, 619-638, doi:10.1029/94PA00208.

Tjallingii, R., et al. (2008), Coherent high- and low-latitude control of the northwest African hydrological balance, *Nat. Geosci.*, 1, 670-675.

Toucanne, S., et al. (2009), Timing of massive “Fleuve Manche” discharges over the last 350 kyr: insights into the European ice-sheet oscillations and the European drainage network from

MIS 10 to 2, *Quat. Sci. Rev.*, 28(13-14), 1238–1256, doi:10.1016/j.quascirev.2009.01.006.

Van Kreveld, S., et al. (2000), Potential links between surging ice sheets, circulation changes, and the Dansgaard-Oeschger cycles in the Irminger Sea, 60-18 kyr, *Paleoceanography*, 15(4), 425-442.

Vautravers, M. J., N. J. Shackleton, C. Lopez-Martinez, and J. O. Grimalt (2004), Gulf stream variability during marine isotope stage 3, *Paleoceanography*, 19, PA2011, doi:10.1029/2003PA000966.

Venz, K. A., D. A. Hodell, C. Stanton, and D. A. Warnke (1999), A 1.0 myr record of glacial North Atlantic Intermediate Water variability from ODP Site 982 in the northeast Atlantic, *Paleoceanography*, 14, 42-52, doi:10.1029/1998PA900013.

Vidal, L., R. R. Schneider, O. Marchal, T. Bickert, T. F. Stocker, and G. Wefer (1999), Link between the North and South Atlantic during the Heinrich events of the last glacial period, *Climate Dynamics*, 15, 909-919.

Voelker, A. H. L., et al. (2006), Mediterranean outflow strengthening during northern hemisphere coolings: A salt source for the glacial Atlantic?, *Earth Planet. Sci. Lett.*, 245, 39-55.

Waddell, L. M., I. L. Hendy, T. C. Moore, and M. W. Lyle (2009), Ventilation of the abyssal Southern Ocean during the late Neogene: A new perspective from the subantarctic Pacific. *Paleoceanography*, 24, PA3206, doi:10.1029/2008PA001661.

Waelbroeck, C., J.-C. Duplessy, E. Michel, L. Labeyrie, D. Paillard, and J. Duprat (2001), The timing of the last deglaciation in North Atlantic climate records, *Nature*, 412, 724-727.

Waelbroeck, C., et al. (2011), The timing of deglacial circulation changes in the Atlantic, *Paleoceanography*, 26, PA3213, doi:10.1029/2010PA002007.

Wang, L., et al. (1999), East-Asian monsoon climate during the Late Pleistocene: high-resolution sediment records from the South China Sea, *Marine Geology*, 156, 245-284, doi:10.1016/S0025-3227(98)00182-0.

Weaver, P. P. E., L. Carter, and H. Neil (1998), Response of surface water masses and circulation to late Quaternary climate change, east of New Zealand, *Paleoceanography*, 13, 70-83.

Wei, G. J., C. Y. Huang, C. C. Wang, M. Y. Lee, and K. Y. Wei (2006), High-resolution benthic foraminifer $\delta^{13}\text{C}$ records in the South China Sea during the last 150 ka, *Marine Geology*, 232, 227-235.

Weinelt, M. (1993), Veränderungen der Oberflächenzirkulation im Europäischen Nordmeer während der letzten 60.000 Jahre - Hinweise aus stabilen Isotopen, *Berichte aus dem Sonderforschungsbereich 313*, Christian Albrechts Univ., Kiel.

Weinelt, M., A. Rosell-Mele, U. Pflaumann, M. Sarnthein, and T. Kiefer (2003), The role of productivity in the Northeast Atlantic on abrupt climate change over the last 80,000 years, *Z. Dtsch. Geol. Ges.*, 154, 47–66.

Wells, P., and H. Okada (1997), Response of nanoplankton to major changes in sea-surface temperature and movements of hydrological fronts over Site DSDP 594 (south Chatham Rise, southeastern New Zealand), during the last 130 kyr, *Marine Micropaleontology*, 32, 341-363.

Winn, K., M. Sarnthein, and H. Erlenkeuser (1991), $\delta^{18}\text{O}$ Stratigraphy and Chronology of Kiel Sediment Cores from the East Atlantic, *Ber. 45*, Geologisch-Paläontologisches Institut und Museum, Univ. of Kiel.

Woodruff, F., S. M. Savin, and R. G. Douglas (1980), Biological fractionation of oxygen and carbon isotopes by recent benthic foraminifera, *Marine Micropaleontology*, 5, 3-11.

Yamane, M. (2003), Late Quaternary variations in water mass in the Shatsky Rise area, northwest Pacific Ocean, *Marine Micropaleontology*, 48, 205-223.

Zabel, M., T. Bickert, L. Dittert, R. R. Haese (1999), The significance of the sedimentary Al/Ti ratio as indicator for reconstructions of the terrestrial input to the equatorial Atlantic, *Paleoceanography*, 14, 789-799.

Zahn, R., K. Winn, and M. Sarnthein (1986), Benthic foraminiferal $\delta^{13}\text{C}$ and accumulation rates of organic carbon: *Uvigerina peregrina* group and *Cibicidoides wuellerstorfi*, *Paleoceanography*, 1, 27-42.

Zarriess, M., and A. Mackensen (2010), The tropical rainbelt and productivity changes off northwest Africa: A 31,000-year high-resolution record, *Mar. Micropaleontol.*, 76(3-4), 76–91, doi:10.1016/j.marmicro.2010.06.001.

Zarriess, M., and A. Mackensen (2011), Testing the impact of seasonal phytodetritus deposition on $\delta^{13}\text{C}$ of epibenthic foraminifer *Cibicidoides wuellerstorfi*: A 31,000 year high-resolution record from the northwest African continental slope, *Paleoceanography*, 26, PA2202, doi:10.1029/2010PA001944.

Zarriess, M., H. Johnstone, M. Prange, S. Steph, J. Groeneveld, S. Mulitza, and A. Mackensen (2011), Bipolar seesaw in the northeastern tropical Atlantic during Heinrich stadials, *Geophys. Res. Lett.*, 38, L04706, doi:10.1029/2010GL046070.

Zhang, J., P. Wang, Q. Li, X. Cheng, H. Jin, and S. Zhang (2007), Western equatorial Pacific productivity and carbonate dissolution over the last 550 kyr: Foraminiferal and nannofossil evidence from ODP Hole 807A, *Marine Micropaleontology*, 64, 121-140.