

1 Threatened by mining, polymetallic nodules are required to preserve
2 abyssal epifauna

3 **Ann Vanreusel^{1*}⁺, Ana Hilario²⁺, Pedro A. Ribeiro^{3,4+}, Lenaick Menot⁵⁺ and Pedro Martínez Arbizu⁶⁺**

4 ¹Marine Biology, Ghent University, 9000 Belgium.

5 ²Biology Department & Centre for Environmental and Marine Studies, University of Aveiro, 3810 Aveiro,
6 Portugal.

7 ³MARE – Marine and Environmental Sciences Centre, Universidade dos Açores, Departamento de
8 Oceanografia e Pescas, 9901-862 Horta, Açores, Portugal.

9 ⁴IMAR- Center of the University of the Azores, Departamento de Oceanografia e Pescas, 9901-862 Horta,
10 Portugal.

11 ⁵Deep-Sea Environment Laboratory, Ifremer, Centre de Bretagne, 29280 Plouzané, France.

12 ⁶Senckenberg am Meer, Abt. DZMB, 26382 Wilhelmshaven, Germany.

13 *ann.vanreusel@ugent.be

14 ⁺These authors contributed equally

15

17 **Supplementary Materials**

18

19 Table S1-1. Transect data. The nodule coverage is presented in percentage. The age of the experimental
 20 mining simulations and trawl tracks is indicated between brackets for the transects marked with *.
 21
 22

Station	License area	Length (m)	Image width (m)	Nodule coverage % (mean ± sd)
S0239_041	BGR	320	4	15.0 ± 2.6
S0239_064	BGR	870	4	0.3 ± 0.1
S0239_082	IOM	320	2	41.3 ± 5.3
S0239_082	IOM	410	2	44.3 ± 3.6
S0239_101	IOM	380	2	0.5 ± 0.2
S0239_101 *	IOM	250	2.9	0 (20 yr)
S0239_131	GSR	420	2	24.4 ± 2.7
S0239_141	GSR	450	2	21.9 ± 5.6
S0239_141 *	GSR	200	2.6	0 (0.7 yr)
S0239_157	IFREMER	400	2	29.5 ± 2.9
S0239_157 *	IFREMER	200	2.5	0 (37 yr)
S0239_161	IFREMER	450	2	0.6 ± 0.2
S0239_161 *	IFREMER	200	2.2	0 (3 yr)
S0239_189	APEI	400	2	41.6 ± 3.4
S0239_189	APEI	350	2	43.5 ± 5.2
S0239_200	APEI	400	2	85.4 ± 5.0
S0239_200	APEI	400	2	92.1 1.5

23

24

