Oxidative DNA damage and repair in the radioresistant archaeon *Thermococcus gammatolerans*

Barbier Ewa^{1,2}, Lagorce Arnaud^{3,4}, Hachemi Amine³, Dutertre Murielle³, Gorlas Aurore^{1,2}, Morand Lucie^{1,2}, Saint-Pierre Christine^{1,2}, Ravanat Jean-Luc^{1,2}, Douki Thierry^{1,2}, Armengaud Jean⁵, Gasparutto Didier^{1,2}, Confalonieri Fabrice³, Breton Jean^{1,2,*}

¹ Univ. Grenoble Alpes, INAC, LCIB, F-38000 Grenoble, France

² CEA, INAC, SyMMES, F-38000 Grenoble, France

³ Univ. Paris-Sud, Institute for Integrative Biology of the Cell (I2BC), Université Paris Saclay, CEA, CNRS, Orsay, France

⁴ Univ. Perpignan, IHPE – UMR 5244 CNRS/IFREMER/Univ. Montpellier, Montpellier, F- 34095, France

⁵ CEA, DSV-Li2D, Laboratory “Innovative technologies for Detection and Diagnostics”, BP 17171, Bagnols-sur-Cèze, F-30207, France

* Corresponding author : Jean Breton, email address : jean.breton@cea.fr

Abstract :

The hyperthermophilic archaeon *Thermococcus gammatolerans* can resist huge doses of γ -irradiation – up to 5.0 kGy – without loss of viability. The potential to withstand such harsh conditions is probably due to complementary passive and active mechanisms, including repair of damaged chromosomes. In this work, we documented the formation and repair of oxidative DNA lesions in *T. gammatolerans*. The basal level of the oxidized nucleoside, 8-oxo-2'-deoxyguanosine (8-oxo-dGuo), was established at 9.2 ($\pm$ 0.9) 8-oxo-dGuo per 106 nucleosides, a higher level than those usually measured in eukaryotic cells or bacteria. A significant increase in oxidative damage, i.e. up to 24.2 ($\pm$ 8.0) 8-oxo-dGuo/106 nucleosides, was measured for *T. gammatolerans* exposed to a 5.0 kGy dose of γ -rays. Surprisingly, the yield of radiation-induced modifications was lower than those previously observed for human cells exposed to doses corresponding to a few grays. One hour after irradiation, 8-oxo-dGuo levels were significantly reduced, indicating an efficient repair. Two putative base excision repair (BER) enzymes, TGAM_1277 and TGAM_1653, were demonstrated both by proteomics and transcriptomics to be present in the cells without exposure to ionizing radiation. Their transcripts were moderately upregulated after gamma irradiation. After heterologous production and purification of these enzymes, biochemical assays based on electrophoresis and MALDI-TOF (Matrix-Assisted Laser Desorption Ionization – Time Of Flight) mass spectrometry indicated that both have a β -elimination cleavage activity. TGAM_1653 repairs 8-oxo-dGuo whereas TGAM_1277 is also able to remove lesions affecting pyrimidines (1-[2-deoxy- β -D-erythro-pentofuranosyl]-5-hydroxyhydantoin (5-OH-dHyd) and 1-[2-deoxy- β -D-erythro-pentofuranosyl]-5-hydroxy-5-methylhydantoin (5-OH-5-Me-dHyd)). This work showed that, in normal growth conditions or in the presence of a strong oxidative stress, *T. gammatolerans* has the potential to rapidly reduce the extent of DNA oxidation, with at least these two BER enzymes as bodyguards with distinct substrate ranges.

Graphical abstract

Keywords : Radioresistance; archaeon; oxidative DNA damage; base excision repair; glycosylase

INTRODUCTION

Extremophile microorganisms have developed a wide range of mechanisms to survive in harsh environmental conditions. *Thermococcus gammatolerans*, a hyperthermophilic prokaryote described as the most radioresistant archaeon identified to date, was isolated from samples obtained in a hydrothermal vent located in the Gulf of California.¹ This obligatory anaerobic heterotrophic organism has a genome of 2.04 megabases (Mb), shows an optimal growth temperature of 88°C, and fully withstands a 5.0 kiloGrays (kGy) dose of gamma irradiation without loss of viability.² The molecular and biological mechanisms explaining this remarkable resistance of *T. gammatolerans* have not yet been defined. This microorganism probably combines different and complementary strategies to resist such hostile conditions, as is the case for other radiotolerant organisms.³ Ionizing radiation is associated with the production of reactive oxygen species (ROS), passive resistance mechanisms consist of modification of the Mn/Fe intracellular ion ratio and the production of “compatible solutes” composed of chemical agents such as polyols, sugars and amino acids allowing the stabilization of macromolecules.⁴ A more active mechanism involves combating oxidative stress using antioxidant enzymes. Finally, another way to resist ionizing radiation is to repair shattered chromosomal DNA as efficiently as possible. Pulsed field gel electrophoresis has allowed time course analysis of the archaeon chromosomal repair after irradiation. Such experiments have indicated that *T. gammatolerans* chromosomes are fully degraded after a 5.0 kGy dose of γ -rays. DNA strand breaks are fully repaired after 5 h by cells in stationary phase, while 6 h are necessary for cells in exponential growth phase.⁵ Proteomic approaches have demonstrated that *T. gammatolerans* produces key factors of the main DNA repair systems such as base excision repair (BER) and Double-Strand Break Repair.² It is worth noting that some of these proteins are present in the absence of stress,

which may be useful for survival in the harsh conditions that constantly surround these extremophiles, or during drastic environmental changes that are especially unfavorable to life. Despite a growing amount of data regarding radiotolerant organisms, DNA repair mechanisms in Archaea, and particularly in *T. gammatolerans*, remain poorly understood. Most of the DNA alterations resulting from high doses of radiation have never been analyzed in radioresistant archaea. Among these, 8-oxo-2'-deoxyguanosine (8-oxo-dGuo) is a classic biomarker of oxidative stress. Its chemical structure is presented in Fig. 1. The main mechanism of formation of 8-oxo-dGuo upon exposure to ionizing radiation depends on an indirect effect: the radiolysis of water generates the hydroxyl radical $\cdot\text{OH}$ which reacts with C8 of the guanine.⁶ Mutations (mainly G→T transversions) can occur in the genome if 8-oxo-dGuo is not repaired prior to replication.⁷ Sample preparation and measurement of 8-oxo-dGuo are technically challenging to avoid DNA artifactual oxidation, and as a consequence misleading results can be found in the literature.⁶ DNA lesions 5',8-cyclonucleosides (Fig. 1) are radiation-induced, tandem base-sugar alterations usually produced in lower yields than 8-oxo-dGuo but with a more severe biological impact resulting from a drastic alteration of the DNA structure.⁸ The formation of these complex DNA lesions is known to be favored at low oxygen concentrations.⁹ The large radiation dose that can be applied to *T. gammatolerans* cells and their natural anaerobic growth conditions makes the quantitation of 5',8-cyclonucleosides highly relevant.

In the present study, we first quantified oxidative DNA damage in *T. gammatolerans*, both at basal levels and after exposure to high doses of γ -irradiation, by measuring 8-oxo-dGuo and 5',8-cyclonucleosides by HPLC coupled to tandem mass spectrometry (HPLC-MS/MS). We also documented the molecular mechanisms of two putative BER enzymes identified in the genome: TGAM_1277, initially annotated as a putative type III endonuclease (endoIII), and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TGAM_1653, a predicted DNA glycosylase,² phylogenetically close to the 8-oxoguanine glycosylase from *Pyrobaculum aerophilum*.¹⁰ Transcriptomics and proteomics were used to assess their presence in *T. gammatolerans* with and without exposure to irradiation. After their heterologous production, TGAM_1277 and TGAM_1653 proteins were assayed for their cleaving activities on different oxidative DNA lesions by electrophoresis and Matrix-Assisted Laser Desorption Ionization – Time Of Flight (MALDI-TOF) mass spectrometry.

EXPERIMENTAL PROCEDURES

T. gammatolerans growth conditions

The *T. gammatolerans* EJ3 strain was cultivated in serum bottles or in 1 L Schott bottles, under anaerobic conditions at 85°C in complex organic medium (VSM), as described in ².

Cell culture densities were measured by optical microscopy (Olympus BH-29) using a Thoma counting chamber (Microgravure Precis).

Gamma irradiation and DNA extraction

Gamma irradiation

For 8-oxo-dGuo quantification after irradiation, large volumes of exponentially growing cells (600 mL in 1 L Schott bottles, 7×10^7 cells/mL) were filtered (MicraCloth Calbiochem, La Jolla, CA, USA) to remove sulfur compounds from the VSM-S° medium. In anaerobic conditions, cells were concentrated X10 in Hungate tubes for γ -irradiation. Cells were then exposed on ice to a dose of 2.5 kGy or 5.0 kGy (100% survival)⁵ using a ¹³⁷Cs γ -ray source (40 Gy/min, IBL637 CisBio International, Institut Curie, Orsay, France). As a control, the same number of nonirradiated cells was kept on ice during irradiation. Irradiated and nonirradiated cells were either collected immediately or incubated for 60 min at 85°C for recovery in fresh VSM-S° medium on a reciprocal shaker before centrifugation (4000 g, 20 min at 4°C). Cells were immediately frozen and stored at -80°C until further processing. Each culture was performed in independent duplicates.

For gene expression assays, 600 mL exponentially growing cells were filtered (MicraCloth Calbiochem, La Jolla, CA, USA) to remove sulfur compounds from the VSM-S° medium and

concentrated ten-fold by centrifugation. Cells were exposed on ice to a dose of 2.5 kGy or 5.0 kGy using a ^{137}Cs γ -ray source (19 Gy/min, GSR D1 GSM, Institut Curie, Orsay, France). As a control, the same number of nonirradiated cells was kept on ice during irradiation. Irradiated and nonirradiated cells were either collected immediately or incubated for 60 or 120 min at 85°C for recovery in fresh VSM-S° medium on a reciprocal shaker before centrifugation (4000 g, 20 min at 4°C). Cells were immediately frozen and stored at -80°C until further processing. Each culture was repeated twice.

DNA extraction and digestion

T. gammatolerans cells (10 mg) were supplemented with 300 μL lysis buffer containing 100 mM Tris-HCl pH 8.3, 50 mM EDTA, 100 mM NaCl, and 0.15 mM deferoxamine. Thirty μL 10% SDS (v/v) and 30 μL 10% sarkosyl (v/v) were then added to the cell suspension. The samples were incubated with 150 μg RNase A (Sigma) and 3.5 U RNase T1 (Sigma) for 15 min at 50°C. Six hundred μg protease (Qiagen) were added and the incubation was pursued for 1.5 h at 37°C. DNA was precipitated with 600 μL 20 mM EDTA- Na_2 , 7.6 M NaI, 40 mM Tris-HCl, 0.3 mM deferoxamine pH 8, and 1 mL 100% isopropanol. The pellets were washed with 40% isopropanol (v/v) and subsequently with 70% ethanol (v/v), and finally suspended in 50 μL 0.1 mM deferoxamine.

The DNA digestion was performed in two steps. First, 0.025 U phosphodiesterase II, 2.5 U DNase II, 0.5 U nuclease P1 (Sigma) and 2.5 μL buffer (200 mM succinic acid, 100 mM CaCl_2 , pH 6) were added and the samples were incubated for 2 h at 37°C. Then, 6 μL 500 mM Tris-HCl, 1 mM EDTA pH 8, 0.015 U phosphodiesterase I (Worthington) and 2 U alkaline phosphatase (Sigma) were added and the samples were incubated for 2 h at 37°C. The reaction was stopped by adding 3.5 μL 0.1 N HCl. For the cyclonucleoside detection, the first step of the digestion was repeated before adding HCl, as optimized previously.⁹

HPLC-MS/MS quantification of 8-oxo-dGuo and 5',8-cyclonucleosides

8-oxo-dGuo

Digested DNA samples were resolved on an Uptisphere 2x150 mm octadecylsilyl silica gel column (Interchim) operated with an Agilent series 1100 HPLC system. The mobile phase was a gradient of methanol in 2 mM aqueous ammonium formate (2 to 10% (v/v) in 25 min and 10 to 25% (v/v) in the following 10 min). Undamaged nucleosides were monitored with a UV detector set at 260 nm while 8-oxo-dGuo was measured in positive mode with an API 3000 triple quadrupolar mass spectrometer (PerkinElmer/SCIEX) equipped with ElectroSpray Ionisation (ESI). The system was operated in the multiple reaction monitoring mode using a specific transition (m/z 284 $\rightarrow$ 168). DNA damage rates measured in different experimental conditions were compared using the Mann-Whitney test ($p < 0.05$). For each condition, five measurements were performed: two technical replicates for the first biological sample and three technical replicates for the second biological sample.

5',8-cyclonucleosides

Four lesions were measured simultaneously: 5'*R* and 5'*S* diastereomers of 5',8-cyclo-2'-deoxyadenosine (5',8-cyclodAdo) and 5',8-cyclo-2'-deoxyguanosine (5',8-cyclodGuo). Digested DNA samples were resolved on an Uptisphere 2x150 mm column of silica octadecylsilyl (Interchim) operated with an Accela HPLC (Thermo Corporation). Nucleosides were resolved by a linear gradient of acetonitrile (0 to 15% (v/v) gradient in 30 min) in ammonium formate. Undamaged nucleosides were detected at 260 nm with a UV detector while cyclonucleosides were specifically monitored as previously described⁹ with a TSQ Quantum Ultra mass detector equipped with ESI.

Transcriptomic and proteomic analysis of TGAM₁₂₇₇ and TGAM₁₆₅₃

For expression profiling by real-time RT-PCR (Reverse Transcription PCR), total RNAs were extracted with TriReagent solution (Sigma) and genomic DNA was digested by Turbo DNase (Ambion) according to the manufacturer's instructions. The absence of genomic DNA in RNA samples was checked by PCR using 16S rRNA primers (Table S1) and 2 µg total RNA as template. Reverse transcription was performed in a total volume of 20 µL in the presence of 5 µg total RNA, 2 pmol gene-specific reverse primers (Table S1), 0.5 mM each dNTP and 200 units SuperScriptII (Invitrogen, Carlsbad, CA). After a 30 min incubation at 42°C, 200 units SuperScriptII were added and the samples were incubated at 42°C for 30 min. The hydrolysis of the RNA was performed in the presence of 11 µL 1N NaOH for 10 min at 70°C and the solution was neutralized with 11 µL 1 N HCl. cDNAs were desalted with Microcon-G30 (Millipore), dried under vacuum and resuspended in 10 µL water. cDNA templates were diluted 1/100 with water for expression measurements. Standard conditions were used with LightCycler FastStart DNA MasterPLUS SYBR green I (Roche), i.e. DNA denaturation at 95°C for 10 min followed by 40 cycles of 95°C for 10 s, 60°C for 10 s and 72°C for 15 s, using the LightCycler System (Roche Diagnostics). Gene expression was calculated relative to the geometric average expression of four reference genes using the formula $2^{-\Delta Ct}$.^{11, 12} The reference genes used in this work, namely *TGAM_0829*, *TGAM_01054*, *TGAM_1165*, and *TGAM_1926*, encode a TATA box-binding protein (tbp), an elongation factor 1-alpha (tuf), a CDP-alcohol phosphatidyltransferase, and a DNA-directed RNA polymerase subunit B (rpoB),¹³ respectively. Expression of these selected reference genes remains particularly stable in *T. gammatolerans* cells challenged with heavy metal (i.e. cadmium)¹³ or with ionizing radiation, as revealed by genome-wide transcriptomic data (unpublished data). Proteomics information for TGAM₁₂₇₇ and TGAM₁₆₅₃ were extracted from data we had previously

obtained through a shotgun proteomic analysis, now available on the ProteomeXchange Consortium (dataset identifier PXD000402).

Heterologous production and purification of BER enzymes

Bacterial strains and expression plasmids

E. coli strain DH5 α (Life Technologies) was used in all cloning experiments and for plasmid amplifications. The *E. coli* Rosetta (DE3) strain (Novagen) was used for protein production. The pET30a(+) plasmid (Novagen) was used for bacterial expression of N-terminal His₆-tagged proteins.

Cloning and expression of TGAM_1277 and TGAM_1653 glycosylase genes

TGAM_1277 (230 aa) and TGAM_1653 (263 aa) coding DNA sequences (CDS) were first amplified by PCR with genomic DNA as the template using a high fidelity DNA polymerase (Fusion high fidelity, ThermoFisher Scientific), according to the manufacturer's instructions (DNA denaturation at 95°C for 2 min followed by 25 cycles of 95°C for 10 s, 55°C for 20 s and 72°C for 30 s) and using primer pairs containing *EcoRV* and *XhoI* restriction sites at their 5' and 3' ends, respectively, as described in table S2. The PCR fragments were digested with the corresponding restriction enzymes and cloned in pET30a (+) at the same unique restriction sites found in the polylinker. The amplified CDS and their fusion in-frame with the series of tags located in the N-terminal of the recombinant proteins, including a His₆-tag, were confirmed by double-stranded sequencing using T7P and T7TERM primers (Beckman Coulter Genomics). Because of the codon usage bias in archaea, both proteins were expressed in the *E. coli* Rosetta (DE3) strain encoding rare tRNA genes, a strategy which has proved successful for numerous archaeal proteins.^{14, 15} *E. coli* Rosetta (DE3) strain cells were

transformed with the individual expression plasmid and were grown overnight at 37°C with shaking in 5 mL LB media containing 25 µg/mL kanamycin and 30 µg/mL chloramphenicol. The cultures were transferred into 500 mL (1:100 dilution) LB media supplemented with 25 µg/mL kanamycin and 30 µg/mL chloramphenicol in 2 L Erlenmeyer flasks until the OD₆₀₀ reached 0.5-0.6. The His₆-tagged TGAM_1277 and TGAM_1653 recombinant proteins were expressed for 4 h at 37°C after induction with 1 mM IPTG (isopropyl β-D-1-thiogalactopyranoside). Cells were harvested by centrifugation and resuspended in 8 mL buffer A (20 mM Tris-HCl pH 7.5, 1 mM NaCl, 1 mM DTT, 20 mM imidazole), supplemented with 300 µL of a protease inhibitor cocktail for bacterial cells (Sigma Aldrich), and finally stored at -80°C overnight.

Purification of TGAM_1277 and TGAM_1653 recombinant glycosylases

After thawing on ice, *E. coli* cells were lysed with an Eaton Press at a pressure of 10,000 psi, and the resulting extracts were sonicated for 2 x 15 s. The protein extracts were centrifuged at 15,000 g for 30 min at 4°C. The soluble proteins were loaded onto His-trap columns (Macherey Nagel) previously equilibrated with buffer A and washed in the same buffer. The recombinant proteins were purified by affinity FPLC (Fast Protein Liquid Chromatography) (Amersham, GE Healthcare) with increasing steps of imidazole (50 mM, 150 mM, 250 mM) in the same buffer and dialyzed twice (Spectrumlabs, MWCO 15 kDa) overnight at 4°C against 500 mL buffer B (20 mM Tris-HCl pH 7.5, 50% Glycerol v/v, 200 mM NaCl, 1 mM DTT). The purified recombinant proteins were almost 95% pure as judged by SDS-PAGE analysis (Fig.2). The concentrations of TGAM_1653 and TGAM_1277 were 0.33 and 0.22 mg/mL, respectively, as estimated by Bradford assays.

DNA glycosylase activity assays

Oligonucleotides (ODNs)

In order to perform activity tests, ODNs of the following sequences were used: 5'-CAC TTC GG¹A T²CG TGA CTG ATC T-3' (control), where G¹ was replaced by 8-oxo-dGuo, and T² by 1-[2-deoxy-β-D-erythro-pentofuranosyl]-5-hydroxyhydantoin (5-OH-dHyd) or 1-[2-deoxy-β-D-erythro-pentofuranosyl]-5-hydroxy-5-methylhydantoin (5-OH-5-Me-dHyd). All ODNs, except those containing 8-oxo-dGuo (Eurogentec), were synthesized in-house as previously described.^{16, 17} In order to form double stranded ODNs, sequences fully complementary to the control ODN were obtained from Eurogentec, along with three other ODNs presenting G, A or T mismatches opposite the G¹ position.

Radiolabeling of ODNs

ODNs (10 pmol) were [5'-³²P]-labeled at their 5'-ends with 10 μCi [γ-³²P] ATP (10 mCi/mL, Perkin Elmer France) and 1 unit of T4 polynucleotide kinase (10 U/μL, Life Technologies/Invitrogen) in the supplied "forward" reaction buffer. The labeling reaction was performed at 37°C for 1 h. Unincorporated [γ-³²P] ATP was then removed using Illustra MicroSpin G-25 Columns (GE Healthcare) and the [5'-³²P]-labeled strands were finally hybridized with their complementary sequences by incubation at 90°C for 3 min followed by slow cooling until room temperature was reached.

Polyacrylamide gel electrophoresis

The [5'-³²P]-labeled ODN duplexes (100 fmol) were incubated with 2 pmol TGAM₁₆₅₃ or TGAM₁₂₇₇ proteins in the reaction buffer consisting of 20 mM Tris-HCl, buffered at pH 7.8 and containing 1 mM EDTA and 150 mM KCl. After 30 min incubation at 37°C, the temperature was raised progressively to 65°C and the reaction was maintained at this

1
2
3 temperature for 10 min. In parallel, control experiments with *E. coli* formamidopyridine DNA
4
5 *N*-glycosylase (Fpg) (a gift from Dr. B. Castaing, CNRS, Orléans, France) were performed.
6
7 An incubation of 1 h at 37°C was carried out in the reaction buffer for Fpg. The samples were
8
9 then denatured by heating for 3 min at 90°C in formamide solution and resolved by
10
11 electrophoresis on a polyacrylamide gel in denaturing conditions. Signal detection was
12
13 performed by phosphorimaging with the Personal Molecular Imager system (Bio-Rad).
14
15
16
17

18 *MALDI-TOF-MS*

19
20 For MALDI-TOF-MS measurements, ODNs were not radioactively labeled but directly
21
22 hybridized with their complementary sequence. These duplexes (10 pmol) were incubated
23
24 with 20 pmol TGAM_1653, TGAM_1277, Fpg or Endo III proteins, in conditions as
25
26 described above for TGAM_1653 and TGAM_1277. The samples were then desalted with the
27
28 50WX8 100-200 Dowex resin (Sigma-Aldrich) and analyzed with a Microflex instrument
29
30 (Bruker).
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Basal levels of oxidative DNA damage and quantitation after irradiation

We first wanted to obtain quantitative data about oxidative DNA damage in *T. gammatolerans* with and without exposure to ionizing radiation. DNA damage was measured in cells that were neither irradiated nor incubated on ice ("basal levels", fig. 3). Cells were also exposed to a γ -ray dose of either 2.5 or 5.0 kGy. The exposure lasted 62.5 min (2.5 kGy) or 125 min (5.0 kGy), and was performed on ice to avoid a significant increase of temperature. Cells were then either immediately collected after irradiation or incubated in fresh medium for 1h before being processed. For each condition, nonirradiated controls were prepared in parallel and stored on ice during irradiation. Extraction protocols were carefully optimized, on the basis of a protocol developed for mammalian cells,¹⁸ to obtain DNA solutions suitable for HPLC-MS/MS measurement of nucleosides and to avoid artifactual oxidation of DNA, which may lead to an overestimation of damage levels.

In these conditions, 8-oxo-dGuo basal levels in *T. gammatolerans* (9.2 ± 0.9 8-oxo-dGuo/ 10^6 nucleosides) were more abundant than in nonirradiated cells stored on ice, which presented a minimum of 4.2 ± 0.5 8-oxo-dGuo/ 10^6 nucleosides for the 5.0 kGy/0 h control and a maximum of 8.4 ± 1.4 8-oxo-dGuo/ 10^6 nucleosides for the 2.5 kGy/1 h control. Irradiated cells exhibited higher levels of DNA lesions than the corresponding controls (Fig. 3), particularly at 5.0 kGy, with an increase from 4.2 ± 0.5 8-oxo-dGuo/ 10^6 nucleosides for controls to 24.2 ± 8.0 8-oxo-dGuo/ 10^6 nucleosides for irradiated cells ($p < 0.05$, Mann-Whitney). The 1 h incubation of irradiated cells in a fresh medium was associated with a decrease in 8-oxo-dGuo mean values, which was only significant after 5.0 kGy treatment,

falling to 9.7 +/- 2.9 8-oxo-dGuo/10⁶ nucleosides. In control cells, incubation in fresh medium was associated with a slight but statistically significant increase of oxidized purines: from 5.1 +/- 1.2 8-oxo-dGuo/10⁶ nucleosides to 8.4 +/- 1.4 8-oxo-dGuo/10⁶ nucleosides for 2.5 kGy controls and from 4.2 +/- 0.5 8-oxo-dGuo/10⁶ nucleosides to 6.5 +/- 1.1 8-oxo-dGuo/10⁶ nucleosides for 5.0 kGy controls.

The 5',8-cyclonucleosides (5'*R* and 5'*S* diastereomers of 5',8-cyclodAdo and 5',8-cyclodGuo) were also measured in the same experimental conditions but chromatograms resulting from these analyses did not show detectable peaks, indicating the absence of these DNA alterations or rates below the limit of detection of our methods, which, during this experiment, was estimated at 1.0 lesion/10⁶ nucleosides.

Proteomic detection of TGAM_1277 and TGAM_1653, and transcriptional expression of the corresponding genes.

We focused on two putative BER enzymes identified in the genome of *T. gammatolerans*: TGAM_1277 and TGAM_1653, which have been annotated as a putative endoIII and a predicted 8-oxoG glycosylase, respectively. We analyzed their occurrence and possible regulation in cells by combining proteomic and transcriptomic approaches. A recent shotgun proteomic survey has been conducted for *T. gammatolerans* cells subjected to a 5.0 kGy γ -irradiation dose and the corresponding control without irradiation.¹⁹ We extracted from the data deposited to the ProteomeXchange Consortium (dataset identifier PXD000402) the spectral counts for the TGAM_1277 and TGAM_1653 proteins in order to evaluate the presence of both proteins. The former was detected and validated through two proteotypic peptides ([134-CANIVLAYGFGR-145] and [215-ELCPYAKGLVR-225]) while the latter was detected by means of a unique peptide ([217-AEETGIPPLHIDSILWPVLGGK-239]) in

1
2
3 this shotgun survey. As estimated by the normalized spectral abundance factor (NSAF)
4
5 obtained by normalizing spectral counts with the molecular weight of each protein,²⁰
6
7 TGAM_1277 (NSAF 0.22) and TGAM_1653 (NSAF 0.10) were slightly less abundant
8
9 compared to TGAM_1679, the reverse gyrase (NSAF 0.95), and in any case present in fairly
10
11 low amounts compared to the whole set of detected proteins.
12

13
14 We monitored by real-time RT-PCR the expression of their corresponding genes,
15
16 *TGAM_1277* and *TGAM_1653*, in control conditions and following γ -irradiation. For γ -
17
18 irradiation, exponentially growing cells were exposed to 2.5 or 5.0 kGy and collected
19
20 immediately (T0) or incubated post irradiation in fresh VSM medium at 85°C for 60 or 120
21
22 min (T1, T2, see methods). At both doses, 100% survival had previously been observed.⁵
23
24 Expression profiles of both genes in these conditions are shown in Fig. 4. For *TGAM_1277*,
25
26 we observed a comparable relative expression in control conditions and in cells immediately
27
28 collected after irradiation (T0; 0.03). A slight increase (two-fold) of the relative expression
29
30 could be detected 1 and 2 h (T1, T2) after a dose of 2.5 kGy. Interestingly, exposure to a
31
32 higher dose (5.0 kGy) led to a significant upregulation (three-fold) as measured 2 h after
33
34 irradiation (T2). For *TGAM_1653*, a similar relative expression was revealed when
35
36 nonirradiated cells and cells collected immediately after irradiation were compared (0.11-
37
38 0.12). Our data show that the basal relative expression of *TGAM_1653* (0.11-0.12) was higher
39
40 than with *TGAM_1277* (0.03). However, *TGAM_1653* relative expression remained stable
41
42 following γ -irradiation at a 2.5 kGy dose and only a weak upregulation could be measured
43
44 after 2 h (T2) for a 5.0 kGy irradiation.
45
46
47
48
49

50 51 52 **Activity of recombinant TGAM_1653 and TGAM_1277** 53 54 55 56 57 58 59 60

To characterize their biochemical activities, the two glycosylases were produced in *Escherichia coli* in fusion with a His₆-tag in the N-terminal domain of the protein, as several glycosylases contain a zinc finger motif in their C-terminal domain. We decided to not remove the 48-amino-acid N-terminal extension by proteolysis as our data showed that it does not disrupt protein activities (see hereafter). The excision activities of the recombinant TGAM_1653 and TGAM_1277 proteins were assessed at a temperature increasing from 37°C to 65°C by means of an electrophoretic cleavage assay using double-stranded and ³²P-labeled ODNs. DNA lesions inserted into the substrates were: i) 8-oxo-dGuo which is a model of oxidized purine, ii) 5-OH-dHyd, and iii) 5-OH-5-Me-dHyd, the latter two being models of oxidized pyrimidines. For this experimental set-up, the bacterial Fpg enzyme, which is known to have a glycosylase activity toward 8-oxo-dGuo, 5-OH-dHyd and 5-OH-5-Me-dHyd, was used as a positive control.

TGAM_1653 was only able to cleave probes containing 8-oxo-dGuo, whereas TGAM_1277 showed a broader substrate specificity by reacting with the three lesions (Fig. 5A). The effect of the opposite base on glycosylase activity toward 8-oxoG (8-oxo-7,8-dihydroguanine, the oxidized base) was also evaluated using four probes with the four different bases in front of the altered position (Fig. 5B). As already shown by Tchou *et al.*,²¹ Fpg activity is partially inhibited by an 8-oxoG:A mismatch. Once again, TGAM_1277 presented a broad activity in contrast to TGAM_1653, the latter being less efficient when confronted with DNA containing 8-oxo-G:A and 8-oxoG:G pairs whereas TGAM_1277 activity was only weakly affected by the opposite base of the lesion.

The gels in Fig. 5A display different sizes of short excision products. For a given enzyme tested for different lesions this phenomenon is simply explained by the different localizations of DNA modifications within the probe sequences. However, this difference in migration was

also noted for one given lesion cleaved by different enzymes: fragments generated by TGAM_1277 with hydantoin lesions exhibited higher molecular weights than others produced with Fpg. Regarding 8-oxo-dGuo, fragments obtained with TGAM_1653 and TGAM_1277 were also of higher molecular weight than the cleavage products of Fpg. As this observation could probably be explained by differences in excision mechanisms, we used MALDI-TOF MS to obtain further insight. This methodology allows a precise determination of the size of the cleavage products, hence providing complementary details about repair mechanisms.²² Regarding the excision process of both *T. gammatolerans* enzymes, our hypothesis was that a β -elimination occurs while, in contrast, Fpg is known to act as a β,δ -elimination enzyme.²³ Both possible elimination processes are described in Fig. 6.

In our experiments, Fpg and endoIII were used as controls for β,δ -elimination of 8-oxo-dGuo and β -elimination of 5-OH-5-Me-dHyd, respectively. Experimental MALDI-TOF MS spectra acquired on the products of Fpg, and recombinant TGAM_1653 and TGAM_1277 excision, are shown in Figs. 7 and 8. 8-oxo-dGuo repair by Fpg was first characterized by the absence of peak **1** corresponding to 8-oxo-dGuo-containing strands. This was indicative of a total cleavage of the substrate by the enzyme. The presence of peaks **3** ([M-H]⁻ = 4332.9 Da; calculated = 4332.8 Da) and **4** ([M-H]⁻ = 2135.6 Da; calculated = 2135.4 Da) is characteristic of the β,δ -elimination process. For TGAM_1653 tested on 8-oxo-dGuo substrates, the absence of peak **1** proves a complete cleavage whereas the presence of peaks **3** ([M-H]⁻ = 4332.5 Da) and **5** ([M-H]⁻ = 2233.9 Da; calculated = 2233.4 Da) confirms a β -elimination mechanism. The MALDI-TOF MS spectrum of TGAM_1277 activity revealed that peak **1** was clearly visible, indicating that in our experimental conditions, this enzyme did not cut all the available substrate. TGAM_1277 was therefore considered as less active than Fpg and TGAM_1653, confirming the results obtained with electrophoretic gels (Fig. 5). Considering

the cleavage products, we were able to detect peak **3** but not peak **4**, thus characterizing a β -elimination process. Actually, the presence of peak **6** ($[M-H]^- = 2251.9$ Da; calculated = 2251.5 Da) probably reveals a hydration product of the α - β -unsaturated ketone, which should have been represented by fragment **5**.²⁴ We can therefore assume that TGAM_1277 uses a β -elimination process.

The same principle was applied to the cleavage of ODNs containing 5-OH-5-Me-dHyd (Fig. 8). The MALDI-TOF MS profile corresponding to our control endoIII confirms its β -elimination activity with the presence of peak **3** ($[M-H]^- = 3714.9$ Da; calculated = 3715.4 Da) and hydrated product **6** ($[M-H]^- = 2893.7$ Da; calculated = 2893.9 Da). The β,δ -elimination exhibited by Fpg is demonstrated again by the presence of peaks **3** ($[M-H]^- = 3715.6$ Da; calculated = 3715.4 Da) and **4** ($[M-H]^- = 2776.5$ Da; calculated = 2777.8 Da). The MS profile of TGAM_1277 is very close to that obtained with endoIII: product **3** at 3714.8 Da and product **6** at 2893.5 Da indicating β -elimination. The only difference with endoIII is the absence of peak **1**, indicating a more efficient cleavage. TGAM_1653 is not associated with short fragments, a characteristic that again indicates its absence of activity on 5-OH-5-Me-dHyd.

In conclusion, TGAM_1653 and TGAM_1277 both exhibit a β -elimination cleavage activity. TGAM_1653 is only able to cleave 8-oxo-dGuo whereas TGAM_1277 is also able to cut ODNs containing pyrimidine lesions (5-OH-dHyd and 5-OH-5-Me-dHyd). It is worth noting that ODN substrates containing 8-oxodA and 5',8-cyclonucleosides were also tested but did not show detectable excisions (data not shown).

DISCUSSION

***T. gammatolerans* shows a basal level of 8-oxo-dGuo that is enhanced by high doses of γ -irradiation, but rapidly decreases post exposure**

Measurement of oxidative base damage in cellular DNA is a rather commonly used biomarker of oxidative stress but has for a long time been associated with questionable results due to possible overestimation related to technical artifacts. By using the most appropriate preanalytical precautions, as well as sensitive and specific methods such as HPLC-electrochemical detection and, more recently, HPLC-MS/MS, it is now accepted that background levels of 8-oxo-dGuo in human cells such as lymphocytes are below the threshold of 1 lesion per 10^6 nucleosides (4 lesions per 10^6 guanines).^{25, 26} This level corresponds to approximately 10,000 8-oxo-dGuo per single nucleus.²⁷ Using HPLC-MS/MS, 8-oxo-dGuo measurements in *E. coli* DNA carried out by our team showed basal rates of 2.6 (+/- 0.9) 8-oxo-dGuo per 10^6 nucleosides (unpublished data). In the present study, basal levels measured in *T. gammatolerans* grown in laboratory experimental conditions were 9.2 (+/- 0.9) 8-oxo-dGuo per 10^6 nucleosides, which represents approximately 18 8-oxo-dGuo per chromosome (*T. gammatolerans* is a polyploid organism for which the mean number of chromosome copies has not been determined yet). While an overwhelming proportion of published studies reporting 8-oxo-dGuo rates have been performed using mammalian cells, the only relevant work comparable with ours was carried out using the archaeon *Halobacterium salinarum*.²⁸ In this halophilic organism, a basal level of 2.1 8-oxoG (oxidized base) per 10^6 bases was observed by means of gas chromatography coupled to mass spectrometry. Two other oxidative purine lesions were assessed in the same study: 4,6-diamino-5-formamidopyrimidine (FapyAde) and 2,6-diamino-4-hydroxy-5-formamidopyrimidine (FapyGua), which showed concentrations of 0.3 and 0.9 lesion per 10^6 bases, respectively.

Elevated rates of 8-oxo-dGuo found in *T. gammatolerans* could be related to the growth conditions, and more specifically to the temperature. Indeed, the optimal growth temperature for this hyperthermophilic organism (85°C) was chosen for the experimental work presented here, while lower temperatures were used in other studies: 37°C for human cells and 42°C for *H. salinarum*. Heat-induced formation of oxidized guanine, at least *in vitro*, has been reported.²⁹ Because nonirradiated cells stored on ice presented only 4.2 8-oxo-dGuo lesions per 10⁶ nucleosides, this hypothesis of heat-induced damage formation is reinforced. This was confirmed by the result obtained with nonirradiated cells incubated for 1 h at 85°C in culture medium, which also showed a significant increase in their 8-oxo-dGuo content. It can be highlighted here that another type of DNA damage, abasic sites, was also found at elevated levels in *Pyrococcus abyssi*: between 25 (exponential phase) and 42 (stationary phase) abasic sites per 100,000 base pairs in the genome of this hyperthermophilic archaeon versus 2 and 4, respectively, abasic sites per 100,000 base pairs in *E. coli*.³⁰

In order to study DNA damage in the context of exposure to ionizing radiation, *T. gammatolerans* samples were subjected to high doses of γ -radiation. Levels of 8-oxo-dGuo measured in cells harvested immediately after treatment significantly increased. The corresponding yields of radiation-induced modifications are 2.8 8-oxo-dGuo/10¹² nucleosides/Gy at 2.5 kGy and 4.0 8-oxo-dGuo/10¹² nucleosides/Gy at 5.0 kGy. Again, measurements of 8-oxo-dGuo after irradiation have previously been measured mainly using eukaryotic cells,^{25, 31-33} and with doses lower than 500 Gy. In the THP1 malignant monocyte cell line, HPLC-MS/MS analysis indicated a damage yield reaching 20 8-oxo-dGuo/10⁹ nucleosides/Gy ($\sim$ 200 to 300 8-oxo-dGuo / cell / Gy).³¹ Kish *et al.*²⁸ analyzed the oxidized base, 8-oxoG, after exposure of the archaeon *H. salinarum* at 0, 2.5, 5.0 and 7.5 kGy. A plateau was reached at 2.5 kGy, for which the levels were 2.1 8-oxoG/10⁶ bases and 3.1 8-oxoG/10⁶ bases at 0 and 7.5 kGy, respectively (⁶⁰Co gamma source), corresponding to less

than 1 lesion/10¹² bases/Gy. Therefore, this archaeon also showed yields of oxidized guanine formation far below those observed in human cells. These very low yields may be explained by a cellular redox environment and low O₂ levels that prevent the formation of oxidative DNA damage, or by detoxification systems.⁴ In the same halophilic organism, Robinson *et al.*³⁴ observed an increase of FapyGua levels at 2.5 kGy but no significant progression of 8-oxoG.

Apart from a potential role of DNA repair enzymes, low levels of oxidative DNA damage observed in *T. gammatolerans* after irradiation may be linked to mechanisms such as highly constitutively expressed detoxification systems. For example, the genome of this archaeon is containing sequences related to proteins known to decrease oxidative stress such as a thioredoxin reductase (TGAM_0180), a glutaredoxin-like protein (TGAM_1302) and 2 peroxiredoxins (TGAM_1253 and TGAM_1220).^{2, 4}

T. gammatolerans cells exposed to 5.0 kGy radiation showed a significant decrease of 8-oxo-dGuo after 1 h recovery in culture medium, whereas this phenomenon was not significant at the lower levels of oxidative DNA damage observed at 2.5 kGy. Studies performed with *H. salinarum* indicated a decrease of FapyGua, FapyAde and 8-oxoG rates after a 2.5 kGy irradiation and 2 h recovery compared with nonexposed organisms (rates after 1 h recovery were not measured).²⁸ This decrease of 8-oxo-dGuo one hour after the end of irradiation may contribute to the resistance of *T. gammatolerans*. It may be explained by DNA repair systems or other factors limiting the extent of DNA damage.

Cyclonucleoside formation in *T. gammatolerans* DNA is questionable

Cyclonucleosides (in this study, cyclopurines) are classified as complex oxidative DNA lesions presenting, in the same nucleotide, a covalent bond between the C8 of the base and the C5' of the 2-deoxyribose. *In vitro* studies show that these bulky lesions are mutagenic³⁵ and

interfere with replication and transcription.³⁶⁻³⁸ Contrary to small oxidative DNA damage such as oxidized bases, it also appears that cyclopurines are repaired by the nucleotide excision repair system and not by BER enzymes. In parallel to these mechanistic results, measurements of cyclopurine rates in DNA extracted from eukaryotic cultured cells and tissues have been performed by a limited number of research teams. In spite of similar methods, i.e. isotope dilution technique in combination with HPLC coupled to mass spectrometry, discrepancies can be noted among the concentrations reported in the literature. In THP1 human monocytes, Belmadoui *et al.*⁹ did not detect 5'*R* and 5'*S* diastereomers of 5',8-cyclodAdo above the detection limit of the method, which was 0.1 lesion per 10⁹ nucleosides. In rodent tissues, the team of Wang measured basal levels of 5'*R* and 5'*S* diastereomers of 5',8-cyclodAdo and 5',8-cyclodGuo superior to 0.1 lesion per 10⁶ nucleosides.³⁹⁻⁴¹ In the work reported here, we did not show the presence of detectable cyclopurines in *T. gammatolerans* cells in any of the conditions tested. Indeed, even exposure to 5.0 kGy radiation did not lead to a signal above our limit of detection, estimated at 1 lesion/10⁶ nucleosides for this experiment (the limit of detection is dependent on the quantity of DNA available for the analysis). In our case, another favorable factor for the detection of cyclonucleosides should be the anaerobic growth conditions of *T. gammatolerans*. Cyclonucleosides are indeed more prone to formation at low O₂ concentrations.⁹ Taken together, these data indicate that cyclonucleoside formation in living cells, *T. gammatolerans* in our case, may occur at very low levels, if at all.

TGAM_1653 and TGAM_1277 are slightly upregulated after heavy radiation exposure

Data extracted from proteomic results indicated the presence at low levels of BER enzymes TGAM_1653 and TGAM_1277 in nonirradiated and irradiated cells. No difference between

these two conditions was noted in terms of protein abundance, but the sensitivity of the MS shotgun approach is not sufficient to detect moderate variations. The expression of the corresponding genes was confirmed by RT-PCR. For TGAM_1277, the main difference appeared 2 h after a 5.0 kGy irradiation, with an increase of the relative expression from 0.03 (T0) to 0.09. TGAM_1653 basal mRNA levels were higher than those of TGAM_1277, but the regulation of the transcription of this gene was less sensitive to irradiation. The expression of BER enzymes before exposure to ionizing radiation is not surprising since extremophiles need constitutive protective mechanisms to deal with their harsh growth conditions. The DNA damage analysis reported here shows the presence of an elevated amount of oxidized bases in basal conditions that may require a sufficient amount of DNA repair enzymes. The weak or absent upregulation of several DNA repair genes after an exposure to genotoxic stress has already been observed in other archaea. *Pyrococcus furiosus* subjected to 2.5 kGy ^{60}Co γ -rays did not show changes in mRNA levels for BER glycosylases but presented a moderate upregulation of *radA* (DNA repair by homologous recombination) and of a DNA repair gene cluster possibly involved in translesion synthesis.⁴² UV irradiation of hyperthermophilic *Sulfolobus* archaea did not lead to overexpression of DNA repair genes, including BER genes.⁴³

It is also recognized that a majority of DNA repair factors are constitutively expressed in eukaryotic cells.⁴⁴ At the protein level, the expression of BER factors is not massively induced, even after significant exposure to genotoxic agents.⁴⁵ We also observed that 8-oxo-dGuo produced after a γ -ray exposure are repaired within the first hour of recovery, whereas maximum levels of TGAM_1277 and TGAM_1653 transcripts were measured at 2 h. We can suggest that newly synthesized enzymes are produced (i) to repair other types of more persistent oxidative DNA damage or (ii) to reconstitute the pool of constitutive enzymes that has probably been, at least partly, altered by irradiation. This degradation of BER factors by

1
2
3 γ -rays has already been shown to occur in human cells⁴⁶ at much lower doses (up to 60 Gy)
4
5 compared to those in the present study. Finally, it is worth mentioning that our observations
6
7 are mainly based on mRNA levels. It would be interesting to complement them with further
8
9 data related to proteins, as well as their post-translational modifications, which are known to
10
11 play an important role in the regulation of BER in eukaryotes.⁴⁷
12
13

14
15
16 **TGAM_1653 excises 8-oxoG, TGAM_1277 excises 8-oxoG and hydantoin lesions. Both**
17
18 **possess a β -elimination activity.**
19
20

21
22
23 According to our study, TGAM_1277 and TGAM_1653 are bifunctional BER enzymes: they
24
25 are able (i) to recognize and remove oxidized bases by cleaving the *N*-glycosyl bond between
26
27 the deoxyribose and the damaged base, and (ii) to cleave the resulting abasic site leading to a
28
29 single-strand break (lyase activity). More precisely, we demonstrated that both enzymes
30
31 cleave the abasic site *via* a β -elimination process resulting in a 3'-phospho- α,β -unsaturated
32
33 aldehyde and in a 5'-phosphate extremity (Fig. 6).
34
35

36 TGAM_1277 was annotated as a putative type III endonuclease.² Indeed, we have shown here
37
38 that this enzyme shares the common features of the bacterial BER enzyme, i.e.
39
40 bifunctionality, a β -elimination process and the capacity to repair hydantoin lesions (5-
41
42 OHdHyd and 5-OH-5-Me-dHyd).^{22, 48} By contrast with bacterial EndoIII, we found that
43
44 TGAM_1277 is also able to cleave 8-oxo-dGuo, particularly when mispaired with T or G.
45
46 This variety of substrates is also found for the *E. coli* enzyme Fpg, which presents a β,δ -
47
48 elimination mechanism.
49
50

51
52 When compared with TGAM_1277, TGAM_1653 shows a more restrictive panel of
53
54 substrates as it does not cleave oxidized pyrimidines and is not as efficient as TGAM_1277 in
55
56 cleaving 8-oxo-dGuo when mispaired, particularly to A and G. This profile of substrates is
57
58
59
60

close to that noted in other studies for the human BER enzyme, hOGG1 (human 8-oxoG DNA Glycosylase I).^{49, 50}

As for TGAM_1277 and TGAM_1653, the 8-oxoG DNA glycosylase from the sulfate-reducing archaeon *Archeoglobus fulgidus* (Afogg) presents a bifunctional and β -elimination mechanism.⁵¹ This protein is able to repair 8-oxo-dGuo when associated with C and G, and to a lesser extent when mispaired with T and A. *Pyrobaculum aerophilum* 8-oxoG DNA glycosylase (Pa-AGOG), a formative member of a new DNA glycosylase family, also demonstrated a bifunctional and β -elimination process.¹⁰ Structural studies indicated that this enzyme has an HhH-GPD motif (helix-hairpin-helix associated with a glycine-proline-rich sequence, followed by a conserved aspartate) that is significantly different in terms of amino acid sequence compared to that of hOGG1, but similar in terms of tridimensional structure.

An alignment of the Pa-AGOG, TGAM_1277 and TGAM_1653 polypeptides is shown in Fig. 9. Pa-AGOG and TGAM_1653 display only 25% sequence identity and 43% similarity, but *P. aerophilum* and *T. gammatolerans* belong to two separate phyla of Archaea (Crenarcheota vs Euryarcheota). However, residues playing an important role for catalytic activity (K140, D172) or substrate recognition (Q31, K147, D218, W222) in Pa-AGOG^{10, 52} are conserved in TGAM_1653 and not in TGAM_1277 (Fig. 9). This suggests that Pa-AGOG and TGAM_1653 may belong to the same glycosylase family, even if some discrepancies are revealed. Pa-AGOG is able to repair 8-oxo-dGuo when paired with C and A but presented a stronger efficiency for 8-oxoG:G mismatches.⁵² We showed here that this strong capacity to cleave 8-oxoG:G pairs was also shared by TGAM_1277, TGAM_1653 and Afogg. It could be interesting for hyperthermophilic organisms to cleave 8-oxoG:G lesions generated by the incorporation of free 8-oxodGMP opposite to a G, this phenomenon probably being favored by high temperatures.^{10, 53} Neither Afogg nor Pa-AGOG were tested for their capacity to repair damaged pyrimidines, making the comparison with *T. gammatolerans* enzymes

difficult to extend further. We also tested cyclonucleosides as potential substrates: TGAM_1277 and TGAM_1653 did not show cleavage activity. Previously published experiments, based on eukaryote whole-cell extracts or purified enzymes, have already demonstrated that BER does not deal with these complex oxidative DNA lesions.⁵⁴⁻⁵⁷ In order to obtain a comprehensive picture of BER in *T. gammatolerans*, complementary experiments have to be performed on other putative BER factors highlighted by our previous genomic and proteomic screenings.² At least seven candidates, annotated as “Uracil-DNA glycosylase”, “Base excision DNA repair protein, HhH-GPD superfamily”, “AP endonuclease family 2” or “endonuclease V”, have been identified. Study of these archaeal enzymes can lead to the discovery of original DNA repair activities, as revealed by the recent characterization of EndoQ found in *Pyrococcus furiosus* and *T. kodakarensis*.⁵⁸ This endonuclease is able to repair deaminated bases generated for example by ionizing radiation or elevated temperatures. Glycosylases and lyases are the very first enzymes involved in the BER system. It may also be interesting to study other BER proteins involved in the later stages of the pathway. For example, it would be relevant to focus attention on BER factors processing single strand breaks as these latter lesions are also generated by ionizing radiation.⁴⁷ Finally, a further characterization of enzymatic activities including parameters such as thermostability or influence of pH, would be interesting, particularly for a potential use of these proteins as new tools for research and applied biotechnologies laboratories.⁵⁹

Conclusion

This work established that the basal level of the oxidative DNA damage, 8-oxo-dGuo, in *T. gammatolerans* is higher than that reported for mammalian cells. As expected, this level of damage increased with high doses of ionizing radiation. Remarkably, the yield of lesions

generated was much lower than those observed in human cell lines. One hour post irradiation, the rate of oxidative DNA damage was the same as the control background, indicating that the repair process is rather rapid. TGAM_1277 and TGAM_1653 BER enzymes were slightly upregulated after heavy radiation exposure. As shown here, these enzymes are able to initiate DNA repair of oxidized pyrimidines (TGAM_1277) and guanines (TGAM_1277 and TGAM_1653). Therefore, *T. gammatolerans* is at any time prepared to counteract DNA oxidation with this protein arsenal. While the search for cyclonucleosides showed no significant levels above technical background, other original DNA lesions such as clusters of damage^{33, 60} would be interesting to identify in radiotolerant organisms such as *T. gammatolerans*. Regarding DNA repair proteins such as TGAM_1277 and TGAM_1653, structural studies with the enzymes alone and with their DNA substrates would provide additional data about recognition and cleaving mechanisms. Further molecular studies on proteins from *T. gammatolerans*, one of the most radiotolerant organisms currently known, should bring valuable information on DNA repair in archaea. Moreover, deciphering the mechanisms explaining its extraordinary resistance to extreme conditions may have potential applications in biotechnology.⁶¹

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FUNDING INFORMATION

This study was supported by the Agence Nationale de la Recherche (ANR-12-BSV6-0012-01).

ACKNOWLEDGEMENTS

We would like to thank Dr. B. Castaing (Orléans, France) for providing Fpg and advice. We also thank Dr. V. Favaudon (Institut Curie, Orsay) for access to the ¹³⁷Cs gamma irradiation source.

SUPPORTING INFORMATION

Table S1: list of oligonucleotides used for quantitative real-time RT-PCR. Table S2: cloning and expression of TGAM_1277 and TGAM_1653 - primer pairs used to amplify coding sequences from total genomic DNA. This material is available free of charge via the Internet at <http://pubs.acs.org>.

ABBREVIATIONS

5',8-cyclodAdo, 5',8-cyclo-2'-deoxyadenosine; 5',8-cyclodGuo, 5',8-cyclo-2'-deoxyguanosine; 5-OH-5-Me-dHyd, 1-[2-deoxy-β-D-erythro-pentofuranosyl]-5-hydroxy-5-methylhydantoin; 5-OH-dHyd, 1-[2-deoxy-β-D-erythro-pentofuranosyl]-5-hydroxyhydantoin; 8-oxo-dGuo, 8-oxo-2'-deoxyguanosine; 8-oxoG, 8-oxo-7,8-dihydroguanine; Afogg, 8-oxoG DNA glycosylase from the sulfate-reducing archaeon *Archeoglobus fulgidus*; BER, Base Excision Repair; CDS, Coding DNA Sequence; Endo III, type III endonuclease; ESI, ElectroSpray Ionisation; FapyAde, 4,6-diamino-5-formamidopyrimidine; FapyGua, 2,6-diamino-4-hydroxy-5-formamidopyrimidine; Fpg, formamidopyridine DNA N-glycosylase; FPLC, Fast Protein Liquid Chromatography; HhH-GPD motif, helix-hairpin-helix associated

1
2
3 with a glycine-proline-rich sequence, followed by a conserved aspartate; hOGG1, human 8-
4
5 oxoG DNA Glycosylase 1; HPLC-MS/MS, HPLC coupled to tandem mass spectrometry;
6
7 kGy, kiloGray; MALDI-TOF, Matrix-Assisted Laser Desorption Ionization – Time Of Flight;
8
9 Mb, megabase; NSAF, normalized spectral abundance factor; ODN, oligodeoxynucleotide;
10
11 Pa-AGOG, *Pyrobaculum aerophilum* 8-oxoG DNA glycosylase; radA, DNA repair by
12
13 homologous recombination; ROS, Reactive Oxygen Species; rpoB, DNA-directed RNA
14
15 polymerase subunit B; RT-PCR, Reverse Transcription PCR; Tbp, TATA box-binding protein
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

- (1) Jolivet, E., L'Haridon, S., Corre, E., Forterre, P., and Prieur, D. (2003) *Thermococcus gammatolerans* sp. nov., a hyperthermophilic archaeon from a deep-sea hydrothermal vent that resists ionizing radiation. *Int. J. Syst. Evol. Microbiol.* 53, 847-851.
- (2) Zivanovic, Y., Armengaud, J., Lagorce, A., Leplat, C., Guerin, P., Dutertre, M., Anthouard, V., Forterre, P., Wincker, P., and Confalonieri, F. (2009) Genome analysis and genome-wide proteomics of *Thermococcus gammatolerans*, the most radioresistant organism known amongst the Archaea. *Genome Biology* 10, R70.
- (3) Confalonieri, F., and Sommer, S. (2011) Bacterial and archaeal resistance to ionizing radiation. *Cost Chemistry Cm0603-Melusyn Joint Meeting: Damages Induced in Biomolecules by Low and High Energy Radiations* 261, 012005.
- (4) Webb, K. M., and DiRuggiero, J. (2012) Role of Mn²⁺ and compatible solutes in the radiation resistance of thermophilic bacteria and archaea. *Archaea* 2012, 845756.
- (5) Tapias, A., Leplat, C., and Confalonieri, F. (2009) Recovery of ionizing-radiation damage after high doses of gamma ray in the hyperthermophilic archaeon *Thermococcus gammatolerans*. *Extremophiles* 13, 333-343.
- (6) Cadet, J., Douki, T., and Ravanat, J. L. (2011) Measurement of oxidatively generated base damage in cellular DNA. *Mutat. Res.* 711, 3-12.
- (7) Michaels, M. L., and Miller, J. H. (1992) The GO system protects organisms from the mutagenic effect of the spontaneous lesion 8-hydroxyguanine (7,8-dihydro-8-oxoguanine). *J. Bacteriol.* 174, 6321-6325.
- (8) Chatgililoglu, C., Ferreri, C., and Terzidis, M. A. (2011) Purine 5',8-cyclonucleoside lesions: chemistry and biology. *Chem. Soc. Rev.* 40, 1368-1382.

- (9) Belmadoui, N., Boussicault, F., Guerra, M., Ravanat, J. L., Chatgililoglu, C., and Cadet, J. (2010) Radiation-induced formation of purine 5',8-cyclonucleosides in isolated and cellular DNA: high stereospecificity and modulating effect of oxygen. *Org. Biomol. Chem.* 8, 3211-3219.
- (10) Sartori, A. A., Lingaraju, G. M., Hunziker, P., Winkler, F. K., and Jiricny, J. (2004) Pa-AGOG, the founding member of a new family of archaeal 8-oxoguanine DNA-glycosylases. *Nucleic Acids Res.* 32, 6531-6539.
- (11) Pfaffl, M. W. (2001) A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* 29, e45.
- (12) Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., and Speleman, F. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* 3, RESEARCH0034.
- (13) Lagorce, A., Fourcans, A., Dutertre, M., Bouyssiere, B., Zivanovic, Y., and Confalonieri, F. (2012) Genome-wide transcriptional response of the archaeon *Thermococcus gammatolerans* to cadmium. *Plos One* 7, e41935.
- (14) Armengaud, J., Fernandez, B., Chaumont, V., Rollin-Genetet, F., Finet, S., Marchetti, C., Myllykallio, H., Vidaud, C., Pellequer, J. L., Gribaldo, S., Forterre, P., and Gans, P. (2003) Identification, purification, and characterization of an eukaryotic-like phosphopantetheine adenylyltransferase (coenzyme A biosynthetic pathway) in the hyperthermophilic archaeon *Pyrococcus abyssi*. *J. Biol. Chem.* 278, 31078-31087.
- (15) Gabant, G., Auxilien, S., Tuszynska, I., Locard, M., Gajda, M. J., Chaussinand, G., Fernandez, B., Dedieu, A., Grosjean, H., Golinelli-Pimpaneau, B., Bujnicki, J. M., and Armengaud, J. (2006) THUMP from archaeal tRNA:m22G10 methyltransferase, a genuine autonomously folding domain. *Nucleic Acids Res.* 34, 2483-2494.

- (16) Gasparutto, D., Ait-Abbas, M., Jaquinod, M., Boiteux, S., and Cadet, J. (2000) Repair and coding properties of 5-hydroxy-5-methylhydantoin nucleosides inserted into DNA oligomers. *Chem. Res. Toxicol.* *13*, 575-584.
- (17) Muller, E., Gasparutto, D., Lebrun, C., and Cadet, J. (2001) Site-specific insertion of the (5R*) and (5S*) diastereoisomers of 1-[2-deoxy-beta-D-erythro-pentofuranosyl]-5-hydroxyhydantoin into oligodeoxyribonucleotides. *European J. Org. Chem.* *2001*, 2091-2099.
- (18) Ravanat, J. L., Douki, T., Duez, P., Gremaud, E., Herbert, K., Hofer, T., Lasserre, L., Saint-Pierre, C., Favier, A., and Cadet, J. (2002) Cellular background level of 8-oxo-7,8-dihydro-2'-deoxyguanosine: an isotope based method to evaluate artefactual oxidation of DNA during its extraction and subsequent work-up. *Carcinogenesis* *23*, 1911-1918.
- (19) Yang, Y. S., Fernandez, B., Lagorce, A., Aloin, V., De Guillen, K. M., Boyer, J. B., Dedieu, A., Confalonieri, F., Armengaud, J., and Roumestand, C. (2015) Prioritizing targets for structural biology through the lens of proteomics: the archaeal protein TGAM_1934 from *Thermococcus gammatolerans*. *Proteomics* *15*, 114-123.
- (20) Christie-Oleza, J. A., Fernandez, B., Nogales, B., Bosch, R., and Armengaud, J. (2012) Proteomic insights into the lifestyle of an environmentally relevant marine bacterium. *ISME J* *6*, 124-135.
- (21) Tchou, J., Kasai, H., Shibutani, S., Chung, M. H., Laval, J., Grollman, A. P., and Nishimura, S. (1991) 8-oxoguanine (8-hydroxyguanine) DNA glycosylase and its substrate-specificity. *Proc. Natl. Acad. Sci. U. S. A.* *88*, 4690-4694.
- (22) D'Ham, C., Romieu, A., Jaquinod, M., Gasparutto, D., and Cadet, J. (1999) Excision of 5,6-dihydroxy-5,6-dihydrothymine, 5,6-dihydrothymine, and 5-hydroxycytosine

- from defined sequence oligonucleotides by *Escherichia coli* endonuclease III and Fpg proteins: kinetic and mechanistic aspects. *Biochemistry* 38, 3335-3344.
- (23) Bhagwat, M., and Gerlt, J. A. (1996) 3'- and 5'-strand cleavage reactions catalyzed by the Fpg protein from *Escherichia coli* occur via successive beta- and delta-elimination mechanisms, respectively. *Biochemistry* 35, 659-665.
- (24) Darwanto, A., Farrel, A., Rogstad, D. K., and Sowers, L. C. (2009) Characterization of DNA glycosylase activity by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry. *Anal. Biochem.* 394, 13-23.
- (25) Frelon, S., Douki, T., Ravanat, J. L., Pouget, J. P., Tornabene, C., and Cadet, J. (2000) High-performance liquid chromatography--tandem mass spectrometry measurement of radiation-induced base damage to isolated and cellular DNA. *Chem. Res. Toxicol.* 13, 1002-1010.
- (26) Gedik, C. M., and Collins, A. (2005) Establishing the background level of base oxidation in human lymphocyte DNA: results of an interlaboratory validation study. *FASEB J.* 19, 82-84.
- (27) Ohno, M., Miura, T., Furuichi, M., Tominaga, Y., Tsuchimoto, D., Sakumi, K., and Nakabeppu, Y. (2006) A genome-wide distribution of 8-oxoguanine correlates with the preferred regions for recombination and single nucleotide polymorphism in the human genome. *Genome Res.* 16, 567-575.
- (28) Kish, A., Kirkali, G., Robinson, C., Rosenblatt, R., Jaruga, P., Dizdaroglu, M., and DiRuggiero, J. (2009) Salt shield: intracellular salts provide cellular protection against ionizing radiation in the halophilic archaeon, *Halobacterium salinarum* NRC-1. *Environ. Microbiol.* 11, 1066-1078.

- (29) Bruskov, V. I., Malakhova, L. V., Masalimov, Z. K., and Chernikov, A. V. (2002) Heat-induced formation of reactive oxygen species and 8-oxoguanine, a biomarker of damage to DNA. *Nucleic Acids Res.* 30, 1354-1363.
- (30) Palud, A., Villani, G., L'Haridon, S., Querellou, J., Raffin, J. P., and Henneke, G. (2008) Intrinsic properties of the two replicative DNA polymerases of *Pyrococcus abyssi* in replicating abasic sites: possible role in DNA damage tolerance? *Mol. Microbiol.* 70, 746-761.
- (31) Pouget, J. P., Frelon, S., Ravanat, J. L., Testard, I., Odin, F., and Cadet, J. (2002) Formation of modified DNA bases in cells exposed either to gamma radiation or to high-LET particles. *Radiat. Res.* 157, 589-595.
- (32) Douki, T., Ravanat, J. L., Pouget, J. P., Testard, I., and Cadet, J. (2006) Minor contribution of direct ionization to DNA base damage induced by heavy ions. *Int. J. Radiat. Biol.* 82, 119-127.
- (33) Regulus, P., Duroux, B., Bayle, P. A., Favier, A., Cadet, J., and Ravanat, J. L. (2007) Oxidation of the sugar moiety of DNA by ionizing radiation or bleomycin could induce the formation of a cluster DNA lesion. *Proc. Natl. Acad. Sci. U. S. A.* 104, 14032-14037.
- (34) Robinson, C. K., Webb, K., Kaur, A., Jaruga, P., Dizdaroglu, M., Baliga, N. S., Place, A., and Diruggiero, J. (2011) A major role for nonenzymatic antioxidant processes in the radioresistance of *Halobacterium salinarum*. *J. Bacteriol.* 193, 1653-1662.
- (35) Yuan, B., Wang, J., Cao, H., Sun, R., and Wang, Y. (2011) High-throughput analysis of the mutagenic and cytotoxic properties of DNA lesions by next-generation sequencing. *Nucleic Acids Res.* 39, 5945-5954.
- (36) Abraham, J., and Brooks, P. J. (2011) Divergent effects of oxidatively induced modification to the C8 of 2'-deoxyadenosine on transcription factor binding: 8,5'(S)-

- cyclo-2'-deoxyadenosine inhibits the binding of multiple sequence specific transcription factors, while 8-oxo-2'-deoxyadenosine increases binding of CREB and NF-kappa B to DNA. *Environ. Mol. Mutagen.* 52, 287-295.
- (37) Jasti, V. P., Das, R. S., Hilton, B. A., Weerasooriya, S., Zou, Y., and Basu, A. K. (2011) (5'S)-8,5'-Cyclo-2'-deoxyguanosine is a strong block to replication, a potent pol V-dependent mutagenic lesion, and is inefficiently repaired in *Escherichia coli*. *Biochemistry* 50, 3862-3865.
- (38) Walmacq, C., Wang, L., Chong, J., Scibelli, K., Lubkowska, L., Gnat, A., Brooks, P. J., Wang, D., and Kashlev, M. (2015) Mechanism of RNA polymerase II bypass of oxidative cyclopurine DNA lesions. *Proc. Natl. Acad. Sci. U. S. A.* 112, E410-419.
- (39) Wang, J., Clauson, C. L., Robbins, P. D., Niedernhofer, L. J., and Wang, Y. (2012) The oxidative DNA lesions 8,5'-cyclopurines accumulate with aging in a tissue-specific manner. *Aging Cell* 11, 714-716.
- (40) Tilstra, J. S., Robinson, A. R., Wang, J., Gregg, S. Q., Clauson, C. L., Reay, D. P., Nasto, L. A., St Croix, C. M., Usas, A., Vo, N., Huard, J., Clemens, P. R., Stolz, D. B., Guttridge, D. C., Watkins, S. C., Garinis, G. A., Wang, Y., Niedernhofer, L. J., and Robbins, P. D. (2012) NF-kappaB inhibition delays DNA damage-induced senescence and aging in mice. *J. Clin. Invest.* 122, 2601-2612.
- (41) Mitra, D., Luo, X., Morgan, A., Wang, J., Hoang, M. P., Lo, J., Guerrero, C. R., Lennerz, J. K., Mihm, M. C., Wargo, J. A., Robinson, K. C., Devi, S. P., Vanover, J. C., D'Orazio, J. A., McMahon, M., Bosenberg, M. W., Haigis, K. M., Haber, D. A., Wang, Y., and Fisher, D. E. (2012) An ultraviolet-radiation-independent pathway to melanoma carcinogenesis in the red hair/fair skin background. *Nature* 491, 449-453.

(42) Williams, E., Lowe, T. M., Savas, J., and DiRuggiero, J. (2007) Microarray analysis of the hyperthermophilic archaeon *Pyrococcus furiosus* exposed to gamma irradiation. *Extremophiles* 11, 19-29.

(43) Dorazi, R., Gotz, D., Munro, S., Bernander, R., and White, M. F. (2007) Equal rates of repair of DNA photoproducts in transcribed and non-transcribed strands in *Sulfolobus solfataricus*. *Mol. Microbiol.* 63, 521-529.

(44) Christmann, M., and Kaina, B. (2013) Transcriptional regulation of human DNA repair genes following genotoxic stress: trigger mechanisms, inducible responses and genotoxic adaptation. *Nucleic Acids Res.* 41, 8403-8420.

(45) Parsons, J. L., and Dianov, G. L. (2013) Co-ordination of base excision repair and genome stability. *DNA Repair (Amst)* 12, 326-333.

(46) Jeong, H. G., Youn, C. K., Cho, H. J., Kim, S. H., Kim, M. H., Kim, H. B., Chang, I. Y., Lee, Y. S., Chung, M. H., and You, H. J. (2004) Metallothionein-III prevents gamma-ray-induced 8-oxoguanine accumulation in normal and hOGG1-depleted cells. *J. Biol. Chem.* 279, 34138-34149.

(47) Dianov, G. L., and Hubscher, U. (2013) Mammalian base excision repair: the forgotten archangel. *Nucleic Acids Res.* 41, 3483-3490.

(48) Gasparutto, D., Muller, E., Boiteux, S., and Cadet, J. (2009) Excision of the oxidatively formed 5-hydroxyhydantoin and 5-hydroxy-5-methylhydantoin pyrimidine lesions by *Escherichia coli* and *Saccharomyces cerevisiae* DNA N-glycosylases. *Biochim. Biophys. Acta* 1790, 16-24.

(49) Morland, I., Rolseth, V., Luna, L., Rognes, T., Bjoras, M., and Seeberg, E. (2002) Human DNA glycosylases of the bacterial Fpg/MutM superfamily: an alternative pathway for the repair of 8-oxoguanine and other oxidation products in DNA. *Nucleic Acids Res.* 30, 4926-4936.

- (50) Leipold, M. D., Workman, H., Muller, J. G., Burrows, C. J., and David, S. S. (2003) Recognition and removal of oxidized guanines in duplex DNA by the base excision repair enzymes hOGG1, yOGG1, and yOGG2. *Biochemistry* 42, 11373-11381.
- (51) Chung, J. H., Suh, M. J., Park, Y. I., Tainer, J. A., and Han, Y. S. (2001) Repair activities of 8-oxoguanine DNA glycosylase from *Archaeoglobus fulgidus*, a hyperthermophilic archaeon. *Mutat. Res.* 486, 99-111.
- (52) Lingaraju, G. M., Protá, A. E., and Winkler, F. K. (2009) Mutational studies of Pa-AGOG DNA glycosylase from the hyperthermophilic crenarchaeon *Pyrobaculum aerophilum*. *DNA Repair (Amst)* 8, 857-864.
- (53) Thivyanathan, V., Somasunderam, A., Hazra, T. K., Mitra, S., and Gorenstein, D. G. (2003) Solution structure of a DNA duplex containing 8-hydroxy-2'-deoxyguanosine opposite deoxyguanosine. *J. Mol. Biol.* 325, 433-442.
- (54) Brooks, P. J., Wise, D. S., Berry, D. A., Kosmoski, J. V., Smerdon, M. J., Somers, R. L., Mackie, H., Spoonde, A. Y., Ackerman, E. J., Coleman, K., Tarone, R. E., and Robbins, J. H. (2000) The oxidative DNA lesion 8,5'-(S)-cyclo-2'-deoxyadenosine is repaired by the nucleotide excision repair pathway and blocks gene expression in mammalian cells. *J. Biol. Chem.* 275, 22355-22362.
- (55) Kuraoka, I., Bender, C., Romieu, A., Cadet, J., Wood, R. D., and Lindahl, T. (2000) Removal of oxygen free-radical-induced 5',8-purine cyclodeoxynucleosides from DNA by the nucleotide excision-repair pathway in human cells. *Proc. Natl. Acad. Sci. U. S. A.* 97, 3832-3837.
- (56) Corne, C., Fiche, J. B., Gasparutto, D., Cunin, V., Suraniti, E., Buhot, A., Fuchs, J., Calemczuk, R., Livache, T., and Favier, A. (2008) SPR imaging for label-free multiplexed analyses of DNA N-glycosylase interactions with damaged DNA duplexes. *Analyst* 133, 1036-1045.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(57) Pande, P., Das, R. S., Sheppard, C., Kow, Y. W., and Basu, A. K. (2012) Repair efficiency of (5'S)-8,5'-cyclo-2'-deoxyguanosine and (5'S)-8,5'-cyclo-2'-deoxyadenosine depends on the complementary base. *DNA Repair (Amst)* 11, 926-931.

(58) Shiraishi, M., Ishino, S., Yamagami, T., Egashira, Y., Kiyonari, S., and Ishino, Y. (2015) A novel endonuclease that may be responsible for damaged DNA base repair in *Pyrococcus furiosus*. *Nucleic Acids Res.* 43, 2853-2863.

(59) Gogos, A., and Clarke, N. D. (1999) Characterization of an 8-oxoguanine DNA glycosylase from *Methanococcus jannaschii*. *J. Biol. Chem.* 274, 30447-30450.

(60) Ravanat, J. L., Breton, J., Douki, T., Gasparutto, D., Grand, A., Rachidi, W., and Sauvaigo, S. (2014) Radiation-mediated formation of complex damage to DNA: a chemical aspect overview. *Br. J. Radiol.* 87, 20130715.

(61) Grasso, S., and Tell, G. (2014) Base excision repair in Archaea: back to the future in DNA repair. *DNA Repair (Amst)* 21, 148-157.

(62) Corpet, F. (1988) Multiple sequence alignment with hierarchical-clustering. *Nucleic Acids Res.* 16, 10881-10890.

FIGURE LEGENDS

Figure 1.

Chemical structures of the DNA lesions mentioned in this work. 8-oxo-dGuo, 5-OH-dHyd and 5-OH-5-Me-dHyd represent single oxidation products of 2'-deoxyguanosine, 2'-deoxycytidine and thymidine, respectively. 5'R-cyclo-dAdo represents an example of 5',8-cyclonucleosides that are considered as a family of complex oxidative DNA damage. 8-oxo-dGuo and 5',8-cyclonucleosides were measured in *T. gammatolerans* DNA at basal levels and after irradiation. 8-oxo-dGuo, 5-OH-dHyd, 5-OH-5-Me-dHyd and 5',8-cyclonucleosides were tested as potential substrates for the *T. gammatolerans* enzymes, TGAM_1653 and TGAM_1277.

Figure 2.

Purification of recombinant TGAM_1277 and TGAM_1653 by FPLC affinity chromatography on His-Trap columns after expression in *E. coli*. TGAM_1277 (4.8 µg) and TGAM_1653 (5.1 µg) were resolved on a 10% SDS-PAGE gel and stained with Instant Blue (Expedeon).

Figure 3.

Levels of 8-oxo-dGuo in *T. gammatolerans*. The “basal level” of 8-oxo-dGuo was determined with cells that were not irradiated neither kept on ice. For gamma irradiation experiments performed at 2.5 kGy or 5.0 kGy: cells were harvested immediately after treatment (0h) or after one hour of recovery in culture media (1h). Nonirradiated controls were stored on ice while the other cells were irradiated. Vertical bars: +/- SD. For each condition, five measurements were performed: two technical replicates for the first biological replicate and three technical replicates for the second biological replicate.

* statistically significant difference, according to the Mann-Whitney test, with $p < 0.05$.

Figure 4.

Expression profiles of TGAM_1277 and TGAM_1653 in *T. gammatolerans* cells exposed to γ -rays at a dose of 2.5 and 5.0 kGy. TGAM_1277 and TGAM_1653 expression were quantified by real-time RT-PCR using RNA extracted from control cells (not irradiated: NI), cells collected immediately after irradiation (T0) and cells incubated for 1 h or 2 h post irradiation (T1; T2). TGAM_1277 and TGAM_1653 expression were calculated relative to the transcript levels of constitutively expressed genes (TGAM_0829, TGAM_01054, TGAM_1165, TGAM_1926) according to the formula: $2^{-\Delta Ct} = 2^{-(Ct_{\text{geneX}} - Ct_{\text{RefGenes}})}$. Each data point is the average of two biological replicates. The standard deviation is indicated by error bars. The TGAM_1277 and TGAM_1653 genes were expressed in all conditions tested.

Figure 5.

A. Base excision activities of Fpg, TGAM_1653 and TGAM_1277 assessed with three oxidative DNA lesions: 8-oxo-dGuo, 5-OH-dHyd and 5-OH-Me-dHyd. 22mer ODNs sequence: 5' -CAC TTC GG¹A T²CG TGA CTG ATC T-3' (control). In modified ODNs: G¹ was replaced by 8-oxo-dGuo, and T² by 5-OH-dHyd or 5-OH-5-Me-dHyd. 100 fmol ODNs were incubated with 2 pmol enzyme. For Fpg, the cleavage was carried out at 37°C. For TGAM_1653 and TGAM_1277, the incubation was initiated at 37°C and the temperature was progressively increased to 65°C. The different sizes of the cleavage products are explained by the positions of the damaged bases in the ODN sequences and by the enzymes presenting different excision processes. B. Base excision activities of Fpg, TGAM_1653 and TGAM_1277 on probes containing 8-oxo-dGuo. The influence of the opposite nucleoside was

evaluated by using, for each enzyme, a set of four ODNs with dC, dT, dA or dG paired with the oxidative lesion.

Figure 6.

Excision reaction catalyzed by bifunctional glycosylases *via* β or β,δ -elimination mechanisms. Successive cleavages of 3' and 5' - strands on each side of the abasic site lead to deoxyribose residue excision. Numbers 1 to 6 were attributed to each reaction product to match with fragments obtained by MALDI-TOF mass spectrometry analysis (Figures 7 and 8). Number 2 (not represented) stands for the complementary strand.

Figure 7.

MALDI-TOF MS spectra resulting from the excision by Fpg, TGAM_1653 or TGAM_1277 of ODNs containing 8-oxo-dGuo. Ten pmol ODNs were incubated with 20 pmol enzyme. The incubation was initiated at 37°C and the temperature was progressively increased to 65°C. The nature of the detected fragments (from 1 to 6) is described in Figure 6. “bis” indicates a secondary ion presenting two charges.

Figure 8.

MALDI-TOF MS spectra resulting from the excision by Fpg, Endo III, TGAM_1653 or TGAM_1277 of ODNs containing 5-OH-5-Me-dHyd. Ten pmol ODNs were incubated with 20 pmol enzyme. The incubation was initiated at 37°C and the temperature was progressively increased to 65°C. The nature of the detected fragments (from 1 to 6) is described in Figure 6. “bis” indicates a secondary ion presenting two charges.

Figure 9.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Sequence alignment of *Pyrobaculum aerophilum* 8-oxoG DNA glycosylase (Pa-AGOG), TGAM_1653 and TGAM_1277. Identical residues are in red, residues shared by two proteins are in blue. Residues playing an important role for catalytic activity or 8-oxo-dGuo recognition in Pa-AGOG (Q31, K140, K147, D172, D218 and W222) are indicated in green. The alignment was performed using the MultAlin tool⁶² followed by manual editing.

FIGURES

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Pa-AGOG	1	MAAESQLKRVISTLRRLGIEEVKLERRDPQYFAVNVVKKRGETV	Q31
TGAM_1653	1	MTLDRFVRIKYEDNERVNRVVEILRELGLDCARTIEEVLDLQFDALENLRENKODEL	
TGAM_1277	1	MSLDGFTFEESWDEKRRRAEKIVEILNRTHPREKLLIGDPYRTLNCIISQRMDEV	
Consensus		m.l.d.f.....e.....t.!EiLrtrlg..e.l.ie...Dpq..a..n.....#ev	
Pa-AGOG	47	GSRLAMLNALI SYRLTGKGEERWEYFGKYFSQLEVIDLCRDFLKYIETS PFLKI QVEARK	
TGAM_1653	60	FIRLVIANALVSYQLSGKGEDWWEFSEYFSENPPEDIVEYSSFLPNSKTNRRLVAGKL	
TGAM_1277	58	TYRVWEELEFKYGDIEIANTPVDEMREFLRKGVGLWKTG	
Consensus		..tl...nal.sy.l.gkge..Weef.kyts#.e.id.....sk....lv..k.	
Pa-AGOG	107	KRALKACDYVPNLEDLGLTLRQLSHIVGARREOKTIVFTIKI LNYAYMCS	K140 K147
TGAM_1653	120	KRIERVEPFLSPLSISEIRDYYFNGMERLDELARVGAARSAKTIVFAVMFPGYAGRIA	
TGAM_1277	100	EWIVKASRIILEKYGGKVPDDIKELMKLPGIGRKCANIVLAYGFGRAIPVD	
Consensus		krl.k.....l.....y.....#L...ga.R..ktzVfa.k...yA....	
Pa-AGOG	157	ROMNRLPFDIPVVDYRVARLTWCAGLIDFPPEALERYEAVQKIWDNAVARETGIPPLH	D172
TGAM_1653	180	FSAFVYPMAIEIPDDVRINAYTKRPTSEPPVSFWGRIAETGIPPLH	
TGAM_1277	152	THVNRISKRGLAPFVPEKVEEYLMELIPKEKVIYVSHANVDRGRSICRPIEPKCE	
Consensus		..vnr..p..i.iP.dvr...t...l...pP.e.....wgt.a.etgiPplh	
Pa-AGOG	217	LDTLWLAGRAVLYGENLHGVPEVIALFQWRGGCRPPSE	D218 W222
TGAM_1653	228	IDSILWFLGGKGEVLRRLKHOGKAERILELRDL	
TGAM_1277	210	SCPLKELCPYAKGLVRDEDIK	
Consensus		.d.llw..l.gkg.v.e..k.....t.....	