

Study Seeks Candidates For Blue Shrimp Polyculture In New Caledonia



The outdoor culture system was set up at Saint Vincent station in Boulouparis, New Caledonia, to help find suitable fish for polyculture with shrimp that could contribute to sustaining the area's shrimp industry.

Dr. Trung Cong Luong
Nha Trang University
Nha Trang, Vietnam

Dr. Sébastien Hochard
ADECAL/Technopole/Programme
ZoNéCo
Nouméa, New Caledonia

Florence Royer
IFREMER Lead
Nouméa, New Caledonia

Dr. Hugues Lemonnier
IFREMER Lead
B. P. 2059
98846 Nouméa, New Caledonia
hugues.lemonnier@ifremer.fr

Pr. Yves Letourneur
Université de la Nouvelle-Calédonie
Laboratoire Live
Nouméa, New Caledonia

Summary:

Seasonal pathologies reduce the profitability and sustainability of the shrimp-farming industry in New Caledonia. A study was therefore conducted to estimate the effects of polyculture of blue shrimp with goldline rabbitfish or mullet on production performance and environmental quality. The fish did not affect shrimp production, and the combined shrimp/fish yields were significantly greater than the yield from shrimp monoculture. Changes in environmental quality in all treatments were few and minor throughout the culture period.

The farming of blue shrimp, *Litopenaeus stylirostris*, has been practiced for over 30 years in New Caledonia, where semi-intensive growout techniques produce an annual average production of around 2,000 mt. The shrimp industry, however, is affected by seasonal mortalities caused by "syndrome 93" during the cold seasons or "summer syndrome" dur-

ing the warmer months. These pathologies reduce profitability and affect the sustainability of the industry.

Blue Shrimp/Fish Polyculture

Shrimp/fish polyculture has demonstrated itself as an ecologically and economically sound method of sustainable shrimp culture and an effective choice for solving and/or minimizing some of the problems the shrimp industry currently faces. Studies on polyculture of penaeid shrimp with omnivorous fish have found a remarkable improvement in water quality. The fish help to reduce organic wastes and selective phytoplankton, clean the pond bottom and increase nutrient uptake into cultured animals.

Based on a literature review, herbivorous siganids and omnivorous mullets could be excellent candidates for polyculture with shrimp because the fish help decrease the impacts of disease, prevent deterioration of the environment and increase the production in the ponds.

The authors therefore conducted an experiment on polyculture of blue shrimp with goldline rabbitfish, *Siganus lineatus*, or *Mugil* species mullet in a mesocosm system to evaluate the effects of polycul-

ture on production performance and environmental quality. Finding suitable fish for polyculture with shrimp would contribute to sustaining the shrimp industry in New Caledonia.

Study Setup

With grant support from the South Province of New Caledonia Agreement C.486-10, the study was conducted in an outdoor system that consisted of 16 circular, 1.7-m² fiberglass tanks. Each tank contained a 15-cm layer of sediment and a 75-cm-high water column with 1,275-L volume. A daily water exchange of around 10% was applied.

Shrimp fry weighing approximately 2.9 g each were randomly stocked into the experimental tanks at density of 15/m². One month later, rabbitfish or mullet fingerlings were added to the shrimp tanks to form polyculture treatments.

Hatchery-sourced rabbitfish weighing 25.5 g were stocked into the shrimp tanks at either 1.2 fish/m² or 2.4 fish/m². In other tanks, 20.5-g wild-caught mullet were added at 1.2 fish/m². A control treatment had no fish. All treatments were randomly distributed among tanks with four replicates per treatment.

Shrimp were fed a commercial pel-



Mullet and rabbitfish are potential species for polyculture with blue shrimp.

leted feed at 3 to 5% of shrimp biomass daily. At the beginning and end of 12 weeks of culture, shrimp and fish were sampled and individually weighted to evaluate the growth performance. Environmental parameters were regularly sampled and analyzed to estimate environmental variations.

Results

At 62.5%, shrimp survival was lowest in mullet polyculture, while the 80.8% survival in the high-density rabbitfish polyculture treatment was highest. In addition, the standard deviation of shrimp survival rate decreased from 20.7% in the control to 7.0% in the high-density rabbitfish treatment.

Although shrimp yields were not significantly different among treatments, the values of 145.4-170.3 g/m² for rabbitfish polyculture were greater than the 143.2 g/m² yield for the control. The mullet treatment had a slightly lower value of 130.3 g/m².

These results showed the presence of the fish had no negative effect on shrimp growth. The shrimp growth obtained in this study was also similar to the results recorded in other research on blue shrimp monoculture in earthen ponds in New Caledonia.

The 100% fish survival and growth rates were similar in all polyculture treatments. The fish yield, however, was significantly higher in the high-density treatment with rabbitfish. The total combined shrimp and fish yields were higher in the polyculture treatments than the shrimp production in the control (Figure 1).

The overall feed-conversion ratio (FCR) was significantly lower in the




Figure 1. Production yields in experimental treatments. Bars with different letters are significantly different ($P < 0.05$).




Figure 2. Feed conversion in experimental treatments. Bars with different letters are significantly different ($P < 0.05$).

high-density rabbitfish polyculture than those in the mullet polyculture and the control (Figure 2). Furthermore, rabbitfish polyculture decreased FCR by 31.6% and 47.4% in the low- and high-density rabbitfish treatments, respectively, compared with the control.

Water Quality

All the results indicated that there were few changes in the environmental quality in all treatments throughout the culture period. The 22.4-26.5 °C temperature and 5.5-9.4 mg/L dissolved oxygen, salinity and 8.2 pH levels were similar in all treatments and generally within suitable ranges for shrimp and fish growth.

Turbidity, total dissolved nitrogen and total ammonia nitrogen values were only slightly different among treatments. Although the mean soluble reactive phosphorus concentration was significantly higher in the high-density rabbitfish polyculture than those in the other treatments, its values maintained low in all treatments (0.15 to 0.26 μ).

No significant difference was observed in all sediment parameters among polyculture treatments and the control. Values for

pH ranged 6.8 to 6.9. Trends in redox potential were similar across all treatments.

Perspectives

Rabbitfish have fast growth, tolerance of crowding and a range of environmental conditions, good flavor, and high demand and market prices. The results of this study indicated that polyculture of blue shrimp with goldline rabbitfish produced an excellent secondary crop and also improved food utilization without adverse effects on shrimp production and additional environmental degradation in the culture system.

Although shrimp/mullet polyculture did not get high results for production performance and environmental improvement, mullets are also potential species for polyculture with shrimp. The omnivorous fish could help control organic matter in polyculture systems.

Further research needs to be conducted to find the optimum density and biomass of fish and shrimp to achieve consistently high production and maximum environmental efficiency. Studies of pathological conditions could test the effects of co-culture on conditions such as vibriosis.