

Supporting Information for

**Zooplankton community response to the winter 2013 deep convection process in the NW Mediterranean Sea**

Katty Donoso<sup>1</sup>, François Carlotti<sup>1</sup>, Marc Pagano<sup>1</sup>, Brian P. V. Hunt<sup>2,3</sup>, Rubén Escribano<sup>4</sup>,  
Léo Berline<sup>1</sup>

<sup>1</sup>Aix-Marseille Univ, Univ Toulon, CNRS, IRD, MIO UM 110, Mediterranean Institute of Oceanography,  
Marseille, France.

<sup>2</sup>University of British Columbia, Department of Earth, Ocean and Atmospheric Sciences, Vancouver, V6T 1 Z4,  
British Columbia, Canada.

<sup>3</sup>Hakai Institute, P.O. Box 309, Heriot Bay, BC, V0P 1H0, Canada.


<sup>4</sup>Millennium Institute of Oceanography (IMO), University of Concepcion, Chile.

**Contents of this file**

Figure S1

**Introduction**

This supporting information provides the averaged vertical profiles (0-250m) from each cluster. This complements the boxplots for environmental variables in Figure 3.



**Figure S1.** Average vertical profile, for temperature, salinity, dissolved oxygen and chlorophyll-a, black line corresponds to the median value for each cluster, colors correspond to quantiles (0.025 to 0.975 range) and clusters of Figure 2.