

Title of article: Rock-type control of Ni, Cr and Co phytoavailability in ultramafic soils  
 Journal: Plant and Soil  
 Authors: Artur Pędziwiatr, Jakub Kierczak, Jarosław Waroszewski, Gildas Ratié, Cécile Quantin, Emmanuel Ponzevera  
 Affiliations: Wrocław University of Environmental and Life Sciences, Institute of Soil Sciences and Environmental Protection, Grunwaldzka 53, 50-537 Wrocław, Poland  
 University of Wrocław, Institute of Geological Sciences, Pl. Maksa Borna 9, 50-204 Wrocław, Poland  
 UMR 8148 GEOPS, Université Paris Saclay - Université Paris Sud - CNRS, Rue du Belvédère 504, 91405 Orsay Cedex, France  
 IFREMER, Centre de Brest, Unité Géosciences Marines, 29280, Plouzané, France

Corresponding author: artur.pedziwiatr@upwr.edu.pl

Online Resource 1. Nickel, Cr and Co concentrations in lechates from the DTPA-CaCl<sub>2</sub> and EDTA extractions.

Horizon	Concentration in lechates (EDTA) <sup>a</sup>			Proportion extracted by EDTA			Concentration in lechates (DTPA-CaCl <sub>2</sub> )	Proportion extracted by DTPA-CaCl <sub>2</sub>
	Ni	Cr	Co	Ni	Cr	Co	Ni	Ni
	mg kg <sup>-1</sup>			%			mg kg <sup>-1</sup>	%
Site 1: Szklary Massif								
A	293.60±21.71	2.22±0.11	18.32±0.92	13.87	0.06	11.17	137.73±0.71	6.51
ABw	247.27±15.09	1.09±0.14	24.31±1.96	10.56	0.04	15.68	77.91±1.43	3.33
R	116.07±9.72	1.25±0.08	2.73±0.20	5.20	0.05	2.79	6.85±0.19	0.31
Site 2: Jordanów								
ABw	42.30±2.40	0.22±0.20	3.10±1.01	5.18	0.01	3.73	32.61±3.80	3.99
Bw	23.76±1.13	0.36±0.23	4.09±0.50	1.52	0.02	3.62	19.30±0.20	1.23
BwC	31.63±0.21	0.11±0.06	4.99±1.20	0.80	0.004	2.97	27.43±0.60	0.70
R	49.79±3.02	0.09±0.03	1.33±1.02	3.16	0.004	1.34	8.09±0.22	0.51

<sup>a</sup> after Kierczak et al. (2016); results reported as mean value ± standard deviation