***Pentaplacodinium saltonense* gen. et sp. nov.**
(Dinophyceae) and its relationship to the cyst-defined genus
Operculodinium* and yessotoxin-producing *Protoceratium
reticulatum

Mertens Kenneth ^{1,*}, Carbonell-Moore Maria Consuelo ², Pospelova Vera ³, Head Martin J. ⁴,
 Highfield Andrea ⁵, Schroeder Declan ^{5,6}, Gu Haifeng ⁷, Andree Karl B. ⁸, Fernandez Margarita ⁸,
 Yamaguchi Aika ⁹, Takano Yoshihito ¹⁰, Matsuoka Kazumi ¹⁰, Nézan Elisabeth ¹¹, Bilien Gwenael ¹¹,
 Okolodkov Yuri ¹², Koike Kazuhiko ¹³, Hoppenrath Mona ¹⁴, Pfaff Maya ¹⁵, Pitcher Grant ¹⁶,
 Al-Muftah Abdulrahman ¹⁷, Rochon André ¹⁸, Lim Po Teen ¹⁹, Leaw Chui Pin ¹⁹, Lim Zhen Fei ¹⁹,
 Ellegaard Marianne ²⁰

¹ Research Unit for Palaeontology, Ghent University, Krijgslaan 281 s8, 9000 Ghent, Belgium

² Oregon State University, Department of Botany and Plant Pathology, College of Agricultural Sciences, 2082 Cordley Hall, Corvallis, OR 97331-2902, USA

³ School of Earth and Ocean Sciences, University of Victoria, OEASB A405, P.O. Box 1700 Box 1700 STN CSC, Victoria, BC, V8W 2Y2, Canada

⁴ Department of Earth Sciences, Brock University, 1812 Sir Isaac Brock Way, St. Catharines, Ontario, L2S 3A1, Canada

⁵ The Marine Biological Association of the United Kingdom, Citadel Hill, Plymouth PL1 2PB, United Kingdom

⁶ School of Biological Sciences, University of Reading, Reading RG6 6AJ, United Kingdom

⁷ Third Institute of Oceanography, SOA, Xiamen 361005, China

⁸ IRTA, Crta. Poble Nou, Km 5.5, 43540 Sant Carles de la Ràpita, Spain

⁹ Kobe University Research Center for Inland Seas, Kobe 657-8501, Japan

¹⁰ Institute for East China Sea Research (ECSER), Nagasaki University, 1551-7, Taira-machi, Nagasaki, 851-2213, Japan

¹¹ Ifremer, LER BO, Station de Biologie Marine, Place de la Croix, BP40537, F-29185 Concarneau Cedex, France

¹² Universidad Veracruzana, Instituto de Ciencias Marinas y Pesquerías, Calle Hidalgo núm. 617, Colonia Río Jamapa, Boca del Río, 94290 Veracruz, Mexico

¹³ Graduate School of Biosphere Science, Hiroshima University, Kagamiyama 1-4-4, Higashi-Hiroshima, Hiroshima 739-8528, Japan

¹⁴ Senckenberg am Meer, Deutsches Zentrum für Marine Biodiversitätsforschung (DZMB), Südstrand 44, D-26382 Wilhelmshaven, Germany

¹⁵ Marine Biology Research Center, Ma-RE Institute, Zoology Department, University of Cape Town, Rondebosch 7701, South Africa

¹⁶ Marine and Coastal Management, Private Bag X2, Rogge Bay 8012, Cape Town, South Africa

¹⁷ Department of Biological and Environmental Sciences, Qatar University, Doha, Qatar

¹⁸ Institut des sciences de la mer de Rimouski (ISMER), Université du Québec à Rimouski, 310 allée des Ursulines, Rimouski, QC, G5L 3A1, Canada

¹⁹ Institute of Ocean and Earth Sciences, University of Malaya, 16310 Bachok, Kelantan, Malaysia

²⁰ Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, DK-1871, Frederiksberg, Denmark

* Corresponding author : Kenneth Neil Mertens, email address : kenneth.mertens29@gmail.com

Abstract :

Strains of a dinoflagellate from the Salton Sea, previously identified as *Protoceratium reticulatum* and yessotoxin producing, have been reexamined morphologically and genetically and *Pentaplacodinium saltonense* n. gen. et sp. is erected to accommodate this species. *Pentaplacodinium saltonense* differs from *Protoceratium reticulatum* (Claparède et Lachmann 1859) Bütschli 1885 in the number of precingular plates (five vs. six), cingular displacement (two widths vs. one), and distinct cyst morphology. Incubation experiments (excystment and encystment) show that the resting cyst of *Pentaplacodinium saltonense* is morphologically most similar to the cyst-defined species *Operculodinium israelianum* (Rossignol, 1962) Wall (1967) and *O. psilatum* Wall (1967). Collections of comparative material from around the globe (including *Protoceratium reticulatum* and the genus *Ceratocorys*) and single cell PCR were used to clarify molecular phylogenies. Variable regions in the LSU (three new sequences), SSU (12 new sequences) and intergenic ITS 1–2 (14 new sequences) were obtained. These show that *Pentaplacodinium saltonense* and *Protoceratium reticulatum* form two distinct clades. *Pentaplacodinium saltonense* forms a monophyletic clade with several unidentified strains from Malaysia. LSU and SSU rDNA sequences of three species of *Ceratocorys* (*C. armata*, *C. gourreti*, *C. horrida*) from the Mediterranean and several other unidentified strains from Malaysia form a well-supported sister clade. The unique phylogenetic position of an unidentified strain from Hawaii is also documented and requires further examination. In addition, based on the V9 SSU topology (bootstrap values >80%), specimens from Elands Bay (South Africa), originally described as *Gonyaulax grindleyi* by Reinecke (1967), cluster with *Protoceratium reticulatum*. The known range of *Pentaplacodinium saltonense* is tropical to subtropical, and its cyst is recorded as a fossil in upper Cenozoic sediments. *Protoceratium reticulatum* and *Pentaplacodinium saltonense* seem to inhabit different niches: motile stages of these dinoflagellates have not been found in the same plankton sample.

Highlights

► *Pentaplacodinium saltonense* gen. et sp. n. is described from the Salton Sea. ► Cover plate, 5 precingulars, contact 1' and Sa, shape, displacement are distinct. ► The genus is well-separated in LSU, SSU and ITS phylogenies. ► The chorate cysts of *P. saltonense*, correspond to *Operculodinium psilatum*. ► Distribution of *Pentaplacodinium saltonense* is considered subtropical to tropical.

Keywords : *Pentaplacodinium*, *Protoceratium*, Precingular plates, Salton Sea, *Ceratocorys*, *Operculodinium*, Cribroperidinioideae

1. Introduction

The dinoflagellate genus *Protoceratium* was erected by Bergh (1881, p. 242) with *Protoceratium aceros* as the type species (fig. 36), which was recovered from Strib, Denmark. Bütschli (1885, p. 1007, plate 52, fig. 2) considered *Peridinium reticulatum* as described earlier by Claparède and Lachmann (1858) from Bergen Fjord, Norway, as a senior synonym, and he proposed the combination *Protoceratium reticulatum*. He also considered *Clathrocysta reticulata* as described by Stein (1883) a junior synonym. The plate formula for *P. reticulatum*, 4', 0a, 6'', 6''', 1p, 1''''', was first provided by Wołoszyńska (1929) through the study of Baltic [Sea](#) specimens. Reinecke (1967) erected the name *Gonyaulax grindleyi* for specimens from Elands Bay in Cape Town, South Africa, with the tabulation 3', 1a, 6'', 6''', 1p, 1'''''. Based on a detailed study of the theca of *Protoceratium reticulatum* from the North Sea, Stosch (1969) considered *G. grindleyi* to be a junior synonym of *P. reticulatum*, although he considered it assignable to the genus *Gonyaulax*. Dodge (1989) agreed with the tabulation of Reinecke (1967), but retained the genus *Protoceratium* because he considered it different from his emendation of the genus *Gonyaulax*, by having only one intercalary plate. Hansen et al. (1997) restudied specimens close to the type locality of *P. aceros*, and based on the plate analysis concluded that *P. reticulatum*, *P. aceros* and *G. grindleyi* were conspecific, and agreed with the tabulation of Wołoszyńska (1929). Paez-Reyes and Head (2013) [reviewed the morphological variability reported for *P. reticulatum* and](#) concurred with Dodge (1989) in maintaining *Protoceratium* as a distinct genus from *Gonyaulax*.

~~Since the early 1900's, s~~Seven other *Protoceratium* species have been described [since the early 1900s](#), ~~and~~ the latest review of these taxa ~~was having been~~ performed by Schiller (1937 p. 322–326). Kofoed (1907) described *P. areolatum* from the tropical Pacific and emended the genus for the first time. Meunier (1910) described a very similar species from

the Kara Sea that he named *Protoceratium splendens*, which is possibly a junior synonym, as suggested by Gómez (2012). Later, Kofoed in Kofoed and Michener (1911) emended *Protoceratium* once more to include several new species from the eastern tropical Pacific that were described without illustration (*P. cancellorum*, *P. globosum*, *P. pellucidum*, *P. pepo*, *P. promissum*), and he suggested a tabulation formula for the genus: 2', 0a, 6" (?c), 6"', 0p, 3''''.

Schiller (1937) transferred *Clathrocysta aculeata* as described by Stein (1883) to *Protoceratium aculeatum*, presumably based on the fact that Bütschli (1885) [had](#) considered the genus *Clathrocysta* described by Stein 1883 as a junior synonym of *Protoceratium*.

Schiller (1937) transferred *Peridinium spinulosum* as described by Murray and Whitting (1899) to the genus *Protoceratium*. Later, Balech (1988) rediscovered this species in the South-~~W~~est Atlantic and suggested yet [another variation on the](#) tabulation for *Protoceratium*, 3', 0a, 6", 6"', 2''''', based on his observations of *Protoceratium spinulosum*.

Protoceratium reticulatum (Claparède et Lachmann) Bütschli 1885 is a very common dinoflagellate found in cold and warm waters, as well as in oceanic and neritic environments (e.g., [as *Operculodinium centrocarpum* in Zonneveld et al., 2013](#)). [-Its resting cyst distribution today reveals a strong link with the North Atlantic Current, an association traceable through the upper Cenozoic fossil record \(Hennissen et al., 2017 and references therein\)](#).

Protoceratium reticulatum is considered potentially toxic because of its production of yessotoxins (e.g. Paz et al., 2008; Sala-Pérez et al., 2016). It has been successfully isolated [and cultured](#) from many parts of the world, and grown into cultures. Cysts of *P. reticulatum* were first observed in cultures established from motile cells from the inner Oslofjord-~~ø~~, Norway by Braarud (1945). This cyst was related by Wall and Dale (1966, 1967, 1968) to the cyst-defined species described from the Miocene of Australia, *Operculodinium centrocarpum* (Deflandre et Cookson 1955) Wall 1967. [That assignation was challenged by Head and Wrenn \(1992\) and Head \(1996a\) on the grounds that *Operculodinium centrocarpum* was](#)

larger and more robust than the cysts recorded by Wall and Dale (1966) from modern sediments. However, a restudy of the holotype of *Operculodinium centrocarpum* refuted this assignment confirmed this, and the name “cyst of *Protoceratium reticulatum*” was recommended (Matsuoka et al., 1997). Wall and Dale (1968) proposed that *P. reticulatum* was also related to the cyst-defined *Operculodinium psilatum* Wall 1967 and furthermore possibly to *Operculodinium israelianum* (Rossignol 1962) Wall 1967. The cyst-defined *Pyxidinosia psilata* (Wall et Dale in Wall et al., 1973) Head 1994 was subsequently also linked to *Protoceratium reticulatum* (Dale, 1996, as *Tectatodinium psilatum*) although this connection was later questioned later (Mertens et al., 2011). Because of uncertainty regarding the links between the cysts produced by *P. reticulatum* (see Head, 2006), and cysts-cyst-defined species named based on from the fossil records, Head (1996a, 1996b) and subsequent authors used the term “*Operculodinium centrocarpum* sensu Wall and Dale, 1966” was used to describe the cysts that had first been observed by Braarud (1945) and Wall and Dale (1966). With the removal of *Pyxidinosia psilata* as a potential cyst of *Protoceratium reticulatum*, Paez-Reyes and Head (2013) argued on the basis of non-overlapping geographic distribution that the “cyst of *Protoceratium reticulatum*” was now unambiguous and should replace the term “*Operculodinium centrocarpum* sensu Wall and Dale, 1966”. That approach is followed here. Recent studies of variation in the process length of cysts of *Protoceratium reticulatum* have been related to variations in sea surface salinity and other parameters (e.g., Mertens et al. 2011; Jansson et al., 2014), and the cyst wall appears to be composed of cellulose glucan (Bogus et al. 2014). Resting cyst production through sexual reproduction has recently been demonstrated by Salgado et al. (2017).

Protoceratium reticulatum was assigned questionably to the subfamily Cribroperidinioideae by Fensome et al. (1993b) based on the presence of six precingular plates, L-type ventral organization and possible dextral torsion, which at the time had not

been documented. This assignation was confirmed by Paez-Reyes and Head (2013). ~~However,~~
~~†~~The description of the ~~very~~ similar cyst-defined *Operculodinium bahamense*, with neutral
torsion and modified L-type ventral organization, ~~which would be placed~~ allowing placement
in the subfamily Leptodinioideae, ~~either challenges~~ challenges such the present subfamilial
classification of the Gonyaulacaceae, or implies that *Operculodinium* is polyphyletic, with
both outcomes being possible -(Paez-Reyes and Head, 2013). Furthermore, molecular
phylogenetics show that *Protoceratium reticulatum* is closely related to the family
Ceratocoryaceae but not to the other extant cribroperidinean, *Lingulodinium polyedra*
(Saldarriaga et al., 2004). It should also be noted that morphological variation and sequencing
of cysts has suggested pseudocryptic speciation in *P. reticulatum* (Mertens et al., 2012a).
Howard et al. (2009) investigated the phylogenetic relationships of yessotoxin-producing
dinoflagellates, including several strains of *P. reticulatum* from different localities. Using
Large Sub Unit (LSU) and Internal Transcribed Spacer (ITS) ribosomal DNA (rDNA)
sequencing, they showed that the *P. reticulatum* strains formed a monophyletic clade in both
phylogenies. ~~However,~~ One particular strain (CCMP404) isolated from the Salton Sea
(California) in 1966 showed significant genetic differences from the other strains in both
phylogenies. Despite these genetic differences, Howard et al. (2009) considered all the strains
to belong to the species *P. reticulatum*.

The Salton Sea is the largest saline lake in California with a surface area of 980 km²
(Reifel et al., 2002). It has a mean depth of 8 m and a maximum depth of 15 m (Ferrari and
Weghorst, 1997). Although originally composed of relatively freshwater, it has become saline
due to a lack of outflow and high evaporation rates. During 1997–1999, the salinity was
between 41 and 45 g l⁻¹ (Watts et al., 2001), while the temperature varied between about 12
and 40°C seasonally (Watts et al., 2001; Holdren and Montaña, 2002). Oxygen at times was
supersaturated due to phytoplankton photosynthesis, but was also often severely depleted,

occasionally even in surface waters (Watts et al., 2001; Holdren and Montaña, 2002). Reifel et al. (2002) reported *P. reticulatum* from the Salton Sea without illustration or description.

In the present study, through reevaluation of the CCMP404 strain originated from the Salton Sea and observations of recently collected plankton samples from the Salton Sea, ~~it is~~we demonstrated~~d~~ that specimens living in the Salton Sea that ~~have had been~~ previously been identified as *P. reticulatum*, have a ~~dissimilar~~different tabulation to that of *P. reticulatum*. To resolve this issue and accommodate these organisms, ~~we have erected a new genus and new species,~~Pentaplacodinium saltonense n. gen. et sp. is erected. From ~~the~~ Salton Sea plankton samples, ~~we describe~~ the morphology of the thecate stage is described, showing significant differences with *P. reticulatum*. Similarly, through incubation of cysts from Salton Sea surface sediments, ~~we describe~~ the corresponding cyst is described. Phylogenetic relationships are explored, including those with several unpublished sequences of *P. reticulatum*, *Ceratocorys armata* (Schütt 1895) Kofoid 1910, *Ceratocorys gourretii* Paulsen 1931, *Ceratocorys horrida* Stein 1883, and several unidentified strains. In addition, both the autecology and fossil record of *Pentaplacodinium saltonense* are examined.

2. Material and Methods

The cyst–theca relationship of *P. saltonense* was established through a germination experiment of a sample from the Salton Sea (CA, USA). To identify differences and similarities between *P. reticulatum* and *P. saltonense*, ~~we compared~~ the morphology of thecate stages of strains present in culture collections and other cells used for sequencing. ~~are were~~ compared (Table 1, Suppl. Table 1). In addition, ~~we studied the~~ phylogenies ~~are were~~ constructed ~~of using~~ LSU, ITS and SSU rDNA based sequences of *P. saltonense* and *P. reticulatum* from several of the same cells or cultured strains, as well as three *Ceratocorys*

species isolated from the Mediterranean, and several unidentified strains from Hawaii and Malaysia (Table 1, Suppl. Table 1).

2.1. Morphological ~~study~~ imaging of cells ~~in~~ from plankton samples and strains present in culture collections ~~with microscopy~~

Plankton samples were obtained from the Salton Sea (California, U.S.A.; 33.50 °N, 115.91 °W) on 24 Oct. 2013 using a plankton net with a 20 µm mesh size. These samples were fixed with ethanol (50% final concentration) and stored cold. Several strains from previously sequenced strains from culture collections established from several other locations were also studied using transmitted light or scanning electron microscopy ([Figure 1](#), Table 1).

For scanning electron microscopy (SEM) of thecate stages by [M.C.C.-M.](#), samples were prepared either by filtering a plankton sample or culture, or isolating a single cell under a Leica™ ~~Inverted-inverted Light-light Microscopemicroscope~~ (Germany). ~~When the~~ Samples were ~~was~~ filtered, ~~by placing an aliquot of ~300 µL aliquot was placed~~ on a Millipore™ 0.25 mm diameter ~~5 µm pore~~ polycarbonate filter at the bottom of a Millipore™ column. Approximately 7 mL of distilled water were added to remove the fixative (ethanol, lugol or formaldehyde) and seawater. A gentle manual vacuum with a 60 cc syringe was used to speed filtration. ~~Individually isolated ce~~ Cells were removed using a glass micropipette under a Leica ~~Inverted-inverted Light m~~ Microscope ~~(Germany) with~~ magnification 10x5x. Individual cells were washed six times with distilled water in double depression microscope slides). After the cells were clean, they were placed on the same kind of filter as for the filtered samples. All filters were air-dried, then ~~adhered~~ affixed to 25 mm diameter aluminium stubs with adhesive tabs (7/16" diameter). The mounted filters were then coated with a mixture of gold-palladium in a Cressington Sputter Coater (U.S.A.) for 60 s.

Observations were performed with a FEI Quanta 3D Dual Beam SEM (Clackamas, Oregon, U.S.A.), at 5 kV. Tilts up to 52° were applied. Digital images were saved in Tiff format (2048 x 1768 pixels). Adobe-Photoshop™ software was used to remove the background while maintaining the integrity of the original image.

For scanning electron microscopy-~~(SEM)~~ of culture CCMP 3243 by K.N.M., the culture was filtered and washed with distilled water and dehydrated in a graded ethanol series (30 to 100% in six steps). The filters were encased in metallic baskets, critical-point dried with CO₂ (CPD Bal-Tec 030), glued onto stubs, sputter coated with platinum/palladium for 90 s (JEOL JFC-2300 HR) and examined in a JEOL 6330F scanning electron microscope (JEOL, Tokyo, Japan) at the University of Copenhagen.

Measurements of ~~thecate-thecae stages~~ of the newly described species were ~~done~~ conducted by ~~M.C.C.-M.~~ under SEM. For each motile cell, the length was measured along the ~~center of the~~ longitudinal axis, the width was measured along the middle of the cingulum, ~~perpendicular to the longitudinal axis~~ from one lateral margin to the other. All motile cell measurements in the species descriptions cite the minimum, average (in parentheses) and maximum values (in µm), in that order. The standard deviation (SD) is also provided where appropriate.

Labelling of tabulation follows a modified Kofoed system that recognizes homologs (e.g., Fensome et al. 1993~~b~~). The sulcal plate labelling ~~is according to~~ with Balech (1980).

2.2. Germination experiment of cysts of *P. saltonense*

Sediment samples were collected from the Salton Sea ~~(CA, U.S.A.)~~ at the same time ~~day~~ of a plankton sample collection ~~on, during the same field campaign~~ on 24 October, 2013, using a Petite Ponar Grab at shallow water depths (<0.5 m). All samples were stored in plastic

bags in a refrigerator at 4°C. *In-situ* sea surface salinities and sea surface temperatures were measured during sampling (Table 1).

About 0.5–1.0 cm³ of wet sediment was immersed in filtered seawater and, after one minute of ultrasonication using an ultrasonic bath, the sediment was rinsed through a 20 µm nylon-mesh sieve using filtered seawater. From this residue, the cyst fraction was separated using the heavy-liquid sodium polytungstate (SPT) at a density of 1.3 g cm⁻¹ (Bolch, 1997). Single cysts were then transferred to Orange Scientific 0.5 mL microwells subjected to an irradiance of 100 µmol photons m⁻² s⁻¹ and 24-hour light, and filled with f/2 medium at room temperature and a salinity of 35 psu. Cysts were regularly checked for germination, and observations were performed under a Leitz DM IL inverted light microscope. Encysted and excysted cysts, as well as motile cells, were photographed and measured using a Leica DM5000B light microscope with 100x oil immersion objectives.

2.3. Morphological study of cysts extracted from surface sediments ~~with~~-using light microscopy and SEM

Surface sediment samples were collected from several ~~sites in the~~ Salton Sea sites were obtained for study ~~of~~ cysts of *Pentaplagodinium saltonense* (Table 1). Palynological techniques were used for processing (e.g., Pospelova et al., 2010; Mertens et al., 2012b). Material was rinsed twice with distilled water to remove salts. The samples were oven-dried at 40°C and then treated with ~~room-temperature~~ 10% hydrochloric acid (HCl) at room temperature to remove calcium carbonate particles. To dissolve ~~siliceous~~-silicate particles, samples were treated with 48–50% ~~room-temperature~~ hydrofluoric acid (HF) at room temperature for two days, and then treated for 10 min with room-temperature HCl (10%) to remove fluorosilicates. The residue was rinsed twice with distilled water, ultrasonicated for

~30 sec and finally collected on a 15 µm mesh. Aliquots of residue were mounted on microscope slides using glycerine jelly.

All measurements and light photomicrographs were obtained by K.N.M. and V.P., respectively using an Olympus BX51 with a Nikon digital sight DS-1L 1 module, and a Nikon Eclipse 80i transmitting light microscope with a DS-L2 module, all with 100x oil immersion objectives.

For each cyst, the lengths of the three longest visible processes with the corresponding widths at their base were measured within the focal plane. Process length was measured from the middle of the process base to the process tip. The average distance between processes was determined by measuring the distance between a process ~~near the center of the cyst~~ on the upper surface of the cyst near the center and the five processes nearest to it, as measured between the middle of the process bases as seen from the surface of the cyst. The central body wall thickness was measured at two to three positions around the cross section of each cyst. The central body maximum and minimum diameters were also measured unless specimens were overly compressed or broken. Fragments representing less than half of a cyst, and cysts with mostly broken processes, were not measured. All cyst measurements in the species descriptions cite the minimum, average (in parentheses) and maximum values (in µm), in that order. The standard deviation (SD) is also provided where appropriate.

For SEM observation of cysts at Geotop (the Université du Québec à Montréal, Canada), single specimens were picked under an inverted microscope with a micropipette, sputter coated with platinum/palladium for 60 s and observed using a scanning electron microscope (Hitachi S-3400N SEM).

2.4. Single-cell polymerase chain reaction (PCR) amplification and sequencing of ~~culture of~~ Salton Sea culture

353

354 Isolated cells were washed three times in serial drops of 0.22 µm filtered and sterilized
355 seawater by micropipette. Each cell was transferred to a 200 µm PCR tube containing 10 µL
356 of Quick Extract FFPE DNA Extraction Solution (Epicentre, Madison, WI, USA) and
357 incubated for 1h at 56°C, then for 2 min at 90°C. The resulting extract was used as a DNA
358 template for the initial PCR amplification. Sequences of SSU and partial LSU rDNA were
359 determined from single cells of *P. saltonense*. The PCR was performed with EconoTaq 2X
360 Master Mix (Lucigen, Middleton, WI, USA) following the manufacture's protocols. The
361 external primers (SR1 and LSU R2) were used for the initial PCR. The first PCR product was
362 used as a DNA template for the second PCR. The following combinations of primer pairs
363 were used separately for the second PCR: SR1 and SR12, 25F1 and LSU R2. Using the
364 second PCR products as the template DNA, the third PCR ~~were~~was performed by the
365 following combinations of primer pairs: SR1b and SR3, SR1b and SR5TAK, SR4 and
366 SR7TAK, SR6 and SR9p, SR8p and SR12, 25F1 and 25R1, D3A and LSU R2. ~~The d~~Details
367 of the primers are described in Takano and Horiguchi (2004) and Yamaguchi et al. (2016).
368 The PCR protocols and sequencing are described in Yamaguchi et al. (2016).

369

370 *2.5. Sequencing of single cells of Protoceratium reticulatum from Elands Bay (South Africa),*
371 *originally described as Gonyaulax grindleyi by Reinecke (1967)*

372

373 Isolated cells were washed three times in serial drops of 0.22 µm filtered and sterilized
374 distilled water and then transferred to a 0.2 mL PCR tube. Cells were subjected to three
375 rounds of heating to 95°C for 5 min~~utes~~ and cooling on ice for 5 min~~utes~~ to induce cellular
376 lysis. 5 µL of the cell lysate was then used as a template for PCR using primers to amplify a
377 168 bp region of the SSU, encompassing the V9 region, V9 For (5'-

GTACACACCGCCCGTC-3') V9 Rev (5'-TGATCCTTCTGCAGGTTACCTAC-3')
(Lane-et al., 1991; Medlin et al., 1988). PCR reactions were carried out in 25 µL volumes
containing 5 µL DNA template, 10 pmol each primer, 1 x buffer, 1 mM MgCl₂, 0.0025 mM
dNTPs, 0.5 Unit Gotaq polymerase (Promega). PCR reactions proceeded with an initial
denaturation at 95°C for 5 min~~utes~~, followed by 35 cycles of denaturation at 95°C for 30
s~~econds~~, annealing at 54°C for 20 s~~econds~~ and extension at 72°C for 20 s~~econds~~ and a final
extension step of 72°C for 5 min~~utes~~. PCR products were sequenced directly in both
directions using the respective primers (Source Bioscience).

2.6. Sequencing of unidentified cultured strains

For strains from Hawaii and Malaysia, single cells were isolated from plankton samples
(Suppl. Table 1) and washed three times with sterilized bi-distillate water and were used as
the template to amplify about 1,430 bp of the LSU rRNA gene (D1-D6 domains), using the
primers D1R (forward, 5'-ACCCGCTGAATTTAAGCATA-3') (Scholin et al., 1994), 28-
1483R (reverse, 5'-GCTACTACCACCAAGATCTGC-3') (Daugbjerg et al., 2000), 1740
bp of the SSU rRNA gene, using the primers SR1(forward, 5'-
TACCTGGTTGATCCTGCCAG-3') and SR12b (reverse, 5'-
CGGAAACCTTGTTACGACTTCTCC-3') (Takano & Horiguchi, 2006), and 600 bp of the
total ITS1–5.8S–ITS2, using the primers ITSA (forward, 5'-CCTCGTAAC
AAGGHTCCGTAGGT-3'), ITSB (reverse, 5'-CAGATGCTTAARTTCAGCRGG)
(Adachi et al., 1996). A 50 µL PCR cocktail containing 0.2 µM forward and reverse primer,
PCR buffer, 50 µM dNTP, 1U of Taq DNA polymerase (Takara, Dalian, China) was
subjected to 35 cycles using a Mastercycler PCR (Eppendorf, Hamburg, Germany). The PCR

reaction procedure was 4 min at 94 °C, followed by 25 cycles of 1 min at 94 °C, 2 min at 45 °C, 3 min at 72 °C, and final extension of 7 min at 72 °C. PCR products were sequenced directly in both directions using the ABI Big-Dye dye-terminator technique (Applied Biosystems, Foster City, CA, USA), according to the manufacturer's recommendations.

DNA extracts from strains collected in Spain (processed at IRTA) were prepared according to the protocol described in Andree et al. (2011). The extracted DNA was used in the amplification of ITS-1, 5.8S, ITS-2 sequences utilizing primers described in Andree et al. (2011), and a partial LSU sequence was amplified utilizing the primers described previously in Hansen et al. (2000). The amplification reactions were carried out in 25µL volume containing: 10 mM Tris-HCl pH 8.3 (at 25 °C), 50 mM KCl, 2 mM MgCl₂, 0.001 % w/v gelatin, 400 pM dNTP's, 1 µM of each primer, and 1 U Taq polymerase. Amplifications were performed using the following parameters: 94 °C for 5 min followed by 35 cycles of 95 °C for 30 s, 50 °C for 45 s, 72 °C for 1 min, and a final extension of 72 °C for 5 min. The PCR products were purified using Qiagen spin columns (Qiagen PCR Purification Kit) and sent for bi-directional sequencing by a commercial company (Sistemas Genomicos, Valencia, Spain) utilizing the same primers as those used in the original amplification. The resulting nucleic acid sequence data was manually proofed using BioEdit (Hall et al., 1999) to confirm the consensus sequence.

The strain 091223-38_M16 from Helgoland (North Sea) was sequenced by M.H.. The Epicentre MasterPure complete DNA & RNA Purification Kit was used for the DNA extraction. ~~We use~~ puReTaq ready-to-go PCR beads ~~are were used~~; annealing temperature was 50°C; 33 cycles; primers: ITS1 (forward) 5' GGTGAACCTGAGGAAGGAT 3'; ITS4 (reverse) 5' TCCTCCGCTTATTGATATGC 3'. The PCR product of the right-correct size was gel isolated (QIAquick Gel Extraction Kit). Sequencing was done by Macrogen with the ITS1 primer.

Strains and single cells from Japan, sequenced by Yoshihito Takano and Kazuhiko Koike, were sequenced using methods mentioned in Mertens et al. (2012a).

DNA was extracted from cultures (strain references K1474, 1476, 1477, 1478, 1479 and 0976) acquired from the NCMA (National Centre for Marine Algae) using the DNeasy DNA extraction kit (Qiagen) according to manufacturers' instructions. The 760 bp region of the LSU rRNA gene was amplified using 2 µL DNA in PCR reactions spanning the D1-D2 variable region D1R (forward, 5'-ACCCGCTGAATTTAAGCATA-3'), D2C (reverse, 5'-GCTTGGTCCGTGTTTCAAGA-3') (Scholin et al., 1994) a 168 bp region of the SSU rRNA gene (V9) V9 For (5'-GTACACACCGCCCGTC-3') V9 Rev (5'-TGATCCTTCTGCAGGTTACCTAC-3') (Lane-et al., 1991; Medlin et al., 1988) and a 710 bp intergenic region ITS1, 5.8S, ITS2, EITS2 For (5'-GTAGGTGAACCTGCVGAAGA-3') EITS2 Rev (5'-TGGGGATCCTGTTTAGTTTC-3') (Guillou et al. 2002). PCR for V9 is detailed in section 2.5. For LSU and ITS, PCR reactions were carried out in 50 µL volumes containing 2 µL DNA, 20 pmol each primer, 1 x buffer, 1.5 mM MgCl₂, 0.0025 mM dNTPs, 1 Unit Gotaq polymerase (Promega). PCR reactions proceeded with an initial denaturation at 95°C for 5 minutes, followed by 35 cycles of denaturation at 95°C for 30 seconds, annealing at 60°C for 45 seconds and extension at 72°C for 1 min and a final extension step of 72°C for 5 minutes. PCR products were sequenced directly in both directions using the respective primers (Source Bioscience) and sequences were manually verified using Chromas (Technelysium Pty Ltd) prior to phylogenetic analysis.

Novel sequences were deposited in Genbank under accession numbers

~~XXXXX~~[MG646283-MG646333](#).

2.7. Sequence alignments and phylogenetic analyses

Multiple sequence alignments were constructed for sequences generated for the variable regions V9 (SSU), D1-D2 (LSU) and partial ITS1, 5.8S, ITS2 (intergenic region), respectively, in BioEdit 7.0 (Hall 1999) using ClustalW along with other available sequences from Genbank. Alignments were trimmed accordingly based on the lengths of the sequences acquired and to allow for a sufficient number of sequences to be included in the phylogeny. Phylogenetic analysis based on neighbour-joining and maximum likelihood was undertaken using MEGA 6 (Tamura et al., 2013) using the default parameters. Bootstrap values were retrieved from 1000 replicates and are indicated on the nodes of the trees.

3. Results

3.1. Study of plankton samples, culture strains, germination experiments, and surface sediments

Investigation of plankton samples from the Salton Sea revealed the presence of a species that is superficially similar to *P. reticulatum* and is here assigned to *Pentaplacodinium saltonense* gen. et sp. nov. n. sp. Three process-bearing cysts (Plate 1) were isolated from surface sediments of the Salton Sea, California, USA (Table 1) and identical morphologies emerged from these cysts (Plate 2). These cells started dividing after germination, and one strain was maintained. The cells were identical in morphology to specimens observed in plankton samples from the Salton Sea (Plate 3), as well as to specimens from several culture strains (Plate 4, Suppl. Table 1), as described below.

3.2. Systematics

477 | Division DINOFLAGELLATA (Bütschli 1885) Fensome et al. 1993**b**
 478 | Class DINOPHYCEAE Pascher 1914
 479 | Subclass PERIDINIPHYCIDAE Fensome et al. 1993**b**
 480 | Order GONYAULACALES Taylor 1980
 481 | Suborder Gonyaulacineae autonym
 482 | Family uncertain
 483 | Genus ***Pentaplacodinium*** Mertens, Carbonell-Moore, Pospelova et Head gen. n. (Plate 3)
 484 | Type: [Plate 3A, the holotype of *Pentaplacodinium saltonense*](#) gen. et sp. nov.
 485 | Diagnosis: A ~~gonyaulacoid~~ [gonyaulacinean](#) genus with roundish to ~~slightly~~ polyhedral thecae
 486 | ~~with bearing~~ heavily reticulated plates without ~~appendices~~ [appendages](#). The tabulation is Po,
 487 | Pt, [4' or 2'+*2'](#), 5'', ~~6C6c~~, ~~6S6s~~, ~~*65'''~~, 1p, 1''', cover plate is oval.
 488 | Etymology: The name is derived from the Greek words *penta* meaning five, *plax* plate, and
 489 | *dino* whirling; with reference to the five precingular plates that characterize this dinoflagellate
 490 | genus.
 491 | ***Pentaplacodinium saltonense*** Mertens, Carbonell-Moore, Pospelova et Head gen. et sp. n.
 492 | (Plates 3, 4, Figs 2A, 3, 4A)
 493 | Synonymy:
 494 | 1970 *Protoceratium reticulatum* (Claparède et Lachmann); Steidinger and Williams, p. 62,
 495 | plate 38, fig. 140a–c.
 496 | 1991 *Protoceratium reticulatum* (Claparède et Lachmann); Al-Muftah, pp. 180–181, figs.
 497 | 246–247.
 498 | ? 2002 *Protoceratium reticulatum* (Claparède et Lachmann); Reifel et al., p. 275.
 499 | 2005 *Gonyaulax grindleyi* Reinecke; Faust et al., p. 110, figs. 2–4.
 500 | ? 2007 *Gonyaulax grindleyi* Reinecke; Tiffany et al., p. 582.

? 2009 "*Protoceratium globosum*" Kofoid ~~et~~ Michener; Morquecho et al., p. 18, 20, figs. 13–17.

Diagnosis: Theca roundish to somewhat polyhedral with tabulation Po, Pt, 2'+*2', 35''+*2'', 66c, 68s, *65'', 1p, 1'', with 3'' interpreted as *(3''+4''). The theca has an L-type ventral organization and dextral torsion. The plates are heavily reticulated with one pore inside each reticulation, although two or more pores might be found in reticulations next to a suture. The ends of the descending cingulum are displaced by ~2.0 widths. The cysts have an approximately ~~roundish-spherical~~ central body with a thin pedium and thicker spongy-fibrous luxuria. ~~Processes-Process distribution apparently have a no intratabular distribution, are~~ Processes fibrous and distally tapering, and have acuminate to minutely expanded distal ends. The archeopyle corresponds to the ~~third~~ *(3''+4'') precingular plate and has a smooth margin with rounded angles. The operculum is free.

Etymology: The specific epithet refers to the type locality for this species.

Type locality: The Salton Sea, California, U.S.A. (station 1 at 33°30.192' N, 115°54.869' W).

Gene sequence: The 28S and 18S gene sequence of the cell isolated from culture 2E3, established from a cyst extracted from surface sediment from station 2 in the Salton Sea (Table 1). —GenBank Accession No. ~~XXXXXX~~ MG646301 (18S) and ~~XXXXXX~~ MG646323 (28S). Several other strains are considered to belong to the same species (Suppl. Table 1).

Holotype: ~~Illustrated on~~ Plate 3A. The specimen illustrated is on an SEM stub (designated CEDiT2017H62) ~~will be deposited-curved~~ at the Senckenberg Research Institute and Natural History Museum, Centre of Excellence for Dinophyte Taxonomy, Germany.

Description: Motile cells observed in the Salton Sea plankton samples (Plate 3, except D).

~~The cell shape varies from~~ Thecae have a roundish to somewhat polyhedral shape (Plate 3A, C). ~~The thecae have and~~ a typical sexiform gonyaulacoid tabulation (sensu Fensome et al.,

1993**b**, ~~t~~Text-Fig. 64B) with an L-type ventral organization (sensu Fensome et al., 1993**b**,
Text-Figs. 82A, C) and dextral torsion (sensu Fensome et al., 1993**b**, Text-Fig. 83C). The
epitheca ~~was~~is often somewhat shorter in length than the hypotheca. The plates are
reticulated with one pore inside each reticulation, although two or more pores ~~could~~may
~~occur~~be found in reticulations next to a suture. All pores each contain ~3 minute pores (small
arrowhead in Plate 3B). The reticulations are faintly expressed on the sulcus and cingulum
(Plate 3C). The cell content is brownish-red owing to the presence of chloroplasts (Plate 2A).
Several red bodies are present (Plate 2A–C).

The apical pore complex consists of a cover plate surrounded by a pore plate (Plate 7F,
H). The oval cover plate, which is often absent (Plate 3B), is relatively broad and is
surrounded by the pore plate. The pore plate is perforated by 5–7 large pores. A low apical
collar may encompass the pore plate and is formed by the raised edges of the first and second
apical plates, and the fourth apical homolog as well (Plate 3F, H). The first and second apical
plates (1' and 2') and the fourth apical homolog (*4') are elongated. The first apical plate (1')
is rectangular, ~~while~~whereas the second apical plate (2') and the fourth apical homolog (*4')
are six-sided and irregularly shaped (Plate 3B). The third apical homolog (*3') is small and
contacts 2' and *4', but in the specimens ~~that were~~ observed it never contacted the apical pore
plates (Plate 3B). There is a large ventral pore located posteriorly between 1' and *4' (Plates
3A, B, 4A–C). The precingular series consists of five large plates, where 2'' is the largest,
*(3''+4'') forms the keystone plate, and *6'' is the smallest. Plates 1'', *(3''+4''), and *5'' are
five-sided, 2'' is four-sided, while *6'' is six-sided (the suture with the anterior right sulcal is
very small (Fig. 3) (Plates 3B, 4A–C). External views of the theca ~~can~~could suggest that
there would be no contact between the anterior sulcal plate and 1' (e.g., Plate 3A, B).
~~However, p~~Properly oriented external views and internal views, however, show ~~however~~a
narrow contact between both plates (Plate 3D). This contact between the anterior sulcal plate

and 1', in combination with the contact between *6'' and 1' therefore results in an insert configuration (sensu Fensome et al. 1993**b**, Text-Fig. 62A). The cingulum is left-handed (descending), lined with narrow lists, and comprises six cingular plates. The ends of the cingulum do not overhang, and are displaced by ~2.0 widths (Plates 3A, 4B).

The sulcus is narrow anteriorly and slightly widens posteriorly. It consists of six plates (Plate 3D, Fig. 3) — the first postcingular plate 1''' is treated as a sulcal and labeled the anterior left sulcal plate (Ssa). The anterior sulcal plate (Sa) is relatively large and anteriorly intruded between plates 1'' and *6'' and barely contacts 1' (Plate 3D). The anterior left sulcal plate (Ssa) is similar in size to the anterior right sulcal plate (Sda). Immediately below these two plates, lay the small posterior right sulcal (Sdp) and a much larger plate, the left posterior sulcal. Finally, there occurs the large posterior sulcal (Sp) ~~is found~~, which presents lines of pores around its sutures with the adjacent non-sulcal plates (Plate 3D, Fig. 3).

The hypotheca is asymmetrical as a consequence of dextral torsion (Plates ~~3~~ 3A). There are five homolog postcingular plates. Plate *2''' is irregularly shaped and the smallest in the series. All other postcingular plates are large, though *6''' is relatively smaller; in addition, they are trapezoidal and four-sided (Plates 3E, 4E). The posterior intercalary plate (1p) bears a conspicuous flange on its right margin (Plates 3A, 4C). The plate overlap is typical for gonyaulacoids, with 3'' (in our case *(3''+4'')) forming the keystone plate (the plate that overlaps all adjacent plates) oin the epitheca, and ~~*4'''~~ forming the keystone plate oin the hypotheca (Fig. 4, Plate 3E).

Cysts from the Salton Sea surface sediments (Plates 1, 5). The central body is approximately spherical~~roundish~~. The wall is thick, consisting of a thin, solid pedium that has a smooth inner surface, and a thicker spongy-fibrous luxuria that appears loosely granular in surface view. Processes are numerous and are solid and fibrous along their entire length, often loosely

fibrous at the base. Process bases are expanded, and larger processes may be concave in lateral profile for at least half of their length. Some closely adjacent processes are joined at the base. Most processes usually have a minute distal expansion, observed under SEM as a concave platform $\sim 1.0\ \mu\text{m}$ or less in diameter with strongly irregular margins that may be approximately perpendicular to the shaft. Alongside these, some processes on most specimens taper to distal points, and such processes occasionally predominate on ~~specimens~~ individual specimens. Processes are mostly of even height, but shorter and thinner processes may be interspersed. The process length/central body ratio is about 0.06. Processes are not evenly spaced, and their parallel alignment and bands devoid of processes observed in many specimens suggest intratabular distribution. ~~However, there is~~ however no clear evidence of tabulation except for the archeopyle and often parallel alignment along the ~~cingulum~~ cingular margins. ~~The archeopyle is formed from the loss of plate $*(3''+4'')$, is reduced, and has slightly rounded angles and straight margins. The usual archeopyle that is moderately wide and reflects the precingular thecal plate $*(3''+4'')$, whereas the operculum is released as a single piece, and has well defined to moderately rounded angles and straight margins, as illustrated on Plate 1G-H. -A reentrant angle along the anterior margin of the archeopyle, signaling the fusion of plates 3'' and 4'', was not seen in the thecal or cyst tabulation of *P. saltonense* although this might not-in fact be expected (see Below, 1987, p. 36, fig. 18a; translated in Fensome et al., 1993a, p. 844). An unusually wide archeopyle that seems to reflect two adjacent precingular thecal plates, $*(3''+4'')$ and 2'', where the operculum is again released as a single piece, is illustrated on Plate 1C-E. If this interpretation is correct, then the component representing 2'' in the archeopyle/operculum is reduced in size, because on the theca the second precingular plate is actually similar or larger in size than the $*(3''+4'')$ plate.~~

601 | *Dimensions:* The ~~holotype~~ cell [illustrated in Plate 3A, the holotype figure](#), is 44 μm in length,
602 | 41 μm in width and 38 μm in depth. Germinated motile cells: length, 48.1 (53.7) 63.4 μm
603 | (SD=6.0, n=5); width, 38.5 (42.4) 47.5 μm (SD= 3.2, n=5). Cells observed in plankton from
604 | St. 2 in the Salton Sea: length, 37.8 (46.1) 59.8 μm (SD = 5.5, n=28); width, 31.0 (39.5) 48.5
605 | μm (SD=4.2, n=28).

606 | Two single cysts germinated to give the identifiable thecae: maximum central body diameter,
607 | 52.3 (53.5) 54.7 μm (SD=1.7, n=2); minimum central body diameter, 51.1 (52.2) 53.3 μm
608 | (SD=1.6, n=2); average length of three randomly chosen processes per cyst, 2.4 (3.0) 3.6 μm
609 | (SD=0.4, n=6); process width at base 1.4 (2.2) 2.7 (SD=0.6, n=6) and wall thickness 1.3 (1.7)
610 | 2.1 (SD=0.3, n=6). Palynologically treated cysts from surface sediments of the Salton Sea:
611 | maximum central body diameter, 48.6 (56.3) 70.9 μm (SD=5.3, n=23); minimum central body
612 | diameter, 45.7 (52.1) 61.4 μm (SD=3.8, n=22); average length of three processes per cyst, 1.0
613 | (3.1) 5.7 μm (SD=1.2, n=66); process width at base 1.0 (2.2) 3.9 (SD=0.6, n=66) and wall
614 | thickness 0.9 (1.6) 2.4 (SD=0.4, n=66).

615 | *Comments:* *Pentaplacodinium saltonense* n. gen et sp. is defined [primarily](#) from the characters
616 | of the motile stage, these distinguishing it from species of the genus *Protoceratium*. The
617 | morphology of several thecae observed from off Yucatan (Gulf of Mexico), the Indian River
618 | Lagoon (Florida, USA), and off Qatar (Persian Gulf) (Table 1; Plate 2) and from cultures
619 | established from cells from Biscayne Bay (Florida, USA) (CCMP1720, CCMP1721), the
620 | Indian River Lagoon (Florida, USA) (CCMP3241, CCMP3243) and the Salton Sea
621 | (California, USA) (CCMP404) ([Suppl. Table 13](#), Plate 1) agree with the description of *P.*
622 | *saltonense* given above. Cysts formed from cultures established from a strain from the Indian
623 | River Lagoon (Florida, USA) (CCMP3243) have the same morphologies (Plate 6). The
624 | observed cysts correspond [most closely](#) to the fossil based ~~taxon-species~~ [Operculodinium](#)
625 | [israelianum](#) (Rossignol 1962) Wall 1967 described from the Pleistocene of Israel, and

Operculodinium psilatum Wall 1967 described from the postglacial (Holocene) of the Caribbean, p. 111–112, Plate 6, figs. 6–8. However, *Operculodinium israelianum* has longer processes (6–10 µm; Rossignol, 1964), and *O. psilatum* has a psilate surface interrupted by minute and sparsely distributed processes, and a pronounced cingulum (Wall, 1967). Both have archeopyles that are less wide than for the cyst of *P. saltonense*.

3.3. Phylogenetic position of *P. saltonense* and other studied strains

The SSU rDNA sequences for all *P. reticulatum* strains analysed were identical, forming a distinct clade separated from the *P. saltonense* sequences which were identical to the unidentified Malaysian sequences (Fig. 5). *P. reticulatum* and *P. saltonense* sequences shared 92% nucleotide identity for the V9 region analysed.

For the LSU rDNA V4 analysis ([Figure 6](#)), *P. reticulatum* sequences were identical apart from a couple of sporadic nucleotide substitutions which were ~~called identified~~ as ambiguous bases by the sequencing software. The unidentified strain from Hawaii had 12 nucleotide substitutions across the 570 bp multiple sequence alignment compared to *P. reticulatum*. *P. saltonense* sequences shared more similarity with the unidentified GgSm strains from Malaysia (96%) compared to that of *P. reticulatum* (94%).

The ITS (intergenic region between ITS1 and 2) was the only marker to resolve intraspecific diversity within the *P. reticulatum* species, with strain E12 (Baffin Bay, Arctic) sharing 98% nucleotide similarity with strain VG0757 isolated from Spain. The phylogeny separates *P. reticulatum* into two large subclades: subclade 1A that regroups several strains from warmer waters, and subclade 1B that regroups several strains from colder waters (Fig 7).

The three phylogenies ([Figures 5–7](#)) show that strains identified as *Protoceratium reticulatum* form a monophyletic group (Clade 1), as well as strains identified as *P.*

saltonense that form a clade with the unidentified GgSm strains from Malaysia (Clade 2), as well as the *Ceratocorys* species that form a clade with PrTT strains from Malaysia (Clade 3) supported by high bootstrap values (>70). The unidentified strain from Hawaii does not group with the *Protoceratium reticulatum* or *Pentaplacodinium saltonense* clades. The topology of the trees ~~are-is~~ not consistent between the three phylogenies (i.e. the relatedness between clades), ~~however-but~~ the three clades identified are consistently formed. The trees furthermore highlight the unexplored diversity within this group of dinoflagellates, and further incubation and plankton studies from these locations should reveal whether the unidentified strains are new species or not.

In addition, the phylogenies show that V9 SSU sequences from cells from Elands Bay (South Africa); (bootstrap values >80%); that have been previously identified as *G. grindleyi* by Reinecke (1967), clusters with *Protoceratium reticulatum* (Fig-~~ure~~ 5).

The three studied species of *Ceratocorys* (*C. armata*, *C. gourreti*, *C. horrida*) share high nucleotide similarity for the SSU (100%) and LSU sequences (>99% identity) (~~Figure~~ 6).

4. Discussion

4.1. Comparison of ~~the cell~~-*theca* of *P. saltonense*

Pentaplacodinium saltonense differs from *Protoceratium reticulatum* because it bears five precingular plates, whereas *P. reticulatum* has six ~~precingular~~ plates. Furthermore, *P. saltonense* has a larger ~~cingulum-cingular~~ displacement (2 widths vs. 1 width respectively) and an oval cover plate, as opposed to a sigmoidal cover plate in *P. reticulatum*. In addition, the theca of *P. saltonense* is mostly roundish, whereas in *P. reticulatum* it is always polyhedral. Both species have an insert configuration, but in *P. saltonense* the contact

between Sa and 1' is very narrow ~~whilst~~ whereas in *P. reticulatum* this contact is wide — this causes an ~~evident~~ conspicuous separation between 1'' and 6'' in *P. reticulatum*, when in *P. saltonense* there is an ~~apparent~~ almost a small point of contact between those two plates (Plate ~~4D3D~~). *Gonyaulax grindleyi* Reinecke 1967 is ~~here considered~~ shown to be a synonym of *P. reticulatum*, as ~~it has been~~ already suggested by von Stosch (1969) and Hansen et al. (1997), and ~~is~~ now confirmed by the LSU rDNA phylogeny in this study (see below).

Several other *Protoceratium* species have been described (e.g. Schiller, 1937, p. 322–326). *Protoceratium splendens* Meunier 1910 from the Kara Sea has six precingular plates; it is possibly a junior synonym of *Protoceratium reticulatum*, as suggested by Gómez (2012). *Protoceratium aculeatum* (Stein 1883) Schiller 1937 bears antapical spines and an apical horn. *Protoceratium areolatum* Kofoid 1907 and *Protoceratium spinulosum* (Murray and Whitting 1899) Schiller 1937 have fewer reticulations in both the epitheca and hypotheca than *P. saltonense*. Of the five species described by Kofoid and Michener (1911~~–~~), *Protoceratium cancellorum*, *Protoceratium pellucidissimum*, *Protoceratium pepo*, *Protoceratium globosum* and *Protoceratium promissum*, none ~~have~~ has illustrations and it is therefore ~~it is~~ impossible to compare them to *P. saltonense*.

Pentaplacodinium saltonense differs from *Ceratocorys anacantha* Carbonell-Moore 1996 because it is not as polyhedral. In addition, in contrast to the insert epithecal configuration of *P. saltonense*, *C. anacantha* has an episert type I epithecal configuration, meaning that 1' does not contact the anterior sulcal plate and that 1'' and 6'' are in contact (Paez-Reyes and Head, 2013).

4.2. Comparison of the cyst of *P. saltonense*

The cyst of *Pentaplacodinium saltonense* ~~corresponds to~~compares with *Operculodinium psilatum* because its cysts display an alignment of processes along the cingulum, it bears short processes (1.0–5.7 μm ~~vs. 2 μm (Wall, 1967) or 0.0–2.9 μm (Head, 1996)~~), its body diameter is of similar size (45.7–70.9 μm vs. 50–60 μm ~~(, Wall, 1967);~~ or 62–79 μm ~~(, Head, 1996b)~~) and its wall thickness (0.9–2.4 vs. 1.4–2.2 μm ~~(; Head, 1996b)~~) is similar. *Operculodinium psilatum* differs, however, in having processes that in general are shorter (2 μm , Wall, 1967; 0.0–2.9 μm , Head, 1996b) and sparsely distributed. The cingulum and sulcus are also more conspicuously expressed in *Operculodinium psilatum* (Wall, 1967; Head et al., 1996b), and *O. psilatum* lacks the wide archeopyle of *P. saltonense*.

The cyst of *P. saltonense* is also similar to *Operculodinium israelianum* (Rossignol 1962) Wall 1967 as described by Rossignol (1964, as *Baltisphaeridium israelianum*); ~~however,~~ although the processes of the latter species are longer (6–10 μm) than ~~of the specimens from Salton Sea for *P. saltonense* (1.0–5.7 μm).~~ ~~At this time, we do not~~ It is not presently known whether variation in process length is related to variations in ecology, ~~similar~~ as demonstrated for the cysts of *Lingulodinium polyedra* (= *Lingulodinium machaerophorum*) (Mertens et al., 2009), cysts of *Protoceratium reticulatum* (Mertens et al., 2011) and cysts of *Pyrodinium bahamense* (= *Polysphaeridium zoharyi*) (Mertens et al., 2015). ~~In addition,~~ The process distribution appears to be ~~is~~ intratabular for the cysts of *P. saltonense* and often shows parallel alignments, which is not readily visible and this is likely to be the case also for ~~on~~ *O. israelianum* (e.g., *O. cf. israelianum* of Head, 1997, fig. 17.2), although the study of topotype material will be needed for confirmation. The rounded angles of the archeopyle in *O. israelianum*, *O. psilatum* and the cysts of *Pentaplacodinium saltonense*, and the shared presence of a spongy-fibrous to fibroreticulate luxuria, accentuate the overall similarities between these cysts, although the relatively wider archeopyle in *P. saltonense* cysts distinguishes them from these other species.

~~We t~~Therefore, ~~it is not considered~~ it not to be a morphological variant of *Operculodinium israelianum*. There is reasonable doubt that *Operculodinium israelianum* is related to another species: Wall and Dale (1968) (and subsequently Dale, 1983) have suggested that *O. israelianum* can be related to *Protoceratium reticulatum*. *Operculodinium israelianum* ~~is very similar to~~resembles the Miocene *Operculodinium centrocarpum* (Deflandre et Cookson 1955) Wall 1967, which ~~is also~~ has a spongy-fibrous luxuria, although it is somewhat larger (54–80 μm) (~~;~~ Deflandre and Cookson, 1955) and has longer processes. Head (1996b) noted an intergradation in size and process length between *O. israelianum* and *O. centrocarpum* in Pleistocene assemblages of eastern England. ~~It is of interest to remark that specimens from the Ludham borehole intergrade between *O. israelianum* and *O. centrocarpum*~~ Deflandre et Cookson 1955 (Head, 1996). A ~~re~~study of ~~both holotype~~topotype material ~~as well as the~~is needed to confirm the range of variability within each species ~~would be needed to resolve this issue~~.

The cysts of *P. saltonense* differs from ~~the cyst~~those of *P. reticulatum*, ~~due to the in~~ having a thick spongy-fibrous luxuria ~~wall~~ (vs. thin, fibrous luxuria), less developed distal ends of the processes, larger central ~~its~~-body size-diameter (48.6–70.9 μm vs. 33–48 μm ; Rochon et al., 1999), and generally shorter process length (1.0–5.7 μm vs. typically 7–14 μm ; Rochon et al., 1999) (48.6 (56.3) 70.9 μm (SD=5.3, n=23)) for *P. saltonense*, as opposed to 33–48 μm for *P. reticulatum* (Rochon et al., 1999) and its process length (1.0 (3.1) 5.7 μm (SD=1.2, n=66)) for *P. saltonense*, as opposed to typical 7–14 μm for *P. reticulatum* (Rochon et al., 1999); although the cysts of *P. reticulatum* ~~shows very wide variation~~vary widely sizes, with ~~formation of~~some being completely bald ~~eysts~~ (e.g., Mertens et al., 2012a; Jansson et al., 2014).

~~Several~~ Numerous other *Operculodinium* species have been described and a detailed
comparison is given by~~we refer to~~ Marret and Kim (2009)~~for their detailed comparison, as;~~
none of these closely resembles the cysts of *P. saltonense*.

4.3. Phylogenetics, evolution and position and relationships of *Protoceratium*, *Pentaplacodinium*, and *Ceratocorys*

~~There are s~~Several morphological characteristics of the theca ~~that~~ are important ~~to~~ in
understanding the evolution of *Protoceratium*, *Pentaplacodinium*, and *Ceratocorys* ~~the three~~
~~genera in question~~ (Plate 7). The shape of the cover plate of *Ceratocorys* is more similar to
that of the cover plate of *Pentaplacodinium*, but less similar to the sigmoidal cover plate of
Protoceratium.

Pentaplacodinium and *Ceratocorys* can be considered closer to *Gonyaulax* than
Protoceratium, because the anterior intercalary is always well-separated from the apical pore
plates, ~~whilst~~ whereas in *Protoceratium reticulatum* it is closer and has even been suggested
to contact the apical pore plates (Hansen et al. 1997). ~~However, It should~~ however be noted,
however, that ~~*Protoceratium reticulatum*~~ and *Gonyaulax* have six precingular plates, whereas
~~whilst~~ *Ceratocorys* and *Pentaplacodinium* have five ~~precingulars~~. So it is not surprising that
in the molecular phylogenies, *Pentaplacodinium* has an intermediate position between
Ceratocorys and *Protoceratium* (Figs. 5–7); the relation to other gonyaulacoids at this time is
unclear and further molecular studies of related genera are required, particularly to understand
how to resolve the position of *Protoceratium* at family level. Another issue regards a conflict
in the dual nomenclature: the cyst of *P. reticulatum* and *P. saltonense* both are considered to
belong to the cyst-defined genus *Operculodinium*, whereas ~~whilst~~ the thecate stages belong to

two different genera; further cyst–theca experiments within this group of related species should help to understand how the genus-generic concepts can be made-conformationalized.

In addition, the ITS marker was able to separate two large subclades within *P. reticulatum*: strains that are predominantly associated with warmer waters (Sub-clade 1A), and other strains largely associated with colder waters (Sub-clade 1B) (Fig. 7). Do these subclades reflect pseudocryptic speciation in *Protoceratium reticulatum* as previously suggested by Mertens et al. (2012a)?

Other morphological characteristics of the theca are conserved in *Protoceratium*, *Pentaplacodinium*, and *Ceratocorys* and other gonyaulacoids. For instance, there is no difference in overlap pattern between *Protoceratium*, *Pentaplacodinium*, *Ceratocorys*, *Gonyaulax* and *Lingulodinium* (Fig. 4).

4.4. Biogeography and ecology of *P. saltonense*

According to ~~the~~ plankton observations, *P. saltonense* can be found in tropical to subtropical regions. ~~We have not observed~~ *P. saltonense* and *P. reticulatum* have not been observed in the same samples, which suggests that both-these species possibly inhabit different niches, where *P. saltonense* has a preference for higher temperatures and salinities, and *P. reticulatum* ~~has a preference~~ for somewhat lower temperatures and salinities. This difference would need to be established-quantified through culture experiments.

4.5. Toxicity

Strains identified as *Pentaplacodinium saltonense* (CCMP404, CCMP1720 and CCMP1721), have been identified as yessotoxin producers using fluorescence HPLC (Paz et al., 2004). A

later toxin analysis by LC-MS of the same strains was negative (Paz et al., 2007), and the authors considered that these strains had lost their toxicity after a number of years in culture. The ~~produced~~ toxins produced by these strains of *Pentaplacodinium saltonense* are similar to toxins produced by strains ~~we~~ identified as *Protoceratium reticulatum*, all of which are yessotoxin producers, such as strains from Chile (Alvarez et al., 2011), Jervis Inlet, British Columbia, Canada (Cassis, 2005), German Bight, ~~the~~ North Sea (Röder et al., 2011, 2012), Okkirai Bay, Japan (Koike et al., 2006) and Spain (Paz et al., 2007, 2013). ~~The presence of *P. saltonense* in the Salton Sea has been considered a potential causative agent of mortality events in the Salton Sea indicates that it is potentially toxic~~ (Reifel et al., 2002, whom identified it as *Protoceratium reticulatum*). ~~However, t~~However, there have not however not been ~~no~~ reports of toxic events knowingly involving *P. saltonense*.

Several other studies have investigated the toxicity of strains ~~that~~ they designate as *Protoceratium reticulatum*, but for which ~~we could not verify~~ the identifications could not be verified (e.g., Satake et al., 1999; Ciminiello et al., 2003; Samdal et al., 2004; Finch et al., 2005; Eiki et al., 2005; Mitrovic et al., 2005; Guerrini et al., 2007; Suzuki et al., 2007).

5. Conclusions

Pentaplacodinium saltonense gen. et sp. nov. is described from the Salton Sea (CA, USA). The distinct cover plate (similar to *Ceratocorys*, but sigmoidal in *Protoceratium*), five precingular plates (also five as in *Ceratocorys*, but six in *Protoceratium*), the very narrow contact between 1' and Sa (wide contact in *Protoceratium*, no contact in *Ceratocorys*), a more roundish-rounded cell-the cal shape, the displacement of the cingulum by two widths (~~vs. one~~ width in *Protoceratium*), as well as the clear separation and distances seen in the three phylogenies, justifies the creation of a new genus. The chorate cysts of *P. saltonense* bear

~~short processes often with parallel alignments. These cysts correspond to the cyst-defined species-genus *Operculodinium* Wall 1967, and are most similar to *O. israelianum* and *O. psilatum* with short processes and often parallel alignments. The geographic distribution of~~
Motile stages of *Pentaplagodinium saltonense* ~~wasare confirmed in~~from four widely dispersed locations ~~and its distribution is thereforeis considered,~~ suggesting a subtropical to tropical ~~distribution for this species., and does not overlap with that of~~ *Protoceratium reticulatum* and *Pentaplagodinium saltonense* are not known to inhabit the same environments. ~~Similar to~~As with the ~~yessotoxin-producing~~ *Protoceratium reticulatum*, *Pentaplagodinium saltonense* is potentially a yessotoxin producer, as shown by previous studies.

~~Note added: While this paper has been going through the process of final acceptance to Harmful Algae, another study was accepted (Salgado et al., accepted) that addresses similar scientific questions.~~

Acknowledgements

V.P. and M.J.H. ~~each~~ acknowledge support from a Discovery Grant of the Natural Sciences and Engineering Research Council of Canada (NSERC). K.B.A. and M.F. ~~acknowledge were supported from by the~~ Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria of the Spanish Government (project RTA2005-00109-00-00). ~~We thank~~ CINVESTAV, Merida ~~is thanked~~ for supplying information about the Yucatan samples. Beatriz Paz is acknowledged for interesting discussions on toxicity. ~~We would like to thank~~ Anke Kremp, Maija Hutunnen, Jacob Larsen, Paul Hargraves and Ximena Vivanco ~~are thanked for kindly providing-provided~~ plankton samples. Nancy Lewis and Margaret Beaton ~~are thanked~~

~~for generously sharing~~ shared information on strains from Nova Scotia, Canada. ~~We are~~
~~grateful~~ Thanks to ~~to~~ Captain Brown, crew of the MSV Strickland, Ms. Sarah Thornton, and
EOS313-2010 (University of Victoria) students are all thanked for their participation in
sediment sample collection. Paul Hargraves ~~is acknowledged for~~ kindly providing ~~provided~~
SEM images of *P. saltonense* from cultures established from the Indian River Lagoon.

References

Adachi, M., Sako, Y., Ishida, Y., 1996. Analysis of *Alexandrium* (Dinophyceae) species using
sequences of the 5.8S ribosomal DNA and internal transcribed spacer regions. J. Phycol. 32,
424–432.

Al-Muftah, A.R., 1991. Dinoflagellates of Qatari ~~Waters~~ waters (PhD thesis), vol. 2.
University College of North Wales, Bangor University, pp. 261.

Alvarez, G., Uribe, E., Díaz, R., Braun, M., Mariño, C., Blanco, J., 2011. Bloom of the
Yessotoxin producing dinoflagellate *Protoceratium reticulatum* (Dinophyceae) in ~~Northern~~
northern Chile. J. Sea Res. 64, 427–434.

Andree, K.B., Quijano-Scheggia, S., ~~et al.~~ Fernández, M., Elandalousi, L.M., Garcés, E., Camp,
J., Diogene, J., ~~et al.~~ (2011.) Quantitative PCR ~~Coupled~~ coupled with ~~Melt~~ melt ~~Curve~~ curve
~~Analysis~~ analysis for ~~Detection~~ detection of ~~Selected~~ selected *Pseudonitzschia* spp.
(Bacillariophyceae) from the ~~Northwestern~~ northwestern Mediterranean Sea. App. and Env.
Micro, 77(5) 1651–1659.

872 | Balech, E., 1980. On thecal morphology of dinoflagellates with special ~~emphases~~emphasis on
873 | circular and sulcal plates. Anales del Centro de Ciencias del Mar y Limnología, UNAM 7(1),
874 | 57–68.

875

876 | Balech, E., 1988. Los dinoflagellados del Atlantico sudoccidental. Publicaciones Especiales
877 | Instituto Español de Oceanografía 1, 1–310.

878 |

879 | Below, R., 1987. Evolution und Systematik von Dinoflagellaten-Zysten aus der Ordnung
880 | Peridinales. I. Allgemeine Grundlagen und Subfamilie Rhaetogonyaulacoideae (Familie
881 | Peridiniaceae). Palaeontographica, Abteilung B 205, 1–164.

882

883 | Bergh, R.S., 1881. Der Organismus der Cilioflagellaten. Eine phylogenetische Studie.
884 | Morphologisches Jahrbuch 7(2), 177–288.

885

886 | Bogus, K. Mertens, K.N., Lauwaert, J., Harding, I.C., Vrielinck, H., Zonneveld, K.A.F.,
887 | Versteegh, G.J.M., 2014. Differences in the chemical composition of organic-walled
888 | dinoflagellate resting cysts from phototrophic and heterotrophic dinoflagellates. J. Phycol. 50,
889 | 254–266.

890

891 | Bolch, C.J.S., 1997. The use of polytungstate for the separation and concentration of living
892 | dinoflagellate cysts from marine sediments. Phycologia 37, 472–478.

893

894 | Braarud, T., 1945. Morphological observations on marine dinoflagellate cultures (*Porella*
895 | *perforata*, *Goniaulax tamarensis*, *Protoceratium reticulatum*). Avh. Utgit. Nor. Vidensk.
896 | Akad. Oslo Mat.–Naturvidensk. Kl. 11, 1–18.

897

898 Bütschli, O. 1885. Erster Band. Protozoa. 3. Unterabtheilung (Ordnung) Dinoflagellata. In:
899 Dr. H.G. Bronn's Klassen und Ordnungen des Thier-Reichs, wissenschaftlich dargestellt in
900 Wort und Bild. C.F. Winter'sche Verlagshandlung, Leipzig und Heidelberg. p. 906–1029.

901

902 Cassis, D., 2005. The effect of harmful algae on the summer mortality of juvenile Pacific

903 ~~Oysters~~ oysters (*Crassostrea gigas*). MSc. Thesis, University of British Columbia, 73 pp.

904

905 Ciminiello, P., Dell' Aversano, C., Fattorusso, E., Forino, M., Magno, S., Guerrini, F.,

906 Pistocchi, R., Boni, L. 2003. Complex yessotoxins profile in *Protoceratium reticulatum* from

907 north-western Adriatic sea revealed by LC-MS analysis. Toxicon 42, 7–14.

908

909 Claparède, E., Lachmann, J., 1858. Etudes sur les infusoires et les rhizopodes. Mem. Inst. Nat.
910 Genev. 6, 261–482.

911

912 Dale, B., 1983. Dinoflagellate resting cysts: "benthic plankton". In: Fryxell, G.A. (ed.),

913 Survival Strategies of the Algae, Cambridge University Press, Cambridge, U.K., pp.69–136.

914

915 Dale, B., 1996. Dinoflagellate cyst ecology: modelling and geological applications. In:

916 Jansonius, J., McGregor, D.C. (eds.), Palynology: Principles and Applications. Volume 3.

917 American Association of Stratigraphic Palynologists Foundation, Dallas, Texas, pp. 1249–

918 1275.

919

920 Daugbjerg, N., Hansen, G., Larsen, J., Moestrup, Ø., 2000. Phylogeny of some of the major

921 genera of dinoflagellates based on ultrastructure and partial LSU rDNA sequence data,

including the erection of three new genera of unarmoured dinoflagellates. *Phycologia* 39,
302–317.

Deflandre, G., Cookson, I.C., 1955. Fossil microplankton from Australian Late Mesozoic and
Tertiary sediments. *Australian Journal of Marine and Freshwater Research* 6, 242–313.

Dodge, J.D., 1989. Some revisions of the family Gonyaulaceae (Dinophyceae) based on a
scanning electron microscope study. *Bot. Mar.* 32, 275–298.

Eiki, K., Satake, M., Koike, K., Ogata, T., Mitsuya, T., Oshima, Y., 2005. Confirmation of
yessotoxin production by the dinoflagellate *Protoceratium reticulatum* in Mutsu Bay. *Fish.*
Sci. 71, 633–638.

Faust, M.A., Litaker, R.W., Vandersea, M.W., Kibler, S.R., Tester, P.A. 2005. Dinoflagellate
diversity and abundance in two Belizean coral-reef mangrove lagoons: a test of Margalef's
mandala. *Atoll Research Bulletin* 534, 105-132.

Fensome, R.A., Gocht, H., Stover, L.E., Williams, G.L., 1993a. The Eisenack Catalog of
Fossil Dinoflagellates. New Series. Volume 2. p.829–1461; E. Schweizerbart'sche
Verlagsbuchhandlung, Stuttgart, Germany.

Fensome, R., Taylor, F., Norris, G., Sarjeant, W., Wharton, D., Williams, G., 1993b. A
classification of fossil and living dinoflagellates. *Micropaleontology Special Publication* 7, 1–
245.

947 Ferrari, R.L., Weghorst, P., 1997. Salton Sea 1995 hydrographic GPS survey. U.S. Bureau of
 948 Reclamation, Water Resources Services. Denver, Colorado, 23 pp.
 949
 950 Finch, S.C., Wilkins, A.L., Hawkes, A.D., Jensen, D.J., MacKenzie, A.L., Beuzenberg, V.,
 951 Quilliam, M.A., Olseng, C.D., Samdal, S.A., Aasen, J., Selwood, A.I., Cooney, J.M., Sandvik,
 952 M., Miles, C.O., 2005. Isolation and identification of (44-R,S)-44,55-dihydroxyessotoxin
 953 from *Protoceratium reticulatum*, and its occurrence in extracts of shellfish from New
 954 Zealand, Norway and Canada. *Toxicon* 46, 160–170.
 955
 956 Gómez, F., 2012. A checklist and classification of living dinoflagellates (Dinoflagellata,
 957 Alveolata). *CICIMAR Océanides* 27(1), 65–140.
 958
 959 Guerrini, F., Ciminiello, P., Dell’Aversano, C., Tartaglione, L., Fattorusso, E., Boni, L.,
 960 Pistocchi, R., 2007. Influence of temperature, salinity and nutrient limitation on yessotoxin
 961 production and release by the dinoflagellate *Protoceratium reticulatum* in batch-cultures.
 962 *Harmful Algae* 6, 707–717.
 963
 964 | Guillou, L., Nezan, E., Cueff-Gauchard, V., Barbier, G., 2002. Genetic diversity and
 965 | molecular detection of three toxic dinoflagellate genera (*Alexandrium*, *Dinophysis* and
 966 | *Karenia*) from French coasts. *Protist*. 153, 223-238
 967
 968 | Hall, T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis
 969 | program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.* 41, 95–98.
 970

Hansen, G., Moestrup, Ø. and Roberts, K.R., 1997. Light and [Electron-electron Microscopical](#)
[microscopical Observations-observations](#) on *Protoceratium reticulatum* (Dinophyceae). Arch.
Protistenk. 147, 381–391.

Hansen, G., Daugbjerg, N., Henriksen, P., 2000. Comparative study of *Gymnodinium*
mikimotoi and *Gymnodinium aureolum*, comb. nov. (= *Gyrodinium aureolum*) based on
morphology, pigment composition, and molecular data. J. Phycol., 36, 394– 410.

[Head, M.J., Wrenn, J.H. \(eds.\), 1992. A forum on Neogene–Quaternary dinoflagellate cysts:
The edited transcript of a round table discussion held at the Second Workshop on Neogene
dinoflagellates. In: Head, M.J., Wrenn, J.H. \(eds.\), *Neogene and Quaternary Dinoflagellate
Cysts and Acritarchs*. American Association of Stratigraphic Palynologists Foundation,
Dallas, Texas, pp. 1–31.](#)

Head, M.J., 1996a. Modern dinoflagellate cysts and their biological affinities. In: Jansonius,
J., McGregor, D.C. (eds.), *Palynology: principles and applications*. American Association of
Stratigraphic Palynologists Foundation, Dallas, Texas, vol. 3, pp. 1197–1248.

[Head, M.J., 1996b. Late Cenozoic dinoflagellates from the Royal Society borehole at
Ludham, Norfolk, eastern England. J. Paleontol. 70, 543–570.](#)

Head, M.J., [1997. Thermophilic dinoflagellate assemblages from the mid Pliocene of eastern
England. J. Paleontol. 71, 165–193.](#)

995 [Hennissen, J.A.I., Head, M.J., De Schepper, S., Groeneveld, J., 2017. Dinoflagellate cyst](#)
 996 [paleoecology during the Pliocene–Pleistocene climatic transition in the North Atlantic.](#)
 997 [Palaeogeogr., Palaeoclimatol., Palaeoecol. 470, 81–108.](#)
 998
 999 Holdren, G.C., Montaña, A., 2002. Chemical and physical characteristics of the Salton Sea,
 1000 California. *Hydrobiol.* 473, 1–21.
 1001
 1002 Howard, M.D.A., Smith, G.J., Kudela, R.M., 2009. Phylogenetic relationship of yessotoxin-
 1003 producing dinoflagellates based on the Large Subunit and Internal Transcribed Spacer
 1004 Ribosomal DNA ~~Domains~~[domains](#). *Appl. Environ. Microbiol.* 75, 54–63.
 1005
 1006 Jansson, I.-M., Mertens, K.N., Head, M.J. with contributions from de Vernal, A., Londeix, L.,
 1007 Marret, F., Matthiessen, J., Sangiorgi, F., 2014. Statistically assessing the correlation between
 1008 salinity and morphology in cysts produced by the dinoflagellate *Protoceratium reticulatum*
 1009 from surface sediments of the North Atlantic Ocean, Mediterranean–Marmara–Black Sea
 1010 region, and Baltic–Kattegat–Skagerrak estuarine system. *Palaeogeogr. Palaeoclimatol.*
 1011 *Palaeoecol.* 399, 202–213.
 1012
 1013 Kofoed, C.A., 1907. Reports on the scientific results of the expedition to the eastern tropical
 1014 Pacific, in charge of Alexander Agassiz, by the U.S. Fish Commission steamer “~~Albatross~~”,
 1015 ~~”~~, from October, 1904, to March, 1905, Lieut.~~.-~~[_](#)Commander L.M. Garrett, U.S.N.,
 1016 commanding. IX. New species of dinoflagellates. *Bull. Mus. Comp. Zool.* 50(6), 163–207.
 1017
 1018 Kofoed, C.A., Michener, J.R., 1911. Reports on the Scientific Results of the Expedition to the
 1019 Eastern Tropical Pacific, in Charge of Alexander Agassiz, by the U.S. Fish Commission

1020 | Steamer “~~Albatross~~~~ALBATROSS~~”, from October 1904, to March, 1906, Lieut. L.M.
1021 | Garrett, U.S.N., Commanding. XXII. New genera and species of
1022 | ~~Dinoflagellates~~~~dinoflagellates~~. Bull. Mus. Comp. Zool. 54(7), 267–302.
1023 |
1024 | Koike, K., Horie, Y., Suzuki, T., Kobiyama, A., Kurihara, K., Takagi, K., Kaga, S.N.,
1025 | Oshima, Y., 2006. *Protoceratium reticulatum* in northern Japan: environmental factors
1026 | associated with seasonal occurrence and related contamination of yessotoxin in scallops. J.
1027 | Plankton Res. 28, 103–112.
1028 |
1029 | Lane, D. J., 1991. 16S/23S sequencing. In: Nucleic Acid Technologies in Bacterial
1030 | Systematic. Stackebrandt E, Goodfellow M (Eds) pp. 115–175, Wiley, NY.
1031 |
1032 | Marret, F., Kim, S.-Y., 2009. *Operculodinium aguinawense* sp. nov., a dinoflagellate cyst
1033 | from the late Pleistocene and recent sediments of the east Equatorial Atlantic ocean.
1034 | Palynology 33, 125–139.
1035 |
1036 | Matsuoka, K., McMinn, A., Wrenn, J.H., 1997. Restudy of the Holotype of *Operculodinium*
1037 | *centrocarpum* (Deflandre & Cookson) Wall (Dinophyceae) from the Miocene of Australia,
1038 | and the ~~Taxonomy~~~~taxonomy~~ of ~~Related-related~~~~Species~~~~species~~. Palynology 21, 19–33.
1039 |
1040 | Medlin, L. K, Elwood, H. J., Stickel, S., Sogin, M. L., 1988. The characterization of
1041 | enzymatically amplified eukaryotic 16S-like rRNA-coding regions. Gene, 71, 491–499.
1042 |
1043 | Mertens, K.N., Ribeiro, S., Bouimetarhan, I., Caner, H., Combourieu Nebout, N., Dale, B., de
1044 | Vernal, A., Ellegaard, M., Filipova, M., Godhe, A., Goubert, E., Grøsfjeld, K., Holzwarth, U.,

1045 Kotthoff, U., Leroy, S.A.G., Londeix, L., Marret, F., Matsuoka, K., Mudie, P.J., Naudts, L.,
 1046 Peña-Manjarrez, J.L., Persson, A., Popescu, S.-M., Pospelova, V., Sangiorgi, F., van der
 1047 Meer, M., Vink, A., Zonneveld, K.A.F., Vercauteren, D., Vlassenbroeck, J., Louwye, S.,
 1048 2009. Process length variation in cysts of a dinoflagellate, *Lingulodinium machaerophorum*,
 1049 in surface sediments: Investigating its potential as salinity proxy. Mar. Micropaleontol. 70,
 1050 54–69.
 1051
 1052 Mertens, K.N., Dale, B., Ellegaard, M., Jansson, I.-M., Godhe, A., Kremp, A., Louwye, S.,
 1053 2011. Process length variation in cysts of the dinoflagellate *Protoceratium reticulatum*, from
 1054 surface sediments of the Baltic-Kattegat-Skagerrak estuarine system: a regional salinity
 1055 proxy. Boreas 40, 242–255.
 1056
 1057 Mertens, K.N., Bringué, M., Van Nieuwenhove, N., Takano, Y., Pospelova, V., Rochon, A.,
 1058 de Vernal, A., Radi, T., Dale, B., Patterson, R.T., Weckström, K., Andrén, E., Louwye, S.,
 1059 Matsuoka, K., 2012a. Process length variation of the cyst of the dinoflagellate *Protoceratium*
 1060 *reticulatum* in the North Pacific and Baltic-Skagerrak region: calibration as an annual density
 1061 proxy and first evidence of pseudo-cryptic speciation. J. Quaternary Sci. 27, 734–744.
 1062
 1063 Mertens, K.N., Price, A., Pospelova, V., 2012b. Determining the absolute abundance of
 1064 dinoflagellate cysts in recent marine sediments II: further tests of the *Lycopodium* marker-
 1065 grain method. Rev. Palaeobot. Palynol. 184, 74–81.
 1066
 1067 Mertens, K.N., Wolny, J., Carbonell-Moore, C., Bogus, K., Ellegaard, M., Limoges, A., de
 1068 Vernal, A., Gurdebeke, P., Omura, T., Mohd. A Al-Muftah, A., Matsuoka, K., 2015.
 1069 Taxonomic re-examination of the toxic armoured dinoflagellate *Pyrodinium bahamense* Plate

1070 1906: can morphology or LSU sequencing separate *P. bahamense* var. *compressum* from var.
1071 *bahamense*? Harmful Algae 41, 1–24.

1072

1073 Meunier A., 1910. Microplancton des mers de Barents & de Kara. In: Duc d'Orléans
1074 Campagne Arctique de 1907. Bulens, Bruxelles, 355 pp.

1075

1076 Mitrovic, S.M., Hamilton, B., McKenzie, L., Furey, A., James, K.J., 2005. Persistence of
1077 yessotoxin under light and dark conditions. Mar. Environ. Res. 60, 397–401.

1078

1079 [Morquecho, L., Góngora-González, D.T. & Okolodkov, Y.B. 2009. Cyst-theca relationships](#)
1080 [of Gonyaulacales and Peridinales \(Dinophyceae\) from Bahía Concepción, Gulf of California.](#)
1081 [Acta Botanica Mexicana 88, 9–29.](#)

1082

1083 Murray, G., Whitting, F.G., 1899. New Peridiniaceae from the Atlantic. Trans. Linn. Soc.
1084 Lond., Bot. 5, 321–342.

1085

1086 Paez-Reyes, M., Head, M.J., 2013. The Cenozoic ~~Gonyaulacacean~~ ~~gonyaulacacean~~
1087 ~~Dinoflagellate~~ ~~dinoflagellate~~ ~~Genera~~ ~~genera~~ *Operculodinium* Wall, 1967 and *Protoceratium*
1088 Bergh, 1881 and ~~Their~~ ~~their~~ ~~Phylogenetic~~ ~~phylogenetic~~ ~~Relationships~~ ~~relationships~~. J. Paleo.
1089 87(5), 786–803.

1090

1091 Paz, B., Riobó, P., Fernández, M.L., Fraga, S., Franco, J.M., 2004. Production and release of
1092 yessotoxins by the dinoflagellates *Protoceratium reticulatum* and *Lingulodinium polyedrum*
1093 in culture. Toxicon 44, 251–258.

1094

1095 | [Paz, B., Riobó, P., Ramilo, I., Franco, J.M., 2007. Yessotoxins profile in strains of](#)
1096 | [Protoceratium reticulatum](#) from Spain and USA. *Toxicon* 50, 1–17.

1097 |

1098 | Paz, B., Daranas, A.H., Norte, M., Riobó, P., Franco, J.M., Fernández, J.J., 2008.

1099 | Yessotoxins, a ~~Group-group~~ of ~~Marine-marine Polyether-polyether Toxinstoxins~~: an
1100 | ~~Overviewoverview~~. *Mar. Drugs* 6, 73–102.

1101 |

1102 | Paz, B., Blanco, J., Franco, J.M., 2013. Yessotoxins production during the culture of
1103 | *Protoceratium reticulatum* strains isolated from Galician Rias Baixas (NW Spain). *Harmful*
1104 | *algae* 21–22, 13–19.

1105 |

1106 | Pospelova, V., Esenkulova S., Johannessen S.C., ~~O'Brien-O'Brien~~ M. C. & Macdonald
1107 | R.W., 2010. Organic-walled dinoflagellate cyst production, composition and flux from 1996
1108 | to 1998 in the central Strait of Georgia (BC, Canada): a sediment trap study. *Mar.*
1109 | *Micropaleontol.* 75, 17–37.

1110 |

1111 | Reifel, K.M., McCoy, M.P., Roche, T.E., Tiffany, M.A., Hurlbert, S.H., Faulkner, D.J., 2002.

1112 | Possible importance of algal toxins in the Salton Sea, California. *Hydrobiologia* 473, 275–
1113 | 292.

1114 |

1115 | Reinecke, P., 1967. *Gonyaulax grindleyi* sp. nov.: a dinoflagellate causing a red tide at Elands
1116 | Bay, Cape Province in December 1966. *S. Afr. J. Bot.* 33, 157–160.

1117 |

1118 | Rochon, A., de Vernal, A., Turon, J.-L., Matthiessen, J., Head, M.J., 1999. Distribution of
1119 | ~~Recent-recent~~ dinoflagellate cysts in surface sediments from the North Atlantic Ocean and

1120 adjacent areas in relation to sea-surface parameters. American Association of Stratigraphic
1121 Palynologists, Contributions Series, no. 35, 146 p.
1122
1123 Röder, K., Fritz, N., Gerds, G., Luckas, B., 2011. Accumulation and ~~Depuration~~[depuraton](#)
1124 of ~~Yessotoxin~~[yessotoxin](#) in ~~Two two~~ [Bivalves](#)[bivalves](#). J. Shellfish Res. 30, 167–175.
1125
1126 Röder, K., Hantzsch, F.M., Gebühr, C., Miene, C., Helbig, T., Krock, B., Hoppenrath, M.,
1127 Luckas, B., Gerds, G., 2012. Effects of salinity, temperature and nutrients on growth, cellular
1128 characteristics and yessotoxin production of *Protoceratium reticulatum*. Harmful Algae 15,
1129 59–70.
1130
1131 Rossignol, M., 1964. Hystrichosphères du Quaternaire en Méditerranée orientale, dans les
1132 sédiments Pléistocènes et les boues marines actuelles. Revue de micropaléontologie 7, 83–99.
1133
1134 Sala-Pérez, M., Alpermann, T.J., Krock, B., Tillmann, U., 2016. Growth and bioactive
1135 secondary metabolites of arctic *Protoceratium reticulatum* (Dinophyceae). Harmful Algae 55,
1136 85–96.
1137
1138 Saldarriaga, J.F., Taylor, F.J.R., Cavalier-Smith, T., Menden-Deuer, S., Keeling, P.J., 2004.
1139 Molecular data and the evolutionary history of dinoflagellates. Eur. J. Protistol. 40, 85–111.
1140
1141 Salgado, P., Figueroa, R.I., Ramilo, I., Bravo, I., 2017. The life history of the toxic marine
1142 dinoflagellate *Protoceratium reticulatum* (Gonyaulacales) in culture. Harmful Algae 68, 67–
1143 81.
1144

1145 [Salgado, P., Fraga, S., Rodríguez, F., Riobó, P., accepted. *Ceratocorys mariaovidiorum* sp.](#)
1146 [nov. \(Gonyaulacales\), a new dinoflagellate species previously reported as *Protoceratium*](#)
1147 [*reticulatum*. Journal of Phycology, accepted.](#)
1148
1149 Samdal, I.A., Naustvoll, L.J., Olseng, C.D., Briggs, L.R., Miles, C.O., 2004. Use of ELISA to
1150 identify *Protoceratium reticulatum* as a source of yessotoxin in Norway. Toxicon 44, 75–82.
1151
1152 Satake, M., Ichimura, T., Sekiguchi, K., Yoshimatsu, S., Oshima, Y. 1999. Confirmation of
1153 Yessotoxin and 45,46,47-Trinoyessotoxin production by *Protoceratium reticulatum* collected
1154 in Japan. Natural toxins 7, 147–150.
1155
1156 Schiller, J., 1937. Dinoflagellatae (Peridineae) in monographischer Behandlung. In: Dr. L.
1157 Rabenhorst's Kryptogamen-Flora von Deutschland, Österreich und der Schweiz. Bd. 10(3).
1158 Teil 2(3), pp. 321–480.
1159
1160 Scholin, C.A., Herzog, M., Sogin, M., Anderson, D.M. 1994. Identification of group- and
1161 strain-specific genetic markers for globally distributed *Alexandrium* (Dinophyceae). II.
1162 Sequence analysis of a fragment of the LSU rRNA gene. J. Phycol. 30, 999–1011.
1163
1164 [Steidinger K.A. & J. Williams \(1970\) Dinoflagellates. Memoirs of the Hourglass Cruises 2,](#)
1165 [1–251. Published by Marine Research Laboratory, Florida Department of Natural Resources,](#)
1166 [St. Petersburg, Florida.](#)
1167

1168 Stein, F. von, 1883. Der Organismus der Infusionsthier. 3. Abt. Der Organismus der
 1169 Arthrodelen Flagellaten nach eigenen Forschungen in systematischer Reihenfolge bearbeitet.
 1170 2. Hälfte. Einleitung und Erklärung der Abbildungen, W. Engelmann, Leipzig, pp. 1–30.
 1171
 1172 Stosch, H.A. von, 1969. Dinoflagellaten aus der Nordsee I. Über *Cachonina niei* Loeblich
 1173 (1968), *Gonyaulax grindleyi* Reinecke (1967) und eine Methode zur Darstellung von
 1174 Peridineenpanzern. Helgoländ. Wiss. Meer. 19, 558–568.
 1175
 1176 Suzuki, T., Horie, Y., Koike, K., Satake, M., Oshima, Y., Iwataki, M., Yoshimatsu, S., 2007.
 1177 Yessotoxin analogues in several strains of *Protoceratium reticulatum* in Japan determined by
 1178 liquid chromatography–hybrid triple quadrupole/linear ion trap mass spectrometry. Journal of
 1179 Chromatography A 1142, 172–177.
 1180
 1181 Takano, Y., Horiguchi, T., 2004. Surface ultrastructure and molecular phylogenetics of four
 1182 unarmoured heterotrophic dinoflagellates, including the type species of the genus *Gyrodinium*
 1183 (Dinophyceae). Phycol. Res. 52, 107–116.
 1184
 1185 Takano, Y., Horiguchi, T. 2006. Acquiring scanning electron microscopical, light
 1186 microscopical and multiple gene sequence data from a single dinoflagellate cell. J. Phycol. 42,
 1187 251–256.
 1188
 1189 Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S. 2013 MEGA6: Molecular
 1190 Evolutionary Genetics Analysis Version 6.0. Mol Biol Evol. 30: 2725–2729
 1191

1192 [Tiffany, M.A., González, M.R., Swan, B.K., Reifel, K.M., Watts, J.M., Hurlbert, S.H. 2007.](#)
1193 [Phytoplankton dynamics in the Salton Sea, California, 1997-1999. Lake and Reservoir](#)
1194 [Management 23, 582–605.](#)
1195
1196 Wall, D., 1967. Fossil microplankton in deep-sea cores from the Caribbean Sea.
1197 Palaeontology 10, 95–123.
1198
1199 Wall, D., Dale, B., 1966. “Living fossils” in western Atlantic plankton. Nature 211 (5053),
1200 1025–1026.
1201
1202 Wall, D., Dale, B., 1968. Modern dinoflagellate cysts and evolution of the Peridiniales.
1203 Micropaleontology 14, 265–304.
1204
1205 Watts, J.M., Swan, B.K., Tiffany, M.A., Hurlbert, S.H., 2001. Thermal, mixing and oxygen
1206 regimes of the Salton Sea, California, 1997–1999. Hydrobiol. 162, 159–176.
1207
1208 Wołoszyńska, J., 1929. Dinoflagellatae Polskiego Bałtyku i blot nad Piasnica. Archivum
1209 Hydrobiologii i Rybactwa 3, 153–278 [In Polish].
1210
1211 Yamaguchi, A., Yoshimatsu, S., Hoppenrath, M., Wakeman, K.C., Kawai, H. 2016.
1212 Molecular phylogeny of the benthic dinoflagellate genus *Amphidiniopsis* and its relationships
1213 with the family Protoperidiniaceae. Protist. 167, 568–583.
1214
1215 Zonneveld, K.A.F., Marret, F., Versteegh, G.J.M., Bogus, K., Bonnet, S., Bouimetarhan, I.,
1216 Crouch, E., de Vernal, A., Elshanawany, R., Edwards, L., Esper, O., Forke, S., Grøsfjeld, K.,

1217 Henry, M., Holzwarth, U., Kielt, J.-F., Kim, S.-Y., Ladouceur, S., Ledu, D., Chen, L.,
1218 Limoges, A., Londeix, L., Lu, S.-H., Mahmoud, M.S., Marino, G., Matsouka [sic], K.,
1219 Matthiessen, J., Mildenhall [sic], D.C., Mudie, P., Neil, H.L., Pospelova, V., Qi, Y., Radi, T.,
1220 Richerol, T., Rochon, A., Sangiorgi, F., Solignac, S., Turon, J.L., Verleye, T., Wang, Y.,
1221 Wang, Z., Young, M., 2013. Atlas of modern dinoflagellate cyst distribution based on 2405
1222 datapoints. *Rev. Palaeobot. Palynol.* 191, 1–197.
1223

Figure captions

Figure 1. Sites of studied plankton samples and cultured strains containing thecate stages of *Pentaplacodinium saltonense* (in red) and *Protoceratium reticulatum* (in blue). The locations are listed in Table 1.

Figure 2. Line drawings of extant members of the subfamily *Cribroperidinioideae* in dorsal view to show the dextral torsion typical of these gonyaulacoids. A. *Pentaplacodinium saltonense*. B. *Protoceratium reticulatum*. C. *Lingulodinium polyedra*. Labeling of tabulation follows a modified Kofoid system that recognizes homologs.

Figure 3. Line drawing of sulcal area of *Pentaplacodinium saltonense*. FP: flagellar pore; Sa: anterior sulcal plate; Sda: right anterior sulcal plate; Sdp: right posterior sulcal plate; Ssa: anterior left sulcal plate; Ssp: posterior left sulcal plate; Sp: posterior sulcal plate; c: cingular plates.

Figure 4. Line drawings of epithecal overlapping plate patterns of gonyaulacoids discussed in this paper. Arrows indicate direction of overlap. A. *Pentaplacodinium saltonense*. B. *Protoceratium reticulatum*. C. *Lingulodinium polyedra*. D. *Ceratocorys horrida*. E. *Gonyaulax spinifera*.

Figure 5. Neighbour-joining tree of *P. reticulatum*, *P. saltonense* and related strains sequenced in this study and sequences from Genbank based on an 80 bp alignment of the V9 region of the SSU gene. Bootstrap values were retrieved from 1000 replicates and those >70% are indicated at the nodes for neighbour-joining and maximum likelihood respectively.

Strain names are indicated followed by their geographic origin and accession number (Genbank).

Figure 6. Neighbour-joining tree of *P. reticulatum*, *P. saltonense* and related strains sequenced in this study and sequences from Genbank based on a 571 bp alignment of the V4 region of the LSU gene. Bootstrap values were retrieved from 1000 replicates and those >70% are indicated at the nodes for neighbour-joining and maximum likelihood respectively. Strain names are indicated followed by their geographic origin and accession number (Genbank).

Figure 7. Neighbour-joining tree of *P. reticulatum*, *P. saltonense* and related strains sequenced in this study and sequences from Genbank based on a 356 bp alignment of the ITS 1–2 region. Bootstrap values were retrieved from 1000 replicates and those >70% are indicated at the nodes for neighbour-joining and maximum likelihood respectively. Strain names are indicated followed by their geographic origin and accession number (Genbank).

Plate Captions

Plate 1. Light microscope images of *Pentaplacodinium saltonense* based on cyst-theca experiment from the Salton Sea. A. Living cyst from the Salton Sea St. 1. B–F. Germinated cyst from St. 2 (culture 2E3 used for single-cell PCR). B. Cross section, showing attached operculum. C. Focus on elongated simple operculum reflecting plates $2''+*(3''+4'')$. D–E. Focus on archeopyle, after removal of operculum. F. Cross section showing processes. G–I. Germinated cyst from St. 2 (culture 1A7 used for single-cell PCR). G. Focus on precingular

1273 archeopyle reflecting plate *(3''+4''), showing attached operculum. H. Focus on operculum. I.
1274 Cross section, showing opened operculum. Scale bars = 20 µm.

1275

1276 **Plate 2.** Light microscope images of cyst-theca experiment from the Salton Sea. A–I. Images
1277 of living cells of *Pentaplacodinium saltonense* germinated from cyst depicted in Plate 1, Figs.
1278 B–F (culture 2E3). A. Globular cell. B. Angular cell. C. Fusiform cell. D. Epitheca. E.
1279 Hypotheca. F. Ventral view showing configuration of apical plates. G–I. Sulcal plates. Scale
1280 bars = 20 µm.

1281

1282 **Plate 3.** Scanning electron microscope images of *Pentaplacodinium saltonense*, all different
1283 cells from the Salton Sea, except D. A. Holotype. Ventral view. Arrowhead points to ventral
1284 pore between plates 1' and *4'. Arrow shows flange on plate 1p. B. Apical view, missing the
1285 cover plate. Small arrowhead points to small pores inside the thecal pores. Large arrowhead
1286 points to ventral pore between plates 1' and *4'. Small arrowhead points to the three minute
1287 pores inside most pores. C. Dorsal view, showing dextral torsion. Note the cell roundness. D.
1288 Sulcal plates of a cell from culture SSCAP K-1479 ([Indian River Lagoon, Florida](#)).
1289 Arrowhead shows the narrow point of contact between the Sa and 1' plates. E. Antapical
1290 view. Scale bars A–C, E = 10 µm; D = 5 µm.

1291

1292 **Plate 4.** Scanning electron microscope images of *Pentaplacodinium saltonense* from the
1293 Indian River Lagoon. A. Apical view of a cell from culture SSCAP K-1479. Arrowhead
1294 points to ventral pore between plates 1' and *4'. B. Same specimen as in A. Ventral view.
1295 Arrowhead points to ventral pore between plates 1' and *4'. C. A different cell from a
1296 plankton sample courtesy of Paul Hargraves. Ventral view. Arrowhead points to ventral pore
1297 between plates 1' and *4'. Arrow shows flange on plate 1p. D. Ventral view of a cell from a

1298 culture established by Paul Hargraves. E. Antapical view of a cell from the same culture as in
1299 D. E. Apical view of a cell from the same culture as in D. D–F: SEMs by Paul Hargraves.
1300 Scale bars = 10 µm.

1301
1302 **Plate 5.** Scanning electron microscope images of cysts of *Pentaplacodinium saltonense*
1303 extracted from Salton Sea sediment (St. 2) using palynological methods. A–C. Views showing
1304 shape of archeopyle, reflecting plate *(3''+4''). D. Specimen that is torn along the cingulum.
1305 E. Specimen showing alignment of processes along the cingulum. F. Specimen with relatively
1306 large openings in cyst wall. G. Specimen with distinct intratabular processes. H. Specimen
1307 with relatively coarsely reticulated wall surface. I. Internal view of smooth cyst wall. Scale
1308 bars = 10 µm.

1309
1310 **Plate 6.** Scanning electron microscope images of cysts of *Pentaplacodinium saltonense*
1311 formed in culture of strain 3243 (Indian River Lagoon). A. Specimen showing preformed
1312 archeopyle and margins of principal archeopyle suture with reduced ornament. B. Specimen
1313 with attached thecal plate. C. Specimen with partly developed processes. D. Specimen with
1314 processes clearly reflecting tabulation. E. Specimen with preformed archeopyle. F. Specimen
1315 showing reflection of the sulcus. G–H. Specimen with well-developed wall texture. I. Wall
1316 texture of specimen with ‘spider-web’ microreticulation. Scale bars = 10 µm, except H, I,
1317 scale bars = 1 µm.

1318
1319 **Plate 7.** Scanning electron microscope images of *Protoceratium reticulatum* cells and of the
1320 apical pore plates of the gonyaulacoids discussed in this study. A. *Protoceratium reticulatum*.
1321 Cell from Greenland, ventral view. B. Same cell in apical view. C. *Protoceratium reticulatum*.
1322 Cell from Elands Bay, South Africa. Dorsal view, note the dextral torsion. D. Apical pore

1323 plates of a different cell of *Protoceratium reticulatum* (Greenland). E. Apical pore plates of a
1324 cell of *Ceratocorys horrida* [\(Central equatorial Pacific\)](#). F. Apical pore plates of a cell of
1325 *Pentaplacodinium saltonense* from culture SSCAP K-1479 [\(Indian River Lagoon\)](#). G. Apical
1326 pore plates of a cell of *Ceratocorys gourretii*. H. Apical pore plates of another cell of
1327 *Pentaplacodinium saltonense* from culture SSCAP K-1479 [\(Indian River Lagoon\)](#). Scale bars
1328 A–C = 10 μm ; D–H = 5 μm .

1329

Figure 1
[Click here to download high resolution image](#)

Figure 2
[Click here to download high resolution image](#)

Figure 3
[Click here to download high resolution image](#)

Figure 4
[Click here to download high resolution image](#)

Figure 5
[Click here to download high resolution image](#)

Figure 5

Figure 6

[Click here to download high resolution image](#)

Figure 6

Figure 7
[Click here to download high resolution image](#)

Figure 7

Table 1

Table 1. Site location of plankton samples investigated, location mark on Figure 1, sampling date, latitude, longitude, sea surface salinity (psu), sea surface temperature (°C), sampling device, used fixative, and name of persons who did the sampling.

Sampling site	Location mark on Figure 1	Sampling date	Latitude (°)	Longitude (°)	SSS (psu)	SST (°C)	Sampling device	Fixative used	Species present	Sampled by
Salton Sea, St. 1, California, USA	A	24-Oct-09	33,50	-115,91	>40.2 (62)	23,1	Plankton net > 20 µm	Ethanol 100%	PS	KM, VP, MH, MCCM
Salton Sea, St. 2, California, USA	A	24-Oct-09	33,50	-115,91	>40.2 (64)	25,5	Plankton net > 20 µm	Ethanol 100%	PS	KM, VP, MH, MCCM
Salton Sea, St. 3, California, USA	A	24-Oct-09	33,50	-115,92	>40.2 (65)	25,5	Plankton net > 20 µm	Ethanol 100%	PS	KM, VP, MH, MCCM
Salton Sea, St. 4, California, USA	A	24-Oct-09	33,50	-115,91	>40.2 (56)	30,3	Plankton net > 20 µm	Ethanol 100%	PS	KM, VP, MH, MCCM
Salton Sea, St. 5, California, USA	A	24-Oct-09	33,50	-115,91	>40.2 (65)	~30	Plankton net > 20 µm	Ethanol 100%	PS	KM, VP, MH, MCCM
Off Yucatan, St. 4, Gulf of Mexico, Mexico	B	19-May-09	21,39	-88,08	31,0	24,1	Plankton tow >20 µm	Formaldehyde	PS	YO
Off Yucatan, Gulf of Mexico, Mexico	B	19-May-09	21,40	-88,84	31,0	24,1	Plankton tow >20 µm	Formaldehyde	PS	YO
Indian River Lagoon, St. TR, Florida, USA	C	28-May-08	27,50	-80,34	37,4	29,5	Plankton tow >20 µm	Formalin 2%	PS	PH
Off Qatar, Persian Gulf	D	Sept. 1991	25,29	51,54	38-43	20-35	Plankton net	Formalin 5%	PS	AA
North Sea, Helgoland, Germany	E	3-Jun-02	54,19	7,90	32,0	12,5	Plankton net >20 µm	Lugol	PR	MHO
Kattegat, St. Central, Denmark	F	NA	56,92	11,28	NA	NA	Plankton net	Lugol & Formalin	PR	JL
Kattegat, St. 431, Denmark	F	22-Jun-05	55,77	12,75	NA	NA	Plankton net	Lugol	PR	JL
Kattegat, St. 925, Denmark	F	17-Aug-00	56,08	11,02	NA	NA	Plankton net	Lugol	PR	JL
Baltic Sea, St. F64, Finland	G	20-Aug-10	60,18	19,13	5,7	17,6	Plankton net	Lugol	PR	AK, MHU
Gulf of Bothnia, Baltic Sea, St. US5B, Finland	H	20-Aug-10	62,58	19,98	5,1	16,0	Plankton net	Lugol	PR	AK, MHU
Western Greenland, St. 516, Denmark	I	30-Jul-12	69,20	-54,10	33,2	6,9	Water Bottle-CTD	Formaldehyde	PR	UT
Baffin Bay, St. 2008-029-0039/9/0039A, Canada	J	6-Sep-08	76,57	-73,96	31,0	2,2	Plankton net >20 µm	Formaldehyde	PR	AR
Baffin Bay, St. 2008-029-027A, Canada	J	5-Sep-08	77,29	-74,34	30,8	2,2	Plankton net >20 µm	Formaldehyde	PR	AR
Baffin Bay, St. 2008-029-0043A, Canada	K	7-Sep-08	75,58	-78,63	31,0	2,0	Plankton net >20 µm	Formaldehyde	PR	AR
Baffin Bay, St. 2008-029-0035A, Canada	J	6-Sep-08	76,33	-71,43	31,0	4,5	Plankton net >20 µm	Formaldehyde	PR	AR
Off Puerto Aguirre, Chile	L	21-May-06	-44,99	-73,53	28,8	10,0	Plankton net > 26 µm	Formaldehyde	PR	XV
Elands Bay, South Africa	M	16-Mar-13	-32,31	18,32	NA	NA	Plankton net	Formaldehyde	PR	MP
Off Cape Town, South Africa	M	NA	-33,89	18,42	NA	NA	Plankton net	Lugol + formalin	PR	JL
Mutsu Bay, Aomori, Japan	N	5-Apr-10	40,92	141,12	32,7	5,9	Plankton net >20 µm	1% Glutaraldehyde	PR	KK
Okkirai Bay, Iwate, Japan	O	16-Jul-03	39,08	141,85	32,6	15,8	Plankton net >20 µm	5% Glutaraldehyde	PR	KK
Okkirai Bay, Iwate, Japan	O	28-Aug-03	39,08	141,85	32,3	20,0	Plankton net >20 µm	5% Glutaraldehyde	PR	KK
Okkirai Bay, Iwate, Japan	O	23-Jun-04	39,08	141,85	32,7	16,0	Plankton net >20 µm	5% Glutaraldehyde	PR	KK
Okkirai Bay, Iwate, Japan	O	12-Aug-04	39,08	141,85	33,5	23,0	Plankton net >20 µm	5% Glutaraldehyde	PR	KK
Omura Bay, Nagayo-ura, Japan	P	9-May-11	32,85	129,87	32,7	20,2	Plankton net >20 µm	Formalin	PR	KMA
Omura Bay, Inoura, Japan	P	12-May-11	33,05	129,74	25,6	18,5	Plankton net >20 µm	Formalin	PR	KMA
Omura Bay, Togitsu Port, Japan	P	23-May-11	32,85	129,87	32,6	19,5	Plankton net >20 µm	Formalin	PR	KMA
Kagoshima Bay, St. 1, Japan	P	20-Jun-11	31,55	130,57	23,8	22,9	Plankton net >20 µm	Formalin	PR	KMA
Saroma Lake (Lagoon), St. 1, Japan	Q	22-Jul-11	44,12	143,82	32,2	17,1	Plankton net	Ethanol	PR	KM

Saanich Inlet, St. S2, BC, Canada	R	13-Jul-10	48,55	-123,53	27,8	14,8	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. S5, BC, Canada	R	13-Jul-10	48,71	-123,46	27,8	13,6	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. S3, BC, Canada	R	13-Jul-10	48,59	-123,48	27,8	14,8	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. S4.5, BC, Canada	R	14-Jul-10	48,67	-123,49	28,0	13,7	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. S11, BC, Canada	R	14-Jul-10	48,73	-123,54	28,0	13,7	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. S4, BC, Canada	R	13-Jul-10	48,63	-123,49	27,4	15,1	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. Pat UVic 262, BC, Canada	R	10-Aug-11	48,65	-123,44	29,0	17,8	Plankton net	Formaldehyde	PR	VP
Saanich Inlet, St. Pat UVic 264, BC, Canada	R	21-Aug-11	48,65	-123,44	31,0	17,9	Plankton net	Formaldehyde	PR	VP

Abbreviations: NA = Not available, PS = *Pentaplecodinium saltonense*, PR = *Protoceratium reticulatum*, AK = Anke Kremp, AR = André Rochon, AA = Abdulrahman Al-Muftah, MCCM = Consuelo Carbonell-Moore, JL = Jacob Larsen, KK = Kazuhiko Koike, KM = Kenneth Neil Mertens, KMA = Kazumi Matsuoka, MH= Martin J. Head, MHO = Mona Hoppenrath, MHU=Maia Huttunen, MP = Maya Pfaff, PH = Paul Hargreaves, UT = Urban Tillmann, VP = Vera Pospelova, XV = Ximena Vivanco, YO = Yuri B. Okolodkov

Suppl. Table 1. Culture strains and cells or cyst picked for SEM and/or phylogenetic analysis.

CODI (Strain ID)	Identified here by its morphology as	Geographic Origin	Latitude (°N)	Longitude (°E)	Isolation date	Isolated by	LSU (28S) Genbank	ITS Genbank	SSU (18S) Genbank	Sequenced in this study by
CCMP404 = ALO011	<i>P. saltonense</i>	Salton Sea (California)	33.375 33°	-116.0	1966	Dodson, A.	EU532476	FJ489629	FJ489629***	Aika Yamaguchi
Salton Sea 2E3 *	<i>P. saltonense</i>	Salton Sea (California)	30.122'	-115° 54.879'	19/11/2013	Mertens, K.	MG646301	Not sequenced	MG646323	
CCMP1720 = K-1475 = ALO013	<i>P. saltonense</i>	Biscayne Bay (Florida)	25.8	-80.3333	15/02/1995	Hargraves, P.	FJ155820	FJ489628	FJ489628***	
CCMP1721 = K-1480 = ALO012	<i>P. saltonense</i>	Biscayne Bay (Florida)	25.8	-80.3333	01/06/1994	Hargraves, P.	EU165321	Not sequenced	Not sequenced	
CCMP3241 = K-1479	<i>P. saltonense</i>	Indian River Lagoon (Florida)	27.83	-80.45	11/06/2008	Hargraves, P.	Not sequenced	MG646297	MG646322	
CCMP3243 = K-1476	<i>P. saltonense</i>	Indian River Lagoon (Florida)	27.83	-80.45	05/06/2008	Hargraves, P.	Not sequenced	MG646296	MG646321	Andrea Highfield / Declan Schroeder
CCMP3031 = K-1474	<i>P. reticulatum</i>	Off coast Florida, Gulf of Mexico	25.0167	-81.4003	NA	NA	Not sequenced	MG646287	MG646314	Andrea Highfield / Declan Schroeder
Victoria-no.4_5_6 *	<i>P. reticulatum</i>	Brentwood Bay, Saanich Inlet	48.57	123.47	Oct. 2011	Mertens, K.	AB727656	AB727656	AB727656	
CCMP1889 = K-0634 = CCCM535	NA	Friday Harbor, San Juan Island, Washington USA	48.544	-123.01	1983	Taylor, F.J.R.	EU532475	EU532484	FJ626858***	Andrea Highfield / Declan Schroeder
CCMP2776 = K-1477	<i>P. reticulatum</i>	Gulf of Mexico, U.S.A.	25,0167	-81,4003	NA	Sinigalliano, C.	EU165290	MG646288	MG646315	Andrea Highfield / Declan Schroeder
CCMP3113 = K-1478	<i>P. reticulatum</i>	Marquesa Keys, Florida Keys	-24.58	-82.1	NA	NA	Not sequenced	MG646286	MG646316	Andrea Highfield / Declan Schroeder
VGO757	NA	Ebro Delta (Catalonia, Spain)	NA	NA	NA	Fernandez-Tejedor, M.	FJ155819	DQ990371	DQ990371***	
VGO758	<i>P. reticulatum**</i>	Alfacs Bay, Ebro Delta (Catalonia Spain)	NA	NA	NA	Fernandez-Tejedor, M.	FJ155817	FJ489624	FJ489624***	Andrea Highfield / Declan Schroeder
VGO903 = ALO014	<i>P. reticulatum**</i>	Ría de Pontevedre (Bueu)	NA	NA	NA	Fernandez-Tejedor, M.	FJ155821	FJ489625	FJ489625***	
VGO904	NA	NA	NA	NA	NA	Fernandez-Tejedor, M.	FJ155822	FJ489630	FJ489630***	
VGO905	NA	NA	NA	NA	NA	Fernandez-Tejedor, M.	FJ155823	FJ489627	FJ489627***	
IRTA015	<i>P. reticulatum</i>	Ebro Delta (Catalonia Spain)	NA	NA	NA	Fernandez-Tejedor, M.	EF642980	EF642972	EF642972***	
20	NA	Catalan Coast	NA	NA	NA	Fernandez-Tejedor, M.	Not sequenced	EU327363	EU327363***	
GG1AM	NA	La Atunara (Cádiz, Spain)	NA	NA	NA	Fernandez-Tejedor, M.	FJ155824	FJ489626	FJ489626***	
17	NA	Catalan Coast	NA	NA	NA	Fernandez-Tejedor, M.	Not sequenced	EU327362	EU327362***	
4	NA	Catalan Coast	NA	NA	NA	Fernandez-Tejedor, M.	Not sequenced	EU327361	EU327361***	
Katte-B3 *	<i>P. reticulatum</i> (cyst-based)	Kattegat, Swedish coast	57.5	11.8	Nov. 2011	Mertens, K.	AB727655	AB727655	AB727655	Karin Röder
091223-38_M16_Protocer3-ITS1	<i>P. reticulatum</i>	Helgoland, Germany	54,19	7,9	avr-03	Hoppenrath, M.	Not sequenced	MG646294	Not sequenced	
Lake Saroma *	<i>P. reticulatum</i> (cyst-based)	Lake Saroma, Japan	44.12	143.87	Aug. 2011	Mertens, K.	AB727654	AB727654	AB727654	Andrea Highfield / Declan Schroeder
K-0976	<i>P. reticulatum</i>	Attu, Greenland	67.924068	-53.649824	21.08.2005	Moestrup, Ø.	Not sequenced	MG646292	MG646313	
Arctic E12 = K1-1-1 - K-1-3 *	<i>P. reticulatum</i>	Station 323, Northern Baffin Bay	74.12	79.45	02-mai-11	Mertens, K.	MG646300	MG646293	MG646333	Yoshihito Takano
K-0485	<i>P. reticulatum</i>	Southern Kattegat, Bouy St.	56.20	12.04	03-avr-89	Hansen, G.	AF260386	Not sequenced	Not sequenced	
PRSH6 (NRC Halifax)	<i>P. reticulatum</i>	Ship harbour, NS, Canada	NA	NA	NA	Ferrell, J. F.	Not sequenced	EU927572	EU927572***	Andrea Highfield / Declan Schroeder
PRSH3 (NRC Halifax)	<i>P. reticulatum</i>	Ship harbour, NS, Canada	NA	NA	NA	Ferrell, J. F.	Not sequenced	EU927569	EU927569***	
PRSH4 (NRC Halifax)	<i>P. reticulatum</i>	Ship harbour, NS, Canada	NA	NA	NA	Ferrell, J. F.	Not sequenced	EU927570	EU927570***	
PRSH1 (NRC Halifax)	<i>P. reticulatum</i>	Ship harbour, NS, Canada	NA	NA	NA	Ferrell, J. F.	Not sequenced	EU927568	EU927568***	
CTCC 01	<i>P. reticulatum</i>	Southern Benguela upwelling region	NA	NA	NA	NA	EU532477	EU532486	EU532486***	
Elands Bay 2013 (EB2) *	<i>P. reticulatum</i>	Elands Bay, South Africa	32°18.618'	18°19.267	16 March 2013	Carbonell-Moore, C.	Not sequenced	Not sequenced	MG646319	

JHW0007-6	NA	Jinhae Bay, Korea	NA	NA	juil-00	Kim, K.-Y.	EF613362	Not sequenced	AY421790	
CBA-1	NA	Adriatic Sea, Ancona, Italy	NA	NA	NA	Totti C.	Not sequenced	AM183800	AM183800***	
020717-OK-PR7	<i>P. reticulatum</i>	Okkirai Bay, Iwate, Japan	39.08	141.85	17-Jul-02	Koike, K.	Not sequenced	MG646283	Not sequenced	Kazuhiko Koike
030707-YB-PR1	<i>P. reticulatum</i>	Yamada Bay, Iwate, Japan	39.46	141.97	7-Jul-03	Koike, K.	Not sequenced	MG646284	Not sequenced	Kazuhiko Koike
030624-OK-PR1	<i>P. reticulatum</i>	Okkirai Bay, Iwate, Japan	39.08	141.85	24-Jun-03	Koike, K.	Not sequenced	MG646285	Not sequenced	Kazuhiko Koike
Omura-no.7_no.8 *	<i>P. reticulatum</i>	Omura Bay, Japan	32.85	129.87	2 may 2011	Takano, Y.	Not sequenced	MG646291	MG646332	Yoshihito Takano
Nagayo-sc1 *	<i>P. reticulatum</i>	Nagayo, Japan	32.85	129.87	24 May 2011	Takano, Y.	MG646299	MG646290	MG646331	Yoshihito Takano
Togitsu-sc2 *	<i>P. reticulatum</i>	Togitsu Port, Japan	32.85	129.87	23 May 2011	Takano, Y.	Not sequenced	MG646289	Not sequenced	Yoshihito Takano
777	<i>P. reticulatum</i>	Concarneau large, France	47.83	-3.95	16 July 2008	Nézan, E.	Not sequenced	Not sequenced	MG646317	Gwenael Bilien
778	<i>P. reticulatum</i>	Concarneau large, France	47.83	-3.95	16 July 2008	Nézan, E.	Not sequenced	Not sequenced	MG646318	Gwenael Bilien
16-034	<i>Ceratocorys armata</i>	Bouée 7, Arcachon, France	44.54	-1.26	18 Dec 2015	Nézan, E.	MG646311	Not sequenced	MG646329	Gwenael Bilien
16-246	<i>Ceratocorys gourretii</i>	Station B70, Banyuls-sur-mer, French Mediterranean	42.48	3.18	26 Sept 2016	Nézan, E.	MG646310	Not sequenced	MG646328	Gwenael Bilien
16-245	<i>Ceratocorys horrida</i>	Station B70, Banyuls-sur-mer, French Mediterranean	42.48	3.18	26 Sept 2016	Nézan, E.	MG646312	Not sequenced	MG646330	Gwenael Bilien
CCMP157	<i>Ceratocorys horrida</i>	Banda Sea, South East Asia	-5,00	130,00	22 April 1975	Sweeney, B.	Not sequenced	EU927577	AF022154	
SCCAP K-1504	<i>Thecadinium kofoidii</i>	Helgoland, Germany	54.19	7.9	Aug. 2002	Hoppenrath, M.	GU295207***	Not sequenced	GU295204	
JH0210	<i>Alexandrium affine</i>	Jinhae Bay, South Korea	NA	NA	NA	NA	Not sequenced	Not sequenced	AY775286	
JHW0004-12	<i>Alexandrium tamarense</i>	Korea	NA	NA	NA	NA	Not sequenced	Not sequenced	AY421777	
CCMP116	<i>Alexandrium tamarense</i>	Ria de Vigo, Spain	42.23	-8.8	1 June 1984	Yentsch, C.M.	HM483868	Not sequenced	Not sequenced	
CCMP1738	<i>Lingulodinium polyedra</i>	Gulf of Mexico, USA	27.8	-97.13	NA	Buskey, E.	Not sequenced	Not sequenced	EF492507	
DRW0208	<i>Lingulodinium polyedra</i>	Korea	NA	NA	Aug. 2001	NA	Not sequenced	Not sequenced	AY421788	
BLACK1	<i>Lingulodinium polyedra</i>	Black Sea, Ukraine	49.90	30.29	2011	Takano, Y.	AB678400	AB678399***	AB693195***	
CASP1	<i>Lingulodinium polyedra</i>	Caspian Sea, Iran	37.51	49.91	2011	Takano, Y.	AB678402	AB678401***	AB693194***	
SANPEDRO1	<i>Lingulodinium polyedra</i>	San Pedro Harbor, California, USA	33.74	-118.24	2011	Takano, Y.	AB678404	AB678403***	AB693196***	
GS0209	<i>Gonyaulax polygramma</i>	Gunsan, South Korea	NA	NA	NA	NA	Not sequenced	Not sequenced	AY775287	
11A1	<i>Protoperidinium tricingulatum</i>	Wadden Sea, Netherlands	53,60	6,58	2006	Kawami, H.	Not sequenced	Not sequenced	AB716917	
CCAP1140/3	<i>Parvodinium inconspicuum</i>	Kl. Ukleisee, Schleswig-Holstein, Germany	NA	NA	NA	Meyer	Not sequenced	Not sequenced	FR865631	
hwi d1	Unidentified	Hawaii	21,59	-158,10	04-mars-14	Anne de Vernal and Geneviève Vautour	MG646298	MG646295	MG646320	Haifeng Gu
GgSm10R	Unidentified	Malaysia	1,60	110,32	17-janv-13	Bao Juan Kam	MG646305	Not sequenced	MG646326	Haifeng Gu
GgSm11R	Unidentified	Malaysia	1,60	110,32	17-janv-13	Bao Juan Kam	MG646306	Not sequenced	MG646327	Haifeng Gu
GgSm01R	Unidentified	Malaysia	1,60	110,32	10-août-10	Toh Hii Tan	MG646302	Not sequenced	MG646324	Haifeng Gu
GgSm03R	Unidentified	Malaysia	1,60	110,32	22-sept-10	Toh Hii Tan	MG646303	Not sequenced	Not sequenced	Guat Ru Liow
GgSm07R	Unidentified	Malaysia	1,60	110,32	17-janv-13	Toh Hii Tan	MG646304	Not sequenced	MG646325	Zhen Fei Lim
PrTT02R	Unidentified	Malaysia	1,92	109,77	28-mars-13	Sing Tung Teng	MG646308	Not sequenced	Not sequenced	Haifeng Gu
PrTT03R	Unidentified	Malaysia	1,92	109,77	28-mars-13	Sing Tung Teng	MG646309	Not sequenced	Not sequenced	Haifeng Gu
PrTT01R	Unidentified	Malaysia	1,92	109,77	28-mars-13	Sing Tung Teng	MG646307	Not sequenced	Not sequenced	Haifeng Gu

					sequenced	
--	--	--	--	--	-----------	--

* = single cells or cysts sequenced through single-cell PCR; Accession numbers in bold denote sequences from this study. ** = these cultures showed presence of 5 precingular plates and are considered aberrant. *** = sequence not used in phylogenies. NA = Not acknowledged.