

Published online: 03 August 2018

OPEN Author Correction: Intensified summer monsoon and the urbanization of Indus Civilization in northwest India

Yama Dixit^{1,2,9}, David A. Hodell¹, Alena Giesche¹, Sampat K. Tandon³, Fernando Gázquez^{1,4}, Hari S. Saini⁵, Luke C. Skinner¹, Syed A. I. Mujtaba⁵, Vikas Pawar⁶, Ravindra N. Singh⁷ & Cameron A. Petrie 608

Correction to: Scientific Reports https://doi.org/10.1038/s41598-018-22504-5, published online 09 March 2018

In Figure 4 the Ostracod curve does not align with δD (orange) and $\delta^{18}O$ (blue) curves. The correct Figure 4 appears below as Figure 1.

¹Godwin Laboratory for Palaeoclimate Research, Department of Earth Sciences, University of Cambridge, Cambridge, CB2 3EQ, United Kingdom. ²IFREMER, Unité de Recherche Géosciences Marines, Z.I. Pointe du Diable, BP 70, 29280, Plouzané, France. ³Department of Earth and Environmental Sciences, IISER Bhopal, India. ⁴School of Earth and Environmental Sciences, University of St. Andrews, St. Andrews, UK. ⁵Geological Survey of India, Faridabad, India. ⁶Department of History, Maharshi Dayanand University, Rohtak, Haryana, India. ⁷Department of AIHC and Archaeology, Banaras Hindu University, Varanasi, India. ⁸Department of Archaeology, University of Cambridge, Cambridge, CB2 3DZ, United Kingdom. ⁹Present address: Earth Observatory of Singapore, Nanyang Technological University, 50 Nanyang Avenue, 639798, Singapore. Correspondence and requests for materials should be addressed to Y.D. (email: ydixit@ntu.edu.sq)




Figure 1. Correlation of climatic variability recorded in the lithostratigraphy, δD (orange), $\delta^{18}O$ (blue), of paleolake Karsandi water and ostracod abundance with Indus cultural changes. The calibrated radiocarbon ages (ka BP) are shown in black with red arrows pointing to their respective depths. OSL dates and depth of sand collection for dating are shown in blue. Grey bands denote the nearly pure gypsum deposits indicating periods of relatively lower rainfall and blue band denotes wetter periods. Roman numerals denote lithologic units. The Early phase of the Indus Civilization developed during increased monsoon intensity as indicated by lower GHW isotopes and high shell abundance after $\sim 5.1 \pm 0.2$ ka BP. The Mature Harappan phase and peak in urbanism coincides with the lowest GHW isotopes and highest shell abundance between ~ 5.0 and ~ 4.4 ka BP. Note that the subsequent decline in urbanism and disappearance of Post-urban Harappan sites in this region is coincident with drying conditions suggested by reappearance of massive gypsum with increasing GHW isotopes and complete absence of ostracod and gastropod shells.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit https://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2018