
Dynamics and control of a periodically forced microalgae culture

Martínez Carlos ^{1,2}, Mairet Francis ⁴, Martinon Pierre ³, Bernard Olivier ^{1,2}

¹ Université Côte d'Azur, Inria, INRA, CNRS, UPMC Univ Paris 06, BIOCORE team, France

² LOV-UPMC Sorbonne-CNRS, UMR 7093, Station Zoologique, B.P. 28, 06234 Villefranche-sur-mer, France

³ INRIA, Saclay, École Polytechnique, CMAP 91128 Palaiseau, France

⁴ IFREMER, PBA, Nantes, France

Email addresses : carlos.martinez@inria.fr ; francis.mairet@ifremer.fr ; pierre.martinon@inria.fr ; olivier.bernard@inria.fr

Abstract :

Microalgae cultivation with wastewater is a promising way of reducing the energetic needs for wastewater treatment and the costs of biofuel production. However, the very turbid medium is not favorable for the development of microalgae. Indeed, light, the key element for photosynthesis, rapidly vanishes along depth due to absorption and scattering. Therefore it is crucial to understand the effects of the depth on turbid cultures. In this work, we study theoretically the long-term behavior of a continuous culture of microalgae exposed to a periodic source of light. By allowing periodic variations of the depth and the hydraulic retention time, we show that the microalgae population is forced to a periodic regime. Finally, we address numerically the problem of determining the optimal variations of the depth and the hydraulic retention time for maximizing the productivity of the culture in the periodic regime.

Keywords : Dynamics control, Industrial biotechnology, Wastewater, Biomass productivity, Microalgae

Microalgae grown as a by-product of wastewater treatment in high rate algal ponds (HRAPs) represents a great opportunity for reducing costs of biofuel production (Park et al. (2011); Christenson and Sims (2011)). Moreover, HRAPs show a great potential for nutrient removal in wastewater treatment (Abdel-Raouf et al. (2012)). One problem with the cultivation of algae in wastewater systems is that light, one of the main factors affecting microalgae growth, rapidly becomes limiting after its decrease due to absorption and scattering by algal cells and the high content of particulate matter and colored substances (Morel and Bricaud (1981); Borowitzka (1998)). We refer to the light extinction due to all non-microalgae components as background turbidity. In the standard analyses of microalgae cultures, background turbidity is neglected. However, Martínez et al. (2018) showed theoretically that background turbidity results in a reduction of the productivity when increasing the depth of the system. Indeed, according to Larsdotter (2006), depths of the HRAP between 15 and

The objective of this paper is to give some insights in the optimization of biomass productivity by playing with the depth of the culture and considering a natural light source. In our approach, we combine the modelling approaches of Droop (1968) and Huisman et al. (2002) to build a model accounting for light and substrate limitation. Our model is periodically forced by the incident light with a period of one day. We argue that by providing periodic input and output flows to the system, any solution of the model reaches a 1-day periodic regime i.e. approaches asymptotically a periodic solution. Thus, we study the productivity of the system in a periodic regime by controlling the input and output flow rates, or equivalently, the depth of the system and the hydraulic retention time.

This article is organized as follows. In section 2, we introduce the model of a microalgae culture given by a system of nonautonomous differential equations. In section 3, we study the dynamics of the model. The main result of this section (Theorem 3) gives sufficient conditions to ensure that any solution approaches asymptotically to a periodic solution characterized by the presence of microalgae (i.e. different from the washout). This result follows from reducing the model to a cooperative system and following some ideas of Smith (1997). In section 4, we state an optimal control problem for maximizing the microalgae productivity per day in the periodic regime. Periodic optimal

* This work was supported by the CONICYT doctoral grant (Carlos Martínez), and by the Phycover (ANR-14-CE04-0011) and IPL-Algae in silico projects.

¹ INRIA Sophia Antipolis, 2004, route des Lucioles BP 93, 06902 Sophia Antipolis Cedex, France (e-mail {carlos.martinez,olivier.bernard}@inria.fr)

² pierre.martinon@inria.fr

³ francis.mairet@ifremer.fr

problems have already been studied by Groganard et al. (2014); Bayen et al. (2015). Here, we present some numerical solutions obtained with the software BOCOP. Finally, we illustrate the asymptotic behavior of the model when repeating each day the 1-day periodic optimal control.

2. MODEL DESCRIPTION

Let us consider a perfectly mixed microalgae culture of depth L illuminated from above with an incident light I_{in} . We assume that light is attenuated exponentially according to the Lambert-Beer law i.e. at a distance $z \in [0, L]$ from the illuminated surface, the corresponding light intensity $I(x, z)$ satisfies

$$I(x, z) = I_{in}e^{-(kx+K_{bg})z}, \quad (1)$$

with x the microalgae concentration, $k > 0$ the specific light attenuation coefficient of microalgae, and $K_{bg} \geq 0$ the background turbidity. We assume that microalgae growth is limited by light and a substrate s . Based on the model of Droop (1968), we assume that the specific growth rate μ depends on an internal cell quota q . The latter corresponds to an internal pool of nutrient per unit of biomass. The light intensity effects are described by a Monod type function. Thus, the specific growth rate is taken to be

$$\mu(I, q) = \mu_{max} \left(1 - \frac{q_0}{q}\right) \frac{I}{K_I + I} - r, \quad (2)$$

where $r > 0$ is the respiration rate, $q_0 > 0$ represents the value of q at which growth ceases, $K_I > 0$ is a half-saturation constant, and $\mu_{max} > 0$ is the maximal specific growth rate. Following Huisman et al. (2002), we compute the average growth rate in the reactor, denoted $\bar{\mu}$, by integrating the local growth rates over all the reactor

$$\bar{\mu}(\cdot) := \frac{1}{L} \int_0^L \mu(I(x, z), q) dz. \quad (3)$$

A simple computation leads to

$$\bar{\mu}(\cdot) := \frac{\mu_{max}}{(kx + K_{bg})L} \ln \left(\frac{K_I + I_{in}}{K_I + I_{out}(x)} \right) \left(1 - \frac{q_0}{q}\right) - r, \quad (4)$$

with $I_{out}(x) = I(x, L)$ the light intensity at the bottom of the culture.

Cell quota q decreases with cell growth and increases with nutrient uptake. The nutrient uptake rate is

$$\rho(q, s) = \begin{cases} \rho_{max} \frac{s}{K_s + s} d(q) & \text{if } q \leq q_L, \\ 0 & \text{if } q > q_L, \end{cases} \quad (5)$$

where ρ_{max} is the maximal uptake rate of nitrogen, q_L is the hypothetical maximal quota, and K_s is a half-saturation constant. $d(q) \in [0, 1]$ is a down regulation term. While $d(q)$ is usually taken as a linear function (Bougaran et al. (2010); Morel (1987)), here we take $d(q)$ as

$$d(q) = \left(\frac{q_L - q}{q_L - q_0} \right)^2 \left(\frac{q_L - q}{q_L - q_0} + 3 \frac{q - q_0}{q_L - q_0} \right) \quad (6)$$

We note that d is the unique cubic function satisfying

$$d(q_0) = 1, d(q_L) = 0, \text{ and } d'(q_0) = d'(q_L) = 0. \quad (7)$$

The volume (V) of the culture varies with the supply flow rate (Q_{in}) and the withdrawal rate (Q_{out}) according to

$$\dot{V} = Q_{in} - Q_{out}. \quad (8)$$

The transversal area (A) of the culture is constant in time and along the depth of the culture. After dividing both sides in Eq.(8) by A , we obtain the following equation for the depth of the culture

$$\dot{L} = F_{in} - F_{out}, \quad (9)$$

with $F_{in} := Q_{in}/A$ and $F_{out} := Q_{out}/A$ linear flows. Mass balances in the culture give the following equations (see Chapter 2 in Dochain (2013) and Bougaran et al. (2010))

$$\begin{aligned} \dot{x} &= [\bar{\mu}(t, x, q, L) - \frac{F_{in}}{L}]x, \\ \dot{q} &= \rho(q, s) - \bar{\mu}(t, x, q, L)q, \\ \dot{s} &= \frac{F_{in}}{L}(s_{in} - s) - \rho(q, s)x, \end{aligned} \quad (10)$$

where s_{in} is the (constant) substrate influent concentration.

3. PERIODIC SYSTEM AND ASYMPTOTIC BEHAVIOR.

We assume that the incident light is variable in time according to

$$I_{in}(t) = I_{max} \max\{0, \sin(2\pi t/\omega)\}^2, \quad (11)$$

with $\omega > 0$ the length of a day. Thus, by assuming that F_{in} and F_{out} vary periodically with a period ω , the system (9)-(10) becomes an ω -periodic system.

We use the following notation for the time-average of an ω -periodic function ⁴

$$\langle f \rangle := \frac{1}{\omega} \int_0^\omega f(t) dt. \quad (12)$$

By integrating Eq.(9) we obtain that

$$L(t) = L(0) + \int_0^t [F_{in}(s) - F_{out}(s)] ds. \quad (13)$$

We assume that

$$\langle F_{in}(t) - F_{out}(t) \rangle = 0.$$

Then L is an ω -periodic function. From now on, we assume that:

$$\text{there is } Q > 0 \text{ such that } \langle \bar{\mu}(t, 0, Q) \rangle > 0. \quad (14)$$

From a biological point of view, (14) states that there is a quota such that a very small population can grow.

⁴ A function f is ω -periodic if $f(t + \omega) = f(t)$ for all t

We define the total amount of limiting element both in the substrate and in the biomass by means of

$$S = s + xq. \quad (15)$$

A simple calculation shows that S satisfies the differential equation

$$\dot{S} = D(t)(s_{in} - S), \quad (16)$$

with $D(t) : \frac{F_{in}(t)}{L(t)}$ known as the dilution rate. We can rapidly verify that $S(t) \rightarrow s_{in}$ as $t \rightarrow \infty$ for any value of $S(0)$. Thus, the solutions of Eq.(10) approach the surface

$$s_{in} = s + xq.$$

We take advantage of this by dropping the equation for s and replacing s in Eq.(10) by $s = s_{in} - xq$ in the second equation. This results in the two-dimensional system

$$\begin{aligned} \dot{x} &= [\bar{\mu}(t, x, q) - D(t)]x, \\ \dot{q} &= \rho(q, s_{in} - xq) - \bar{\mu}(t, x, q)q. \end{aligned} \quad (17)$$

In the following we study the asymptotic behavior of Eq.(17). Care must be exercised in extrapolating results obtained from the analysis of Eq.(17) to analogous results for Eq.(9)-(10). A discussion about this can be found in Smith (1997).

As s must be non-negative, initial conditions for Eq. (17) must satisfy $x(0)q(0) \leq s_{in}$. We note that if $x(0) = 0$ (resp. $x(0) > 0$), then $x(t) = 0$ (resp. $x(t) > 0$) for all $t \geq 0$ (see Zanolin (1992) to find rigorous arguments). We also note that $q(0) \geq q_0$ implies $q(t) \geq q_0$ for all $t \geq 0$. Therefore we are only interested in solutions starting on the set

$$P := \{(x, q) ; x > 0, q \geq q_0, \text{ and } xq \leq s_{in}\}. \quad (18)$$

A solution (x, q) of Eq.(17) will be called an ω -periodic solution provided each component is ω -periodic. There is a trivial periodic solution corresponding to the absence of microalgae in the chemostat. Putting $x = 0$ in the second equation of (17) results in

$$\dot{q} = \rho(q, s_{in}) - \bar{\mu}(t, 0, q)q. \quad (19)$$

The following proposition guarantees the existence and uniqueness of the washout.

Proposition 1. Eq. (19) has a unique periodic solution q^* . Moreover, if $g : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a continuous function such that $g(t) \rightarrow 0$ as $t \rightarrow 0$, then any solution to $\dot{q} = \rho(q, s_{in} - qg(t)) - \bar{\mu}(t, g(t), q)q$ starting on P approaches asymptotically to q^* .

Proof. Let $h : \mathbb{R}_+^2 \times [q_0, \infty) \rightarrow \mathbb{R}$ be the function defined by $h(t, x, q) = \frac{\rho(q, s_{in} - xq)}{q} - \bar{\mu}(t, x, q)$. Now, let $\hat{h} : \mathbb{R}_+^3 \rightarrow \mathbb{R}$ be the continuous extension of h on $\mathbb{R}_+^2 \times [q_0, \infty)$ defined as $\hat{h}(t, x, q) = \rho_{max} \frac{s_{in} - xq}{K_n + s_{in} - xq} + r$ for $q \in [0, q_0]$. We can verify that $\int_0^\omega \hat{h}(t, 0, 0)dt > 0$. From Eq.(14), there exists $Q > 0$ such that $\int_0^\omega \hat{h}(t, 0, Q)dt < 0$. Since $q \mapsto \hat{h}(t, 0, q)$ is strictly decreasing, we can apply Proposition 5 in Zanolin (1996) and conclude that the Kolmogorov periodic equation $dq/dt = qh(t, 0, q)$ (*) admits a unique positive ω -periodic solution q^* . Since $[q_0, \infty)$ is positively invariant with respect to (*), we

conclude that $q_*(t) \geq q_0$. Thus, q^* is the unique ω -periodic solution of (19). We note that $|h(t, g(t), q) - h(t, 0, q)| \rightarrow 0$ uniformly for t . It can be shown that the solutions of the non-autonomous Kolmogorov equation $dq/dt = qh(t, g(t), q)$ (***) are uniformly bounded. Using the Theory of Asymptotically Periodic Semiflows (Theorems 2.4 and 3.1 in Zhao (1996)) and the fact that $[q_0, \infty)$ is positively invariant with respect to (**), we conclude that any solution of (***) is asymptotic to q^* . \square

Thus, the trivial periodic solution of (17) is given by $(0, q^*)$. Now we show that any solution of (17) is asymptotic to an ω -periodic solution.

Proposition 2. Any solution to (17) starting on P approaches to an ω -periodic solution.

Proof. Let (\bar{x}, \bar{q}) be a solution of (17) with $(\bar{x}(0), \bar{q}(0)) \in P$. The change of variables $x_s := qx$ leads the system (17) to

$$\begin{aligned} \dot{x} &= [\bar{\mu}(t, x, x_s/x) - D(t)]x \\ \dot{x}_s &= \rho(x_s/x, s_{in} - x_s)x - D(t)x_s. \end{aligned} \quad (20)$$

Eq. (20) is a cooperative system i.e. $\frac{\partial \dot{x}}{\partial x_s}, \frac{\partial \dot{x}_s}{\partial x} \geq 0$. Since $\bar{x}_s = \bar{x}\bar{q}$ satisfies the differential equation

$$\frac{dx_s}{dt} = x_s \left(\frac{\rho(q, s_{in} - x_s)}{q} - D \right), \quad (21)$$

we have that $\bar{x}_s(t) = \bar{x}(t)\bar{q}(t) \leq s_{in}$ for all $t \geq 0$. Thus, $\bar{x}(t) \leq s_{in}/q_0$ for all $t \geq 0$. Consequently, (\bar{x}, \bar{x}_s) is bounded. Following the proof of Theorem 4.2 in Chapter 7 in the Book (Smith and Waltman, 1995), we have that the sequences $x_n := \bar{x}(n\omega)$ and $x_{sn} = \bar{x}_s(n\omega)$ are convergent. Let $l := \lim_{n \rightarrow \infty} x_n$ and $l_s := \lim_{n \rightarrow \infty} x_{sn}$. If $l > 0$, then $l_s > 0$ and $q_n := q(n\omega) \rightarrow l/l_s$. Then, (\bar{x}, \bar{q}) approaches a periodic solution. If $l = 0$, then it can be shown that $\bar{x}(t) \rightarrow 0$ as $t \rightarrow \infty$. Applying Proposition 1, we conclude that \bar{q} approaches asymptotically to q^* , and therefore (\bar{x}, \bar{q}) approaches asymptotically to the washout. \square

We state now the main result of this section. We give conditions to avoid solutions of (17) to approach to the washout.

Theorem 3. If $\langle \bar{\mu}(t, 0, q^*(t)) - D(t) \rangle$ is greater than 0 and D is continuous, then any solution of Eq. (17) starting in P is asymptotic to an ω -periodic solution of Eq.(17) with positive x -component.

Proof. We use the Floquet theory (see Chapter 7 in the book of Sideris (2013)) to determine the stability of the washout. The characteristic multipliers of the washout $(0, q^*)$ are given by $\rho_1 = \exp(\omega \langle a(t) \rangle)$ and $\rho_2 = \exp(-\omega \langle c(t) \rangle)$ with

$$a(t) = \bar{\mu}(t, 0, q^*(t)) - D(t), \quad (22)$$

and

$$\begin{aligned} c(t) &= -\frac{\partial \rho}{\partial q}(q^*(t), s_{in}) + \bar{\mu}(t, 0, q^*(t)) \\ &+ q^*(t) + \frac{\partial \bar{\mu}}{\partial q}(t, 0, q^*(t)) > 0. \end{aligned} \quad (23)$$

Thus, $\rho_1 > 1$ and $\rho_2 < 1$, and the washout is not stable. This implies that not all the solutions of Eq.(17) starting on P are asymptotic to the washout. From Proposition

2, any solution converges to an ω -periodic solution, then there exists an ω -periodic solution different from the washout that attracts some solutions. Following the same arguments as in the proof of Proposition 1.3 in Smith (1997), we can conclude that this is true for any solution starting on P . \square

Note that Theorem 3 does not say anything about the number of periodic solutions with positive x -component.

4. OPTIMAL CONTROL PROBLEM

We define the *areal biomass harvest* on the interval of time $[t_i, t_f]$ as

$$H = \int_{t_i}^{t_f} F_{out}(t)x(t)dt. \quad (24)$$

H represents the quantity of biomass that is harvested on the interval of time $[t_i, t_f]$ per unit of area.

Here we consider the case where the chemostat is operated in long term, with a periodic daily biomass production from the reactor outlet. The problem that we consider is the maximization of the biomass production over a single day. The control variables are F_{in} (u_1) and F_{out} (u_2). The problem can be stated as

$$\begin{aligned} \max H(u_1, u_2) &= \int_0^\omega u_2(t)x(t)dt, \\ \text{s.t. } \dot{x} &= [\bar{\mu}(t, x, q, L) - u_1/L]x, \\ \dot{q} &= \rho(q, s_{in} - xq) - \bar{\mu}(t, x, q, L)x, \\ \dot{L} &= u_1 - u_2 \\ x(0) &= x(\omega) > 0, q(0) = q(\omega), L(0) = L(\omega) \\ u_1(t) &\in [0, F_{in,max}], u_2(t) \in [0, F_{out,max}] \end{aligned} \quad (25)$$

In the formulation of our problem we use Eq. (17) together with Eq. (9).

4.1 Numerical results

We solve numerically the problem (25), with a direct method implemented in the software BOCOP (Bonnans et al. (2011)) (version 2.10). The problem is discretized by a two-stage Gauss-Legendre method of order 4 with 200 time steps. We consider a constant initialization, and the tolerance for IPOPT NLP solver is set at 10^{-12} .

We consider two microalgae cultures; *culture 0* without background turbidity i.e. $K_{bg} = 0$, and *culture 1* with a background turbidity $K_{bg} = 10 m^{-1}$. For both cultures the rest of parameters are given in Table 1. Huisman et al. (2002) reported a background turbidity of $7.2 m^{-1}$ for a nutrient-rich medium used in a laboratory culture.

Figures 1 and 2 show the results. In both cultures, algae concentration increases during half or almost all the day (first half of the period), then decreases at night (see Figure 1A). We note that in culture 1 the microalgae concentration is higher, and consequently the nutrient in the medium is lower (see Figure 1C). However, the daily

microalgae productivity is $1.7 gC m^{-2}$ in contrast to the culture 0 that reaches a daily microalgae productivity of $4.0 gC m^{-2}$ (i.e. $8.0 gDW m^{-2}$).

The control structures are Bang-Singular (u_1) and Bang-Bang (u_2) for both cultures (see Figure 2A). These controls result in a different depth for each culture. Figure 2B shows that there is a difference of almost 40 cm during all the period of 24 hours with a lower depth for the culture that accounts for background turbidity.

Table 1. Parameters

| Parameter | Value | Unit | Reference |
|---------------|---------|-------------------------|------------------------|
| μ_{max} | 1.72 | d^{-1} | Huisman et al. (2002) |
| K_I | 107 | $\mu mol m^{-2} s^{-1}$ | Huisman et al. (2002) |
| k | 0.2 | $m^2 gC^{-1}$ | Béchet et al. (2013) |
| ρ_{max} | 11.2 | $gN gC^{-1} d^{-1}$ | Bougaran et al. (2010) |
| q_L | 0.28 | $gN gC^{-1}$ | Bougaran et al. (2010) |
| q_0 | 0.07 | $gN gC^{-1}$ | Bougaran et al. (2010) |
| K_s | 0.0007 | $gN m^{-3}$ | Bougaran et al. (2010) |
| r | 0.1 | d^{-1} | - |
| s_{in} | 10 | $gN m^{-3}$ | - |
| I_{max} | 2000 | $\mu mol m^{-2} s^{-1}$ | - |
| K_{bg} | 0 or 10 | m^{-1} | - |
| $F_{in,max}$ | 1000 | $L m^{-2} d^{-1}$ | - |
| $F_{out,max}$ | 1000 | $L m^{-2} d^{-1}$ | - |

Consider the controls (u_1, u_2) (see Fig. 2A) obtained from the optimization of one periodic day-night cycle. We apply the respective control to each culture (i.e. to solve Eq.(9)-(17)) repeatedly for a long interval of time. The initial depth for each culture is equal to the initial depth of the respective optimal solution (see Figure 2B). The initial microalgae concentration and the initial quota are $5 gC/m^3$ and $0.28 gN/gC$ respectively for both cultures. Figures 3 and 4 show that the microalgae concentration approaches the optimal periodic microalgae concentration (Fig. 1A) found previously. For the culture 1 the microalgae concentration approaches more slowly to the periodic solution. Indeed, it needs 50 days in contrast to 30 days for the culture 0.

5. CONCLUSIONS AND FUTURE WORK

A first series of numerical simulations using a direct method illustrates the impact of the background turbidity on the choice of the depth of the culture for maximizing the microalgae productivity. We have shown that a background turbidity of $10 m^{-1}$ can result in a reduction of 40 cm of the optimal depth and a productivity loss of 55%. Thus, the background turbidity should not be neglected. The choice of the depth is not trivial and should be adapted to the overall turbidity, in particular when both light and nutrient limitations can occur. This choice should also include thermal considerations, since low depth induces lower thermal inertia. Processes with low depth are likely to overwarm very rapidly in hot seasons (DeLuca et al. (2017)) and a trade-off between light and temperature performances should be found.

We also studied theoretically the long-term behavior of the microalgae population when the system is operated periodically in agreement with the period of the light




Fig. 1. Optimal solutions for problem (25) with two different background turbidities. A. Microalgae concentration. B. Cell quota. C. Substrate concentration in the medium.

source. We established sufficient conditions to ensure that any solution of the model, independently of the initial conditions, is attracted to an oscillatory solution with a positive microalgae concentration.

Photoinhibition (i.e. reduction of the photosynthetic rate due to high light intensities) should be included in a future work. Low depths of the culture could enhance the photoinhibitory process resulting in a lower productivity, and for low initial microalgae concentrations the population could go extinct (see Gerla et al. (2011); Martínez et al. (2017)).

In this work, we emphasized the productivity of microalgae using wastewater streams without paying attention to the aspect of wastewater treatment. In a future work, other optimization problems will be studied. For example, the




Fig. 2. Optimal solutions for problem (25) with two different background turbidities. A. Control variables. B. Depth of the culture.




Fig. 3. Comparison of the optimal 1-day periodic microalgae concentration and the evolution of the microalgae with a low initial concentration when the periodic 1-day optimal periodic control is applied. Culture without background turbidity.




Fig. 4. Comparison of the optimal periodic microalgae concentration and the evolution of the microalgae with a low initial concentration when the periodic optimal control is applied all the time. With background turbidity.

maximization of the removal of nitrogen with restrictions over the volume of wastewater to be treated.

REFERENCES

- Abdel-Raouf, N., Al-Homaidan, A., and Ibraheem, I. (2012). Microalgae and wastewater treatment. *Saudi journal of biological sciences*, 19(3), 257–275.
- Bayen, T., Mairet, F., Martinon, P., and Sebbah, M. (2015). Analysis of a periodic optimal control problem connected to microalgae anaerobic digestion. *Optimal Control Applications and Methods*, 36(6), 750–773.
- Béchet, Q., Shilton, A., and Guieysse, B. (2013). Modeling the effects of light and temperature on algae growth: state of the art and critical assessment for productivity prediction during outdoor cultivation. *Biotechnology advances*, 31(8), 1648–1663.
- Bonnans, J., Grelard, V., and Martinon, P. (2011). Bocop, the optimal control solver, open source toolbox for optimal control problems. URL <http://bocop.org>.
- Borowitzka, M.A. (1998). Limits to growth. In *Wastewater treatment with algae*, 203–226. Springer.
- Bougaran, G., Bernard, O., and Sciandra, A. (2010). Modeling continuous cultures of microalgae colimited by nitrogen and phosphorus. *Journal of theoretical biology*, 265(3), 443–454.
- Christenson, L. and Sims, R. (2011). Production and harvesting of microalgae for wastewater treatment, biofuels, and bioproducts. *Biotechnology advances*, 29(6), 686–702.
- De-Luca, R., Bezzo, F., Béchet, Q., and Bernard, O. (2017). Exploiting meteorological forecasts for the optimal operation of algal ponds. *Journal of Process Control*, 55, 55–65.
- Dochain, D. (2013). *Automatic control of bioprocesses*. John Wiley & Sons.
- Droop, M.R. (1968). Vitamin b 12 and marine ecology. iv. the kinetics of uptake, growth and inhibition in *monochrysis lutheri*. *Journal of the Marine Biological Association of the United Kingdom*, 48(3), 689–733.
- Gerla, D.J., Mooij, W.M., and Huisman, J. (2011). Photoinhibition and the assembly of light-limited phytoplankton communities. *Oikos*, 120(3), 359–368.
- Grognard, F., Akhmetzhanov, A.R., and Bernard, O. (2014). Optimal strategies for biomass productivity maximization in a photobioreactor using natural light. *Automatica*, 50(2), 359–368.
- Huisman, J., Matthijs, H.C., Visser, P.M., Balke, H., Sigon, C.A., Passarge, J., Weissing, F.J., and Mur, L.R. (2002). Principles of the light-limited chemostat: theory and ecological applications. *Antonie van Leeuwenhoek*, 81(1-4), 117–133.
- Larsdotter, K. (2006). Wastewater treatment with microalgae—a literature review. *Vatten*, 62(1), 31.
- Martínez, C., Bernard, O., and Mairet, F. (2017). Maximizing microalgae productivity by shading outdoor cultures. *IFAC-PapersOnLine*, 50(1), 8734–8739.
- Martínez, C., Mairet, F., and Bernard, O. (2018). Theory of turbid microalgae cultures. *Journal of theoretical biology*, 456, 190–200.
- Morel, A. and Bricaud, A. (1981). Theoretical results concerning light absorption in a discrete medium, and application to specific absorption of phytoplankton. *Deep-Sea Res*, 28(11), 1375–1393.
- Morel, F.M. (1987). Kinetics of nutrient uptake and growth in phytoplankton 1. *Journal of phycology*, 23(1), 137–150.
- Park, J., Craggs, R., and Shilton, A. (2011). Wastewater treatment high rate algal ponds for biofuel production. *Bioresour technology*, 102(1), 35–42.
- Sideris, T.C. (2013). Ordinary differential equations and dynamical systems.
- Smith, H.L. and Waltman, P. (1995). *The theory of the chemostat: dynamics of microbial competition*, volume 13. Cambridge university press.
- Smith, H. (1997). The periodically forced droop model for phytoplankton growth in a chemostat. *Journal of Mathematical Biology*, 35(5), 545–556.
- Zanolin, F. (1996). Continuation theorems for the periodic problem via the translation operator. *Rend. Sem. Mat. Univ. Politec. Torino*, 54, 1–23.
- Zanolin, F. (1992). Permanence and positive periodic solutions for kolmogorov competing species systems. *Results in Mathematics*, 21(1-2), 224–250.
- Zhao, X.q. (1996). Asymptotic behavior for asymptotically periodic semiflows with applications. In *MR 97i: 58150*. Citeseer.