

Supplementary Data "Seasonal and plant-part isotopic and biochemical variation in *Posidonia oceanica*"

P. CRESSON, C.F. BOUDOURESQUE, S. RUITTON, L. CASALOT, M. VERLAQUE and M. HARMELIN-VIVIEN

Table S1: Results of ANOVA mean comparison tests for seasonal variation of biochemical and isotopic parameters in each plant part and leaf epibionts (***) p < 0.001, ** p < 0.01, * p < 0.05, ns p > 0.05). Results of post-hoc tests are provided, with letters standing for the season (Sp: Spring, Su: Summer, A: Autumn, W: Winter). For leaves and rhizome, protein is calculated as %N × 4.28.

Plant part	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	%C	%N / Proteins	C/N	IC	SC	Lipids
Rhizomes	F = 16.6 **	F = 10.9 **	F = 2.2 ns	F = 39.6 ***	F = 207.8 ***	F = 0.7 ns	F = 4.2 *	F = 15.1 **
	A < Sp < W = Su	Sp < W = Su < A		W < A = Su < Sp	Sp < Su < A < W		Sp = W < Su = A	W = A < Sp = Su
Juveniles	F = 298.5 ***	F = 183.6 ***	F = 2.3 ns	F = 8.6 **	F = 209.2 ***	F = 1.4 ns	F = 5.6 *	F = 0.1 ns
	W < A < Su = Sp	W < A = Sp < Su		Sp = A < W = Su	W < Su < Sp < A		Sp < W = Su < A	
Intermediate	F = 42.0 ***	F = 108.1 ***	F = 2.3 ns	F = 98.1 ***	F = 461.1 ***	F = 6.3 *	F = 5.8 *	F = 38.0 ***
	W < Sp < Su = A	Sp < W < A < Su		A < Sp = W < Su	Sp = W < Su < A	Sp < W = Su = A	Su = Sp = W < A	W < A = Sp < Su
Adult	F = 381.0 ***	F = 146.1 ***	F = 3.8 ns	F = 161.4 ***	F = 5051.7 ***	F = 17.1 ***	F = 17.3 ***	F = 8.7 **
	Su < Sp < A < W	Sp = W < A < Su		W < A < Su < Sp	Sp = E < A < W	W = Su = Sp < A	Su = W = Sp < A	W = A = Sp < Su
Senescent	F = 56.2 ***	F = 79.1 ***	F = 41.5 ***	F = 633.4 ***	F = 1136.8 ***	F = 4.9 *	F = 7.2 *	F = 47.8 ***
	Su < Sp < A < W	Sp = Su < A = W	A < Sp < W < Su	A < W < Sp < Su	Su < Sp < A < W	Sp < Su = W = A	W < Sp = A = Su	A = W < Sp < Su
Drifting	F = 12.9 **	F = 8.2 **	F = 0.3 ns	F = 3.6 ns	F = 141.7 ***	F = 34.6 ***	F = 6.1 *	F = 15.7 **
	Su < Sp = W	Su < Sp = W			Sp = Su < W	Sp < Su < W	Su = W < Sp	W < Su < Sp
Leaf epibionts	F = 2.4 ns	F = 336.8 ***	F = 0.9 ns	F = 1.1 ns	F = 13.2 ***	F = 114.3 ***	F = 5.1 *	F = 20.6 ***
		Sp < E < A = W			E = Sp < W < A	E = W = A < Sp	E = W < Sp = A	A = W < E < Sp

Medit. Mar. Sci., suppl.data, 20/2, 2019

Table S2: Ash, lipid, nitrogen, protein and carbohydrate compositions of *Posidonia oceanica* leaves and rhizomes. Values in percentage of the dry mass.

Ash	Lipid	Nitrogen	Protein	Soluble carbohydrate	Insoluble carbohydrate	Remarks	Locality or country	Reference
Leaves								
-	-	2.18	13.6 ^k	-	-	?	Provence? France	Moliner and Pellegrini, 1966
23.9 ^f	1.2 ^g	-	14.3	-	-	February-April, 4 m depth	Western Italy	Gallarati Scotti, 1969
31.0 ^f	0.7 ^g	-	10.5	-	-	June-July, 4 m depth	Western Italy	Gallarati Scotti, 1969
34.5 ^f	0.5 ^g	-	9.9	-	-	August-September, 4 m depth	Western Italy	Gallarati Scotti, 1969
-	-	1.65-2.18	-	-	-	October, shallow, short (young?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	1.33-1.36	-	-	-	October, shallow, long (old?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	2.54-3.76	-	-	-	December-Febr., shallow, short (young?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	2.20-2.84	-	-	-	December-Febr., shallow, long (old?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	2.32-2.48	-	-	-	May-June, shallow, short (young?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	1.69-2.32	-	-	-	May-June, shallow, long (old?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	2.12-2.32	-	-	-	August, shallow, short (young?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	1.37-2.07	-	-	-	August, shallow, long (old?) leaves	Le Brusuc, France	Pellegrini, 1971
-	-	1.55-1.95	-	-	-	January-April, 10 m depth	Calvi, Corsica	Bay, 1978
-	-	1.08-1.55	-	-	-	August-November, 10 m depth, young leaves	Calvi, Corsica	Bay, 1978
-	-	0.71-0.95	-	-	-	August-November, 10 m depth, old leaves	Calvi, Corsica	Bay, 1978
-	-	1.89-2.51	-	-	-	January, 20-38 m depth	Calvi, Corsica	Bay, 1978
5.5	-	1.31	6.6	-	-	June? (°), 10 m depth	Port-Cros, France	Augier and Santimone, 1979
4.0	-	1.48	7.0	-	-	June? (°), 20 m depth	Port-Cros, France	Augier and Santimone, 1979
5.1	-	1.30	5.6	-	-	June? (°), 30 m depth	Port-Cros, France	Augier and Santimone, 1979
-	-	0.62	-	-	-	July, 3 m depth, base	Mallorca, Spain	Mouriño <i>et al.</i> , 1981
-	-	1.71-1.85	-	-	-	July, 3 m depth, 5-15 cm above the base	Mallorca, Spain	Mouriño <i>et al.</i> , 1981
-	-	-	-	10-11	-	April, 5 m depth, leaf base	Ischia, Italy	Velimirov <i>et al.</i> , 1981
-	-	-	-	1.9-2.9	-	April, 5 m depth, leaf limb	Ischia, Italy	Velimirov <i>et al.</i> , 1981
-	-	-	-	8-13	-	April, 15 m depth, leaf base	Ischia, Italy	Velimirov <i>et al.</i> , 1981
-	-	-	-	2.5-6.3	-	April, 15 m depth, leaf limb	Ischia, Italy	Velimirov <i>et al.</i> , 1981
-	-	-	-	9-17	-	April, 30 m depth, leaf base	Ischia, Italy	Velimirov <i>et al.</i> , 1981
-	-	-	-	2.8-7.0	-	April, 30 m depth, leaf limb	Ischia, Italy	Velimirov <i>et al.</i> , 1981
5.5	-	1.31	6.6	-	-	June, 10 m depth	Port-Cros, France	Augier <i>et al.</i> , 1982
5.9	-	1.30	5.6	-	-	June, 30 m depth	Port-Cros, France	Augier <i>et al.</i> , 1982
-	-	956:39:1 ¹	-	-	-	36 m depth	Corsica	Atkinson and Smith, 1983
31 ¹	-	-	-	-	-	November, 9-10 m depth	Côte Bleue, France	Maubert and Le Goff, 1984

Medit. Mar. Sci., suppl.data, 20/2, 2019

Ash	Lipid	Nitrogen	Protein	Soluble carbohydrate	Insoluble carbohydrate	Remarks	Locality or country	Reference
23 ^f	-	-	-	-	-	July, 10-12 m depth	El Dabaa, Egypt	Maubert and Le Goff, 1984
18-21	2.3-2.4	-	4.1-4.3	15-19	55-60	July, 2 m depth	Port Cros, France	Lawrence <i>et al.</i> , 1989
19-22	1.9-3.2	-	3.7-4.3	13-15	58-60	July 10 m depth	Port Cros, France	Lawrence <i>et al.</i> , 1989
12	3.9 ^e	1.14 ^e	-	-	-	Spring, 10 m depth	Corsica	Gobert <i>et al.</i> , 1995
12	4.7 ^g	1.02 ^g	-	-	-	Summer, 10 m depth	Corsica	Gobert <i>et al.</i> , 1995
12	4.3 ^h	1.25 ^h	-	-	-	Autumn, 10 m depth	Corsica	Gobert <i>et al.</i> , 1995
12	4.4 ⁱ	2.05 ⁱ	-	-	-	Winter, 10 m depth	Corsica	Gobert <i>et al.</i> , 1995
-	-	1.35	-	-	-	Sept, 10 m fish farm, 22 m depth	Italy	Pérez <i>et al.</i> , 2008
-	-	1.50	-	-	-	Sept, 1 000 m fish farm, 22 m depth	Italy	Pérez <i>et al.</i> , 2008
-	-	1.75	-	-	-	June, 15 m fish farm, 16 m depth	Greece	Pérez <i>et al.</i> , 2008
-	-	1.40	-	-	-	June, 1 200 m fish farm, 16 m depth	Greece	Pérez <i>et al.</i> , 2008
-	-	1.50	-	-	-	Sept, 10 m fish farm, 28 m depth	Spain	Pérez <i>et al.</i> , 2008
-	-	1.40	-	-	-	Sept, 1 000 m fish farm, 28 m depth	Spain	Pérez <i>et al.</i> , 2008
-	-	0.9-1.3 ^a	-	-	-	Sept, 7 m depth, polluted and pristine	Balearic Islands	Terrados and Medina Pons, 2008
-	-	0.7-1.0 ^a	-	-	-	Sept, 17 m depth, polluted and pristine	Balearic Islands	Terrados and Medina Pons, 2008
-	4.05	2.90 ⁱ	60.7 ^j	-	-	Summer, 40 and 68 m (l)	Egypt (Mediterranean coast)	Shams el Din and El-Sherif, 2013

Rhizomes								
Ash	Lipid	Nitrogen	Protein	Soluble carbohydrate	Insoluble carbohydrate	Remarks	Locality or country	Reference
-	-	1.02	-	-	-	October, shallow	Le Brus, France	Pellegrini, 1971
-	-	3.06-3.72	-	-	-	December-February, shallow	Le Brus, France	Pellegrini, 1971
-	-	2.63-2.76	-	-	-	May-June, shallow	Le Brus, France	Pellegrini, 1971
-	-	1.79	-	-	-	August, shallow	Le Brus, France	Pellegrini, 1971
18.5	-	0.51	2.5	-	-	June? (°), 20 m depth	Port-Cros, France	Augier and Santimone, 1979
-	-	1 749:40:1 ⁱ	-	-	-	38 m depth	Corsica	Atkinson and Smith, 1983
52 ^f	-	-	-	-	-	November, 9-10 m depth	Côte Bleue, France	Maubert and Le Goff, 1984
7.8 ^d	-	3.03	-	-	-	August, 15 m depth, 5 last years	Corsica	Calmet <i>et al.</i> , 1986
10.4 ^d	-	1.06-1.11	-	-	-	August, 15 m depth, 10-15 years ago	Corsica	Calmet <i>et al.</i> , 1986
18.9 ^d	-	0.78-0.84	-	-	-	August, 15 m depth, 30-35 years ago	Corsica	Calmet <i>et al.</i> , 1986
5.4	1.3	-	4.5	40	49	July, 2 m	Port Cros, France	Lawrence <i>et al.</i> , 1989
4.8	1.6	-	4.8	47	42	July 10 m	Port Cros, France	Lawrence <i>et al.</i> , 1989
5.9-6.0	-	3.63	-	-	-	March, 10 m depth, last year	Côte Bleue, France	Carlotti <i>et al.</i> , 1992
6.4-8.4	-	0.83-3.03	-	-	-	March, 10 m depth, 2-10 years ago	Côte Bleue, France	Carlotti <i>et al.</i> , 1992

Medit. Mar. Sci., suppl.data, 20/2, 2019

Ash	Lipid	Nitrogen	Protein	Soluble carbohydrate	Insoluble carbohydrate	Remarks	Locality or country	Reference
5.0-5.2	-	1.78	-	-	-	May, 10 m depth, last year	Corsica	Carlotti <i>et al.</i> , 1992
5.3-7.7	-	0.63-1.60	-	-	-	May, 10 m depth, 2-10 years ago	Corsica	Carlotti <i>et al.</i> , 1992
-	-	1.25-1.35	-	-	-	March-July, 10 m depth	Catalonia, Spain	Pérez <i>et al.</i> , 2007
-	-	1.75-1.80	-	-	-	May-July, org. matter enriched, 10 m depth	Catalonia, Spain	Pérez <i>et al.</i> , 2007
-	-	1.55-2.00	-	-	-	May-July, org. matter and N enriched, 10 m depth	Catalonia, Spain	Pérez <i>et al.</i> , 2007
-	-	1.35	-	-	-	Sept, 10 m fish farm, 22 m depth	Italy	Pérez <i>et al.</i> , 2008
-	-	1.25	-	-	-	Sept, 1 000 m fish farm, 22 m depth	Italy	Pérez <i>et al.</i> , 2008
-	-	2.60	-	-	-	June, 15 m fish farm, 16 m depth	Greece	Pérez <i>et al.</i> , 2008
-	-	0.75	-	-	-	June, 1 200 m fish farm, 16 m depth	Greece	Pérez <i>et al.</i> , 2008
-	-	2.30	-	-	-	Sept, 10 m fish farm, 28 m depth	Spain	Pérez <i>et al.</i> , 2008
-	-	1.20	-	-	-	Sept, 1 000 m fish farm, 28 m depth	Spain	Pérez <i>et al.</i> , 2008
-	-	-	-	0.4-0.5 ^b	-	Oct, 7 m depth, within a MPA	French Catalonia	Ferrari <i>et al.</i> , 2008
-	-	-	-	0.5-0.6 ^b	-	Oct, 7 m depth, outside a MPA	French Catalonia	Ferrari <i>et al.</i> , 2008
-	-	-	-	0.5-5.0	4-5 ^c	April, 3-5 m depth, beach replenishment	Andalusia, Spain	González-Correa <i>et al.</i> , 2008

^a Calculated from authors' figure

^b Nonstructural carbohydrate content

^c Starch

^d Calculated from authors' figure

^e Extrapolated from another article of the same authors

^f Ash mass putatively over-evaluated, possibly due to the non-removing of leaf calcareous epibionts

^g Possibly under-evaluated, due to the over-evaluation of ash (see ^f)

ⁱ C:N:P ratio

^j Obviously erroneous values. The authors may have inverted the values of the proteins and total carbohydrates.

^k The authors calculated protein content from total nitrogen, according to an Atwater coefficient of 6.25

References

Atkinson, M.J., Smith, S.V., 1983. C:N:P ratios in benthic marine plants. *Limnology and Oceanography*, 28 (3), 568-574.

Augier, H., Calvert, H., Wollaston, E., Santimone M., 1982. A comparison of the C, H, N, protein and amino acid composition of *Posidonia australis* Hook. F. with that of *Posidonia oceanica* (L.) Delile and several other marine phanerogams. *Aquatic Botany*, 12, 69-80.

Augier, H., Santimone, M., 1979. Contribution à l'étude chimique de la phanérogame marine *Posidonia oceanica* Delile: Composition en cendres, carbone, hydrogène, azote, protéines et acides aminés en milieu exempt de pollution et en fonction de la profondeur dans le Parc national de Port-Cros. *Travaux scientifiques du Parc national de Port-Cros*, 5, 105-123.

Bay D., 1978. Etude *in situ* de la production primaire d'un herbier de posidonies (*Posidonia oceanica* (L.) Delile) de la baie de Calvi -Corse. *Progr. Rep. Stat. océanogr. STARESO*, Univ. Liège, 18, 1-6 + 1-251.

Medit. Mar. Sci., suppl.data, 20/2, 2019

- Calmet, D., Boudouresque, C.F., Meinesz, A., 1988. Memorization of nuclear atmospheric tests by rhizomes and scales of the Mediterranean seagrass *Posidonia oceanica* (Linnaeus) Delile. *Aquatic Botany*, 30, 279-293.
- Carlotti, P., Boudouresque, C.F., Calmet, D., 1992. Mémorisation du cadmium et des radioéléments par les rhizomes et les écailles de *Posidonia oceanica* (Potamogetonaceae). *Travaux scientifiques du Parc naturel Régional et des Réserves Naturelle de Corse*, 36, 1-34.
- Ferrari, B., Raventos, N., Planes, S., 2008. Assessing effects of fishing prohibition on *Posidonia oceanica* seagrass meadows in the Marine Natural Reserve of Cerbère-Banyuls. *Aquatic Botany*, 88, 295-302.
- Gallarati Scotti, G., 1969. Variazioni stagionali del contenuto in principi alimentari ed in alcuni elementi minerali delle foglie di *Posidonia oceanica*. *Revista di Zootecnica Agricoltura Veterinaria*, 7 (4-6), 1-10.
- Gobert, S., Belkhiria, S., Dauby, P., Havelange, S., Soullard, M., Bouquegneau, J.M., 1995. Variations temporelles de la phénologie et de la composition biochimique de la phanérogame marine *Posidonia oceanica* en baie de Calvi. *Bulletin de la Société Royale des Sciences de Liège*, 64 (4-5), 263-284.
- González-Correa, J.M., Fernández Torquemada, Y., Sánchez Lizaso, J.L., 2008. Long-term effect of beach replenishment on natural recovery of shallow *Posidonia oceanica* meadows. *Estuarine, Coastal and Shelf Science*, 76, 834-844.
- Lawrence, J.M., Boudouresque, C.F., Maggiore F., 1989. Proximate constituents, biomass and energy in *Posidonia oceanica* (Potamogetonaceae). *PSZN 1 Marine Ecology*, 10 (3), 263-270.
- Maubert, H., Le Goff, A., 1984. Contribution à l'étude des niveaux de pollution des herbiers à *Posidonia oceanica* (Linné). Commissariat à l'énergie atomique, Institut de Protection et de Sureté Nucléaire, Rapport SRcA 84/28, 1-19.
- Mouriño, C., Fernández, A., Niell, F.X., 1981. Comentarios sobre el uso del índice C:N en el estudio de algas bentónicas. *Oecologia aquatica*, 5, 35-42.
- Pellegrini, M., 1971. Contribution à l'étude biochimique des phanérogames marines. Répartition et évolution de l'azote total chez *Posidonia oceanica* Delile. *Bulletin du Museum d'Histoire Naturelle de Marseille*, 31, 197-203.
- Peréz, M., Garcia, T., Invers, O., Ruiz, J.M., 2008. Physiological responses of the seagrass *Posidonia oceanica* as indicators of fish farm impact. *Marine Pollution Bulletin*, 56, 869-879.
- Pérez, M., Invers, O., Ruiz, J.M., Frederiksen, M.S., Holmer, M., 2007. Physiological responses of the seagrass *Posidonia oceanica* to elevated organic matter content in sediments: an experimental assessment. *Journal of Experimental Biology and Ecology*, 344, 149-160.
- Pirc, H., 1985. Growth dynamics in *Posidonia oceanica* (L.) Delile. I. Seasonal changes of soluble carbohydrates, starch, free amino acids, nitrogen and organic anions in different parts of the plant. *PSZN I: Marine Ecology*, 6 (2), 141-165.
- Shams El Din, N.G., El-Sherif Z.M. 2013. Nutritional value of *Cymodocea nodosa* and *Posidonia oceanica* along the western Egyptian Mediterranean coast. *Egyptian Journal of Aquatic Research*, 39, 153-165.

Terrados, J, Medina Pons, F.J. 2008. Epiphyte load on the seagrass *Posidonia oceanica* (L.) Delile does not indicate anthropogenic nutrient loading in Cabrera Archipelago National Park (Balearic Islands, Western Mediterranean). *Scientia Marina* 72, 503-510.

Velimirov, B., Ott, J.A., Novak, R., 1981. Microorganisms on macrophyte debris: biodegradation and its implication in the food web. *Kieler Meeresforschungen Sonderheft*, 5, 333-344.

Medit. Mar. Sci., suppl.data, 20/2, 2019