

SUPPORTING INFORMATION APPENDIX 1: Data Sources

The next 64 pages comprise a reference list for all literary sources given as references for trait scores and/or comments in the sFDvent raw (marked with an asterisk (*) if not then included in recommended) and/or recommended datasets (Tables S4.3 and S4.2, respectively). These references are not in alphabetical order, as the database is a ‘living’ record, so new references will be added and a new number assigned.

In the recommended dataset (Table S4.2), the references are recorded according to the numbers listed below (and in Table S1.1), to ensure that citations are relatively easy for users to carry through when conducting analyses using subsets of the data, for example. If a score in the recommended dataset is supported by more than one reference, multiple reference identifiers are provided and separated by a semi-colon (;).

The references are not provided as numbers / identifiers in the other versions of the dataset, as information is lost during this processing step (e.g., ‘expert opinion’, or 66, replaces comments made by experts in each reference column regarding additional observations, rationale for certainty scores, etc.), which may prove useful for some users. Other versions of the dataset thus maintain raw reference entries for transparency and as potentially useful metadata. We provide a copy of the recommended dataset without the references as numbers (Table S4.2A), in case it is easier for users to cross-reference between the two sheets to seek additional comments for a given data subset of interest.

1. Aguado, M. T., & Rouse, G. W. (2006). First record of Sphaerodoridae (Phyllodocida : Annelida) from hydrothermal vents. *Zootaxa*, 1383, 1 - 21.
2. Ahyong, S. T. (2009). New Species and New Records of Hydrothermal Vent Shrimps from New Zealand (Caridea: Alvinocarididae, Hippolytidae). *Crustaceana*, 82(7), 775 - 794.
3. *Aktipis, S. W., Boehm, E., & Giribet, G. (2011). Another step towards understanding the slit-limpets (Fissurellidae, Fissurelloidea, Vetigastropoda, Gastropoda): a combined five-gene molecular phylogeny. *Zoologica Scripta*, 40(3), 238 - 259.
4. Alberti, G., & Meyer-Rochow, V. B. (2002). Ultrastructural investigations of testes and spermiogenesis in two species of halacarid mites (Halacaridae, Actinedida, Actinotrichida): Thalassarachna basteri from the Baltic Sea and Halacarellus thomasi from McMurdo Sound (Antarctica). *Arthropod Structure & Development*, 30(4), 315-328. doi:10.1016/S1467-8039(02)00008-7
5. Allen, C. E., Tyler, P. A., & Van Dover, C. L. (2001). Lipid composition of the hydrothermal vent clam *Calyptogena pacifica* (Mollusca: Bivalvia) as a trophic indicator. *Journal of the Marine Biological Association of the United Kingdom*, 81(5), 817 - 821. doi:10.1017/S0025315401004647

6. *Allen, J. A. (1993). A new deep-water hydrothermal species of Nuculana (Bivalvia, Protobranchia) from the Guaymas Basin. *Malacologia*, 35(1), 141 - 151.
7. Anderson, M. E., Somerville, R., & Copley, J. T. (2016). A new species of *Pachycara* Zugmayer, 1911 (Teleostei: Zoarcidae) from deep-sea chemosynthetic environments in the Caribbean Sea. *Zootaxa*, 4066(1), 71 - 77.
8. *Arellano, S. M., & Young, C. M. (2009). Spawning, development, and the duration of larval life in a deep-sea cold-seep mussel. *Biological Bulletin*, 216(2), 149 - 162.
9. Audzijonyte, A., Krylova, E. M., Sahling, H., & Vrijenhoek, R. C. (2012). Molecular taxonomy reveals broad trans-oceanic distributions and high species diversity of deep-sea clams (Bivalvia: Vesicomyidae: Pliocardiinae) in chemosynthetic environments. *Systematics and Biodiversity*, 10(4), 403 - 415. doi:10.1080/14772000.2012.744112
10. Ax, P. (2000). Leptostraca — Eumalacostraca. In P. Ax (Ed.), *Multicellular Animals* (Vol. The Phylogenetic System of the Metazoa. Volume II, pp. 161 - 162). Berlin, Heidelberg: Springer.
11. Baba, K. (1995). A new squat lobster (Decapoda: Anomura: Galatheidae) from an active thermal vent area in the North Fiji Basin, SW Pacific. *Crustacean Research*, 24, 188 - 193.
12. Baba, K. (2005). Deep-sea chirostylid and galatheid Crustaceans (Decapoda: Anomura) from the Indo-Pacific, with a list of species. *Galathea Report*, 20, 1 - 317.
13. Baba, K., & de Saint Laurent, M. (1992). Chirostylid and Galatheid Crustaceans (Decapoda: Anomura) from active thermal vent areas in the southwest Pacific. *Scientia Marina*, 56(4), 321 - 332.
14. Baba, K., & Türkay, M. (1992). *Munida magniatennulata*, a new deep sea decapod crustacean from active thermal vent areas of Valu-Fa-Ridge in the Lau Basin, SW-Pacific (Anomura: Galatheidae). *Senckenbergiana Maritima*, 22, 203 - 210.
15. *Baeza, J. A. (2011). Squat lobsters as symbionts and in chemo-autotrophic environments. In G. C. B. Poore, S. T. Ahyong, & J. Taylor (Eds.), *The Biology of Squat Lobsters* (pp. 249 - 270): CSIRO Publishing: Melbourne and CRC Press: Boca Raton.
16. *Barnard, J. L. (1954a). Amphipoda of the family Ampeliscidae collected in the Eastern Pacific Ocean by the *Velero III* and *Velero IV*. *Allan Hancock Pac. Expeds.*, 18(1), 1 - 137.
17. *Barnard, J. L. (1954b). Marine Amphipoda of Oregon. *Oregon State Monographs*,

Studies in Zoology, 8, 1 - 103.

18. Barnard, J. L. (1960a). The amphipod family Phoxocephalidae in the eastern Pacific Ocean, with analyses of other species and notes for a revision of the family. *Hancock Pacific Expeditions*, 18(3), 175 - 368.
19. *Barnard, J. L. (1960b). New bathyal and sublittoral ampeliscid amphipods from California, with an illustrated key to *Ampelisca*. *Pacific Naturalist*, 1(16), 1 - 36.
20. Barnard, J. L., & Ingram, C. (1990). Lysianassoid Amphipoda (Crustacea) from deep-sea thermal vents. *Smithsonian Contributions to Zoology*, 499, 1 - 80.
21. *Barnard, K. H. (1960). New species of South African marine gastropods. *Journal of Conchology London*, 24, 438 - 442.
22. Barry, J. P., Buck, K. R., Kochevar, R. K., Nelson, D. C., Fujiwara, Y., Goffredi, S. K., & Hashimoto, J. (2002). Methane-based symbiosis in a mussel, *Bathymodiolus platifrons*, from cold seeps in Sagami Bay, Japan. *Invertebrate Biology*, 121(1), 47 - 54.
23. Barry, J. P., Buck, K., Goffredi, S. K., & Hashimoto, J. (2000). Ultrastructure studies of two chemosynthetic invertebrate bacterial symbioses (*Lamellibrachia* sp. and *Acharax* sp.) from the Hatsushima cold seeps in Sagami Bay, Japan. *Jamstec Journal of Deep Sea Research*, 16, 91 -99.
24. Barry, J. P., & Kochevar, R. E. (1999). *Calyptogena diagonalis*, a new vesicomyid bivalve from subduction zone cold seeps in the Eastern North Pacific. *The Veliger*, 42(2), 117 - 123.
25. Barry, J. P., Kochevar, R. E., & Baxter, C. H. (1997). The influence of pore-water chemistry and physiology on the distribution of vesicomyid clams at cold seeps in Monterey Bay: Implications for patterns of chemosynthetic community organization. *Limnology and Oceanography*, 42(2), 318-328. doi:10.4319/lo.1997.42.2.0318
26. *Bartels-Hardege, H. D., & Zeeck, E. (1990). Reproductive behaviour of *Nereis diversicolor* (Annelida: Polychaeta). *Mar Biol*, 106(3), 409 - 412.
27. Bartsch, I. (1994). Halacarid mites (Acari) from hydrothermal deep-sea sites: new records. *Cahiers de Biologie Marine*, 35(4), 479 - 490.
28. Bates, A. E. (2007). Feeding strategy, morphological specialisation and presence of bacterial epibionts in lepetodrilid gastropods from hydrothermal vents. *Marine Ecology Progress Series*, 347, 87 - 99.

29. Bates, A. E., Harmer, T. L., Roeselers, G., & Cavanaugh, C. M. (2011). Phylogenetic Characterization of Episymbiotic Bacteria Hosted by a Hydrothermal Vent Limpet (Lepetodrilidae, Vetigastropoda). *Biological Bulletin*, 220(2), 118 - 127.
30. Bates, A. E., Tunnicliffe, V., & Lee, R. W. (2005). Role of thermal conditions in habitat selection by hydrothermal vent gastropods. *Marine Ecology Progress Series*, 305, 1 - 15.
31. *Bayer, S. R., Mullineaux, L. S., Waller, R. G., & Solow, A. R. (2011). Reproductive traits of pioneer gastropod species colonizing deep-sea hydrothermal vents after an eruption. *Marine Biology*, 158(1), 181 - 192.
32. Beaulieu, S. E., Sayre-McCord, R. T., Mills, S. W., Pradillon, F., & Watanabe, H. (2015). Swimming speeds of polychaete larvae collected near deep-sea hydrothermal vents. *Marine Ecology*, 36(Suppl. 1), 133 - 143.
33. *Beck, L. A. (1989a). *Symmetromphalus hageni* sp. n., a new neomphalid gastropod (Prosobranchia: Neomphalidae) from hydrothermal vents at the Manus Back-Arc Basin (Bismarck Sea, Papua New Guinea). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 93(B), 243 - 257.
34. *Beck, L. A. (1989b). Two new neritacean limpets (Gastropoda: Prosobranchia: Neritacea: Phenacolepadidae) from active hydrothermal vents at Hydrothermal Field 1 "Wienerwald" in the Manus Back-Arc Basin (Bismarck Sea, Papua-New Guinea). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 93(B), 259 - 275.
35. Beck, L. A. (1990 / 1991). Morphological and anatomical studies on a new lepetodrilacean limpet (Gastropoda, Prosobranchia) from hydrothermal vents at the Manus Back-Arc Basin (Bismarck Sea, Papua New Guinea). *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*, 94 / 95(B), 167 - 179.
36. Beck, L. A. (1996). Systematic position and relationship of *Phymorhynchus hyifluxi* n. sp., a further new turrid gastropod species associated with hydrothermal vent sites in the north Fiji Basin. *Archiv für Molluskenkunde*, 126, 109 - 115.
37. *Beedessee, G., Watanabe, H., Ogura, T., Nemoto, S., Yahagi, T., Nakagawa, S., . . . Marie, D. E. P. (2013). High Connectivity of Animal Populations in Deep-Sea Hydrothermal Vent Fields in the Central Indian Ridge Relevant to Its Geological Setting. *PLoS One*, 8(12, e81570), 1 - 11.
38. *Beinart, R. A., Sanders, J. G., Faure, B., Sylva, S. P., Lee, R. W., Becker, E. L., . . . Girguis, P. R. (2012). Evidence for the role of endosymbionts in regional-scale habitat partitioning by hydrothermal vent symbioses. *PNAS*, E3241 - E3250.

39. *Bell, J. B., Reid, W. D. K., Pearce, D. A., Glover, A. G., Sweeting, C. J., Newton, J., & Woulls, C. (2016). Hydrothermal activity lowers trophic diversity in Antarctic sedimented hydrothermal vents. *Biogeosciences Discussions*. doi:10.5194/bg-2016-318
40. *Bell, K. L. C., Brennan, M. L., & Raineault, N. A. (2015). New frontiers in ocean exploration: The E/V *Nautilus* 2014 Gulf of Mexico and Caribbean field season. *Oceanography*, 28(1, supplement), 60.
41. Bell, K. L. C., Flanders, J., Bowman, A., & Raineault, N. A. (2017). New frontiers in ocean exploration: The E/V *Nautilus*, NOAA Ship *Okeanos Explorer*, and R/V *Falkor* 2016 field season. *Oceanography*, 30(1), 94pp.
42. Bellan-Santini, D. (2005). European species of *Haustorius* (Crustacea: Amphipoda: Haustoriidae), with description of a new Mediterranean species. *Journal of Natural History*, 39(15), 1101 - 1110.
43. Bellan-Santini, D. (2007). New amphipods of hydrothermal vent environments on the Mid-Atlantic Ridge, Azores Triple junction zone. *Journal of Natural History*, 41(9 - 12), 567 - 596.
44. Bellan-Santini, D., & Kaim-Malka. (1977). In T. Horton, J. Lowry, C. De Broyer, D. Bellan-Santini, C. O. Coleman, L. Corbari, M. Daneliya, J. C. Dauvin, C. Fišer, R. Gasca, M. Grabowski, J. M. Guerra-García, E. Hendrycks, L. Hughes, D. Jaume, K. Jazdzewski, Y. H. Kim, R. King, T. Krapp-Schickel, S. LeCroy, A. N. Lötz, T. Mamos, A. R. Senna, C. Serejo, B. Sket, J. F. Souza-Filho, A. H. Tandberg, J. Thomas, M. Thurston, W. Vader, R. Väinölä, R. Vonk, K. White, & W. Zeidler (Eds.), World Amphipoda Database. Retrieved from: <http://www.marinespecies.org/amphipoda> on 2017-12-07.
45. Bellan-Santini, D., & Thurston, M. H. (1996). Amphipoda of the hydrothermal vents along the mid-Atlantic Ridge. *Journal of Natural History*, 30(5), 685 - 702.
46. *No reference is associated with this number (as the reference originally included here was a duplicate). This number can therefore be used as a reference code in future versions of sFDvent.*
47. Bennett, S. A., Van Dover, C. L., Breier, J. A., & Coleman, M. (2015). Effect of depth and vent fluid composition on the carbon sources at two neighboring deep-sea hydrothermal vent fields (Mid-Cayman Rise). *Deep Sea Research Part I: Oceanographic Research Papers*, 104, 122 - 133.
48. Bergquist, D. C., Eckner, J. T., Urcuyo, I. A., Cordes, E. E., Hourdez, S., Macko, S. A., & Fisher, C. R. (2007). Using stable isotopes and quantitative community characteristics to determine a local hydrothermal vent food web. *Marine Ecology Progress Series*, 330, 49 - 65.

49. Bischoff, M., Almeida, A. J., & Armbruster, J. W. (2004). New Species of *Pachycara* Zugmayer (Pisces: Zoarcidae) from the Rainbow Hydrothermal Vent Field (Mid-Atlantic Ridge). *Copeia*, 2004(3), 562-568. doi:10.1643/CI-03-031R2
50. *Blake, E. A., & Van Dover, C. L. (2005). The reproductive biology of *Amathys lutzi*, an ampharetid polychaete from hydrothermal vents on the Mid-Atlantic Ridge. *Invertebrate Biology*, 124(3), 254 - 264.
51. Blake, J. A. (1991). A new species of *Hesiocaeaca* (Polychaeta: Hesionidae) from hydrothermal vents at the Mariana Back-arc basin with notes on other polychaetes. *Proceedings of The Biological Society of Washington*, 104(1), 175 - 180.
52. Blake, J. A. (1985). Polychaeta from the vicinity of deep-sea geothermal vents in the eastern Pacific. 1. Euphrosinidae, Phyllodocidae, Hesionidae, Nereidae, Glyceridae, Dorvilleidae, Orbiniidae, and Maldanidae. *Bulletin of the Biological Society of Washington*, 6, 67 - 101.
53. Blake, J. A., & Hilbig, B. (1990). Polychaeta from the Vicinity of Deep-sea Hydrothermal Vents in the Eastern Pacific. II. New Species and Records from the Juan de Fuca and Explorer Ridge Systems. *Pacific Science*, 44(3), 219 - 253.
54. Böggemann, M. (2009). Polychaetes (Annelida) of the abyssal SE Atlantic. *Organisms Diversity & Evolution*, 9(4), 251 - 428. doi: 10.1016/j.ode.2009.10.001
55. Borda, E., Kudenov, J. D., Chelvadonné, P., Blake, J. A., Desbruyères, D., Fabri, M. C., . . . Rouse, G. W. (2013). Cryptic species of *Archinome* (Annelida: Amphionomida) from vents and seeps. *Proceedings of the Royal Society B: Biological Sciences*, 280(1770), 1 - 9.
56. Boschen, R. E., Rowden, A. A., Clark, M. R., Barton, S. J., Pallentin, A., & Gardner, J. P. A. (2015). Megabenthic assemblage structure on three New Zealand seamounts: implications for seafloor massive sulfide mining. *Marine Ecology Progress Series*, 523, 1 - 14.
57. Boschen, R. E., Rowden, A. A., Clark, M. R., & Gardner, J. P. A. (2015). Limitations in the Use of Archived Vent Mussel Samples to Assess Genetic Connectivity Among Seafloor Massive Sulfide Deposits: A Case Study with Implications for Environmental Management. *Frontiers in Marine Science*, 2(105), 1 - 14.
58. *Boschen, R. E., Rowden, A. A., Clark, M. R., Pallentin, A., & Gardner, J. P. A. (2016). Seafloor massive sulfide deposits support unique megafaunal assemblages: Implications for seabed mining and conservation. *Marine Environmental Research*, 115, 78 - 88.
59. Boss, K. J., & Turner, R. D. (1980). The giant white clam from the Galapagos Rift,

Calyptogena magnifica species novum. *Malacologia*, 20, 161 - 194.

60. Bowden, D. A., Rowden, A. A., Leduc, D., Beaumont, J., & Clark, M. R. (2016). Deep-sea seabed habitats: Do they support distinct mega-epifaunal communities that have different vulnerabilities to anthropogenic disturbance? *Deep Sea Research Part I: Oceanographic Research Papers*, 107(Supplement C), 31 - 47. doi: 10.1016/j.dsr.2015.10.011
61. *Boyle, E. A., Thaler, A. D., Jacobson, A., Plouviez, S., & Van Dover, C. L. (2013). Characterization of 10 polymorphic microsatellite loci in *Munidopsis lauensis*, a squat-lobster from the southwestern Pacific. *Conservation Genetics Resources*, 5(3), 647 - 649.
62. Brescia, L. A., & Tunnicliffe, V. (1998). Population biology of two pycnogonid species (Amnotheidae) at hydrothermal vents in the northeast Pacific, in: *Proceedings of the First International Symposium on Deep-Sea Hydrothermal Vent Biology: Funchal, Madeira, Portugal 20-24 October 1997. Cahiers de Biologie Marine*, 39(3 - 4), 233 - 236.
63. *Breusing, C., Biastoch, A., Drews, A., Metaxas, A., Jollivet, D., Vrijenhoek, R. C., . . . Reusch, T. B. H. (2016). Biophysical and Population Genetic Models Predict the Presence of "Phantom" Stepping Stones Connecting Mid-Atlantic Ridge Vent Ecosystems. *Current Biology*, 26, 2257 - 2267.
64. Breusing, C., Johnson, S. B., Tunnicliffe, V., & Vrijenhoek, R. C. (2015). Population structure and connectivity in Indo-Pacific deep-sea mussels of the *Bathymodiolus septemdierum* complex. *Conservation Genetics*, 16(6), 1415-1430. doi:10.1007/s10592-015-0750-0
65. *Breusing, C., Vrijenhoek, R. C., & Reusch, T. B. H. (2017). Widespread introgression in deep-sea hydrothermal vent mussels. *BMC Evolutionary Biology*, 17(1), 13. doi:10.1186/s12862-016-0862-2Britayev, T. A., Martin, D., Krylova, E. M., Von Cosel, R., & Aksiuk, T. S. (2007). Life-history traits of the symbiotic scale-worm *Branchipolynoe seepensis* and its relationships with host mussels of the genus *Bathymodiolus* from hydrothermal vents, *Marine Ecology*, 28(1), 36 - 48.
66. **Expert opinion is assigned this value (3380 records in the recommended dataset).**
67. Bronnimann, P., Van Dover, C. L., & Whittaker, J. E. (1989). *Abyssotherma pacifica*, n. gen., n. sp., a Recent remaneicid (Foraminiferida, Remaneicacea) from the East Pacific Rise. *Micropaleontology*, 35(2), 142 - 149.
68. Buckeridge, J. S. (2000). *Neolepas osheai* sp . nov., a new deep-sea vent barnacle (Cirripedia: Pedunculata) from the Brothers Caldera, south-west Pacific Ocean. *New Zealand Journal of Marine and Freshwater Research*, 34(3), 409 - 418.

69. Buckeridge, J. S., Linse, K., & Jackson, J. A. (2013). *Vulcanolepas scotiaensis* sp. nov., a new deep-sea scalpelliform barnacle (Eolepadidae: Neolepadinae) from hydrothermal vents in the Scotia Sea, Antarctica. *Zootaxa*, 3745(5), 551 - 568.
70. Calder, D. R., & Vervoort, W. (1998). Some hydroids (Cnidaria: Hydrozoa) from the Mid-Atlantic Ridge, in the North Atlantic Ocean. *Zoologische verhandelingen / uitgegeven door het Rijksmuseum van Natuurlijke Historie te Leiden (Ministerie van Cultuur, Recreatie en Maatschappelijk Werk)*, 319(29), 1 - 65.
71. Campbell, B. J., Stein, J. L., & Cary, S. C. (2003). Evidence of Chemolithoautotrophy in the Bacterial Community Associated with *Alvinella pompejana*, a Hydrothermal Vent Polychaete. *Applied and Environmental Microbiology*, 69(9), 5070 - 5078.
72. *Carey, S., Ballard, R., Bell, K. L. C., Bell, R. J., Connally, P., Dondin, F., . . . Smart, C. (2014). Cold seeps associated with a submarine debris avalanche deposit at Kick'em Jenny volcano, Grenada (Lesser Antilles). *Deep Sea Research Part I: Oceanographic Research Papers*, 93(Supplement C), 156 - 160. doi:10.1016/j.dsr.2014.08.002
73. Cary, S. C., Cottrell, M. T., Stein, J. L., Camacho, F., & Desbruyères, D. (1997). Molecular Identification and Localization of Filamentous Symbiotic Bacteria Associated with the Hydrothermal Vent Annelid *Alvinella pompejana*. *Applied and Environmental Microbiology*, 63(3), 1124 - 1130.
74. *Cavanaugh, C. M., Gardiner, S. L., Jones, M. L., Jannasch, H. W., & Waterbury, J. B. (1981). Prokaryotic Cells in the Hydrothermal Vent Tube Worm *Riftia pachyptila* Jones: Possible Chemoautotrophic Symbionts. *Science*, 213(4505), 340 - 342.
75. Cavanaugh, C. M., Wirsén, C. O., & Jannasch, H. W. (1992). Evidence for methylotrophic symbionts in a hydrothermal vent mussel (Bivalvia: Mytilidae) from the Mid-Atlantic Ridge. *Applied and Environmental Microbiology*, 58(12), 3799 - 3803.
76. Cavanaugh, C. M., McKiness, Z. P., Newton, I. L. G., & Stewart, F. J. (2006). Marine Chemosynthetic Symbioses *The Prokaryotes* (3rd Edition ed., pp. 475-507). New York: Springer.
77. *Chaffee, C., & Lindberg, D. R. (1986). Larval biology of early Cambrian molluscs: The implications of small body size. *Bulletin of Marine Science*, 39(2), 536 - 549.
78. Chao, L. S.-L., Davis, R. E., & Moyer, C. L. (2007). Characterization of bacterial community structure in vestimentiferan tubeworm *Ridgeia piscesae* trophosomes. *Marine Ecology*, 28(1), 72 - 85.
79. Chelvadonné, P., & Olu, K. (1996). Occurrence of anomuran crabs (Crustacea: Decapoda) in hydrothermal vent and cold-seep communities: a review. *Proceedings of*

The Biological Society of Washington, 109(2), 286 - 298.

80. Chen, C., Copley, J. T., Linse, K., & Rogers, A. D. (2015). Low connectivity between 'scaly-foot gastropod' (Mollusca: Peltospiridae) populations at hydrothermal vents on the Southwest Indian Ridge and the Central Indian Ridge. *Organisms Diversity and Evolution*, 15(4), 663 - 670.
81. Chen, C., Copley, J. T., Linse, K., Rogers, A. D., & Sigwart, J. D. (2015). The heart of a dragon: 3D anatomical reconstruction of the 'scaly-foot gastropod' (Mollusca: Gastropoda: Neomphalina) reveals its extraordinary circulatory system. *Frontiers in Zoology*, 12(13), 1 - 16.
82. Chen, C., Linse, K., Copley, J. T., & Rogers, A. D. (2015). The 'scaly-foot gastropod': a new genus and species of hydrothermal vent-endemic gastropod (Neomphalina: Peltospiridae) from the Indian Ocean. *Journal of Molluscan Studies*, 81(3), 322 - 334.
83. Chen, C., Linse, K., Roterman, C. N., Copley, J. T., & Rogers, A. D. (2015). A new genus of large hydrothermal vent-endemic gastropod (Neomphalina: Peltospiridae). *Zoological Journal of the Linnean Society*, 175(2), 319 - 335.
84. Chen, C., Uematsu, K., Linse, K., & Sigwart, J. D. (2017). By more ways than one: Rapid convergence in adaptations to hydrothermal vents shown by 3D anatomical reconstruction of *Gigantopelta* (Mollusca: Neomphalina). *BMC Evolutionary Biology*, 17(62), 1 - 19.
85. *ChEss. (2010). 'Meiofauna from chemosynthetic deep-sea environments', *Report on the ChEss workshop held 17 July 2010 at Ghent University, Belgium* Retrieved from: http://archive.noc.ac.uk/chess/docs/meiofauna_report.pdf
86. Chevreux, E. (1909). Biospeologica XII. Amphipodes (Première série). *Archives de zoologie expérimentale et générale, ser 5(2)*, 27 - 42.
87. *Child, C. A. (1987). *Ammothea verenae* and *Sericosura venticola*, two new hydrothermal vent associated pycnogonids from the Northeast Pacific. *Proceedings of The Biological Society of Washington*, 100(4), 892 - 901.
88. Child, C. A., & Segonzac, M. (1996) *Sericosura heteroscela* and *S. cyrtoma*, new species, and other Pycnogonida from Atlantic and Pacific hydrothermal vent, with notes on habitat and environment. *Proc. Biol. Soc. Wash.*, 109(4).
89. *Child, C. A. (2000). *Sericosura dissita*, n. sp., a Third Hydrothermal Vent Pycnogonida Described from the Northeast Pacific, and Other Known Vent Species. *Species Diversity*, 5, 1 - 6.

90. *Chu, J. W. F., & Reiswig, H. M. (2014). Mechanisms of propagule release in the carnivorous sponge *Asbestopluma occidentalis*. *Invertebrate Biology*, 133(2), 109 - 120.
91. *Clark, M. R., & O'Shea, S. (2001). Hydrothermal vent and seamount fauna from the southern Kermadec Ridge, New Zealand. *InterRidge News*, 10(2), 14 - 17.
92. Coan, E. V., Scott, P. V., & Bernard, F. R. (2000). *Bivalve Seashells of Western North America: Marine Bivalve Mollusks from Arctic Alaska to Baja California*. Santa Barbara, California, USA: Santa Barbara Museum of Natural History.
93. *Cohen, D. M., Inada, T., Iwamoto, T., & Scialabba, N. (1990). FAO species catalogue. Vol. 10. Gadiform fishes of the world (Order Gadiformes). An annotated and illustrated catalogue of cods, hakes, grenadiers and other gadiform fishes known to date. *FAO Fisheries Synopsis*, 125(10), 442.
94. Cohen, D. M., Rosenblatt, R. H., & Moser, H. G. (1990). Biology and description of a bythitid fish from deep-sea thermal vents in the tropical eastern Pacific. *Deep Sea Research Part A: Oceanographic Research Papers*, 37(2), 267 - 283.
95. Colaço, A., Desbruyères, D., & Guezennec, J. (2007) Polar lipid fatty acids as indicators of trophic associations in a deep-sea vent system community. *Marine Ecology*, 28(1), 15 - 24.
96. Colaço, A., Dehairs, F., & Desbruyères, D. (2002). Nutritional relations of deep-sea hydrothermal fields at the Mid-Atlantic Ridge: a stable isotope approach. *Deep Sea Research Part I: Oceanographic Research Papers*, 49(2), 395 - 412.
97. *No reference is associated with this number (as the reference originally included here was a duplicate). This number can therefore be used as a reference code in future versions of sFDvent.*
98. Colaço, A., Desbruyères, D., Comtet, T., & Alayse, A-M. (1998) Ecology of the Menez Gwen hydrothermal vent field (Mid-Atlantic Ridge / Azores triple junction), *Cahiers de Biologie Marine*, 39, 237 - 240.
99. *Collins, P. C., Kennedy, R., & Van Dover, C. L. (2012). A biological survey method applied to seafloor massive sulphides (SMS) with contagiously distributed hydrothermal-vent fauna. *Marine Ecology Progress Series*, 452(89 - 107).
100. *Comtet, T., Le Pennec, M., & Desbruyères, D. (1999). Evidence of a sexual pause in *Bathymodiolus azoricus* (Bivalvia: Mytilidae) from hydrothermal vents of the Mid-Atlantic Ridge. *Journal of the Marine Biological Association of the U.K.*, 79(6), 1149 - 1150.

101. Connelly, D. P., Copley, J. T., Murton, B. J., Stansfield, K., Tyler, P. A., German, C. R., . . . Wilcox, S. (2012). Hydrothermal vent fields and chemosynthetic biota on the world's deepest seafloor spreading centre. *Nature Communications*, 3, 620. doi:10.1038/ncomms1636
102. Copley, J. T., Marsh, L., Glover, A. G., Huhnerbach, V., Nye, V. E., Reid, W. D., . . . Wiklund, H. (2016). Ecology and biogeography of megafauna and macrofauna at the first known deep-sea hydrothermal vents on the ultraslow-spreading Southwest Indian Ridge. *Scientific Reports*, 6, 39158. doi:10.1038/srep39158
103. Copley, J. T. P., Flint, H. C., Ferrero, T. J., & Van Dover, C. L. (2007). Diversity of meiofauna and free-living nematodes in hydrothermal vent mussel beds on the northern and southern East Pacific Rise. *Journal of the Marine Biological Association of the U.K.*, 87(5), 1141 - 1152.
104. Copley, J. T. P., Jorgensen, P. B. K., & Sohn, R. A. (2007). Assessment of decadal-scale ecological change at a deep Mid-Atlantic hydrothermal vent and reproductive time-series in the shrimp *Rimicaris exoculata*. *Journal of the Marine Biological Association of the U.K.*, 87(4), 859 - 867.
105. *Copley, J. T. P., Tyler, P. A., Van Dover, C. L., & Philp, S. J. (2003). Spatial variation in the reproductive biology of *Paralvinella palmiformis* (Polychaeta: Alvinellidae) from a vent field on the Juan de Fuca Ridge. *Marine Ecology Progress Series*, 255, 171-181.
106. Corbari, L., & Sorbe, J.-C. (2018)¹. First observations of the behaviour of the deep-sea amphipod *Dulichiopsis dianae* sp. nov. (Senticaudata, Dulichiidae) in the TAG hydrothermal vent field (Mid-Atlantic Ridge). *Marine Biodiversity*, 48 (1), 631 - 645.
107. Corbari, L., Durand, L., Cambon-Bonavita, M. A., Gaill, F., & Compère, P. (2012). New digestive symbiosis in the hydrothermal vent amphipoda *Ventiella sulfuris*. *Comptes rendus biologies*, 335(2), 142 -154.
108. Corbera, J., Segonzac, M., & Cunha, M. R. (2008). A new deep-sea genus of Nannastacidae (Crustacea, Cumacea) from the Lucky Strike hydrothermal vent field (Azores Triple Junction, Mid- Atlantic Ridge). *Marine Biology Research*, 4(3), 180 - 192.
109. *Cordes, E. E., Becker, E. L., & Fisher, C. R. (2010). Temporal shift in nutrient input to cold-seep food webs revealed by stable-isotope signatures of associated communities. *Limnology and Oceanography*, 55(6), 2537 - 2548.
110. Cordes, E. E., Becker, E. L., Hourdez, S., & Fisher, C. R. (2010). Influence of foundation species, depth, and location on diversity and community composition at Gulf of Mexico lower-slope cold seeps. *Deep Sea Research Part II: Topical Studies in*

¹ Referred to in some of the dataset versions as Corbari and Sorbe (in press).

Oceanography, 57(21 - 23), 1870 - 1881.

111. *Cordes, E. E., Carney, S. L., Hourdez, S., Carney, R. S., Brooks, J. M., & Fisher, C. R. (2007). Cold seeps of the deep Gulf of Mexico: Community structure and biogeographic comparisons to Atlantic equatorial belt seep communities. *Deep Sea Research Part I: Oceanographic Research Papers*, 54(4), 637 - 653.
112. *Cordes, E. E., Bergquist, D. C., Predmore, B. L., Jones, C., Deines, P., Teleshnicki, G., & Fisher, C. R. (2006). Alternate unstable states: Convergent paths of succession in hydrocarbon-seep tubeworm-associated communities. *Journal of Experimental Marine Biology and Ecology*, 339(2), 159 - 176.
113. Cubelio, S. S., Tsuchida, S., & Watanabe, S. (2008). New species of *Munidopsis* (Decapoda: Anomura: Galatheidae) from hydrothermal vent areas of Indian and Pacific Oceans. *Journal of the Marine Biological Association of the U.K.*, 88(1), 111 - 117.
114. Cubelio, S. S., Tsuchida, S., & Watanabe, S. (2007). Vent Associated *Munidopsis* (Decapoda: Anomura: Galatheidae) from Brothers Seamount, Kermadec Arc, Southwest Pacific, with Description of One New Species. *Journal of Crustacean Biology*, 27(3), 513 - 519.
115. Cuvelier, D., Beesau, J., Ivanenko, V. N., Zeppilli, D., Sarradin, P.-M., & Sarrazin, J. (2014). First insights into macro- and meiofaunal colonisation patterns on paired wood/slate substrata at Atlantic deep-sea hydrothermal vents. *Deep Sea Research Part I: Oceanographic Research Papers*, 87(Supplement C), 70-81.
doi:10.1016/j.dsr.2014.02.008
116. Cuvelier, D., de Busserolles, F., Lavaud, R., Floc'h, E., Fabri, M-C., Sarradin, P-M., and Sarrazin, J. (2012) Biological data extraction from imagery – How far can we go? A case study from the Mid-Atlantic Ridge. *Marine Environmental Research*, 82, 15 - 27.
117. Cuvelier, D., Sarradin, P.-M., Sarrazin, J., Colaço, A., Copley, J. T., Desbruyères, D., . . . Tyler, P. A. (2011). Hydrothermal faunal assemblages and habitat characterisation at the Eiffel Tower edifice (Lucky Strike, Mid-Atlantic Ridge). *Marine Ecology*, 32(2), 243 - 255.
118. Cuvelier, D., Sarrazin, J., Colaço, A., Copley, J., Desbruyères, D., Glover, A. G., . . . Serrão-Santos, R. (2009). Distribution and spatial variation of hydrothermal faunal assemblages at Lucky Strike (Mid-Atlantic Ridge) revealed by high-resolution video image analysis. *Deep Sea Research Part I: Oceanographic Research Papers*, 56(11), 2026 - 2040.
119. No reference is associated with this number. This number can therefore be used as a reference code in future versions of sFDvent.

120. No reference is associated with this number. This number can therefore be used as a reference code in future versions of sFDvent.
121. *Dahlgren, T. G., Glover, A. G., Baco, A., & Smith, C. R. (2004). Fauna of whale falls: systematics and ecology of a new polychaete (Annelida: Chrysopetalidae) from the deep Pacific Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 51(12), 1873 - 1887.
122. *Dahlhoff, E., Schneidemann, S., & Somero, G. N. (1990). Pressure-Temperature Interactions on M4-Lactate Dehydrogenases From Hydrothermal Vent Fishes: Evidence for Adaptation to Elevated Temperatures by the Zoarcid *Thermarces andersoni*, but not by the Bythitid, *Bythites hollisi*. *The Biological Bulletin*, 179(1), 134-139. doi:10.2307/1541747
123. de Burgh, M. E., Juniper, S. K., & Singla, C. L. (1989). Bacterial symbiosis in Northeast Pacific Vestimentifera: a TEM study. *Marine Biology*, 101, 97 - 105.
124. de Burgh, M. E., & Singla, C. L. (1984). Bacterial colonization and endocytosis on the gill of a new limpet species from a hydrothermal vent. *Marine Biology*, 84, 1 - 6.
125. De Busserolles, F., Sarrazin, J., Gauthier, O., Gélinas, Y., Fabri, M. C., Sarradin, P. M., & Desbruyères, D. (2009). Are spatial variations in the diets of hydrothermal fauna linked to local environmental conditions? *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(19), 1649-1664. doi:10.1016/j.dsr2.2009.05.011
126. de Saint Laurent, M., & Macpherson, E. (1997). Une nouvelle espèce du genre *Paralomis* White, 1856, des sources hydrothermales du Sud-ouest Pacifique (Crustacea, Decapoda, Lithodidae). *Zoosystema*, 19, 721 - 727.
127. *de Chaine, E. G., Bates, A. E., Shank, T. M., & Cavanaugh, C. M. (2006). Off-axis symbiosis found: characterization and biogeography of bacterial symbionts of *Bathymodiolus* mussels from Lost City hydrothermal vents. *Environmental Microbiology*, 8(11), 1902 - 1912.
128. *Desbruyères, D., Hashimoto, J., & Fabri, M. C. (2006). Composition and Biogeography of Hydrothermal Vent Communities in Western Pacific Back-Arc Basins. In D. M. Christie, C. R. Fisher, S.-M. Lee, & S. Givens (Eds.), *Back-Arc Systems: Geological, Biological, Chemical, and Physical Interactions*. Washington, D. C.: American Geophysical Union.
129. Desbruyères, D., Segonzac, M., & Bright, M. (2006). *Handbook of Deep-Sea Hydrothermal Vent Fauna* (D. Desbruyères, M. Segonzac, & M. Bright Eds.).
130. Desbruyères, D., Biscoito, M., Caprais, J.-C., Colaço, A., Comtet, T., Crassous, P., . . . Vangriesheim, A. (2001). Variations in deep-sea hydrothermal vent communities on

the Mid-Atlantic Ridge near the Azores plateau. *Deep-sea Research I*, 48, 1325 - 1346.

131. *Desbruyères, D., Almeida, A., Biscoito, M., Comtet, T., Khripounoff, A., Le Bris, N., . . . Segonzac, M. (2000). A review of the distribution of hydrothermal vent communities along the northern Mid-Atlantic Ridge: dispersal vs. environmental controls. *Hydrobiologia*, 440, 201 - 216.
132. *Desbruyères, D., & Hourdez, S. (2000). A new species of scale-worm (Polychaeta: Polynoidae), *Lepidonotopium jouinae* sp.nov., from the Azores Triple Junction on the Mid-Atlantic Ridge. *Cahiers de Biologie Marine*, 41(4), 399 - 406.
133. Desbruyères, D., Alayse-Danet, A.-M., Ohta, S., & the Scientific Parties of biolauand, s. (1994). Deep-sea hydrothermal communities in Southwestern Pacific back-arc basins (the North Fiji and Lau Basins): Composition, microdistribution and food web. *Marine Geology*, 116(1), 227 - 242. doi:10.1016/0025-3227(94)90178-3
134. *Desbruyères, D., & Laubier, L. (1982). *Paralvinella grasslei*, new genus, new species of Alvinellinae (Polychaeta: Ampharetidae) from the Galápagos Rift geothermal vents. *Proceedings of The Biological Society of Washington*, 95(3), 484 - 494.
135. *Di Camillo, C. G., Giordano, G., Bo, M., Betti, F., Mori, M., Puce, S., & Bavestrello, G. (2013). Seasonal patterns in the abundance of *Ectopleura crocea* (Cnidaria: Hydrozoa) on a shipwreck in the Northern Adriatic. *Marine Ecology*, 34(s1), 25 - 32.
136. Distel, D. L., Lane, D. J., Olsen, G. J., Giovannoni, S. J., Pace, B., Pace, N. R., . . . Felbeck, H. (1988). Sulfur-oxidizing bacterial endosymbionts: analysis of phylogeny and specificity by 16S rRNA sequences. *Journal of Bacteriology*, 170(6), 2506 - 2510.
137. Dittel, A. I., Perovich, G., & Epifanio, C. E. (2008). Biology of the Vent Crab *Bythograea thermydron*: A Brief Review. *Journal of Shellfish Research*, 27(1), 63 - 77.
138. *Dittel, A. I., Epifanio, C. E., & Perovich, G. (2005). Food sources for the early life history stages of the hydrothermal vent crab *Bythograea thermydron*: a stable isotope approach. *Hydrobiologia*, 544(1), 339 - 346.
139. *Dixon, D. R., Lowe, D. M., Miller, P. I., Villemin, G. R., Colaço, A., Serrão-Santos, R., & Dixon, L. R. J. (2006). Evidence of seasonal reproduction in the Atlantic vent mussel *Bathymodiolus azoricus*, and an apparent link with the timing of photosynthetic primary production. *Journal of the Marine Biological Association of the U.K.*, 86(6), 1363 - 1371.
140. Dolan, M. F. J., Buhl-Mortensen, P., Thorsnes, T., Buhl-Mortensen, L., Bellec, V. K., & Bøe, R. (2009). Developing seabed nature-type maps offshore Norway: initial results from the MAREANO programme. *Norwegian Journal of Geology*, 89, 17 - 28.

141. *Dubilier, N., Windoffer, R., & Giere, O. (1998). Ultrastructure and stable carbon isotope composition of the hydrothermal vent mussels *Bathymodiolus brevior* and *B. sp. affinis brevior* from the North Fiji Basin, western Pacific. *Marine Ecology Progress Series*, 165, 187 - 193.
142. Duperron, S., Gaudron, S. M., Rodrigues, C. F., Cunha, M. R., Decker, C., & Olu, K. (2013). An overview of chemosynthetic symbioses in bivalves from the North Atlantic and Mediterranean Sea. *Biogeosciences*, 10(5), 3241 - 3267.
143. *Duperron, S. (2010). The diversity of deep-sea mussels and their bacterial symbioses. In S. Kiel (Ed.), *The Vent and Seep Biota: Aspects from Microbes to Ecosystems* (pp. 137 - 167). Dordrecht: Springer.
144. *Duperron, S., Sibuet, M., MacGregor, B. J., Kuypers, M. M. M., Fisher, C. R., & Dubilier, N. (2007). Diversity, relative abundance and metabolic potential of bacterial endosymbionts in three *Bathymodiolus* mussel species from cold seeps in the Gulf of Mexico. *Environmental Microbiology*, 9(6), 1423 - 1438.
145. Duperron, S., Bergin, C., Zielinski, F., Blazejak, A., Pernthaler, A., McKiness, Z. P., . . . Dubilier, N. (2006). A dual symbiosis shared by two mussel species, *Bathymodiolus azoricus* and *Bathymodiolus puteoserpentis* (Bivalvia: Mytilidae), from hydrothermal vents along the northern Mid-Atlantic Ridge. *Environmental Microbiology*, 8(8), 1441 - 1447.
146. *Ellis, J., & Solander, D. (1786). *The Natural History of many curious and uncommon Zoophytes, collected from various parts of the Globe. Systematically arranged and described by the late John Ellis, etc.*. London. 1 - 206.
147. *Encyclopedia of Life*. Retrieved from: <http://www.eol.org>.
148. *Epifanio, C. E., Perovich, G., Dittel, A. I., & Cary, S. C. (1999). Development and behavior of megalopa larvae and juveniles of the hydrothermal vent crab *Bythograea thermydron*. *Marine Ecology Progress Series*, 185, 147 - 154.
149. Erickson, K. L., Macko, S. A., & Van Dover, C. L. (2009). Evidence for a chemoautotrophically based food web at inactive hydrothermal vents (Manus Basin). *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(19 - 20), 1577 - 1585.
150. *Esposito, V., Giacobbe, S., Cosentino, A., Minerva, C. S., Romeo, T., Canese, S., & Andaloro, F. (2015). Distribution and ecology of the tube-dweller *Ampelisca ledoyerii* (Amphipoda: Ampeliscidae) associated with the hydrothermal field off Panarea Island (Tyrrhenian Sea, Mediterranean). *Marine Biodiversity*, 45(4), 763 - 768.
151. Fabri, M. C., Bargain, A., Briand, P., Gebruk, A., Fouquet, Y., Morineaux, M., &

- Desbruyères, D. (2011). The hydrothermal vent community of a new deep-sea field, Ahazze-1, 12°58'N on the Mid-Atlantic Ridge. *Journal of the Marine Biological Association of the U.K.*, 91, 1 - 13.
152. Fauchald, K., & Jumars, P. A. (1979). The diet of worms: a study of polychaete feeding guilds. *Oceanography and Marine Biology, An Annual Review*, 17(193 - 284).
153. Fautin, D. G., & Barber, B. R. (1999). *Maractis rimicarivora*, a new genus and species of sea anemone (Cnidaria: Anthozoa: Actiniaria: Actinostolidae) from an Atlantic hydrothermal vent. *Proceedings of the Royal Society of Washington*, 112(3), 624 - 631.
154. Fautin, D. G., & Hessler, R. R. (1989). *Marianactis bythios*, a new genus and species of actinostolid sea anemone (Coelenterata: Actiniaria) from the Mariana vents. *Proceedings of The Biological Society of Washington*, 102, 815 - 825.
155. Felbeck, H., Childress, J., & Somero, G. (1981). Calvin-Benson cycle and sulphide oxidation enzymes in animals from sulphide-rich habitats. *Nature*, 293, 291 - 293.
156. *Feldman, R. A., Black, M. B., Cary, C. S., Lutz, R. A., & Vrijenhoek, R. C. (1997). Molecular Phylogenetics of Bacterial Endosymbionts and Their Vestimentiferan Hosts. *Molecular Marine Biology and Biotechnology*, 6(3), 268 - 277.
157. Feng, D., Cheng, M., Kiel, S., Qiu, J.-W., Yang, Q., Zhou, H., . . . Chen, D. (2015). Using *Bathymodiolus* tissue stable carbon, nitrogen and sulfur isotopes to infer biogeochemical process at a cold seep in the South China Sea. *Deep Sea Research Part I: Oceanographic Research Papers*, 104, 52 - 59. doi:10.1016/j.dsr.2015.06.011
158. Fiala-Médioni, A., McKiness, Z. P., Dando, P., Boulegue, J., Mariotti, A., Alayse-Danet, A., . . . Cavanaugh, C. (2002). Ultrastructural, biochemical, and immunological characterization of two populations of the mytilid mussel *Bathymodiolus azoricus* from the Mid-Atlantic Ridge: evidence for a dual symbiosis. *Marine Biology*, 141(6), 1035 - 1043.
159. Fiala-Médioni, A., & Metivier, C. (1986). Ultrastructure of the gill of the hydrothermal vents bivalve *Calyptogena magnifica* with considerations on its nutrition. *Marine Biology*, 90, 215 - 222.
160. Fisher, C. R., Childress, J. J., Macko, S. A., & Brooks, J. M. (1994). Nutritional interactions in Galapagos Rift hydrothermal vent communities: inferences from stable carbon and nitrogen isotope analyses *Marine Ecology Progress Series*, 103, 45 - 55.
161. Fisher, C. R. (1990). Chemoautotrophic and methanotrophic symbioses in marine invertebrates. *Reviews in Aquatic Sciences*, 2, 399 - 436.

162. Fleminger, A. (1983). Description and phylogeny of *Isaacsicalanus paucisetus*, n. gen., n. sp., (Copepoda: Calanoida: Spinocalanidae) from an East Pacific hydrothermal vent site (21°N). *Proceedings of The Biological Society of Washington*, 96(4), 605 - 622.
163. Flørenes, H. v. P. (2013). *Taxonomic overview of calcareous sponges (Porifera, Calcarea) in the Jan Mayen vent field at the Mohn Ridge*. (Master of Science in Marine Biology; Marine Biodiversity), The University of Bergen.
164. Fretter, V. (1988). New archaeogastropod limpets from hydrothermal vents; Superfamily Lepetodrilacea. II. Anatomy. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 319(1192), 1 - 82.
165. Froese, R., & Pauly, D. (2017). *FishBase*. Retrieved from: www.fishbase.org
166. *Fry, B., Gest, H., & Hayes, J. M. (1983). Sulphur isotopic compositions of deep-sea hydrothermal vent animals. *Nature*, 306, 51 - 52.
167. Fujikura, K., Kojima, S., Fujiwara, Y., Hashimoto, J., & Okutani, T. (2000). New Distribution Records of Vesicomyid Bivalves from Deep-sea Chemosynthesis-based Communities in Japanese Waters. *Venus*, 59, 103 - 121.
168. Fujikura, K., Okutani, T., & Maruyama, T. (2012). *Deep-sea Life : Biological observations using research submersibles* (2nd ed.). Tokyo: Tokyo University Press.
169. *Fujikura, K., Okutani, T., & Maruyama, T. (2008). *Deep-sea Life—Biological Observations Using Research Submersibles*. Hadano, Japan: Tokai University Press.
170. Fujiwara, Y., Takai, K., Uematsu, K., Tsuchida, S., Hunt, J. C., & Hashimoto, J. (2000). Phylogenetic characterization of endosymbionts in three hydrothermal vent mussels: influence on host distributions. *Marine Ecology Progress Series*, 208, 147 - 155.
171. Gale, K. S. P., Hamel, J.-F., & Mercier, A. (2013). Trophic ecology of deep-sea Asteroidea (Echinodermata) from eastern Canada. *Deep Sea Research Part I: Oceanographic Research Papers*, 80, 25 - 36. doi: 10.1016/j.dsr.2013.05.016
172. *Galea, H. R., Häussermann, V., & Försterra, G. (2007). Cnidaria, Hydrozoa: latitudinal distribution of hydroids along the fjords region of southern Chile, with notes on the world distribution of some species. *Check List*, 3(4), 308 - 320.
173. Galkin, S. V. (1997). Megafauna associated with hydrothermal vents in the Manus Back-Arc Basin (Bismarck Sea). *Marine Geology*, 142(1), 197 - 206. doi: 10.1016/S0025-3227(97)00051-0

174. Gardebrecht, A., Markert, S., Sievert, S. M., Felbeck, H., Thürmer, A., Albrecht, D., . . . Schweder, T. (2011). Physiological homogeneity among the endosymbionts of *Riftia pachyptila* and *Tevnia jerichonana* revealed by proteogenomics. *The Isme Journal*, 6, 766. doi:10.1038/ismej.2011.137
175. Gasca, R., & Shih, C.-T. (2001). Hyperiid amphipods from surface waters of the western Caribbean Sea (1991). *Crustaceana*, 74(5), 489 - 499.
176. Gaudron, S. M., Marqué, L., Thiébaut, E., Riera, P., Duperron, S., & Zbinden, M. (2015). How are microbial and detrital sources partitioned among and within gastropods species at East Pacific Rise hydrothermal vents? *Marine Ecology*, 36(S1), 18 - 34.
177. Gebruk, A. V., Chelvadonné, P., Shank, T., Lutz, R. A., & Vrijenhoek, R. C. (2000). Deep-sea hydrothermal vent communities of the Logatchev area (14°45' N, Mid-Atlantic Ridge): diverse biotopes and high biomass. *Journal of the Marine Biological Association of the U.K.*, 80(3), 383 - 393.
178. Gebruk, A. V., Pimenov, N. V., & Savvichev, A. S. (1993). Feeding specialization of bresiliid shrimps in the TAG site hydrothermal community *Marine Ecology Progress Series*, 98, 247 - 253.
179. Gebruk, A. V., Southward, E. C., Kennedy, H., & Southward, A. J. (2000). Food sources, behaviour, and distribution of hydrothermal vent shrimps at the Mid-Atlantic Ridge. *Journal of the Marine Biological Association of the U.K.*, 80(3), 485 - 499.
180. Geistdoerfer, P. (1994). *Careproctus hyaleius*, nouvelle espèce de poisson cyclopteridae (Liparinae) de l'écosystème hydrothermal de la dorsale de l'Océan Pacifique oriental (13°N). *Cybium*, 18(3), 325 - 333.
181. *Geistdoerfer, P. (1991). Ichthyofaune associée à l'hydrothermalisme océanique et description de *Thermobiotes mytilogeiton*, nouveau genre et nouvelle espèce de Synaphobranchidae (Pisces, Anguilliformes) de l'Océan Pacifique. *Comptes Rendus de l'Académie des Sciences, Paris*, 312(3), 91 - 97.
182. *Gibson, R. N., & Atkinson, R. J. A. (2003). *Oceanography and Marine Biology, An Annual Review* (Vol. 41): CRC Press, Taylor and Francis.
183. *Gibson, R. N., Atkinson, R. J. A., & Gordon, J. D. M. (2010). *Oceanography and Marine Biology*. New York: Chapman and Hall / CRC.
184. Giongo, A., Haag, T., Simão, T. L. L., Medina-Silva, R., Utz, L. R. P., Bogo, M. R., . . . Eizirik, E. (2016). Discovery of a chemosynthesis-based community in the western South Atlantic Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 112(Supplement C), 45-56. doi: 10.1016/j.dsr.2015.10.010

185. Glover, A. G., Goetze, E., Dahlgren, T. G., & Smith, C. R. (2005). Morphology, reproductive biology and genetic structure of the whale-fall and hydrothermal vent specialist, *Bathykurila guaymasensis* (Annelida: Polynoidae). *Marine Ecology*, 26(3 - 4), 223 - 234.
186. Glover, E. A., Taylor, J. D., & Rowden, A. A. (2004). *Bathyaustriella thionipta*, a new lucinid bivalve from a hydrothermal vent on the Kermadec Ridge, New Zealand and its relationship to shallow-water taxa (Bivalvia: Lucinidae). *Journal of Molluscan Studies*, 70(3), 283 - 295. doi:10.1093/mollus/70.3.283
187. Göcke, C., Hestetun, J. T., Uhlir, C., Freiwald, A., Beuck, L., & Janussen, D. (2016). *Cladorhiza corallophila* sp. nov., a new carnivorous sponge (Cladorhizidae, Demospongiae) living in close association with *Lophelia pertusa* and *Madrepora oculata* (Scleractinia). *Zootaxa*, 4168(3), 512 - 524.
188. Goffredi, S. K., & Barry, J. P. (2002a). Energy acquisition and allocation in vesicomyid symbioses. *Cahiers de Biologie Marine*, 43(3 - 4), 345 - 350.
189. Goffredi, S. K., & Barry, J. P. (2002b). Species-specific variation in sulfide physiology between closely related Vesicomyid clams. *Marine Ecology Progress Series*, 225, 227 - 238.
190. Goffredi, S. K., Hurtado, L., Hallam, S., & Vrijenhoek, R. C. (2003). Evolutionary relationships of deep-sea vent and cold seep clams (Mollusca: Vesicomyidae) of the "pacifica/lepta" species complex. *Marine Biology*, 142(2), 311 - 320.
191. *Goffredi, S. K., Johnson, S., Tunnicliffe, V., Caress, D., Clague, D., Escobar, E., . . . Vrijenhoek, R. (2017). Hydrothermal vent fields discovered in the southern Gulf of California clarify role of habitat in augmenting regional diversity. *Proceedings of the Royal Society B: Biological Sciences*, 284(1859), 1 - 10.
192. Goffredi, S. K., Jones, W. J., Erhlich, H., Springer, A., & Vrijenhoek, R. C. (2008). Epibiotic bacteria associated with the recently discovered Yeti crab, *Kiwa hirsuta*. *Environmental Microbiology*, 10(10), 2623 - 2634.
193. Goffredi, S. K., Warén, A., Orphan, V. J., Van Dover, C. L., & Vrijenhoek, R. C. (2004). Novel Forms of Structural Integration between Microbes and a Hydrothermal Vent Gastropod from the Indian Ocean. *Applied and Environmental Microbiology*, 70(5), 3082 - 3090.
194. *Gollner, S., Fontaneto, D., & Martínez Arbizu, P. (2011). Molecular taxonomy confirms morphological classification of deep-sea hydrothermal vent copepods (Dirivultidae) and suggests broad physiological tolerance of species and frequent dispersal along ridges. *Marine Biology*, 158(1), 221 - 231.

195. Gollner, S., Govenar, B., Fisher, C. R., & Bright, M. (2015). Size matters at deep-sea hydrothermal vents: different diversity and habitat fidelity patterns of meio- and macrofauna. *Marine Ecology Progress Series*, 520, 57 - 66.
196. Gollner, S., Govenar, B., Martinez Arbizu, P., Mills, S., Le Bris, N., Weinbauer, M., . . . Bright, M. (2015). Differences in recovery between deep-sea hydrothermal vent and vent-proximate communities after a volcanic eruption. *Deep Sea Research Part I: Oceanographic Research Papers*, 106, 167 - 182.
197. Gollner, S., Ivanenko, V. N., & Martínez Arbizu, P. (2008). A new species of deep-sea Tegastidae (Crustacea: Copepoda: Harpacticoida) from 9°50'N on the East Pacific Rise, with remarks on its ecology. *Zootaxa*, 1866, 323.
198. Gollner, S., Ivanenko, V. N., Martínez Arbizu, P., & Bright, M. (2010). Advances in Taxonomy, Ecology, and Biogeography of Dirivultidae (Copepoda) Associated with Chemosynthetic Environments in the Deep Sea. *PLoS One*, 5(8, e9801), 1 - 13.
199. Gollner, S., Miljutina, M., & Bright, M. (2013). Nematode succession at deep-sea hydrothermal vents after a recent volcanic eruption with the description of two dominant species. *Organisms Diversity & Evolution*, 13(3), 349 - 371.
doi:10.1007/s13127-012-0122-2
200. Gollner, S., Riemer, B., Martínez Arbizu, P., Le Bris, N., & Bright, M. (2010). Diversity of Meiofauna from the 9°50'N East Pacific Rise across a Gradient of Hydrothermal Fluid Emissions. *PLoS One*, 5(8, e12321), 1 - 14.
201. *Gollner, S., Stuckas, H., Kihara, T. C., Laurent, S., Kodami, S., & Martínez Arbizu, P. (2016). Mitochondrial DNA Analyses Indicate High Diversity, Expansive Population Growth and High Genetic Connectivity of Vent Copepods (Dirivultidae) across Different Oceans. *PLoS One*, 11(10, e0163776), 1 - 23.
202. Gollner, S., Zekely, J., Govenar, B., Le Bris, N., Nemeschkal, H. L., Fisher, C. R., & Bright, M. (2007). Tubeworm-associated permanent meiobenthic communities from two chemically different hydrothermal vent sites on the East Pacific Rise. *Marine Ecology Progress Series*, 337, 39 - 49.
203. González, A. F., Guerra, A., Pascual, S., & Briand, P. (1998). *Vulcanoctopus hydrothermalis* gen. et sp. nov. (Mollusca, Cephalopoda): an octopod from a deep-sea hydrothermal vent site. *Cahiers de Biologie Marine*, 39, 169 - 184.
204. Gordon, D. P. (n.d.). *Annual distribution of the giant seep mussel Bathymodiolus tangaroa lineage*. New Zealand's National Aquatic Biodiversity Information System (NABIS): New Zealand Government Retrieved from:
<http://www.nabis.govt.nz/LineageDocuments/Annual%20distribution%20of%20Giant%20seep%20mussel%20Lineage.pdf>.

205. *Govenar, B., Fisher, C. R., & Shank, T. M. (2015). Variation in the diets of hydrothermal vent gastropods *Deep Sea Research Part II: Topical Studies in Oceanography*, 121, 193 - 201.
206. Govenar, B., Le Bris, N., Gollner, S., Glanville, J., Aperghis, A. B., Hourdez, S., & Fisher, C. R. (2005). Epifaunal community structure associated with *Riftia pachyptila* aggregations in chemically different hydrothermal vent habitats. *Marine Ecology Progress Series*, 305, 67 - 77.
207. Grelon, D., Morineaux, M., Desrosiers, G., & Juniper, S. (2006). Feeding and territorial behavior of *Paralvinella sulfincola*, a polychaete worm at deep-sea hydrothermal vents of the Northeast Pacific Ocean. *Journal of Experimental Marine Biology and Ecology*, 329(2), 174 - 186. doi:10.1016/j.jembe.2005.08.017
208. Guinot, D., & Hurtado, L. A. (2003). Two new species of hydrothermal vent crabs of the genus *Bythograea* from the southern East Pacific Rise and from the Galapagos Rift (Crustacea Decapoda Brachyura Bythograeidae). *Comptes rendus biologies*, 326(4), 423 - 439. doi: 10.1016/S1631-0691(03)00126-4
209. Guri, M., Durand, L., Cueff-Gauchard, V., Zbinden, M., Crassous, P., Shillito, B., & Cambon-Bonavita, M. (2012). Acquisition of epibiotic bacteria along the life cycle of the hydrothermal shrimp *Rimicaris exoculata*. *The Isme Journal*, 6, 597 - 609.
210. *Gustafson, R. G., & Lutz, R. A. (1994). Molluscan life history traits at deep-sea hydrothermal vents and cold methane/sulfide seeps. *Reproduction, Larval Biology, and Recruitment of the Deep-sea Benthos*, Columbia University Press, NY, 76 - 97.
211. Harasewych, M. G., & Kantor, Y. I. (2002). *Buccinum thermophilum* (Gastropoda: Neogastropoda: Buccinidae), a new species from the Endeavour vent field of the Juan de Fuca Ridge. *Journal of Molluscan Studies*, 68(1), 39 - 44. doi:10.1093/mollus/68.1.39
212. *Hashimoto, J., Fujikura, K., & Hotta, H. (1989). Deep-sea communities associated with warm seep at the Minami-Ensei Knoll. *Preprint for 6th Symposium on Research by the Shinkai 2000*, 77 - 79.
213. Hashimoto, J., & Furuta, M. (2007). A new species of *Bathymodiolus* (Bivalvia: Mytilidae) from hydrothermal vent communities in the Manus Basin, Papua New Guinea. *Venus*, 66(1 - 2), 57 - 68.
214. Hashimoto, J., & Yamane, T. (2005). A new species of *Gigantidas* (Bivalvia: Mytilidae) from a vent site on the Kaikata Seamount Southwest of the Ogasawara (Bonin) Islands, Southern Japan. *Venus*, 64(1- 2), 1 - 10.
215. *Henry, M. S., Childress, J. J., & Figueroa, D. (2008). Metabolic rates and thermal

tolerances of chemoautotrophic symbioses from Lau Basin hydrothermal vents and their implications for species distributions. *Deep Sea Research Part I: Oceanographic Research Papers*, 55(5), 679 - 695. doi:10.1016/j.dsr.2008.02.001

216. Heptner, M. V. (1968). Description and functional morphology of *Megapontius pleurospinosis* sp. n. from the Pacific with some remarks on the status (Copepoda, Cyclopoida). *Zoologicheskii Zhurnal*, 47(11), 197 - 206.
217. Heptner, M. V., & Ivanenko, V. N. (2002). Copepoda (Crustacea) of hydrothermal ecosystems of the World Ocean. *Arthropoda Selecta*, 11(2), 117 - 134.
218. *Hernández-Ávila, I., Cambron-Bonavita, M.-A., & Pradillon, F. (2016). Morphology of First Zoal Stage of Four Genera of Alvinocaridid Shrimps from Hydrothermal Vents and Cold Seeps: Implications for Ecology, Larval Biology and Phylogeny. *PLoS One*, 10(12), e0144657. doi:10.1371/journal.pone.0144657
219. *Herrera, S., Watanabe, H., & Shank, T. M. (2015). Evolutionary and biogeographical patterns of barnacles from deep-sea hydrothermal vents. *Molecular Ecology*, 24(3), 673 - 689. doi:10.1111/mec.13054
220. Hertz, E., Robinson, J. P. W., Trudel, M., Mazumder, A., & Baum, J. K. (2014). Estimation of predator-prey mass ratios using stable isotopes: sources of errors. *Marine Ecology Progress Series*, 516, 1 - 6.
221. Hessler, R. R. (1984). *Dahlella Caldariensis*, New Genus, New Species: a Leptostracan (Crustacea, Malacostraca) from Deep-sea Hydrothermal Vents. *Journal of Crustacean Biology*, 4(4), 655 - 664.
222. Hessler, R. R., & Martin, J. W. (1989). *Austinograea williamsi*, New Genus, New Species, a Hydrothermal Vent Crab (Decapoda: Bythograeidae) from the Mariana Back-arc Basin, Western Pacific. *Journal of Crustacean Biology*, 9(4), 645 - 661. doi:10.1163/193724089X00647
223. Hestetun, J. T., Dahle, H., Jørgensen, S. L., Olsen, B. R., & Rapp, H. T. (2016). The Microbiome and Occurrence of Methanotrophy in Carnivorous Sponges. *Frontiers in Microbiology*, 7(1781). doi:10.3389/fmicb.2016.01781
224. *Hestetun, J. T., Fourt, M., Vacelet, J., Boury-Esnault, N., & Rapp, H. T. (2013). Cladorhizidae (Porifera, Demospongiae, Poecilosclerida) of the deep Atlantic collected during Ifremer cruises, with a biogeographic overview of the Atlantic species. *Journal of the Marine Biological Association of the United Kingdom*, 95(7), 1311 - 1342. doi:10.1017/S0025315413001100
225. Hestetun, J. T., Tompkins-Macdonald, G., & Rapp, H. T. (2017). A review of carnivorous sponges (Porifera: Cladorhizidae) from the Boreal North Atlantic and

226. *Hestetun, J. T., Vacelet, J., Boury-Esnault, N., Borchiellini, C., Kelly, M., Ríos, P., . . . Rapp, H. T. (2016). The systematics of carnivorous sponges. *Molecular Phylogenetics and Evolution*, 94, 327 - 345. doi: 10.1016/j.ympev.2015.08.022
227. Heywood, J. L., Chen, C., Pearce, D. A., & Linse, K. (2017). Bacterial communities associated with the Southern Ocean vent gastropod, *Gigantopelta chessoia*: indication of intergenerational, horizontal symbiont transfer. *Polar Biology*, 40(11), 2335 - 2342. doi:10.1007/s00300-017-2148-6
228. Hicks, G. R. F. (1988). Harpacticoid copepods from biogenic substrata in offshore waters of New Zealand. 1: New species of *Paradactylopodia*, *Stenhelia* (St.) and *Laophonte*. *Journal of the Royal Society of New Zealand*, 18(4), 437 - 452.
229. Hidaka, H., Ogura, T., Watanabe, H., Kano, Y., & Kojima, S. (2015). Population History of a Hydrothermal Vent-Endemic Gastropod *Alviniconcha hessleri* in the Mariana Trough. 325 - 333. doi:10.1007/978-4-431-54865-2_26
230. *Hilário, A., & Cunha, M. R. (2013). Notes on a mating event of the deep-sea crab *Chaecon affinis* in the Gorringe Bank (NE Atlantic). *Deep Sea Research Part II: Topical Studies in Oceanography*, 92(Supplement C), 58 - 62. doi: 10.1016/j.dsr2.2013.01.021
231. Hilário, A., Vilar, S., Cunha, M. R., & Tyler, P. A. (2009). Reproductive aspects of two bythograeid crab species from hydrothermal vents in the Pacific-Antarctic Ridge. *Marine Ecology Progress Series*, 378, 153 - 160.
232. *Hilário, A., Young, C. M., & Tyler, P. A. (2005). Sperm Storage, Internal Fertilization, and Embryonic Dispersal in Vent and Seep Tubeworms (Polychaeta: Siboglinidae: Vestimentifera). *The Biological Bulletin*, 208(1), 20 - 28. doi:10.2307/3593097
233. Horton, T., & Thurston, M. H. (2014). A revision of the bathyal and abyssal necrophage genus *Cyclocaris* Stebbing, 1888 (Crustacea: Amphipoda: Cyclocaridae) with the addition of two new species from the Atlantic Ocean. *Zootaxa*, 3796(3), 507 - 527.
234. Hourdez, S., & Desbruyères, D. (2003). A new species of scale-worm (Polychaeta: Polynoidae), *Levensteiniella iris* sp. nov., from the Rainbow and Lucky Strike vent fields (Mid-Atlantic Ridge). *Cahiers Biologie Marine*, 44, 13 - 21.
235. *Hourdez, S., & Jouin-Toulmond, C. (1998). Functional anatomy of the respiratory system of *Branchipolynoe* species (Polychaeta, Polynoidae), commensal with *Bathymodiolus* species (Bivalvia, Mytilidae) from deep-sea hydrothermal vents. *Zoomorphology*, 118(4), 225 - 233. doi:10.1007/s004350050071

236. Hügler, M., Petersen, J. M., Dubilier, N., Imhoff, J. F., & Sievert, S. M. (2011). Pathways of Carbon and Energy Metabolism of the Epibiotic Community Associated with the Deep-Sea Hydrothermal Vent Shrimp *Rimicaris exoculata*. *PLoS One*, 6(1), e16018. doi:10.1371/journal.pone.0016018
237. Humes, A. G. (1999a). *Collocherides brychius*, a new species (Copepoda: Siphonostomatoida: Astrocheridae) from a deep-water hydrothermal site in the northeastern Pacific. *Proceedings of The Biological Society of Washington*, 112(1), 181 - 188.
238. Humes, A. G. (1999b). Copepoda (Cyclopinidae and Misophriidae) from a deep-sea hydrothermal site in the northeastern Pacific. *Journal of Natural History*, 33(7), 961 - 978. doi:10.1080/002229399300038.
239. Humes, A. G. (1996). Deep-Sea Copepoda (Siphonostomatoida) from Hydrothermal Sites on the Mid-Atlantic Ridge at 23° and 37°N. *Bulletin of Marine Science*, 58(3), 609 - 653.
240. *Humes, A. G. (1991). Siphonostomatoid copepods from a deep-water hydrothermal zone in the Lau Basin, South Pacific. *Bulletin du Muséum National d'Histoire Naturelle, Paris*, 4(13, sect. A (1-2)), 121 - 134.
241. *Humes, A. G. (1990). Copepods (Siphonostomatoida) from a deep-sea hydrothermal vent at the Mariana Back-Arc Basin in the Pacific, including a new genus and species. *Journal of Natural History*, 24(2), 289 - 304.
242. Humes, A. G. (1989). Copepoda from deep-sea hydrothermal vents at the East Pacific Rise. *Bulletin du Muséum National d'Histoire Naturelle. Section A, Zoologie, Biologie et Ecologie Animales*, 11(4), 829 - 849.
243. Humes, A. G. (1987). Copepoda from deep-sea hydrothermal vents. *Bulletin of Marine Science*, 41(3), 645 - 788.
244. Humes, A. G., & Lutz, R. A. (1994). *Aphotopontius acanthinus*, New Species (Copepoda: Siphonostomatoida), from Deep-Sea Hydrothermal Vents on the East Pacific Rise. *Journal of Crustacean Biology*, 14(2), 337 - 345.
245. Humes, A. G., & Segonzac, M. (1998). Copepoda from deep-sea hydrothermal sites and cold seeps: description of a new species of *Aphotopontius* from the East Pacific Rise and general distribution. *Cahiers de Biologie Marine*, 39, 51 - 62.
246. Hurtado, L. A., Lutz, R. A., & Vrijenhoek, R. C. (2004). Distinct patterns of genetic differentiation among annelids of eastern Pacific hydrothermal vents. *Molecular Ecology*, 13(9), 2603 - 2615.

247. Hurtado, L. A., Mateos, M., Lutz, R. A., & Vrijenhoek, R. C. (2003). Coupling of Bacterial Endosymbiont and Host Mitochondrial Genomes in the Hydrothermal Vent Clam *Calyptogena magnifica*. *Applied and Environmental Microbiology*, 69(4), 2058 - 2064.
248. Husson, B., Sarradin, P.-M., Zeppilli, D., & Sarrazin, J. (2017). Picturing thermal niches and biomass of hydrothermal vent species. *Deep Sea Research Part II: Topical Studies in Oceanography*, 137, 6 - 25. doi:10.1016/j.dsr2.2016.05.028
249. Ikuta, T., Takaki, Y., Nagai, Y., Shimamura, S., Tsuda, M., Kawagucci, S., . . . Yoshida, T. (2015). Heterogeneous composition of key metabolic gene clusters in a vent mussel symbiont population. *The Isme Journal*, 10, 990. doi:10.1038/ismej.2015.176
250. Imhoff, J. F., Sahling, H., Süling, J., & Kath, T. (2003). 16S rDNA-based phylogeny of sulphur-oxidising bacterial endosymbionts in marine bivalves from cold-seep habitats. *Marine Ecology Progress Series*, 249, 39 - 51.
251. *Ise, Y., & Vacelet, J. (2010). New Carnivorous Sponges of the Genus Abyssocladia (Demospongiae, Poecilosclerida, Cladorhizidae) from Myojin Knoll, Izu-Ogasawara Arc, Southern Japan. *Zoological Science*, 27(11), 888 - 894. doi:10.2108/zsj.27.888
252. *Ishibashi, J., Okino, K., & Sunamura, M. (2015). *Subseafloor Biosphere Linked to Hydrothermal Systems*. Tokyo: Springer.
253. *Ivanenko, V. N., Defaye, D., Segonzac, M., Khripounoff, A., Sarrazin, J., & Ferrari, F. D. (2011). A new species of *Exrima*, synonymy of four species of *Aphotopontius*, *Stygiopontius* and *Rhogobius*, and record of first copepodid stage of Dirivultidae (Copepoda: Siphonostomatoida) from deep-sea hydrothermal vents of the East Pacific Rise (13°N). *Journal of the Marine Biological Association of the United Kingdom*, 91(7), 1547 - 1559. doi:10.1017/S0025315411000178
254. *Ivanenko, V. N., Martínez Arbizu, P., & Stecher, J. (2007). Lecithotrophic nauplius of the family Dirivultidae (Copepoda; Siphonostomatoida) hatched on board over the Mid-Atlantic Ridge (5°S). *Marine Ecology*, 28(1), 49 - 53. doi:10.1111/j.1439-0485.2007.00142.x
255. *Japan Agency for Marine-Earth Science and Technology. (2009 onwards). Biological Information System for Marine Life. Retrieved from: <http://www.godac.jamstec.go.jp/bismal/>
256. Jan, C., Petersen, J. M., Werner, J., Teeling, H., Huang, S., Glöckner, F. O., . . . Cambon-Bonavita, M.-A. (2014). The gill chamber epibiosis of deep-sea shrimp *Rimicaris exoculata*: an in-depth metagenomic investigation and discovery of Zetaproteobacteria. *Environ Microbiol*, 16(9), 2723 - 2738. doi:10.1111/1462-2920.12406

257. Jeanthon, C. (2000). Molecular ecology of hydrothermal vent microbial communities. *Antonie van Leeuwenhoek*, 77, 117 - 133.
258. *Jenkins, C. D., Ward, M. E., Turnipseed, M., Osterberg, J., & Van Dover, C. L. (2002). The digestive system of the hydrothermal vent polychaete *Galapagomystides aristata* (Phyllodocidae): evidence for hematophagy? *Invertebrate Biology*, 121(3), 243 - 254. doi:10.1111/j.1744-7410.2002.tb00064.x
259. Johnson, S. B., Warén, A., Tunnicliffe, V., Van Dover, C. L., Wheat, C. G., Schultz, T. F., & Vrijenhoek, R. C. (2015). Molecular taxonomy and naming of five cryptic species of *Alviniconcha* snails (Gastropoda: Abyssochrysoidea) from hydrothermal vents. *Systematics and Biodiversity*, 13(3), 278 - 295.
260. Johnson, S. B., Warén, A., & Vrijenhoek, R. C. (2008). DNA Barcoding of *Lepetodrilus* Limpets Reveals Cryptic Species. *Journal of Shellfish Research*, 27(1), 43 - 51. doi:10.2983/0730-8000(2008)27[43:DBOLLR]2.0.CO;2.
261. Johnson, S. B., Won, Y.-J., Harvey, J. B. J., & Vrijenhoek, R. C. (2013). A hybrid zone between *Bathymodiolus* mussel lineages from eastern Pacific hydrothermal vents. *BMC Evolutionary Biology*, 13(21), 1 - 18. doi:10.1186/1471-2148-13-21.
262. *Jollivet, D., Desbruyères, D., Ladrat, C., & Laubier, L. (1995). Evidence for differences in the allozyme thermostability of deep-sea hydrothermal vent polychaetes (Alvinellidae): a possible selection by habitat. *Marine Ecology Progress Series*, 123, 125 - 136.
263. Jones, W. J., & Macpherson, E. (2007). Molecular Phylogeny of the East Pacific Squat Lobsters of the Genus *Munidopsis* (Decapoda: Galatheidae) with the Descriptions of Seven New Species. *Journal of Crustacean Biology*, 27(3), 477 - 501. doi:10.1651/S-2791.1
264. Jumars, P. A., Dorgan, K. M., & Lindsay, S. M. (2015). Diet of Worms Emended: An Update of Polychaete Feeding Guilds. *Annual Review of Marine Science*, 7, 497 - 520.
265. Juniper, S. K. (1994). Ecology and biogeochemistry of *Paralvinella sulfincola* at northeast Pacific hydrothermal vents: review and comparison with *Alvinella* spp. of the east Pacific rise. *Mémoires du Muséum national d'histoire naturelle*, 162, 453 - 462.
266. Juniper, S. K., Jonasson, I. R., Tunnicliffe, V., & Southward, A. J. (1992). Influence of a tube-building polychaete on hydrothermal chimney mineralization. *Geology*, 20, 895 - 898.
267. Juniper, S. K., Tunnicliffe, V., & Southward, E. C. (1992). Hydrothermal vents in turbidite sediments on a Northeast Pacific spreading centre: organisms and substratum at an ocean drilling site. *Canadian Journal of Zoology*, 70(9), 1792 - 1809.

doi:10.1139/z92-247

268. *Kamenev, G. M. (2009). North Pacific Species of the Genus *Solemya* Lamarck, 1818 (Bivalvia: Solemyidae), with notes on *Acharax johnsoni* (Dall, 1891). *Malacologia*, 51(2), 233 - 261.
269. Kamenev, G. M., Nadtochy, V. A., & Kuznetsov, A. P. (2001). *Conchocele bisecta* (Conrad, 1849) (Bivalvia : Thyasiridae) from cold-water methane-rich areas of the Sea of Okhotsk. *Veliger*, 44, 84 - 94.
270. Kano, Y., Kimura, S., Kimura, T., & Warén, A. (2012). Living Monoplacophora: morphological conservatism or recent diversification? *Zoologica Scripta*, 41(5), 471 - 488.
271. Karaseva, N. P., Rimskaya-Korsakova, N. N., Galkin, S. V., & Malakhov, V. V. (2016). Taxonomy, geographical and bathymetric distribution of vestimentiferan tubeworms (Annelida, Siboglinidae). *Biology Bulletin*, 43(9), 937 - 969.
doi:10.1134/S1062359016090132
272. Karl, D. M., Christian, J. R., Dore, J. E., Hebel, D. V., Letelier, R. M., Tupas, L. M., & Winn, C. D. (1996). Seasonal and interannual variability in primary production and particle flux at Station ALOHA. *Deep Sea Research Part II: Topical Studies in Oceanography*, 43(2 - 3), 539 - 568.
273. Kelley, D., & Shank, T. (2010). Hydrothermal Systems: A Decade of Discovery in Slow Spreading Environments. In P. A. Rona, C. W. Devey, J. Dyment, & B. J. Murton (Eds.), *Diversity of Hydrothermal Systems on Slow Spreading Ocean Ridges*. Washington, D.C.: American Geophysical Union.
274. *Kelly, N. E., & Metaxas, A. (2006). Recruitment patterns of invertebrates at anhydrite hydrothermal vents on the Juan de Fuca Ridge, NE Pacific. *Cahiers de Biologie Marine*, 47, 435 - 441.
275. *Kenk, V. C., & Wilson, B. R. (1985). A new mussel (Bivalvia, Mytilidae) from hydrothermal vents in the Galapagos Rift zone. *Malacologia*, 26(1 - 2), 253 - 271.
276. *Kharlamenko, V. I., Kamenev, G. M., Kalachev, A. V., Kiyashko, S. I., & Ivin, V. V. (2016). Thyasirid bivalves from the methane seep community off Paramushir Island (Sea of Okhotsk) and their nutrition. *Journal of Molluscan Studies*, 82(3), 391 - 402.
doi:10.1093/mollus/eyw004
277. Kiel, S. (2010). *The Vent and Seep Biota: Aspects from Microbes to Ecosystems* (Vol. 33). Dordrecht: Springer.

278. Kikuchi, T., & Ohta, S. (1995). Two Caridean Shrimps of the Families Bresiliidae and Hippolytidae from a Hydrothermal Field on the Iheya Ridge, off the Ryukyu Islands, Japan. *Journal of Crustacean Biology*, 15(4), 771 - 785.
279. *Kim, E. S., Sakai, H., Hashimoto, J., Yanagisawa, F., & Ohta, S. (1989). Sulfur isotopic ratios of hydrothermal vent-animals at Ogasawara Arc and Mid-Okinawa Trough-evidence for microbial origin of hydrogen sulfide at low-temperature submarine hydrothermal areas. *Geochemical Journal*, 23(4), 195 - 208.
doi:10.2343/geochemj.23.195
280. Kim, Y. W., Yasuda, M., Yamagishi, A., Oshima, T., & Ohta, S. (1995). Characterization of the endosymbiont of a deep-sea bivalve, *Calyptogena soyoae*. *Applied and Environmental Microbiology*, 61(2), 823 - 827.
281. King, N. J., Bagley, P. M., & Priede, I. G. (2006). Depth zonation and latitudinal distribution of deep-sea scavenging demersal fishes of the Mid-Atlantic Ridge, 42 to 53°N. *Marine Ecology Progress Series*, 319, 263 - 274.
282. *Knowles, J. D., Wenink, E., Schult, N., Tunnicliffe, V., & McHugh, D. (2005). Molecular analysis indicates gene flow among populations of *Paralvinella pandorae* Desbruyeres and Laubier 1986 (Alvinellidae, Terebellida), a polychaete annelid endemic to hydrothermal vents of the northeast Pacific. *Marine Ecology*, 26(3-4), 216 - 222. doi:10.1111/j.1439-0485.2005.00063.x.
283. Kojima, S., Tsuchida, E., Numanami, H., Fujikura, K., & Okutani, T. (2006) Synonymy of *Calyptogena solidissima* with *Calyptogena kawamurai* (Bivalvia: Vesicomyidae) and its population structure revealed by mitochondrial DNA sequences. *Zool. Sci.*, 23, 835 - 842.
284. No reference is associated with this number (as the reference originally included here was a duplicate). This number can therefore be used as a reference code in future versions of sFDvent.
285. Kojima, S., Fujikura, K., & Okutani, T. (2004). Multiple trans-Pacific migrations of deep-sea vent/seep-endemic bivalves in the family Vesicomyidae. *Molecular Phylogenetics and Evolution*, 32(1), 396 - 406.
286. *Kojima, S., & Ohta, S. (1997). *Calyptogena okutani* n. sp., a sibling species of *Calyptogena soyoae* Okutani, 1957 (Bivalvia: Vesicomyidae). *Venus*, 56(3), 189 - 195.
287. Kojima, S., Ohta, S., Yamamoto, T., Miura, T., Fujiwara, Y., Fujikura, K., & Hashimoto, J. (2002). Molecular taxonomy of vestimentiferans of the western Pacific and their phylogenetic relationship to species of the eastern Pacific. *Mar Biol*, 141(1), 57 - 64.
doi:10.1007/s00227-002-0818-5

288. *Kojima, S., Segawa, R., Fujiwara, Y., Fujikura, K., Ohta, S., & Hashimoto, J. (2001). Phylogeny of Hydrothermal-Vent–Endemic Gastropods *Alviniconcha* spp. from the Western Pacific Revealed by Mitochondrial DNA Sequences. *The Biological Bulletin*, 200(3), 298 - 304. doi:10.2307/1543511
289. Kojima, S., & Watanabe, H. (2015). Vent Fauna in the Mariana Trough. In J.-i. Ishibashi, K. Okino, & M. Sunamura (Eds.), *Subseafloor Biosphere Linked to Hydrothermal Systems: TAIGA Concept* (pp. 313-323). Tokyo: Springer Japan.
290. Komai, T., & Segonzac, M. (2008). Taxonomic review of the hydrothermal vent shrimp genera *Rimicaris* Williams and Rona and *Chorocaris* Martin and Hessler (Crustacea: Decapoda: Caridea: Alvinocarididae). *Journal of Shellfish Research*, 27(1), 21 - 41.
291. Komai, T., & Segonzac, M. (2005). A revision of the genus *Alvinocaris* Williams and Chace (Crustacea: Decapoda: Caridea: Alvinocarididae), with descriptions of a new genus and a new species of *Alvinocaris*. *Journal of Natural History*, 39(15), 1111 - 1175.
292. Komai, T., & Segonzac, M. (2004). A new genus and species of alvinocaridid shrimp (Crustacea: Decapoda: Caridea) from hydrothermal vents on the North Fiji and Lau Basins, south-western Pacific. *Journal of the Marine Biological Association of the U.K.*, 84, 1179 - 1188.
293. *Komai, T., & Tsuchida, S. (2015). New records of Alvinocarididae (Crustacea: Decapoda: Caridea) from the southwestern Pacific hydrothermal vents, with descriptions of one new genus and three new species. *Journal of Natural History*, 49(29 - 30), 1789 - 1824.
294. Komai, T., Tsuchida, S., & Segonzac, M. (2012). Records of species of the hippolytid genus *Lebbeus* White, 1847 (Crustacea: Decapoda: Caridea) from hydrothermal vents in the Pacific Ocean, with descriptions of three new species. *Zootaxa*, 3241, 35 - 63.
295. Komai, T., Martin, J. W., Zala, K., Tsuchida, S., & Hashimoto, J. (2006). A new species of *Mirocaris* (Crustacea: Decapoda: Caridea: Alvinocarididae) associated with hydrothermal vents on the Central Indian Ridge, Indian Ocean. *Scientia Marina*, 70, 109 - 119.
296. Kongsrud, J. A., Eilertsen, M. H., Alvestad, T., Kongshavn, K., & Rapp, H. T. (2017). New species of Ampharetidae (Annelida: Polychaeta) from the Arctic Loki Castle vent field. *Deep Sea Research Part II: Topical Studies in Oceanography*, 137, 232 - 245.
297. Kongsrud, J. A., & Rapp, H. T. (2012). *Nicomache (Loxochona) lokii* sp. nov. (Annelida: Polychaeta: Maldanidae) from the Loki's Castle vent field: an important structure builder in an Arctic vent system. *Polar Biology*, 35(2), 161 - 170.

298. *Kornicker, L. S. (1991). *Myodocopid Ostracoda of Hydrothermal Vents in the Eastern Pacific Ocean* (Vol. 516). Washington D.C.: Smithsonian Institution Press.
299. Kornicker, L. S., & Harrison-Nelson, E. (2005). Two new species of Ostracoda from hydrothermal vents of *Riftia pachyptila* aggregations on the East Pacific Rise (Halocypridina; Cladocopina). *Zootaxa*(1071), 19 - 38.
300. Kouris, A., Limén, H., Stevens, C. J., & Juniper, S. K. (2010). Blue mats: faunal composition and food web structure in colonial ciliate (*Folliculinopsis* sp.) mats at Northeast Pacific hydrothermal vents. *Marine Ecology Progress Series*, 412, 93 - 101.
301. Krylova, E. M., Sahling, H. & Janssen, R. (2010). *Abyssogena*: a new genus of the family Vesicomyidae (Bivalvia) from deep-water vents and seeps. *Journal of Molluscan Studies*, 76(2), 107 - 132.
302. Krylova, E. M., & Janssen, R. (2006). Vesicomyidae from Edison Seamount (South West Pacific): Papua New Guinea: New Ireland fore-arc basin) (Bivalvia: Glossoidea). *Archiv für Molluskenkunde*, 135(2), 231 - 261.
303. *Krylova, E. M., Kamenev, G. M., Vladychenskaya, I. P., & Petrov, N. B. (2015). Vesicomyinae (Bivalvia: Vesicomyidae) of the Kuril–Kamchatka Trench and adjacent abyssal regions. *Deep Sea Research Part II: Topical Studies in Oceanography*, 111, 198 - 209. doi:10.1016/j.dsrr.2014.10.004
304. Krylova, E. M., & Moskalev, L. I. (1996). *Ectenagena extenta*, a new species of vesicomyid bivalve from Monterey Bay, California. *Ruthenica*, 6(1), 1 - 10.
305. Krylova, E. M., & Sahling, H. (2006). Recent bivalve molluscs of the genus *Calyptogena* (Vesicomyidae). *Journal of Molluscan Studies*, 72, 359 - 395.
306. Kuwahara, H., Yoshida, T., Takaki, Y., Shimamura, S., Nishi, S., Harada, M., . . . Maruyama, T. (2007). Reduced Genome of the Thioautotrophic Intracellular Symbiont in a Deep-Sea Clam, *Calyptogena okutanii*. *Current Biology*, 17(10), 881 - 886. doi:10.1016/j.cub.2007.04.039
307. *Le Pennec, M., & Beninger, P. G. (2000). Reproductive characteristics and strategies of reducing-system bivalves. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 126(1), 1 - 16. doi: 10.1016/S0742-8413(00)00100-6
308. Le Pennec, M., Beninger, P. G., & Herry, A. (1995). Feeding and digestive adaptations of bivalve molluscs to sulphide-rich habitats. *Comparative Biochemistry and Physiology*, 111A, 183 - 189.

309. Leavitt, W. D., Venceslau, S. S., Pereira, I. A. C., Johnston, D. T., & Bradley, A. S. (2016). Fractionation of sulfur and hydrogen isotopes in *Desulfovibrio vulgaris* with perturbed DsrC expression. *FEMS Microbiology Letters*, 363(20), 1 - 8.
310. Ledru, S. (2000). *Trophic niche and nutritional biochemistry of invertebrates inhabiting East Pacific Rise hydrothermal vent sites*. (Magistère de physico-chimie Moléculaire), Université Paris-Sud.
311. *Lee, R. W., Robert, K., Matabos, M., Bates, A. E., & Juniper, S. K. (2015). Temporal and spatial variation in temperature experienced by macrofauna at main endeavour Hydrothermal vent field. *Deep Sea Research Part I: Oceanographic Research Papers*, 106, 154 - 166.
312. *Lee, W., & Huys, R. (1999). *Bathylaophonte* gen. nov. from deep-sea hydrothermal vents and the polyphyly of Paronychocamptus (Copepoda: Harpacticoida). *Cahiers de Biologie Marine*, 40(4), 293 - 328.
313. Lelievre, Y., Legendre, P., Matabos, M., Mihaly, S., Lee, R. W., Sarradin, P. M., . . . Sarrazin, J. (2017). Astronomical and atmospheric impacts on deep-sea hydrothermal vent invertebrates. *Proceedings. Biological Sciences*, 284(1852). doi:10.1098/rspb.2016.2123
314. Léveillé, R. J., Levesque, C., & Juniper, S. K. (2005). Biotic interactions and feedback processes in deep-sea hydrothermal vent ecosystems. In E. Kristensen, R. R. Haese, & J. E. Kostka (Eds.), *Interactions Between Macro- and Microorganisms in Marine Sediments* (Vol. 60, pp. 299-321). Washington DC: American Geophysical Union.
315. Levesque, C., Kim Juniper, S., & Limén, H. (2006). Spatial organization of food webs along habitat gradients at deep-sea hydrothermal vents on Axial Volcano, Northeast Pacific. *Deep Sea Research Part I: Oceanographic Research Papers*, 53(4), 726 - 739. doi:10.1016/j.dsr.2006.01.007
316. Levesque, C., Juniper, S. K., & Marcus, J. (2003). Food resource partitioning and competition among alvinellid polychaetes of Juan de Fuca Ridge hydrothermal vents. *Marine Ecology Progress Series*, 246, 173 - 182.
317. Levin, L. A., Mendoza, G. F., Konotchick, T., & Lee, R. (2009). Macrobenthos community structure and trophic relationships within active and inactive Pacific hydrothermal sediments. *Deep Sea Research Part II: Topical Studies in Oceanography*, 56, 1632 - 1648.
318. *Levinsen, G. M. R. (1887). Kara-Havets Svampe (Porifera). *Djympha-Togtets zoologisk-botaniske Udbytte*, 1, 339 - 372.
319. Lewis, K. B., & Marshall, B. A. (1996). Seep faunas and other indicators of methane-

rich dewatering on New Zealand convergent margins. *New Zealand Journal of Geology and Geophysics*, 39(2), 181 - 200.

320. Li, X. (2015). Report on two deep-water caridean shrimp species (Crustacea: Decapoda: Caridea: Alvinocarididae, Acanthephyridae) from the northeastern South China Sea. *Zootaxa*, 3911(1), 130 - 138.
321. Limén, H., & Juniper, S. K. (2006). Habitat controls on vent food webs at Eifuku Volcano, Mariana Arc. *Cahiers de Biologie Marine*, 47(4), 449 - 455.
322. Limén, H., Juniper, S. K., Tunnicliffe, V., & Clément, M. (2006). Benthic community structure on two peaks of an erupting seamount: Northwest Rota-1 Volcano, Mariana Arc, western Pacific. *Cahiers de Biologie Marine*, 47, 457 - 464.
323. Limén, H., Levesque, C., & Juniper, S. K. (2007). POM in macro-/meiofaunal food webs associated with three flow regimes at deep-sea hydrothermal vents on Axial Volcano, Juan de Fuca Ridge. *Marine Biology*, 153(2), 129 - 139.
324. Limén, H., Stevens, C. J., Bourass, Z., & Juniper, S. K. (2008). Trophic ecology of siphonostomatoid copepods at deep-sea hydrothermal vents in the northeast Pacific. *Marine Ecology Progress Series*, 359, 161 - 170.
325. *Lin, H. Y., Lin, P. Y., Chang, N. N., Shiao, J. C., & Kao, S. J. (2014). Trophic structure of megabenthic food webs along depth gradients in the South China Sea and off northeastern Taiwan. *Marine Ecology Progress Series*, 501, 53 - 66.
326. *Lisin, S. E., Hannan, E. E., Kockevar, R. E., Harrold, C., & Barry, J. P. (1997). Temporal variation in gametogenic cycles of vesicomyid clams. *Invertebrate Reproduction & Development*, 31, 307 - 318.
327. López-García, P., Gaill, F., & Moreira, D. (2002). Wide bacterial diversity associated with tubes of the vent worm *Riftia pachyptila*. *Environmental Microbiology*, 4(4), 204 - 215. doi:10.1046/j.1462-2920.2002.00286.x
328. López-Gonzalez, P. J., Rodríguez, E., Gili, J.-M., & Segonzac, M. (2003). New records on sea anemones (Anthozoa: Actiniaria) from hydrothermal vents and cold seeps. *Zoologische verhandelingen / uitgegeven door het Rijksmuseum van Natuurlijke Historie te Leiden (Ministerie van Cultuur, Recreatie en Maatschappelijk Werk)*, 345, 215 - 243.
329. Lorion, J., Kiel, S., Faure, B., Kawato, M., Ho, S. Y. W., Marshall, B., . . . Fujiwara, Y. (2013). Adaptive radiation of chemosymbiotic deep-sea mussels. *Proceedings of the Royal Society B: Biological Sciences*, 280(1770).

330. Lundbeck, W. (1905). Porifera. (Part II.) Desmacidonidae (pars.). *The Danish Ingolf-Expedition*, 6(2), 183 - 185.
331. Lundsten, L., Reiswig, H. M., & Austin, W. C. (2014). Four new species of Cladorhizidae (Porifera, Demospongiae, Poecilosclerida) from the Northeast Pacific. *Zootaxa*, 3786, 101 - 123.
332. *Lutz, R. A., Jablonski, D., & Turner, R. D. (1984). Larval Development and Dispersal at Deep-Sea Hydrothermal Vents. *Science*, 226(4681), 1451 - 1454.
333. Macpherson, E., Jones, W. J., & Segonzac, M. (2005). A new squat lobster family of Galatheoidea (Crustacea, Decapoda, Anomura) from the hydrothermal vents of the Pacific-Antarctic Ridge. *Zoosystema*, 27(4), 709 - 723.
334. Maddocks, R. F. (2005). Three New Species of Podocopid Ostracoda from Hydrothermal Vent Fields at 9°50'N on the East Pacific Rise. *Micropaleontology*, 51(5), 345 - 372.
335. Mah, C., Linse, K., Copley, J., Marsh, L., Rogers, A., Clague, D., & Foltz, D. (2015). Description of a new family, new genus, and two new species of deep-sea Forcipulatacea (Asteroidea), including the first known sea star from hydrothermal vent habitats. *Zoological Journal of the Linnean Society*, 174(1), 93 - 113.
336. Marcus, J. (2003). *Community ecology of hydrothermal vents at Axial Volcano, Juan de Fuca Ridge, northeast Pacific*. (Doctor of Philosophy Doctor of Philosophy), University of Victoria, Victoria, Canada.
337. Marcus, J., & Tunnicliffe, V. (2002). Living on the edges of diffuse vents on the Juan de Fuca Ridge. *Cahiers de Biologie Marine*, 43, 263 - 266.
338. Marques, A., Porteiro, F., & Schaefer, S. A. (2000). Hydrothermal Vent Mussel Bathymodiolus sp. (Mollusca: Mytilidae): Diet Item of Hydrolagus affinis (Pisces: Chimaeridae). *Copeia*, 2000(3), 806 - 807. doi:10.1643/0045-8511(2000)000[0806:HVMBSM]2.0.CO;2
339. *Marsh, A. G., Mullineaux, L. S., Young, C. M., & Manahan, D. T. (2001). Larval dispersal potential of the tubeworm *Riftia pachyptila* at deep-sea hydrothermal vents. *Nature*, 411, 77 - 80.
340. Marsh, L., Copley, J. T., Huvenne, V. A., Linse, K., Reid, W. D., Rogers, A. D., . . . Tyler, P. A. (2012). Microdistribution of faunal assemblages at deep-sea hydrothermal vents in the Southern Ocean. *PLoS One*, 7(10), e48348. doi:10.1371/journal.pone.0048348

341. Marsh, L., Copley, J. T., Tyler, P. A., Thatje, S., & Webb, T. (2015). In hot and cold water: differential life-history traits are key to success in contrasting thermal deep-sea environments. *Journal of Animal Ecology*. doi:10.1111/1365-2656.12337
342. Martell, K. A., Tunnicliffe, V., & Macdonald, I. R. (2002). Biological features of a buccinid whelk (Gastropoda, Neogastropoda) at the Endeavour ventfields of Juan de Fuca Ridge, northeast Pacific. *Journal of Molluscan Studies*, 68, 45 - 53.
343. Martin, J. W., & Christiansen, J. C. (1995). A new species of the shrimp genus *Chorocaris* Martin & Hessler, 1990 (Crustacea: Decapoda: Bresiliidae) from hydrothermal vent fields along the Mid-Atlantic Ridge. *Proceedings of The Biological Society of Washington*, 108(2), 220 - 227.
344. Martin, J. W., France, S. C., & Van Dover, C. L. (1993). *Halice hesmonectes*, a new species of pardaliscid amphipod (Crustacea, Peracarida) from hydrothermal vents in the eastern Pacific. *Canadian Journal of Zoology*, 71, 1724 - 1732.
345. Martin, J. W., & Haney, T. A. (2005). Decapod crustaceans from hydrothermal vents and cold seeps: a review through 2005. *Zoological Journal of the Linnean Society*, 145(4), 445 - 522. doi:10.1111/j.1096-3642.2005.00178.x
346. *Martin, J. W., & Hessler, R. R. (1990). *Chorocaris vandoverae*, a new genus and species of hydrothermal vent shrimp (Crustacea, Decapoda, Bresiliidae) from the Western Pacific. *Contributions in Science*, 417, 1 - 11.
347. *Martin, J. W., & Pettit, G. (1998). *Caprella bathytatos* New Species (Crustacea, Amphipoda, Caprellidae), from the Mouthparts of the Crab Macroregonia macrochira Sakai (Brachyura, Majidae) in the Vicinity of Deep-sea Hydrothermal vents off British Columbia. *Bulletin of Marine Science*, 63(1), 189 - 198.
348. Matabos, M., Plouviez, S., Hourdez, S., Desbruyères, D., Legendre, P., Warén, A., . . . Thiébaut, E. (2011). Faunal changes and geographic crypticism indicate the occurrence of a biogeographic transition zone along the southern East Pacific Rise. *Journal of Biogeography*, 38(3), 575 - 594.
349. Matabos, M., & Thiebaut, E. (2010). Reproductive biology of three hydrothermal vent peltospirid gastropods (*Nodopelta heminoda*, *N. subnoda* and *Peltospira operculata*) associated with Pompeii worms on the East Pacific Rise. *Journal of Molluscan Studies*, 76(3), 257 - 266. doi:10.1093/mollus/eyq008
350. Matabos, M., Thiébaut, E., Le Guen, D., Sadosky, F., Jollivet, D., & Bonhomme, F. (2008). Geographic clines and stepping-stone patterns detected along the East Pacific Rise in the vetigastropod *Lepetodrilus elevatus* reflect species crypticism. *Marine Biology*, 153(4), 545 - 563.

351. McHugh, D. (1989). Population structure and reproductive biology of two sympatric hydrothermal vent polychaetes, *Paralvinella pandorae* and *P. palmiformis*. *Mar Biol*, 103, 95-106.
352. McHugh, D., & Tunnicliffe, V. (1994). Ecology and reproductive biology of the hydrothermal vent polychaete *Amphisamytha galapagensis* (Ampharetidae). *Marine Ecology Progress Series*, 106, 111 - 120.
353. *McKinney, Z. P., & Cavanaugh, C. M. (2005). The ubiquitous mussel: *Bathymodiolus* aff. *brevior* symbiosis at the Central Indian Ridge hydrothermal vents. *Marine Ecology Progress Series*, 295, 183 - 190.
354. *McLaughlin, P. A., Komai, T., Lemaitre, R., & Rahayu, D. L. (2010). Annotated checklist of anomuran decapod crustaceans of the world (exclusive of the Kiwaoidea and families Chirostylidae and Galatheidae of the Galatheoidea). Part I-Lithodoidea, Lomisoidea and Paguroidea. *The Raffles Bulletin of Zoology, Supplements*, 23, 5 - 107.
355. McLay, C. (2007). New crabs from hydrothermal vents of the Kermadec Ridge submarine volcanoes, New Zealand: *Gandalfus* gen. nov. (Bythograeidae) and *Xenograpsus* (Varunidae) (Decapoda: Brachyura). *Zootaxa*, 1524, 1 - 22.
356. McLean, J. H. (1987). Taxonomic descriptions of cocculinid limpets (Mollusca, Archaeogastropoda): two new species and three rediscovered species. *Zoologica Scripta*, 16(4), 325 - 333.
357. *McLean, J. H. (1988). New Archaeogastropod Limpets from Hydrothermal Vents; Superfamily Lepetodrilacea I. Systematic Descriptions. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 319, 1 - 32.
358. *McLean, J. H. (1989). New slit-limpets (Scissurellacea and Fissurellacea) from hydrothermal vents. Part 1. Systematic Descriptions and comparisons based on shell and radular characters. *Contributions in Science*, 407, 1 - 29.
359. McLean, J. H. (1993). New Species and Records of *Lepetodrilus* (Vetigastropoda: Lepetodrilidae) from Hydrothermal Vents. *The Veliger*, 36(1), 27-35.
360. *McLean, J. H., & Haszprunar, G. (1987). Pyropeltidae, a new family of cocculiniform limpets from hydrothermal vents. *Veliger*, 30, 196 - 205.
361. Métivier, B., & Von Cosel, R. (1993). *Acharax alinae* n. sp., Solemyidae (Mollusca: Bivalvia) géante du bassin de Lau. *Comptes Rendus de l'Académie des Sciences Série III*, 316, 229 - 237.

362. Miljutin, D., Miljutina, M., & Messié, M. (2015). Changes in abundance and community structure of nematodes from the abyssal polymetallic nodule field, Tropical Northeast Pacific. *Deep Sea Research Part I: Oceanographic Research Papers*, 106, 126 - 135.
363. *Mincks, S. L., Dyal, P. L., Paterson, G. L. J., Smith, C. R., & Glover, A. G. (2009). A new species of *Aurospio* (Polychaeta, Spionidae) from the Antarctic shelf, with analysis of its ecology, reproductive biology and evolutionary history. *Marine Ecology*, 30(2), 181 - 197.
364. Miura, T. (1986) Japanese polychaetes of the genera *Eunice* and *Euniphysa*: Taxonomy and branchial distribution patterns. *Publications of the Seto Marine Biological Laboratory*, 31(3), 269 - 325.
365. Miura, T., & Hashimoto, J. (1996). Nautiliniellid polychaetes living in the mantle cavity of bivalve molluscs from cold seeps and hydrothermal vents around Japan. *Publications of the Seto Marine Biological Laboratory*, 37(3 - 6), 257 - 274.
366. Miura, T., & Kojima, S. (2006). Two new species of vestimentiferan tubeworm (Polychaeta: Siboglinidae aka Pogonophora) from the Brothers Caldera, Kermadec Arc, South Pacific Ocean. *Species Diversity*, 11(3), 209 - 224.
367. *Miura, T., & Ohta, S. (1991). Two polychaete species from the deep-sea hydrothermal vent in the Middle Okinawa Trough. *Zoological Science, Tokyo*, 8, 383 - 387.
368. *Miyake, H., Kitada, M., Tsuchida, S., Okuyama, Y., & Nakamura, K.-i. (2007). Ecological aspects of hydrothermal vent animals in captivity at atmospheric pressure. *Marine Ecology*, 28(1), 86 - 92. doi:10.1111/j.1439-0485.2006.00115.x
369. *Miyazaki, J., Kawagucci, S., Makabe, A., Takahashi, A., Kitada, K., Torimoto, J., . . . Chen, C. (2017). Deepest and hottest hydrothermal activity in the Okinawa Trough: the Yokosuka site at Yaeyama Knoll. *Royal Society Open Science*, 4(12), 1 - 20.
370. Miyazaki, J-I., Martins, L. de O., Fujita, Y., Matsumoto, H., & Fujiwara, Y. (2010) Evolutionary process of deep-sea *Bathymodiolus* mussels. *PLOS ONE*, 5(4), e10363.
371. Moore, J. A., Auster, P. J., Calini, D., Heinonen, K., Barber, K., & Hecker, B. (2008). False Boarfish *Neocytthus helgae* in the Western North Atlantic. *Bulletin of the Peabody Museum of Natural History*, 49(1), 31 - 41. doi:10.3374/0079-032X(2008)49[31:FBNHIT]2.0.CO;2
372. Moore, J. P. (1903). Polychaeta from the Coastal Slope of Japan and from Kamchatka and Bering Sea. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 55, 401 - 490.

373. Morineaux, M., Grelon, D., & Juniper, S. K. (2002). Utilization of nutritional resources and energy budgets for populations of the hydrothermal vent polychaete *Paralvinella sulfincola* on Axial Volcano, Juan de Fuca Ridge (Northeast Pacific). *Cahiers de Biologie Marine (Proceedings of the 2nd International Symposium on Deep-Sea Hydrothermal Vent Biology)*, 43, 241 - 244.
374. Morineaux, M., Nishi, E., Ormos, A., & Mouchel, O. (2010). A new species of *Phyllochaetopterus* (Annelida: Chaetopteridae) from deep-sea hydrothermal Ashadze-1 vent field, Mid-Atlantic Ridge: taxonomical description and partial COI DNA sequence. *Cahiers de Biologie Marine*, 51(3), 239-248.
375. Munroe, T. A., Tyler, J., & Tunnicliffe, V. (2011). Description and biological observations on a new species of deepwater symphurine tonguefish (Pleuronectiformes: Cynoglossidae: *Sympfururus*) collected at Volcano-19, Tonga Arc, West Pacific Ocean. *Zootaxa*, 3061, 53 - 66.
376. Myers, A. A., & Cunha, M. R. (2004). New and little known corophiid amphipods from the 'Lucky Strike' hydrothermal vent, Mid-Atlantic Ridge. *Journal of the Marine Biological Association of the U.K.*, 84(5), 1019 - 1025.
377. Nakagawa, S., Shimamura, S., Takaki, Y., Suzuki, Y., Murakami, S.-i., Watanabe, T., . . . Takai, K. (2013). Allying with armored snails: the complete genome of gammaproteobacterial endosymbiont. *The Isme Journal*, 8, 40. doi:10.1038/ismej.2013.131
378. Nakamura, K., & Takai, K. (2015) Indian Ocean hydrothermal systems: seafloor hydrothermal activities, physical and chemical characteristics of hydrothermal fluids, and vent-associated biological communities. In: Ishibashi, J., Okino, K., & Sunamura, M. (eds.) *Subseafloor Biosphere Linked to Hydrothermal Systems*. Springer, Toyko.
379. *Nakamura, K., Watanabe, H., Miyazaki, J., Takai, K., Kawagucci, S., Noguchi, T., . . . Tamaki, K. (2012). Discovery of New Hydrothermal Activity and Chemosynthetic Fauna on the Central Indian Ridge at 18°–20°S. *PLoS One*, 7(3), e32965. doi:10.1371/journal.pone.0032965
380. *Nakamura, M., Chen, C., & Mitarai, S. (2015). Insights into life-history traits of *Munidopsis* spp. (Anomura: Munidopsidae) from hydrothermal vent fields in the Okinawa Trough, in comparison with the existing data. *Deep Sea Research Part I: Oceanographic Research Papers*, 100, 48 - 53.
381. Naraoka, H., Naito, T., Yamanaka, T., Tsunogai, U., & Fujikura, K. (2008). A multi-isotope study of deep-sea mussels at three different hydrothermal vent sites in the northwestern Pacific. *Chemical Geology*, 255(1 - 2), 25 - 32.
382. *Newman, W. A., & Jones, W. J. (2011). Two Northeast Pacific deep-water barnacle populations (Cirripedia: Calanticidae and Pachylasmatidae) from seamounts of the

Juan de Fuca Ridge; "insular" endemics stemming from Tethys, or by subsequent dispersal from the Western Pacific center of distribution? *Zootaxa*, 2789, 49 - 68.

383. *Newman, W. A., & Ross, A. (1971). *Antarctic Cirripedia: Monographic Account Based on Specimens Collected Chiefly Under the United States Antarctic Research Program, 1962-1965* (Vol. 14): American Geophysical Union.
384. Newman, W. A., & Yamaguchi, T. (1995). A new sessile barnacle (Cirripedia, Brachylepadomorpha) from the Lau Back-Arc Basin, Tonga; first record of a living representative since the Miocene. *Bulletin du Muséum national d'histoire naturelle, Section A, Zoology, biologie et écologie animales*, 17, 221 - 244.
385. Newton, I. L. G., Woyke, T., Auchtung, T. A., Dilly, G. F., Dutton, R. J., Fisher, M. C., . . . Cavanaugh, C. M. (2007). The *Calyptogena magnifica* Chemoautotrophic Symbiont Genome. *Science*, 315(5814), 998.
386. Ng, N. K., Suzuki, H., Shih, H.-T., Dewa, S., & Ng, P. K. L. (2014). The hydrothermal crab, *Xenograpsus testudinatus* Ng, Huang & Ho, 2000 (Crustacea: Decapoda: Brachyura: Grapsidae) in southern Japan. *Proceedings of The Biological Society of Washington*, 127(2), 391 - 399.
387. *Nielsen, J. G., Cohen, D. M., Markle, D. F., & Robins, C. R. (1999). FAO species catalogue. Volume 18. Ophidiiform fishes of the world (Order Ophidiiformes). An annotated and illustrated catalogue of pearlfishes, cusk-eels, brotulas and other ophidiiform fishes known to date. *FAO Fisheries Synopsis*, 18(125), 178.
388. *Nishijima, M., Lindsay, D. J., Hata, J., Nakamura, A., Kasai, H., Ise, Y., . . . Maruyama, T. (2010). Association of Thioautotrophic Bacteria with Deep-Sea Sponges. *Marine Biotechnology*, 12(3), 253 - 260. doi:10.1007/s10126-009-9253-7
389. *Nye, V. E., & Copley, J. T. (2014). Reproductive ecology of a hippolytid shrimp, *Lebbeus virentova* (Caridea: Hippolytidae) at the Von Damm Vent Field, Mid-Cayman Spreading Centre. *Marine Biology*, 161(10), 2371 - 2380.
390. Nye, V., Copley, J., & Plouviez, S. (2012). A new species of *Rimicaris* (Crustacea: Decapoda: Caridea: Alvinocarididae) from hydrothermal vent fields on the Mid-Cayman Spreading Centre, Caribbean. *Journal of the Marine Biological Association of the U.K.*, 92(5), 1057 - 1072.
391. Nye, V., Copley, J., Plouviez, S., & Van Dover, C. L. (2013). A new species of *Lebbeus* (Crustacea: Decapoda: Caridea: Hippolytidae) from the Von Damm Vent Field, Caribbean Sea. *Journal of the Marine Biological Association of the U.K.*, 93(3), 741 - 751.
392. Nye, V., Copley, J. T., & Tyler, P. A. (2013). Spatial Variation in the Population

Structure and Reproductive Biology of *Rimicaris hybisae* (Caridea: Alvinocarididae) at Hydrothermal Vents on the Mid-Cayman Spreading Centre. *PLoS One*, 8(3, e60319), 1 - 15.

393. O'Mullan, G. D., Maas, P. A. Y., Lutz, R. A., & Vrijenhoek, R. C. (2001). A hybrid zone between hydrothermal vent mussels (Bivalvia: Mytilidae) from the Mid-Atlantic Ridge. *Molecular Ecology*, 10, 2819 - 2831.
394. OBIS. (2017). *Ocean Biogeographic Information System*. Retrieved from: www.iobis.org.
395. Ohta, S., & Kim, D. (2001). Submersible Observations of the Hydrothermal Vent Communities on the Iheya Ridge, Mid Okinawa Trough, Japan. *Journal of Oceanography*, 57(6), 663 - 677. doi:10.1023/A:1021620023610
396. Okutani, T., Hashimoto, J., & Sasaki, T. (2004). New gastropod taxa from a hydrothermal vent (Kairei Field) in the central Indian Ocean. *Venus*, 63(1 - 2), 1 - 10.
397. Okutani, T., Kojima, S., & Ashi, J. (1996). An unusual vesicomyid bivalve, *Calyptogena nankaiensis* n. sp. from bathyal depth off Honshu, Japan. *Japanese Journal of Malacology (Japan)*.
398. Okutani, T., Kojima, S., & Iwasaki, N. (2002). New and known vesicomyid bivalves recently collected from the western and central Nankai Trough off Shikoku and Honshu, by deep sea research systems of Japan Marine Science and Technology Center. *Venus*, 61(3 - 4), 129 - 140.
399. Oliver, P. G., & Holmes, A. M. (2006a). The Arcoidea (Mollusca: Bivalvia): a review of the current phenetic-based systematics. *Zoological Journal of the Linnean Society*, 148(3), 237 - 251.
400. Oliver, P. G., & Holmes, A. M. (2006b). New species of Thyasiridae (Bivalvia) from chemosynthetic communities in the Atlantic Ocean. *Journal of Conchology*, 39(2), 175 - 184.
401. Oliver, P. G., & Rodrigues, C. F. (2017). Thyasiridae (Mollusca: Bivalvia) from the Kemp Caldera hydrothermal site, South Sandwich Islands, Antarctica. *Journal of Conchology*, 42(5), 267 - 282.
402. Olsen, B. R., Troedsson, C., Hadziavdic, K., Pedersen, R. B., & Rapp, H. T. (2014). A molecular gut content study of *Themisto abyssorum* (Amphipoda) from Arctic hydrothermal vent and cold seep systems. *Molecular Ecology*, 23(15), 3877 - 3889. doi:10.1111/mec.12511
403. *Olu-Le Roy, K., Caprais, J. C., Fifis, A., Fabri, M. C., Galéron, J., Budzinsky, H., . . . Sibuet, M. (2007). Cold-seep assemblages on a giant pockmark off West Africa:

Spatial patterns and environmental control. *Marine Ecology*, 28(1), 115 - 130.
doi:10.1111/j.1439-0485.2006.00145.x

404. *Olu-Le Roy, K., Sibuet, M., Fiala-Médioni, A., Gofas, S., Salas, C., Mariotti, A., . . . Woodside, J. (2004). Cold seep communities in the deep eastern Mediterranean Sea: Composition, symbiosis and spatial distribution on mud volcanoes. *Deep-Sea Research Part I: Oceanographic Research Papers*, 51(12), 1915 - 1936.
doi:10.1016/j.dsr.2004.07.004
405. Page, H. M., Fiala-Medioni, A., Fisher, C. R., & Childress, J. J. (1991). Experimental evidence for filter-feeding by the hydrothermal vent mussel, *Bathymodiolus thermophilus*. *Deep Sea Research Part A. Oceanographic Research Papers*, 38(12), 1455 - 1461. doi:10.1016/0198-0149(91)90084-S
406. *Palomares, M. L. D., & Pauly, D. (2017). SeaLifeBase. www.sealifebase.org
407. Paradis, S., Jonasson, I. R., Le Cheminant, G. M., & Watkinson, D. H. (1988). Two zinc-rich chimneys from the plume site, southern Juan de Fuca Ridge. *The Canadian Mineralogist*, 26(3), 637 - 654.
408. Paterson, G. L. J. (1985). The deep-sea Ophiuroidea of the North Atlantic Ocean. *Bulletin of the British Museum, Zoology*, 49(1), 1 - 162.
409. *Paxton, H., & Morineaux, M. (2009). Three Species of Dorvilleidae (annelida: Polychaeta) Associated With Atlantic Deep-Sea Reducing Habitats, With The Description of Ophryotrocha Fabriae, New Species. *Proceedings of The Biological Society of Washington*, 122(1), 14 - 25. doi:10.2988/08-22.1
410. Pedersen, R. B., Rapp, H. T., Thorseth, I. H., Lilley, M. D., Barriga, F. J., Baumberger, T., . . . Jorgensen, S. L. (2010). Discovery of a black smoker vent field and vent fauna at the Arctic Mid-Ocean Ridge. *Nature Communications*, 1, 126.
doi:10.1038/ncomms1124
411. Peek, A. S., Feldman, R. A., Lutz, R. A., & Vrijenhoek, R. C. (1998). Cospeciation of chemoautotrophic bacteria and deep sea clams. *Proc Natl Acad Sci U S A*, 95(17), 9962 - 9966.
412. Peek, A. S., Gustafson, R. G., Lutz, R. A., & Vrijenhoek, R. C. (1997). Evolutionary relationships of deep-sea hydrothermal vent and cold-water seep clams (Bivalvia: Vesicomyidae): results from the mitochondrial cytochrome oxidase subunit I. *Marine Biology*, 130, 151 - 161.
413. *Penry, D. L., & Jumars, P. A. (1990). Gut architecture, digestive constraints and feeding ecology of deposit-feeding and carnivorous polychaetes. *Oecologia*, 82(1), 1 - 11.

414. Perez, M., & Juniper, S. K. (2016). Insights into Symbiont Population Structure among Three Vestimentiferan Tubeworm Host Species at Eastern Pacific Spreading Centers. *Applied and Environmental Microbiology*, 82(17), 5197 - 5205.
415. *Perovich, G. M., Epifanio, C. E., Dittel, A. I., & Tyler, P. A. (2003). Spatial and temporal patterns in development of eggs in the vent crab *Bythograea thermydron*. *Marine Ecology Progress Series*, 251, 211 - 220.
416. Petersen, J. M., Ramette, A., Lott, C., Cambon-Bonavita, M.-A., Zbinden, M., & Dubilier, N. (2010). Dual symbiosis of the vent shrimp *Rimicaris exoculata* with filamentous gamma- and epsilonproteobacteria at four Mid-Atlantic Ridge hydrothermal vent fields. *Environmental Microbiology*, 12(8), 2204 - 2218. doi:10.1111/j.1462-2920.2009.02129.x
417. *Pettibone, M. H. (1990). New species and new records of scaled polychaetes (Polychaeta: Polynoidae) from the Axial Seamount caldera of the Juan de Fuca Ridge in the northeast Pacific and the East Pacific Ocean off northern California. *Proceedings of The Biological Society of Washington*, 103(4), 825 - 838.
418. *Pettibone, M. H. (1989). Polynoidae and Sigalionidae (Polychaeta) from the Guaymas Basin, with descriptions of two new species, and additional records from hydrothermal vents of the Galapagos Rift, 21°N, and seep-sites in the Gulf of Mexico (Florida and Louisiana). *Proceedings of The Biological Society of Washington*, 102(1), 154 - 168.
419. Pettibone, M. H. (1986). Review of the Iphioninae (Polychaeta: Polynoidae) and revision of *Iphione cimex* Quatrefages, *Gattyana deludens* Fauvel and *Harmothoe iphionelloides* Johnson (Harmothoinae). *Smithsonian Contributions to Zoology*, 428, 1 - 43.
420. Pettibone, M. H. (1985a). Additional branchiate scale-worms (Polychaeta: Polynoidae) from Galapagos hydrothermal vent and rift-area off western Mexico at 21°N. *Proceedings of The Biological Society of Washington*, 98(2), 447 - 469.
421. Pettibone, M. H. (1985b). New genera and species of deep-sea Macellicephalinae and Harmothoinae (Polychaeta: Polynoidae) from the hydrothermal rift areas off the Galapagos and western Mexico at 21°N and from the Santa Catalina Channel. *Proceedings of The Biological Society of Washington*, 98(3), 740 - 757.
422. *Pettibone, M. H. (1984a). A new scale worm commensal with deep-sea mussels on the Galapagos hydrothermal vent (Polychaeta: Polynoidae). *Proceedings of The Biological Society of Washington*, 97(1), 226 - 239.
423. *Pettibone, M. H. (1984b). Two new species of *Lepidonotopodium* (Polychaeta: Polynoidae: Lepidonotopodinae) from hydrothermal vents off the Galapagos and East Pacific Rise at 21°N. *Proceedings of The Biological Society of Washington*, 97(4), 849 - 863.

424. Pettibone, M. H. (1976). Revision of the Genus *Macellicephalia* McIntosh and the Subfamily Macellicephalinae Hartmann-Schrader (Polychaeta: Polynoidae). *Smithsonian Contributions to Zoology*, 229, 1 - 71.
425. Pimenov, N. V., Kalyuzhnaya, M. G., Khmelenina, V. N., Mityushina, L. L., & Trotsenko, Y. A. (2002). Utilization of Methane and Carbon Dioxide by Symbiotic Bacteria in Gills of Mytilidae (*Bathymodiolus*) from the Rainbow and Logachev Hydrothermal Fields on the Mid-Atlantic Ridge. [Translated from Mikrobiologiya, Vol. 71, No. 5, 2002, pp. 681–689]. *Microbiology*, 71(5), 587 - 594.
426. Plouviez, S., Jacobson, A., Wu, M., & Van Dover, C. L. (2015). Characterization of vent fauna at the Mid-Cayman Spreading Center. *Deep Sea Research Part I: Oceanographic Research Papers*, 97(Supplement C), 124 - 133. doi:10.1016/j.dsr.2014.11.011
427. Podowski, E. L., Ma, S., Luther, I. G., Wardrop, D., & Fisher, C. R. (2009). Biotic and abiotic factors affecting distributions of megafauna in diffuse flow on andesite and basalt along the Eastern Lau Spreading Center, Tonga. *Marine Ecology Progress Series*, 418, 25 - 45.
428. *Polz, M. F., Robinson, J. J., Cavanaugh, C. M., & Van Dover, C. L. (1998). Trophic ecology of massive shrimp aggregations at a Mid-Atlantic Ridge hydrothermal vent site. *Limnology and Oceanography*, 43(7), 1631 - 1638. doi:10.4319/lo.1998.43.7.1631
429. Portail, M., Olu, K., Dubois, S. F., Escobar-Briones, E., Gelinas, Y., Menot, L., & Sarrazin, J. (2016). Food-Web Complexity in Guaymas Basin Hydrothermal Vents and Cold Seeps. *PLoS One*, 11(9), e0162263), 1 - 33.
430. *Pranal, V., Fiala-Medioni, A., & Guezennec, J. (2009). Fatty acid characteristics in two symbiont-bearing mussels from deep-sea hydrothermal vents of the south-western Pacific. *Journal of the Marine Biological Association of the United Kingdom*, 77(2), 473 - 492. doi:10.1017/S0025315400071812
431. Price, R. E., Breuer, C., Reeves, E., Bach, W., & Pichler, T. (2016). Arsenic bioaccumulation and biotransformation in deep-sea hydrothermal vent organisms from the PACMANUS hydrothermal field, Manus Basin, PNG. *Deep Sea Research Part I: Oceanographic Research Papers*, 117, 95 - 106. doi:10.1016/j.dsr.2016.08.012
432. *Ramil, F., & Vervoort, W. (1992). Report on the Hydroida collected by the "BALGIM" expedition in and around the Strait of Gibraltar. *Zoologische verhandelingen / uitgegeven door het Rijksmuseum van Natuurlijke Historie te Leiden (Ministerie van Cultuur, Recreatie en Maatschappelijk Werk)*, 277(7), 1 - 262.
433. *Ramirez-Llodra, E., & Segonzac, M. (2006). Reproductive biology of *Alvinocaris muricola* (Decapoda: Caridea: Alvinocarididae) from cold seeps in the Congo Basin. *Journal of the Marine Biological Association of the United Kingdom*, 86(6), 1347-1356.

434. *Ramirez-Llodra, E., Tyler, P. A., & Copley, J. T. P. (2000). Reproductive biology of three caridean shrimp, *Rimicaris exoculata*, *Chorocaris chacei* and *Mirocaris fortunata* (Caridea: Decapoda), from hydrothermal vents. *Journal of the Marine Biological Association of the U.K.*, 80, 473 - 484.
435. *Rau, G. H. (1981). Hydrothermal vent clam and tube worm $^{13}\text{C}/^{12}\text{C}$: further evidence of nonphotosynthetic food source. *Science*, 213, 338 - 339.
436. Raulfs, E. C., Macko, S. A., & Van Dover, C. L. (2004). Tissue and symbiont condition of mussels (*Bathymodiolus thermophilus*) exposed to varying levels of hydrothermal activity. *Journal of the Marine Biological Association of the U.K.*, 84(1), 229 - 234.
437. Reid, W. D. K., Sweeting, C. J., Wigham, B. D., McGill, R. A. R., & Polunin, N. V. C. (2016). Isotopic niche variability in macroconsumers of the East Scotia Ridge (Southern Ocean) hydrothermal vents: What more can we learn from an ellipse? *Marine Ecology Progress Series*, 542, 13 - 24.
438. Reid, W. D. K., Sweeting, C. J., Wigham, B. D., Zwirglmaier, K., Hawkes, J. A., McGill, R. A. R., . . . Polunin, N. V. C. (2013). Spatial Differences in East Scotia Ridge Hydrothermal Vent Food Webs: Influences of Chemistry, Microbiology and Predation on Trophodynamics. *PLoS One*, 8(6, e65553), 1 - 11.
439. Reuscher, M., Fiege, D., & Wehe, T. (2011). Terebellomorph polychaetes from hydrothermal vents and cold seeps with the description of two new species of Terebellidae (Annelida: Polychaeta) representing the first records of the family from deep-sea vents. *Journal of the Marine Biological Association of the United Kingdom*, 92(5), 997 - 1012. doi:10.1017/S0025315411000658
440. Reynolds, K. C., Watanabe, H., Strong, E. E., Sasaki, T., Uematsu, K., Miyake, H., . . . Young, C. M. (2010). New Molluscan Larval Form: Brooding and Development in a Hydrothermal Vent Gastropod, *Ifremeria nautilei* (Provannidae). *The Biological Bulletin*, 219(1), 7 - 11. doi:10.1086/BBLv219n1p7
441. *Rhoads, D. C., & Lutz, R. A. (1980). *Skeletal Growth of Aquatic Organisms: Biological Records of Environmental Change*. Springer US: New York, 750.
442. Rhoads, D. C., Lutz, R. A., Revelas, E. C., & Cerrato, R. M. (1981). Growth of Bivalves at Deep-Sea Hydrothermal Vents Along the Galápagos Rift. *Science*, 214(4523), 911 - 913.
443. Riou, V., Duperron, S., Halary, S., Dehairs, F., Bouillon, S., Martins, I., . . . Serrão Santos, R. (2010). Variation in physiological indicators in *Bathymodiolus azoricus* (Bivalvia: Mytilidae) at the Menez Gwen Mid-Atlantic Ridge deep-sea hydrothermal

vent site within a year. *Marine Environmental Research*, 70(3), 264 - 271.
doi:10.1016/j.marenvres.2010.05.008

444. *Rodrigues, C. F., Webster, G., Cunha, M. R., Duperron, S., & Weightman, A. J. (2010). Chemosynthetic bacteria found in bivalve species from mud volcanoes of the Gulf of Cadiz. *FEMS Microbiology Ecology*, 73(3), 486 - 499. doi:10.1111/j.1574-6941.2010.00913.x
445. Roe, K. L., Schneider, R. J., Hansel, C. M., & Voelker, B. M. (2016). Measurement of dark, particle-generated superoxide and hydrogen peroxide production and decay in the subtropical and temperate North Pacific Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, 107(Supplement C), 59 - 69.
doi:/10.1016/j.dsr.2015.10.012
446. Rogers, A. D., Gibson, R., & Tunnicliffe, V. (1996). A new genus and species of monostiliferous hoplonemertean colonizing an inchoate hydrothermal field on Juan de Fuca Ridge. *Deep-Sea Research Part I*, 43(10), 1581 - 1599.
447. Rogers, A. D., Tyler, P. A., Connelly, D. P., Copley, J. T., James, R., Larter, R. D., . . . Zwirglmaier, K. (2012). The discovery of new deep-sea hydrothermal vent communities in the Southern Ocean and implications for biogeography. *PLoS Biology*, 10(1), e1001234. doi:10.1371/journal.pbio.1001234
448. *Rossi, G. (2014). *The reproductive and physiological condition of a deep-sea mussel (Bathymodiolus septendierum Hashimoto & Okutani, 1994) living in extremely acidic conditions*. (MSc), University of Victoria, Victoria, BC.
449. Rossi, G. S., & Tunnicliffe, V. (2017). Trade-offs in a high CO₂ habitat on a subsea volcano: condition and reproductive features of a bathymodioline mussel. *Marine Ecology Progress Series*, 574, 49 - 64.
450. Saito, H., & Okutani, T. (1990). Two new chitons (Mollusca: Polyplacophora) from a hydrothermal vent site of the Iheya Small Ridge, Okinawa Trough, East China Sea. *Venus*, 49, 165 - 179.
451. Saldanha, L., & Biscoito, M. (1997). Fishes from the Lucky Strike and Menez Gwen hydrothermal vent sites (Mid-Atlantic Ridge). *Boletim do Museu Municipal do Funchal*, 49(283), 189 - 206.
452. Sarrazin, J., & Juniper, S. K. (1999). Biological characteristics of a hydrothermal edifice mosaic community. *Marine Ecology Progress Series*, 185, 1 - 19.
453. Sarrazin, J., Legendre, P., de Busserolles, F., Fabri, M.-C., Guilini, K., Ivanenko, V. N., . . . Sarradin, P.-M. (2015). Biodiversity patterns, environmental drivers and indicator species on a high-temperature hydrothermal edifice, Mid-Atlantic Ridge. *Deep Sea*

454. Sars, G. O. (1872). On some remarkable forms of animal life from the great deeps off the Norwegian coast. Part 1, partly from posthumous manuscripts of the late prof. Mich. Sars. University Program for the 1rs half-year 1869. *Brøgger & Christie, Christiania viii*, 65 - 68.
455. Sasaki, T., Ogura, T., Watanabe, H., & Fujikura, K. (2016). Four new species of hycarbina discovered from hydrothermal vents off the Nansei Shoto and cold spring springs. *Venus (Journal of the Malacological Society of Japan)*, 74(1 - 2), 1 - 17.
456. Sasaki, T., Okutani, T., & Fujikura, K. (2006). Anatomy of Bathycmaea secunda Okutani, Fujikura & Sasaki, 1993 (Patellogastropoda: Acmaeidae). *Journal of Molluscan Studies*, 72(3), 295 - 309. doi:10.1093/mollus/eyl007
457. Sasaki, T., Okutani, T., & Fujikura, K. (2005). Molluscs from hydrothermal vents and cold seeps in Japan: a review of taxa recorded in twenty recent years (1984–2004). *Venus*, 64(3 - 4), 87 - 133.
458. *Sato, K., Sasaki, T., & Watanabe, H. (2013). A new species of Solemya (Bivalvia: Protobranchia: Solemyidae) from a hydrothermal vent in the Iheya Ridge in the mid-Okinawa Trough, Japan. *Nautilus*, 127, 93 - 100.
459. *Scarlato, O. A. (1981). Bivalve molluscs of the temperate latitudes of the western part of the Pacific ocean. *Transactions of the U.S.S.R. Academy of Sciences Zoological Institute, Nauka Publishers, Leningrad*, 126(Survey of the Fauna of the U.S.S.R.), 479.
460. Schander, C., Rapp, H. T., Kongsrud, J. A., Bakken, T., Berge, J., Cochrane, S., . . . Pedersen, R. B. (2010). The fauna of hydrothermal vents on the Mohn Ridge (North Atlantic). *Marine Biology Research*, 6(2), 155-171. doi:10.1080/17451000903147450
461. Scheltema, A. H. (2008). Biogeography, Diversity, and Evolution Through Vicariance of the Hydrothermal Vent Aplacophoran Genus *Helicoradomenia* (Aplacophora, Mollusca). *Journal of Shellfish Research*, 27(1), 91 - 96. doi:10.2983/0730-8000(2008)27[91:BDAETV]2.0.CO;2
462. Scheltema, A. H. (2000). Two new hydrothermal vent species, *Helicoradomenia bisquama* and *Helicoradomenia acredema*, from the eastern Pacific Ocean (Mollusca, Aplacophora). *Argonauta*, 14(2), 15 - 25.
463. Scheltema, A. H., & Kuzirian, A. M. (1991). *Helicoradomenia juani* gen. et sp. nov., a Pacific Hydrothermal Vent Aplacophora (Mollusca: Neomeniomorpha). *The Veliger*, 34(2), 195 - 203.

464. Schuchert, P. (2001). Hydroids of Greenland and Iceland (Cnidaria, Hydrozoa) Meddelelser om Grønland. *BioScience*, 53, 1 - 184.
465. Segonzac, M., de Saint Laurent, M., & Casanova, B. (1993). L'éénigme du comportement trophique des crevettes Alvinocarididae des sites hydrothermaux de la dorsale médio-Atlantique. *Cahiers de Biologie Marine*, 34, 535 - 571.
466. Segonzac, M., & Vervoort, W. (1995). First record of the genus *Candelabrum* (Cnidaria, Hydrozoa, Athecata) from the Mid-Atlantic Ridge: a description of a new species and a review of the genus. *Bulletin de Muséum national d'histoire naturelle, Paris, 4 séries*, 17(1 - 2), 31 - 64.
467. Sen, A., Becker, E. L., Podowski, E. L., Wickes, L. N., Ma, S., Mullaugh, K. M., . . . Fisher, C. R. (2013). Distribution of mega fauna on sulfide edifices on the Eastern Lau Spreading Center and Valu Fa Ridge. *Deep Sea Research Part I: Oceanographic Research Papers*, 72, 48 - 60.
468. Sen, A., Kim, S., Miller, A. J., Hovey, K. J., Hourdez, S., Luther III, G. W., & Fisher, C. R. (2016). Peripheral communities of the Eastern Lau Spreading Center and Valu Fa Ridge: community composition, temporal change and comparison to near-vent communities. *Marine Ecology*, 37(3), 599 - 617.
469. Sen, A., Podowski, E. L., Becker, E. L., Shearer, E. A., Gartman, A., Yücel, M., . . . Fisher, C. R. (2014). Community succession in hydrothermal vent habitats of the Eastern Lau Spreading Center and Valu Fa Ridge, Tonga. *Limnology and Oceanography*, 59(5), 1510 - 1528.
470. Shank, T. M., Fornari, D. J., Von Damm, K. L., Lilley, M. D., Haymon, R. M., & Lutz, R. A. (1998). Temporal and spatial patterns of biological community development at nascent deep-sea hydrothermal vents (9°50'N, East Pacific Rise). *Deep Sea Research Part II: Topical Studies in Oceanography*, 45(1), 465 - 515. doi:10.1016/S0967-0645(97)00089-1
471. *Sheader, M. (1986). *Primno evansi* sp. nov. (Amphipoda: Hyperiidea) from the eastern North Atlantic. *Journal of Natural History*, 20, 975 - 980.
472. Sheader, M., Van Dover, C. L., & Shank, T. M. (2000). Structure and function of *Halice hesmonectes* (Amphipoda: Pardaliscidae) swarms from hydrothermal vents in the eastern Pacific. *Marine Biology*, 136, 901 - 911.
473. Sheader, M., Van Dover, C. L., & Thurston, M. H. (2004). Reproductive ecology of *Bouvierella curtirama* (Amphipoda: Eusiridae) from chemically distinct vents in the Lucky Strike vent field, Mid-Atlantic Ridge. *Marine Biology*, 144, 503 - 514.
474. Shillito, B., Jollivet, D., Sarradin, P.-M., Rodier, P., Lallier, F., Desbruyères, D., & Gaill,

- F. (2001). Temperature resistance of *Hesiolyra bergi*, a polychaetous annelid living on deep-sea vent smoker walls. *Marine Ecology Progress Series*, 216, 141 - 149.
475. *Signorelli, J., & Pastorino, G. (2015). A New Species of *Laubiericoncha* (Bivalvia: Vesicomyidae) from Deep Waters Off Argentina. *Malacologia*, 58(1-2), 349 - 360. doi:10.4002/040.058.0213
476. Sigvaldadóttir, E., & Desbruyères, D. (2003). Two new species of Spionidae (Annelida: Polychaeta) from Mid-Atlantic Ridge hydrothermal vents. *Cahiers de Biologie Marine*, 44(3), 219 - 225.
477. *Sloan, L. M., Anderson, S. V., & Pernet, B. (2010). Kilometer-Scale Spatial Variation in Prevalence of the Rhizocephalan *Lernaeodiscus Porcellanae* on the Porcelain Crab *Petrolisthes Cabrilloi*. *Journal of Crustacean Biology*, 30(2), 159 - 166.
478. Smirnov, A. V., Gebruk, A. V., Galkin, S. V., & Shank, T. (2000). New species of holothurian (Echinodermata: Holothuroidea) from hydrothermal vent habitats. *Journal of the Marine Biological Association of the U.K.*, 80(2), 321 - 328.
479. Smith, C. R., & Baco, A. R. (2003). Ecology of whale falls at the deep-sea floor. *Oceanography and Marine Biology: An Annual Review*, 41, 311 - 354.
480. Smith, C. R., Bernardino, A. F., Baco, A., Hannides, A., & Altamira, I. (2014). Seven-year enrichment: macrofaunal succession in deep-sea sediments around a 30 tonne whale fall in the Northeast Pacific. *Marine Ecology Progress Series*, 515, 133 - 149.
481. Smith, C. R., & Rabouille, C. (2004) Reply to comment by Boudreau on: What controls the mixed-layer depth in deep-sea sediments? The importance of POC flux. *Limnology and Oceanography*, 49(2), 623 - 624.
482. *Snelgrove, P. V. R., Grassle, J. F., & Petrecca, R. F. (1994). Macrofaunal response to artificial enrichments and depressions in a deep-sea habitat. *Journal of Marine Research*, 52(2), 345 - 369.
483. Solis-Weiss, V. (1993). *Grassleia hydrothermalis*, a new genus and species of Ampharetidae (Annelida: Polychaeta) from the hydrothermal vents off the Oregon Coast (U.S.A.), at Gorda Ridge. *Proc. Biol. Soc. Wash.*, 106(4), 661 - 665.
484. Soto, L. A. (2009). Stable carbon and nitrogen isotopic signatures of fauna associated with the deep-sea hydrothermal vent system of Guaymas Basin, Gulf of California. *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(19 - 20), 1675 - 1682.
485. Southward, E. C. (2008). The morphology of bacterial symbioses in the gills of

mussels of the genera *Adipicola* and *Idas* (Bivalvia: Mytilidae). *Journal of Shellfish Research*, 27, 139 - 146.

486. *Southward, A. J. (2005). Systematics and ecology of a new species of stalked barnacle (Cirripedia: Thoracica: Scalpellomorpha: Eolepadidae: Neolepadini) from the Pacific-Antarctic Ridge at 38° S. *Senckenbergiana maritima*, 35(2), 147 - 156.
487. *Southward, E. C. (1991). Three new species of Pogonophora, including two vestimentiferans, from hydrothermal sites in the Lau Back-arc Basin (Southwest Pacific Ocean). *Journal of Natural History*, 25(4), 859 - 881.
doi:10.1080/00222939100770571
488. *Southward, E. C., Andersen, A. C., & Hourdez, S. (2011). *Lamellibrachia anaximandri* n. sp., a new vestimentiferan tubeworm (Annelida) from the Mediterranean, with notes on frenulate tubeworms from the same habitat. *Zoosystema*, 33(3), 245 - 279.
doi:10.5252/z2011n3a1
489. Southward, E. C., & Galkin, S. V. (1997). A new vestimentiferan (Pogonophora: Obturata) from hydrothermal vent fields in the Manus Back-arc Basin (Bismarck Sea, Papua New Guinea, Southwest Pacific Ocean). *Journal of Natural History*, 31(1), 43 - 55. doi:10.1080/00222939700770041
490. Southward, E. C., Gebruk, A., Kennedy, H., Southward, A. J., & Chelvadonné, P. (2001). Different energy sources for three symbiont-dependent bivalve molluscs at the Logatchev hydrothermal site (Mid-Atlantic Ridge). *Journal of the Marine Biological Association of the U.K.*, 81(4), 655 - 661.
491. Southward, A. J., & Jones, D. S. (2003). A revision of stalked barnacles (Cirripedia: Thoracica: Scalpellomorpha: Eolepadidae: Neolepadinae) associated with hydrothermalism, including a description of a new genus and species from a volcanic seamount off Papua New Guinea. *Senckenbergiana maritima*, 32(1), 77 - 93.
doi:10.1007/bf03043086
492. Southward, E. C., Schulze, A., & Gardiner, S. L. (2005). Pogonophora (Annelida): form and function. *Hydrobiologia*, 535(1), 227 - 251. doi:10.1007/s10750-004-4401-6
493. *Southward, E. C., Schulze, A., & Tunnicliffe, V. (2002). Vestimentiferans (Pogonophora) in the Pacific and Indian Oceans: a new genus from Lihir Island (Papua New Guinea) and the Java Trench, with the first report of *Arcovestia ivanovi* from the North Fiji Basin. *Journal of Natural History*, 36, 1179 - 1197.
494. Southward, E. C., Tunnicliffe, V., & Black, M. (1995). Revision of the species of *Ridgeia* from northeast Pacific hydrothermal vents, with a redescription of *Ridgeia piscesae* Jones (Pogonophora: Obturata = Vestimentifera). *Canadian Journal of Zoology*, 73(2), 282 - 295. doi:10.1139/z95-033

495. *Southward, E. C., Tunnicliffe, V., Black, M. B., Dixon, D. R., & Dixon, L. R. J. (1996). Ocean-ridge segmentation and vent tubeworms (Vestimentifera) in the NE Pacific. *Geological Society, London, Special Publications*, 118(1), 211 - 224. doi:10.1144/gsl.sp.1996.118.01.13
496. Spiridonova, E. M., Kuznetsov, B. B., Pimenov, N. V., & Tourova, T. P. (2006). Phylogenetic characterization of endosymbionts of the hydrothermal vent mussel *Bathymodiolus azoricus* by analysis of the 16S rRNA, *cbbL*, and *pmoA* genes. *Microbiology*, 75(6), 694 - 701.
497. Stebbing, T. R. R. (1888). Report on the Amphipoda collected by H.M.S. Challenger during the years 1873-1876. *Report on the Scientific Results of the Voyage of H.M.S. Challenger during the years 1873-76, Zoology*, 29(1 - 872). doi:10.5962/bhl.title.6513
498. Stecher, J., Tunnicliffe, V., & Türkay, M. (2003). Population characteristics of abundant bivalves (Mollusca, Vesicomyidae) at a sulphide-rich seafloor site near Lihir Island, Papua New Guinea. *Canadian Journal of Zoology*, 81(11), 1815 - 1824.
499. *Steele, D. H., & Brunel, P. (1968). Amphipoda of the Atlantic and Arctic coasts of North America: *Anonyx* (Lysianassidae). *Journal of Fisheries Research Board of Canada*, 25(5), 943 - 1060.
500. *Stevens, C. J., Limén, H., Pond, D. W., Gélinas, Y., & Juniper, S. K. (2008). Ontogenetic shifts in the trophic ecology of two alvinocaridid shrimp species at hydrothermal vents on the Mariana Arc, western Pacific Ocean. *Marine Ecology Progress Series*, 356, 225 - 237.
501. Stiller, J., Rousset, V., Pleijel, F., Chelvadonné, P., Vrijenhoek, R. C., & Rouse, G. W. (2013). Phylogeny, biogeography and systematics of hydrothermal vent and methane seep *Amphisamytha* (Ampharetidae, Annelida), with descriptions of three new species. *Systematics and Biodiversity*, 11(1), 35 - 65.
502. *Stöhr, S., O'Hara, T. D., & Thuy, B. (2012). Global Diversity of Brittle Stars (Echinodermata: Ophiuroidea). *PLoS One*, 7(3), e31940. doi:10.1371/journal.pone.0031940
503. Stöhr, S., & Segonzac, M. (2006). Two new genera and species of ophiuroid (Echinodermata) from hydrothermal vents in the East Pacific. *Species Diversity*, 11, 7 - 32.
504. Stöhr, S., & Segonzac, M. (2005). Deep-sea ophiuroids (Echinodermata) from reducing and non-reducing environments in the North Atlantic Ocean. *Journal of the Marine Biological Association of the United Kingdom*, 85(2), 383-402. doi:10.1017/S0025315405011318h

505. Streit, K., Bennett, S. A., Van Dover, C. L., & Coleman, M. (2015). Sources of organic carbon for *Rimicaris hybisae*: Tracing individual fatty acids at two hydrothermal vent fields in the Mid-Cayman rise. *Deep Sea Research Part I: Oceanographic Research Papers*, 100, 13 - 20.
506. Sun, Q.-I., Zeng, Z.-g., Chen, S., & Sun, L. (2016). First Comparative Analysis of the Community Structures and Carbon Metabolic Pathways of the Bacteria Associated with *Alvinocaris longirostris* in a Hydrothermal Vent of Okinawa Trough. *PLoS One*, 11(4), e0154359. doi:10.1371/journal.pone.0154359
507. Suzuki, Y., Kojima, S., Sasaki, T., Suzuki, M., Utsumi, T., Watanabe, H., . . . Horikoshi, K. (2006). Host-Symbiont Relationships in Hydrothermal Vent Gastropods of the Genus *Alviniconcha* from the Southwest Pacific. *Applied and Environmental Microbiology*, 72(2), 1388 - 1393.
508. Suzuki, Y., Kojima, S., Watanabe, H., Suzuki, M., Tsuchida, S., Nunoura, T., . . . Horikoshi, K. (2006). Single host and symbiont lineages of hydrothermal-vent gastropods *Ifremeria nautillei* (Provannidae): biogeography and evolution. *Marine Ecology Progress Series*, 315, 167 - 175.
509. Suzuki, Y., Sasaki, T., Suzuki, M., Nogi, Y., Miwa, T., Takai, K., . . . Horikoshi, K. (2005). Novel Chemoautotrophic Endosymbiosis between a Member of the *Epsilonproteobacteria* and the Hydrothermal-Vent Gastropod *Alviniconcha* aff. *hessleri* (Gastropoda: Provannidae) from the Indian Ocean. *Applied and Environmental Microbiology*, 71(9), 5440 - 5450.
510. Suzuki, Y., Sasaki, T., Suzuki, M., Tsuchida, S., Nealson, K. H., & Horikoshi, K. (2005). Molecular phylogenetic and isotopic evidence of two lineages of chemoautotrophic endosymbionts distinct at the subdivision level harbored in one host-animal type: The genus *Alviniconcha* (Gastropoda: Provannidae). *FEMS Microbiology Letters*, 249(1), 105 - 112.
511. Suzuki, Y., Suzuki, M., Tsuchida, S., Takai, K., Horikoshi, K., Southward, A. J., . . . Yamaguchi, T. (2009). Molecular investigations of the stalked barnacle *Vulcanolepas osheai* and the epibiotic bacteria from the Brothers Caldera, Kermadec Arc, New Zealand. *Journal of the Marine Biological Association of the United Kingdom*, 89(4), 727 - 733. doi:10.1017/S0025315409000459
512. Sweetman, A. K., Levin, L. A., Rapp, H. T., & Schander, C. (2013). Faunal trophic structure at hydrothermal vents on the southern Mohn's Ridge, Arctic Ocean. *Marine Ecology Progress Series*, 473, 115 - 131.
513. Taboada, S., Wiklund, H., Glover, A. G., Dahlgren, T. G., Cristobo, J., & Avila, C. (2013). Two new Antarctic *Ophryotrocha* (Annelida: Dorvilleidae) described from shallow-water whale bones. *Polar Biology*, 36(7), 1031 - 1045. doi:10.1007/s00300-013-1326-4

514. *Takahashi, Y., Sasaki, Y., Chikaraishi, Y., Tsuchiya, M., Watanabe, H., Asahida, T., . . . Fujikura, K. (2012). Does the symbiotic scale-worm feed on the host mussel in deep-sea vent fields? *Organic Geochemistry*, 28, 23 - 26.
515. *Takeda, M., & Hashimoto, J. (1990). A new species of the genus *Paralomis* (Crustacea, Decapoda, Lithodidae) from the Minami-Ensei Knoll in the Mid-Okinawa Trough. *Bulletin of the National Science Museum, Tokyo, Series A (Zoology)*, 16(2), 79 - 88.
516. Tanaka, H., & Yasuhara, M. (2016). A New Deep-Sea Hydrothermal Vent Species of Ostracoda (Crustacea) from the Western Pacific: Implications for Adaptation, Endemism, and Dispersal of Ostracodes in Chemosynthetic Systems. *Zoological Science*, 33(5), 555 - 565.
517. Tandberg, A. H. S., Vader, W., Olsen, B. R., & Rapp, H. T. (2018). *Monoculodes bousfieldi* sp. n. from the Arctic hydrothermal vent Loki's Castle. *Marine Biodiversity*, doi:10.1007/s12526-018-0869-6.
518. Tandberg, A. H., Rapp, H. T., Schander, C., & Vader, W. (2013). A new species of *Exitomelita* (Amphipoda: Melitidae) from a deep-water wood fall in the northern Norwegian Sea. *Journal of Natural History*, 47(25 - 28).
519. Tandberg, A. H., Rapp, H. T., Schander, C., Vader, W., Sweetman, A. K., & Berge, J. (2012). *Exitomelita sigynae* gen. et sp. nov.: a new amphipod from the Arctic Loki Castle vent field with potential gill ectosymbionts. *Polar Biology*, 35(5), 705 - 716. doi:10.1007/s00300-011-1115-x
520. Teal, J. M. (1971). Pressure Effects on the Respiration of Vertically Migrating Decapod Crustacea. *Am. Zoologist*, 11, 571 - 576.
521. *Thaler, A. D., Plouviez, S., Saleu, W., Alei, F., Jacobson, A., Boyle, E. A., . . . Van Dover, C. L. (2014). Comparative Population Structure of Two Deep-Sea Hydrothermal-Vent-Associated Decapods (*Chorocaris* sp. 2 and *Munidopsis lauensis*) from Southwestern Pacific Back-Arc Basins. *PLoS One*, 9(7), e101345.
522. Thaler, A. D., Zelnio, K., Saleu, W., Schultz, T. F., Carlsson, J., Cunningham, C., . . . Van Dover, C. L. (2011). The spatial scale of genetic subdivision in populations of *Ifremeria nautillei*, a hydrothermal-vent gastropod from the southwest Pacific. *BMC Evolutionary Biology*, 11(1), 372. doi:10.1186/1471-2148-11-372
523. Thatje, S., Marsh, L., Roterman, C. N., Mavrogordato, M. N., & Linse, K. (2015). Adaptations to Hydrothermal Vent Life in *Kiwa tyleri*, a New Species of Yeti Crab from the East Scotia Ridge, Antarctica. *PLoS One*, 10(6), e0127621, 1 - 20.
524. Thiel, V., Hügler, M., Blümel, M., Baumann, H. I., Gärtner, A., Schmaljohann, R., . . .

Imhoff, J. F. (2012). Widespread Occurrence of Two Carbon Fixation Pathways in Tubeworm Endosymbionts: Lessons from Hydrothermal Vent Associated Tubeworms from the Mediterranean Sea. *Frontiers in Microbiology*, 3, 423. doi:10.3389/fmicb.2012.00423

525. Thubaut, J., Puillandre, N., Faure, B., Cruaud, C., & Samadi, S. (2013). The contrasted evolutionary fates of deep-sea chemosynthetic mussels (Bivalvia, Bathymodiolinae). *Ecology and Evolution*, 3(14), 4748 - 4766. doi:10.1002/ece3.749
526. Thurston, M. H. (1990). Abyssal necrophagous amphipods (Crustacea: Amphipoda) in the northeast and tropical Atlantic Ocean. *Progress in Oceanography*, 24(1), 257 - 274. doi:10.1016/0079-6611(90)90036-2
527. *Todt, C., & Von Salvini-Plawen, L. (2005). The digestive tract of *Helicoradomenia* (Solenogastres, Mollusca), aplacophoran molluscs from the hydrothermal vents of the East Pacific Rise. *Invertebrate Biology*, 124(3), 230 - 253.
528. Tokuda, G., Yamada, A., Nakano, K., Arita, N. O., & Yamasaki, H. (2008). Colonization of *Sulfurovum* sp. on the gill surfaces of *Alvinocaris longirostris*, a deep-sea hydrothermal vent shrimp. *Marine Ecology*, 29(1), 106 - 114.
529. *Tomoyuki, M. (1986). Japanese Polychaetes of the Genera *Eunice* and *Euniphysa*: Taxonomy and Branchial Distribution Patterns. *Publications of the Seto Marine Biological Laboratory*, 31(3 - 6), 269 - 325.
530. Topsent, E. (1909). Étude sur quelques *Cladhoriza* et sur *Euchelipluma pristina* n. g. et n. sp. *Bulletin de l'Institut Océanographique (Fondation Albert I", Prince de Monaco)*, 151, 1 - 23.
531. Topsent, E. (1928). Spongiaires de l'Atlantique et de la Méditerranée provenant des croisières du Prince Albert Ier de Monaco. *Résultats des Campagnes Scientifiques du Prince Albert I Monaco*, 74, 1 - 376.
532. *Trask, J. L., & Van Dover, C. L. (1999). Site-specific and ontogenetic variations in nutrition of mussels (*Bathymodiolus* sp.) from the Lucky Strike hydrothermal vent field, Mid-Atlantic Ridge. *Limnology and Oceanography*, 44(2), 334 - 343. doi:10.4319/lo.1999.44.2.0334
533. Tsuchida, S., & Hashimoto, J. (2002). A new species of Bythograeid crab, *Austinograea rodriguezensis* (Decapoda, Bathyura), associated with active hydrothermal vents from the Indian Ocean. *Journal of Crustacean Biology*, 22(3), 642 - 650. doi:10.1651/0278-0372(2002)022[0642:ANSOBC]2.0.CO;2
534. Tsuchida, S., Suzuki, Y., Fujiwara, Y., Kawato, M., Uematsu, K., Yamanaka, T., . . . Yamamoto, H. (2010). Epibiotic association between filamentous bacteria and the

vent-associated galatheid crab, *Shinkaia crosnieri* (Decapoda: Anomura). *Journal of the Marine Biological Association of the United Kingdom*, 91(1), 23 - 32.
doi:10.1017/S0025315410001827

535. Tsurumi, M. (2003). Diversity at hydrothermal vents. *Global Ecology and Biogeography*, 12, 181 - 190.
536. *Tsurumi, M. (2001). *Ecology of hydrothermal vents on three segments of the Juan de Fuca Ridge, northeast Pacific*. University of Victoria.
537. Tsurumi, M., de Graaf, R. C., & Tunnicliffe, V. (2003). Distributional and biological aspects of copepods at hydrothermal vents on the Juan de Fuca Ridge, north-east Pacific ocean. *Journal of the Marine Biological Association of the UK*, 83, 469 - 477.
538. Tsurumi, M., & Tunnicliffe, V. (2003). Tubeworm-associated communities at hydrothermal vents on the Juan de Fuca Ridge, northeast Pacific. *Deep-sea Research I*, 50(5), 611 - 629. doi:10.1016/s0967-0637(03)00039-6
539. Tunnicliffe, V. (1991). The biology of hydrothermal vents: ecology and evolution. *Oceanography and Marine Biology: An Annual Review*, 29, 319 - 407.
540. *Tunnicliffe, V., Koop, B. F., Tyler, J., & So, S. (2010). Flatfish at seamount hydrothermal vents show strong genetic divergence between volcanic arcs. *Marine Ecology*, 31, 158 - 167. doi:10.1111/j.1439-0485.2010.00370.x
541. Tunnicliffe, V., McArthur, A. G., & McHugh, D. (1998). A Biogeographical Perspective of the Deep-Sea Hydrothermal Vent Fauna. In J. H. S. Blaxter, A. J. Southward, & P. A. Tyler (Eds.), *Advances in Marine Biology* (Vol. 34, pp. 353 - 442): Academic Press.
542. *Tunnicliffe, V., & Southward, A. J. (2004). Growth and breeding of a primitive stalked barnacle *Leucolæpas longa* (Cirripedia: Scalpellomorpha: Eolepadidae: Neolepadinae) inhabiting a volcanic seamount off Papua New Guinea. *Journal of the Marine Biological Association of the United Kingdom*, 84(1), 121 - 132.
doi:10.1017/S0025315404008987h
543. *Tunnicliffe, V., St. Germain, C., & Hilário, A. (2014). Phenotypic Variation and Fitness in a Metapopulation of Tubeworms (*Ridgeia piscesae* Jones) at Hydrothermal Vents. *PLoS One*, 9(10), e110578. doi:10.1371/journal.pone.0110578
544. *Tunnicliffe, V., Tyler, J., & Dower, J. F. (2013). Population ecology of the tonguefish *Syphurus thermophilus* (Pisces; Pleuronectiformes; Cynoglossidae) at sulphur-rich hydrothermal vents on volcanoes of the northern Mariana Arc. *Deep Sea Research Part II: Topical Studies in Oceanography*, 92, 172 - 182.
doi:10.1016/j.dsr2.2013.01.026

545. *Tuset, V. M., Espinosa, D. I., García-Mederos, A., Santana, J. I., & González, J. A. (2011). Egg development and fecundity estimation in deep-sea red crab, *Chaceon affinis* (Geryonidae), off the Canary Islands (NE Atlantic). *Fisheries Research*, 109(2), 373 - 378. doi:10.1016/j.fishres.2011.02.021
546. *Tyler, P. A., Pendlebury, S., Mills, S. W., Mullineaux, L., Eckelbarger, K. J., Baker, M., & Young, C. M. (2008). Reproduction of Gastropods from Vents on the East Pacific Rise and the Mid-Atlantic Ridge. *Journal of Shellfish Research*, 27(1), 107 - 118.
547. *Tyler, P. A., & Young, C. M. (2003). Dispersal at hydrothermal vents: a summary of recent progress. *Hydrobiologia*, 503, 9 - 19.
548. *Tyler, P. A., & Young, C. M. (1999). Reproduction and dispersal at vents and cold seeps. *Journal of the Marine Biological Association of the U.K.*, 79, 193 - 208.
549. Urakawa, H., Dubilier, N., Fujiwara, Y., Cunningham, D. E., Kojima, S., & Stahl, D. A. (2005). Hydrothermal vent gastropods from the same family (Provannidae) harbour ε - and γ -proteobacterial endosymbionts. *Environmental Microbiology*, 7(5), 750 - 754. doi:10.1111/j.1462-2920.2005.00753.x
550. Vacelet, J. (2006). New carnivorous sponges (Porifera, Poecilosclerida) collected from manned submersibles in the deep Pacific. *Zoological Journal of the Linnean Society*, 148(4), 553 - 584.
551. Valdés, Á., & Bouchet, P. (1998). Naked in toxic fluids: A nudibranch mollusc from hydrothermal vents. *Deep Sea Research Part II: Topical Studies in Oceanography*, 45, 319 - 327.
552. *Valdés, F., Sellanes, J., & D'Elía, G. (2012). Phylogenetic Position of Vesicomyid Clams from a Methane Seep off Central Chile ($\sim 36^{\circ}$ S) with a Molecular Timescale for the Diversification of the Vesicomyidae. *Zoological Studies*, 51(7), 1154 - 1164.
553. van der Heijden, K., Petersen, J. M., Dubilier, N., & Borowski, C. (2012). Genetic Connectivity between North and South Mid-Atlantic Ridge Chemosynthetic Bivalves and Their Symbionts. *PLoS One*, 7(7), e39994, 1 - 14.
554. Van Dover, C. L. (2002). Trophic relationships among invertebrates at the Kairei hydrothermal vent field (Central Indian Ridge). *Marine Biology*, 141(4), 761 - 772.
555. Van Dover, C. L., Aronson, J., Pendleton, L., Smith, S., Arnaud-Haond, S., Moreno-Mateos, D., . . . Warner, R. (2014). Ecological restoration in the deep sea: Desiderata. *Marine Policy*, 44, 98 - 106. doi:10.1016/j.marpol.2013.07.006
556. Van Dover, C. L., Desbruyères, D., Segonzac, M, Comtet, T., Saldanha, L., Fiala-

- Medioni, A., & Langmuir, C. (1996) Biology of the Lucky Strike hydrothermal field. *Deep Sea Research Part I: Oceanographic Research Papers*, 43(9), 1509 - 1529.
557. *Van Dover, C. L., Factor, J. R., Williams, A. B., & Berg, C. J., Jr. (1985). Reproductive patterns of decapod crustaceans from hydrothermal vents. *Bulletin of the Biological Society of Washington*, 6, 223 - 227.
558. Van Dover, C. L., & Fry, B. (1994). Microorganisms as food resources at deep-sea hydrothermal vents. *Limnology and Oceanography*, 39(1), 51 - 57.
559. *Van Dover, C. L., & Fry, B. (1989). Stable isotopic compositions of hydrothermal vent organisms. *Marine Biology*, 102(2), 257 - 263. doi:10.1007/BF00428287
560. Van Dover, C. L., Humphris, S. E., Fornari, D., Cavanaugh, C. M., Collier, R., Goffredi, S. K., ..., & Vrijenhoek, R. C. (2001). Biogeography and ecological setting of Indian Ocean hydrothermal vents. *Science*, 294(5543), 818 - 823.
561. Van Dover, C. L., & Trask, J. L. (2000) Diversity at deep-sea hydrothermal vent and intertidal mussel beds. *Marine Ecology Progress Series*, 195, 169 - 178.
562. van Harten, D. (1992). Hydrothermal vent ostracoda and faunal association in the deep sea. *Deep-Sea Research*, 39(6), 1067 - 1070.
563. Vereshchaka, A. L., Kulagin, D. N., & Lunina, A. A. (2015). Phylogeny and New Classification of Hydrothermal Vent and Seep Shrimps of the Family Alvinocarididae (Decapoda). *PLoS One*, 10(7), e0129975. doi:10.1371/journal.pone.0129975
564. Vereshchaka, A. L., Vinogradov, G. M., Lein, A. Y., Dalton, S., & Dehairs, F. (2000). Carbon and nitrogen isotopic composition of the fauna from the Broken Spur hydrothermal vent field. *Mar Biol*, 136(1), 11 - 17. doi:10.1007/s002270050002
565. Vinogradov, G. M. (1995). Colonization of pelagic and hydrothermal vent habitats by gammaridean amphipods: an attempt of reconstruction. *Polish Archives of Hydrobiology*, 42(4), 417 - 430.
566. Vinogradov, G. M. (1993). Amfipody (Crustacea) s "chernykh Kuril'shchikov" iz rajonov gidrotermal'noj aktivnosti vostochnoj chasti Tikhogo Okeana. [Amphipods (Crustacea) from hydrothermal vents of the Eastern Pacific]. *Zoologicheskij Zhurnal*, 72(2), 40 - 52.
567. Vinogradov, G., Hernández, F., Tejera, E., & León, M. E. (2004). Anfípodos pelágicos de las islas de Cabo Verde (Campaña TFMCBM/98, Proyecto-Macaronesia 2000). *VIERAEA*, 32, 7 - 27.

568. Voight, J. R. (2000). A review of predators and predation at deep-sea hydrothermal vents. *Cahiers de Biologie Marine*, 41(2), 155 - 166.
569. *von Cosel, R. (2002). A new species of bathymodioline mussel (Mollusca Bivalvia Mytilidae) from Mauritania (West Africa) with comments on the genus *Bathymodiolus* Kenk & Wilson 1985. *Zoosystema*, 24, 259 - 271.
570. *von Cosel, R., Comtet, T., & Krylova, E. M. (1999). *Bathymodiolus* (Bivalvia: Mytilidae) from hydrothermal vents on the Azores triple junction and the Logatchev hydrothermal field, Mid-Atlantic Ridge,. *The Veliger*, 42(3), 218 - 248.
571. von Cosel, R., & Marshall, B. A. (2010). A new genus and speciesof large mussel (Mollusca: Bivalvia: Mytilidae) from the Kermadec Ridge. *Tuhinga*, 21, 59 - 73.
572. von Cosel, R., & Marshall, B. A. (2003). Two new species of large mussels (Bivalvia: Mytilidae) from active submarine volcanoes and a cold seep off the eastern North Island of New Zealand, with description of a new genus. *The Nautilus*, 117(2), 31 - 46.
573. von Cosel, R., Métivier, B., & Hashimoto, J. (1994). Three New Species of *Bathymodiolus* (Bivalvia: Mytilidae) from Hydrothermal Vents in the Lau Basin and the North Fiji Basin, Western Pacific, and the Snake Pit Area, Mid-Atlantic Ridge. *The Veliger*, 37, 374 - 392.
574. Ward, M. E., Jenkins, C. D., & Van Dover, C. L. (2003). Functional morphology and feeding strategy of the hydrothermal-vent polychaete *Archinome rosacea* (family Archinomidae). *Canadian Journal of Zoology*, 81(4), 582 - 590. doi:10.1139/z03-034
575. Warén, A., & Bouchet, P. (2001). Gastropoda and Monoplacophora from Hydrothermal Vents and Seeps; New Taxa and Records. *The Veliger*, 44(2), 116 - 231.
576. Warén, A., & Bouchet, P. (1993). New records, species, genera, and a new family of gastropods from hydrothermal vents and hydrocarbon seeps. *Zoologica Scripta*, 22(1), 1 - 90.
577. Warén, A., & Bouchet, P. (1989). New gastropods from East Pacific hydrothermal vents. *Zoologica Scripta*, 18(1), 67 - 102.
578. *Warén, A., & Bouchet, P. (1986). Four new species of *Provanna* Dall (Prosobranchia, Cerithiacea?) from East Pacific hydrothermal sites. *Zoologica Scripta*, 15(2), 157 - 164. doi:10.1111/j.1463-6409.1986.tb00218.x
579. No reference is associated with this number (as the reference originally included here was a duplicate). This number can therefore be used as a reference code in future versions of sFDvent.

580. Watabe, H., & Hashimoto, J. (2002). A New Species of the Genus *Rimicaris* (Alvinocarididae: Caridea: Decapoda) from the Active Hydrothermal Vent Field, "Kairei Field," on the Central Indian Ridge, the Indian Ocean. *Zoological Science*, 19(10), 1167 - 1174. doi:10.2108/zsj.19.1167
581. Watanabe, H., & Beedessee, G. (2015). Vent Fauna on the Central Indian Ridge. In J. Ishibashi, K. Okino, & M. Sunamura (Eds.), *Subseafloor Biosphere Linked to Hydrothermal Systems: TAIGA Concept* (pp. 205 - 212). Tokyo: Springer Japan.
582. *Watanabe, K. H., Chen, C., Marie, D. P., Takai, K., Fujikura, K., & Chan, B. K. K. (2018) Phylogeography of hydrothermal vent stalked barnacles: a new species fills a gap in the Indian Ocean 'dispersal corridor' hypothesis. *Royal Society Open Science*, 5(4), 1 - 18.
583. *Watanabe, H., Fujikura, K., Kojima, S., Miyazaki, J.-I., & Fujiwara, Y. (2010). Japan: Vents and Seeps in Close Proximity. In S. Kiel (Ed.), *The Vent and Seep Biota*. (Vol. 33, pp. 379 - 401). Dordrecht: Springer.
584. *Watanabe, H., Kado, R., Tsuchida, S., Miyake, H., Kyo, M., & Kojima, S. (2004). Larval development and intermoult period of the hydrothermal vent barnacle *Neoverruca* sp. *Journal of the Marine Biological Association of the United Kingdom*, 84(4), 743 - 745. doi:10.1017/S0025315404009841h
585. *Watanabe, H., & Kojima, S. (2015). Vent Fauna in the Okinawa Trough. In J. Ishibashi, K. Okino, & M. Sunamura (Eds.), *Subseafloor Biosphere Linked to Hydrothermal Systems: TAIGA Concept* (pp. 449 - 459). Tokyo: Springer Japan.
586. Watanabe, H., Seo, E., Takahashi, Y., Yoshida, T., Kojima, S., Fujikura, K., & Miyake, H. (2012). Spatial distribution of sister species of vesicomyid bivalves *Calyptogena okutanii* and *Calyptogena soyoae* along an environmental gradient in chemosynthetic biological communities in Japan. *Journal of Oceanography*, 69(1), 129 - 134. doi:10.1007/s10872-012-0155-3
587. *Watanabe, H., Yahagi, T., Nagai, Y., Seo, M., Kojima, S., Ishibashi, J., . . . Toyofuku, T. (2016). Different thermal preferences for brooding and larval dispersal of two neighboring shrimps in deep-sea hydrothermal vent fields. *Marine Ecology*, 37(6), 1282 - 1289.
588. Watson, C. (2001). New genus and species of Chrysopetalidae (Polychaeta) from hydrothermal vents (south-western Pacific). . *The Beagle, Records of the Northern Territory Museum of Arts and Sciences*, 17, 57 - 66.
589. *Watsuji, T.-o., Nakagawa, S., Tsuchida, S., Toki, T., Hirota, A., Tsunogai, U., & Takai, K. (2010). Diversity and Function of Epibiotic Microbial Communities on the Galatheid Crab, *Shinkaia crosnieri*. *Microbes Environ.*, 25(4), 288 - 294.

590. Webber, W. R. (2004). A new species of *Alvinocaris* (Crustacea: Decapoda: Alvinocarididae) and new records of alvinocarids from hydrothermal vents north of New Zealand. *Zootaxa*, 444, 1 - 26.
591. *Williams, A. B. (1988). New marine decapod crustaceans from waters influenced by hydrothermal discharge, brine, and hydrocarbon seepage. *Fishery Bulletin*, 86(2), 263 - 287.
592. *Williams, A. B. (1980). A new crab family from the vicinity of submarine thermal vents on the Galapagos Rift (Crustacea: Decapoda: Brachyura). *Proceedings of The Biological Society of Washington*, 93, 443 - 472.
593. Williams, A. B., & Chace, J. F. A. (1982). A New Caridean Shrimp of the Family Bresiliidae From Thermal Vents of the Galapagos Rift. *Journal of Crustacean Biology*, 2(1), 136 - 147. doi:10.2307/1548118
594. Williams, A. B., & Van Dover, C. L. (1983). A New Species Of *Munidopsis* From Submarine Thermal Vents Of The East Pacific Rise At 21 Degrees North Anomura Galatheidae. *Proceedings of The Biological Society of Washington*, 96, 481 - 488.
595. *Windoffer, R., & Giere, O. (1997). Symbiosis of the Hydrothermal Vent Gastropod *Ifremeria nautillei* (Provannidae) With Endobacteria - Structural Analyses and Ecological Considerations. *Biological Bulletin*, 193, 381 - 392.
596. Won, Y.-J., Hallam, S. J., O'Mullan, G. D., Pan, I. L., Buck, K. R., & Vrijenhoek, R. C. (2003). Environmental Acquisition of Thiotrophic Endosymbionts by Deep-Sea Mussels of the Genus *Bathymodiolus*. *Applied and Environmental Microbiology*, 69(11), 6785 - 6792.
597. Won, Y., Hallam, S. J., O'Mullan, G. D., & Vrijenhoek, R. C. (2003). Cytonuclear disequilibrium in a hybrid zone involving deep-sea hydrothermal vent mussels of the genus *Bathymodiolus*. *Molecular Ecology*, 12(11), 3185 - 3190. doi:10.1046/j.1365-294X.2003.01974.x
598. Won, Y.-J., Jones, W. J., & Vrijenhoek, R. C. (2008). Absence of Cospeciation Between Deep-Sea Mytilids and Their Thiotrophic Endosymbionts. *Journal of Shellfish Research*, 27(1), 129 - 138. doi:10.2983/0730-8000(2008)27[129:AOCBDM]2.0.CO;2
599. *Won, Y., Young, C. R., Lutz, R. A., & Vrijenhoek, R. C. (2003). Dispersal barriers and isolation among deep-sea mussel populations (Mytilidae: *Bathymodiolus*) from eastern Pacific hydrothermal vents. *Molecular Ecology*, 12(1), 169 - 184. doi:10.1046/j.1365-294X.2003.01726.x
600. Xiao, N., & Liao, Y. (2013). Three new records of deep-water goniasterids

(Echinodermata: Asteroidea: Goniasteridae) from China seas. *Chinese Journal of Oceanology and Limnology*, 31(5), 1036 - 1044. doi:10.1007/s00343-013-2323-8

601. *Yahagi, T., Kayama Watanabe, H., Kojima, S., & Kano, Y. (2017). Do larvae from deep-sea hydrothermal vents disperse in surface waters? *Ecology*, 98(6), 1524 - 1534. doi:10.1002/ecy.1800
602. Yahagi, T., Watanabe, H., Ishibashi, J., & Kojima, S. (2015). Genetic population structure of four hydrothermal vent shrimp species (Alvinocarididae) in the Okinawa Trough, Northwest Pacific. *Marine Ecology Progress Series*, 529, 159 - 169.
603. Yahagi, T., Watanabe, H., Kojima, S., Beedessee, G., & Komai, T. (2014). First record and a new species of *Alvinocaris* Williams & Chace, 1982 (Crustacea: Decapoda: Caridea: Alvinocarididae) from the Indian Ocean. *Zootaxa*, 3893(1), 101 - 113.
604. Yamaguchi, T., & Newman, W. A. (1997). *Eochionelasmus paquensis*, new species (Cirripedia: Balanomorpha) from 17°25'S, north of Easter Island: First record of a sessile hydrothermal barnacle from the East Pacific Rise. *Journal of Crustacean Biology*, 17(3), 488 - 496.
605. *Yamaguchi, T., & Newman, W. A. (1990). A new and primitive barnacle (Cirripedia: Balanomorpha) from the North Fiji Basin abyssal hydrothermal field, and its evolutionary implications. *Pacific Science*, 44(2), 135 - 155.
606. *Yamaguchi, T., Newman, W. A., & Hashimoto, J. (2004). A cold seep barnacle (Cirripedia: Neolepadinae) from Japan and the age of the vent/seep fauna. *Journal of the Marine Biological Association of the United Kingdom*, 84(1), 111 - 120. doi:10.1017/S0025315404008975h
607. Yamanaka, T., Shimamura, S., Nagashio, H., Yamagami, S., Onishi, Y., Hyodo, A., . . . Mizota, C. (2015). A Compilation of the Stable Isotopic Compositions of Carbon, Nitrogen, and Sulfur in Soft Body Parts of Animals Collected from Deep-Sea Hydrothermal Vent and Methane Seep Fields: Variations in Energy Source and Importance of Subsurface Microbial Processes in the Sediment-Hosted Systems. . In J. Ishibashi, K. Okino, & M. Sunamura (Eds.), *Subseafloor Biosphere Linked to Hydrothermal Systems*. (pp. 105 - 129). Tokyo: Springer.
608. Yang, C.-H., Tsuchida, S., Fujikura, K., Fujiwara, Y., Kawato, M., & Chan, T.-Y. (2016). Connectivity of the squat lobsters *Shinkaia crosnieri* (Crustacea: Decapoda: Galatheidae) between cold seep and hydrothermal vent habitats. *Bulletin of Marine Science*, 92(1), 17 - 31.
609. *Yang, S.-H., Chiang, P.-W., Hsu, T.-C., Kao, S.-J., & Tang, S.-L. (2016). Bacterial Community Associated with Organs of Shallow Hydrothermal Vent Crab *Xenograpsus testudinatus* near Kuishan Island, Taiwan. *PLoS One*, 11(3), e0150597. doi:10.1371/journal.pone.0150597

610. *Yorisue, T., Inoue, K., Miyake, H., & Kojima, S. (2012). Trophic structure of hydrothermal vent communities at Myojin Knoll and Nikko Seamount in the northwestern Pacific: Implications for photosynthesis-derived food supply. *Plankton Benthos Research*, 7(2), 35 - 40.
611. *Yorisue, T., Kado, R., Watanabe, H., Høeg, J. T., Inoue, K., Kojima, S., & Chan, B. K. K. (2013). Influence of water temperature on the larval development of *Neoverruca* sp. and *Ashinkailepas seepiophila*—Implications for larval dispersal and settlement in the vent and seep environments. *Deep Sea Research Part I: Oceanographic Research Papers*, 71, 33 - 37.
612. *Young, C. M., & Eckelbarger, K. J. (1994). *Reproduction, larval biology, and recruitment of the deep-sea benthos*: Columbia University Press.
613. Zarenkov, N. A. (1976). On the fauna of decapods of the waters adjacent to South America. *Biologiya Morya*, 5, 8 - 18.
614. Zbinden, M., Marqué, L., Gaudron, S. M., Ravaux, J., Léger, N., & Duperron, S. (2015). Epsilonproteobacteria as gill epibionts of the hydrothermal vent gastropod *Cyathermia naticoides* (North East-Pacific Rise). *Marine Biology*, 162(2), 435 - 448.
615. Zekely, J., Gollner, S., Van Dover, C. L., Govenar, B., Le Bris, N., Nemeschkal, H. L., & Bright, M. (2006). Nematode communities associated with tubeworm and mussel aggregations on the East Pacific Rise. *Cahiers de Biologie Marine*, 47, 477 - 482.
616. Zekely, J., Sørensen, M. V., & Bright, M. (2006). Three nematode species (Monysteridae) from deep-sea hydrothermal vents. *Meiofauna Mar*, 15, 25 - 42.
617. Zekely, J., Van Dover, C. L., Nemeschkal, H. L., & Bright, M. (2006). Hydrothermal vent meiobenthos associated with mytilid mussel aggregations from the Mid-Atlantic Ridge and the East Pacific Rise. *Deep Sea Research Part I: Oceanographic Research Papers*, 53(8), 1363 - 1378. doi:10.1016/j.dsr.2006.05.010
618. Zelnio, K. A. (2009). *Community structure of hydrothermal vents at the Eastern-Lau Spreading Center*. (Master of Science), The Pennsylvania State University. Retrieved from: https://etda.libraries.psu.edu/files/final_submissions/3338.
619. Zelnio, K. A., & Hourdez, S. (2009). A New Species of *Alvinocaris* (Crustacea: Decapoda: Caridea: Alvinocarididae) from Hydrothermal Vents at The Lau Basin, Southwest Pacific, and a Key to The Species of Alvinocarididae. *Proceedings of The Biological Society of Washington*, 122(1), 52 - 71. doi:10.2988/07-28.1
620. Zelnio, K., Rodriguez, E., & Daly, M. (2009). Hexacorals (Anthozoa: Actiniaria, Zoanthidea) from hydrothermal vents in the south-western Pacific. *Marine Biology*

Research, 5(6), 547 - 571.

621. *Zelnio, K. (2007). From The Desk of Zelnio: *Alviniconcha hessleri*. Deep Sea News. Retrieved from: <http://www.deepseanews.com/2007/03/from-the-desk-of-zelnio-alviniconcha-hessleri/>.
622. Zottoli, R. (1983). *Amphisamytha galapagensis*, a new species of Ampharetid polychaete from the vicinity of abyssal hydrothermal vents in the Galapagos Rift, and the role of this species in rift ecosystems. *Proceedings of The Biological Society of Washington*, 96(3), 379 - 391.
623. Zwirglmaier, K., Reid, W. D., Heywood, J., Sweeting, C. J., Wigham, B. D., Polunin, N. V., . . . Linse, K. (2015). Linking regional variation of epibiotic bacterial diversity and trophic ecology in a new species of Kiwaidae (Decapoda, Anomura) from East Scotia Ridge (Antarctica) hydrothermal vents. *Microbiologyopen*, 4(1), 136 - 150. doi:10.1002/mbo3.227
624. Team, P. (2017). *Polytraits: A database on biological traits of polychaetes*. Retrieved from: <http://polytraits.lifewatchgreece.eu>.
625. *No reference is associated with this number (as the reference originally included here was a duplicate). This number can therefore be used as a reference code in future versions of sFDvent.*
626. Pinho, M. R., Gonçalves, J. M., Martins, H. R., & Menezes, G. M. (2001). Some aspects of the biology of the deep-water crab, *Chaceon affinis* (Milne-Edwards and Bouvier, 1894) off the Azores, *Fisheries Research*, 51(2-3), 283 - 295.
627. Fishwise Professional (2019). *Fishwise Professional*. Retrieved from: <https://www.fishwisepro.com/>.
628. Foxton (1972). Observations on the vertical distribution of the genus *Acanthephyra* (Crustacea: Decapoda) in the eastern North Atlantic, with particular reference to species of the 'purpurea' group. *Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences*, 73, 301 - 313.
629. Wang, Y-R., & Sha, Z-L. (2017). A new species of the genus *Alvinocaris* Williams and Chace, 1982 (Crustacea: Decapoda: Caridea: Alvinocarididae) from the Manus Basin hydrothermal vents, Southwest Pacific. *Zootaxa*, 4226(1).
630. Krylova, E. M., & Sahling, H. (2010). Vesicomyidae (Bivalvia): Current taxonomy and distribution. *PLoS One*, 5(4), e9957.
631. Burreson, E. M. (1981). A new deep-sea leech *Bathybdella sawyeri* n. gen., n. sp., from thermal vent areas on the Galápagos Rift, *Proc. Biol. Soc. Wash.*, 94, 483 - 491.

632. Yamanaka, T., Mizota, C., Fujiwara, Y., Chiba, H., Hashimoto, J., Gamo, T., & Okudaira, T. (2003). Sulphur-isotopic composition of the deep-sea mussel *Bathymodiolus marisindicus* from currently active hydrothermal vents in the Indian Ocean. *Journal of the Marine Biological Association of the United Kingdom*, 83(4), 841 - 848.
633. Yamanaka, T., Mizota, C., Maki, Y., Fujikura, K., & Chiba, H. (2000). Sulfur isotope composition of soft tissues of deep-sea mussels, *Bathymodiolus* spp., in Japanese waters. *Benthos Research*, 55(2), 63 - 68.
634. Pettibone, M. H. (1989). New species of scale-worms (Polychaeta: Polynoidae) from the hydrothermal rift-area of the Mariana back-arc basin in the western Central Pacific, *Proc. Biol. Soc. Wash.*, 102(1), 137 - 153.
635. Reiss, C. S., Walsh, J., & Goebel, M. E. (2015) Winter preconditioning determines feeding ecology of *Euphausia superba* in the Antarctic Peninsula, *Marine Ecology Progress Series*, 519, 89 - 101.
636. Aberle, N., Malzahn, A. M., Lewandowska, A. M., & Sommer, U. (2015). Some like it hot: the protozooplankton-copepod link in a warming ocean. *Marine Ecology Progress Series*, 519, 103 - 113.
637. Shaw, P. (1989). New amphipods from geothermal vent sites of the west coast of Vancouver Island, British Columbia, with a reappraisal of the amhipod family Sebidae. *Canadian Journal of Zoology*, 67, 1882 - 1890.
638. Van Dover, C. L., & Doerries, M. B. (2005). Community structure in mussel beds at Logatchev hydrothermal vents and a comparison of macrofaunal species richness on slow- and fast-spreading mid-ocean ridges, *Marine Ecology*, 26(2), 110 - 120.
639. Daguin, C., & Jollivet, D. (2005). Development and cross-amplification of nine polymorphic microsatellite markers in the deep-sea hydrothermal vent polychaete *Branchipolynoe seepensis*, *Molecular Ecology Resources*, 5(4), 780 - 783.
640. Geistdoerfer, P., Auzende, J-M., Batiza, R., Bideau, D., Cormier, M., Fouquet, ... & Spadea, P. (1995). Hydrothermalisme et communautés animales associées sur la dorsale du Pacifique oriental entre 17°S et 19°S (campagne Naudur, décembre 1993), *Comptes Rendus de l'Academie des Sciences Serie II* (1251-8069) (Gauthier-villars), 320(1), 47 - 54.
641. Roskov Y., Ower G., Orrell T., Nicolson D., Bailly N., Kirk P.M., ... & Penev L., (eds.) (2019). *Species 2000 & ITIS Catalogue of Life*. Retrieved from: www.catalogueoflife.org/col. Species 2000: Naturalis, Leiden, the Netherlands.
642. S. Nakagawa, pers. comm.

643. Fox, M., Juniper, S. K., & Vali, H. (2002). Chemoautotrophy as a possible nutritional source in the hydrothermal vent limpet *Lepetodrilus fucensis*. *Cah. Biol. Mar.*, 43, 371 – 376.
644. Arronte, J. C., Bañón, R. , Quigley, D. T., Pis-Millán, J. A., & Heredia, J. (2011). New data on *Lepidion schmidti* (Gadiformes: Moridae) from the north-east Atlantic Ocean. *Journal of Fish Biology*, 79, 1708 - 1721.
645. Rogers, A.D. (ed.) (2010). RRS *James Cook* Cruise JC042. Cruise Report JC42: 'Cruise Report', accessible online via:
https://www.bodc.ac.uk/data/information_and_inventories/cruise_inventory/report/jc042.pdf
646. ETI Bioinformatics. (n.d.) *Marine Species Identification Portal*. Retrieved from: <http://species-identification.org/index.php>
647. Takai, K. *et al.*, unpublished data.
648. Foxton, P., & Roe, H. S. J. (1974). Observations on the nocturnal feeding of some mesopelagic decapod crustacea. *Marine Biology*, 28(1), 37 - 49.
649. Deevey, G. B. (1968). *Bathyconchoecia*, a new genus of pelagic ostracods (Myodocopa: Halocypridae) with six new species from the deeper waters of the Gulf of Mexico. *Proceedings of the Biological Society of Washington*, 81, 539 - 570.
650. Beaulieu, S. E. (2015). *InterRidge Global Database of Active Submarine Hydrothermal Vent Fields: prepared for InterRidge, Version 3.4*. Retrieved from: <http://vents-data.interridge.org>
651. J. Haywood, pers. comm.
652. Eckner, J. T. (1999). *Analysis of community composition and trophic interactions in a hydrothermal vent vestimentiferan tubeworm community collection*. Honors undergraduate thesis in biology. The Pennsylvania State University.
653. Goroslavskaya, E. I., & Galkin, S. V. (2011). Benthic fauna associated with mussel beds and shrimp swarms at hydrothermal fields on the Mid-Atlantic Ridge, *Oceanology*, 51(1), 69 - 79.
654. Fujikura, K., Sasaki, T., Yamanaka, T., & Yoshida, T. (2009). Turrids whelk, *Phymorhynchus buccinoides* feeds on *Bathymodiolus* mussels at a seep site in Sagami Bay, Japan, *Plankton and Benthos Research*, 4(1), 23 - 30.

655. Rybakova (Goroslavskaya), E., & Galkin, S. (2015). Hydrothermal assemblages associated with different foundation species on the East Pacific Rise and Mid-Atlantic Ridge, with a special focus on mytilids, *Marine Ecology*, 36(S1), 45 - 61.
656. Horton, T., Kroh, A., Bailly, N., Boury-Esnault, N., Brandão, S. N., Costello, M. J., . . . Zeidler, W. (2017). *World Register of Marine Species (WoRMS)*. Retrieved from: WoRMS Editorial Board <http://www.marinespecies.org>.

-9999 Gap filled using Location Information entered and/or the InterRidge database.
-9998 Gap filled using the score for 'Trophic Mode'.
-9997 Gap filled according to the typical score for the genus.
-9996 Gap filled according to the most common (mode) score for the taxonomic family, or higher taxonomic groups.
-9995 Gap filled using score for 'Position of Symbiont'.