

---

## Resistance of the oyster pathogen *Vibrio tasmaniensis* LGP32 against grazing by *Vannella* sp. marine amoeba involves Vsm and CopA virulence factors

Robino Etienne <sup>1,5</sup>, Poirier Aurore C. <sup>6</sup>, Amraoui Hajar <sup>6</sup>, Le Bissonnais Sandra <sup>6</sup>, Perret Angelique <sup>1</sup>, Lopez-joven Carmen <sup>7</sup>, Auguet Jean-Christophe <sup>2</sup>, Rubio Tristan P. <sup>8</sup>, Cazevieille Chantal <sup>3</sup>, Rolland Jean-Luc <sup>1</sup>, Héchard Yann <sup>4</sup>, Destoumieux-garzón Delphine <sup>8</sup>, Charrière Guillaume <sup>6,\*</sup>

<sup>1</sup> IHPE UMR 5244, CNRS, Ifremer, University of Montpellier, University of Perpignan Via Domitia Montpellier, France

<sup>2</sup> MARBEC, University of Montpellier, CNRS, Ifremer, IRD Montpellier, France

<sup>3</sup> INM, University of Montpellier Montpellier, France

<sup>4</sup> EBI UMR CNRS 7267, University of Poitiers Poitiers, France

\* Corresponding author : Guillaume Charrière, email address : [guillaume.charriere@umontpellier.fr](mailto:guillaume.charriere@umontpellier.fr)

---

### Abstract :

ibrios are ubiquitous in marine environments and opportunistically colonize a broad range of hosts. Strains of *Vibrio tasmaniensis* present in oyster farms can thrive in oysters during juvenile mortality events and behave as facultative intracellular pathogen of oyster hemocytes. Herein, we wondered whether *V. tasmaniensis* LGP32 resistance to phagocytosis is specific to oyster immune cells or contributes to resistance to other phagocytes, like marine amoebae. To address this question, we developed an integrative study, from the first description of amoeba diversity in oyster-farms to the characterization of LGP32 interactions with amoebae. An isolate of the *Vannella* genus, *Vannella* sp. AP1411, which was collected from oyster farms, is ubiquitous, and belongs to one clade of *Vannella* that could be found associated with Vibrionaceae. LGP32 was shown to be resistant to grazing by *Vannella* sp. AP1411 and this phenotype depends on some previously identified virulence factors: secreted metalloprotease Vsm and copper efflux p-ATPase CopA, which act at different steps during amoeba-vibrio interactions, whereas some other virulence factors were not involved. Altogether, our work indicate that some virulence factors can be involved in multi-host interactions of *V. tasmaniensis* ranging from protozoans to metazoans, potentially favoring their opportunistic behavior.

**Keywords :** Facultative intracellular pathogen, vibrios, marine amoebae, virulence, oyster

## INTRODUCTION

Vibrios are  $\gamma$ -proteobacteria living in aquatic environments ranging from saline to freshwater.

They are ubiquitous in marine coastal environments and have evolved the capacity to colonize a

broad range of hosts from protozoans to metazoans (Austin, 2010). Vibrios belong to the microbiota of healthy oysters but some species that behave as opportunistic pathogens can thrive in host tissues and cause disease (Lemire *et al.*, 2014; Bruto *et al.*, 2017; Rubio *et al.*, 2019). This occurs mostly as a result of environmental changes such as shifts in water temperature, exposure to high animal densities and stressful farming practices or upon immune suppression of host defenses by other microorganisms such as the OsHV-1 virus (de Lorgeril *et al.*, 2018). Currently, vibrioses are recognized as a major factor limiting the development of aquaculture (Le Roux *et al.*, 2015). In addition, vibrios can cause severe disease outbreaks in human populations, the best-known example being cholera. As they are multi-host pathogens, the ecology of vibrios depends on a series of biotic interactions in the environment that influence their ecology, their evolution and their transmission to animal and human hosts (Constantin de Magny *et al.*, 2008; Takemura *et al.*, 2014).

Bacteriovorous protists, such as free-living amoebae (FLA), are bacterial predators present in natural environments. They can be found in most aquatic environments and their predation through grazing activity shapes bacterial communities (Pernthaler *et al.*, 2005). However, it has been shown that some bacteria are able to resist grazing, persist and even grow within amoebae (Greub and Raoult, 2004). The best-known and described example is the facultative intracellular pathogen *Legionella pneumophila* (Rowbotham, 1980). Predation activity by protists shares cellular and molecular processes with metazoan immune cells antibacterial responses, in particular phagocytosis (Boulais *et al.*, 2010). Hence, amoebae have been hypothesized to act as

“evolutionary training ground” for grazing-resistant bacteria and in particular for intracellular bacterial pathogens, by favoring the selection of virulence factors (Molmeret *et al.*, 2005).

A number of studies have demonstrated that some *Vibrio* species adopt intracellular stages (Rosenberg and Falkovitz, 2004; Ma *et al.*, 2009; Vidal-Dupiol *et al.*, 2011; Ritchie *et al.*, 2012; de Souza Santos and Orth, 2014). In particular, intracellular stages are required for strains of *Vibrio tasmaniensis* to express virulence in oysters (Duperthuy *et al.*, 2011; Rubio *et al.*, 2019). Virulent strains of this species have been isolated from moribund oysters during juvenile mortality events on the French Atlantic coast (Gay *et al.*, 2004; Lemire *et al.*, 2014; Bruto *et al.*, 2017). This species was also found in oyster flesh during juvenile mortalities, as well as in the water column and the sediment of oyster farms in the Mediterranean Thau Lagoon, France (de Lorgeril *et al.*, 2018; Lopez-Joven *et al.*, 2018). Some of the *V. tasmaniensis* LGP32 virulence factors have been described in oysters. They include secreted proteases, such as Vsm (Binesse *et al.*, 2008) or Vsp (Vanhove *et al.*, 2014), antioxidants such as the superoxide dismutase SodA, efflux pumps to detoxify heavy metals such as the copper p-ATPase CopA (Vanhove *et al.*, 2016), and a Type VI secretion system that mediates the intracellular secretion of toxic effectors (Rubio *et al.*, 2019).

Here we wondered whether some virulence factors of the *Vibrio tasmaniensis* LGP32 playing a key role in its interaction with oysters could also be beneficial during other biotic interactions in the environment. With this objective, we first investigated the natural associations between vibrios and marine amoebae. Second, we did comparative cellular biology to study the

Accepted Article

interaction between LGP32 and an environmental free-living marine amoeba (*Vannella* sp strain AP1411) ubiquitous in oyster farms, which was found to establish interactions with vibrios. By performing a combination of flow cytometry, confocal and electron microscopy analyses, we show that LGP32 resists the predation by *Vannella* sp. AP1411. Using mutant strains inactivated for previously characterized virulence factors in oysters, we found that the copper p-ATPase efflux pump CopA and the secreted metalloprotease Vsm are both involved. CopA appears to play a role in the intracellular survival of LGP32 whereas Vsm appears to play a role in the inhibition of amoebae motility during grazing. Altogether, our results indicate that some of the previously identified virulence factors involved in *V. tasmaniensis* LGP32 interaction with oysters, are also involved in resistance to predation by marine amoeba *Vannella* sp. AP1411, thus potentially favoring opportunism.

## RESULTS

### **Marine amoebae belonging to the *Vannellidae* family and present in oyster farming area can interact with vibrios**

Although amoebae represent an important category of benthic bacterial grazers, their diversity remains poorly explored in marine environments, particularly in Mediterranean lagoons, which are exploited for oyster farming. Therefore, we first attempt to identify and isolate free-living amoebae in marine environments hosting oyster farms. Amoebae were isolated from the Thau lagoon at the vicinity of an oyster farming area. Samples were collected from the water column,

oyster gills and sediment at all seasons between the year 2014 and 2015, and at least two cultures of amoebae per fraction and per season were analyzed. Cloning-sequencing of the v7 hypervariable region of 18S rDNA gene was used to identify the isolated amoebae. A total of 110 sequences were analyzed. From 24 different cultures, 103 amoebae sequences showed more than 90% sequence homology to the *Vannella* genus, which belongs to the *Vannellidae* family. The 103 sequences were classified into 11 different Operational Taxonomic Units (OTUs) (Figure 1 and Table S1). In parallel, barcoding of the different amoebae cultures confirmed that only species belonging to the *Vannella* genus were present in the samples (Figure S1). One OTU of *Vannella sp.* was ubiquitous, as it was found in the 3 fractions during four seasons, and referred to the isolate AP1411 (arrow in Figure 1). In an attempt to determine whether marine amoebae could be found associated with *Vibrionaceae* in the environment, a dedicated sampling was performed in 2017 and amoebae were clonally isolated from oyster gills and from water column in another site (Banyuls-sur-mer, France) as an out-group. Then 16s barcoding analyses were performed to characterize the bacteria that remained associated to these *Vannella* clones four weeks after culturing. From all the clones analyzed, four were found associated with *Vibrionaceae* OTUs (Figure 2A). In order to validate this association, the presence of *Vibrionaceae* associated with the four *Vannella* clones was then assessed by FISH using a 16s probe specific for vibrios (Huggett *et al.*, 2008) (Figure 2B). Among the four clones that were analyzed, two showed an association with *Vibrionaceae* (0517-TOEc-2 and 0517-TOEc-4). The

vibrios could be found inside amoebae as well as outside. These results suggest that amoebae of the *Vannella* genus can establish interactions with vibrios in the environment.

**The pathogen *V. tasmaniensis* LGP32 resists grazing by *Vannella* sp. AP1411.**

To study further the interaction between vibrios and *Vannella* amoebae, we chose the *Vannella* sp. AP1411 clone, which belongs to the most ubiquitous OTU of marine amoebae and is not already associated with other vibrios, as shown by FISH (Figure 2B). The trophozoite, planktonic and cystic forms of *Vannella* sp. AP1411 were typical of the *Vannella* genera and very similar to *Vannella plurinucleolus* (Smirnov *et al.*, 2007), which was also the closest 18S matching sequence for this isolate (98% Blastn score). Transmission electron microscopy of trophozoites revealed a typical intracellular organization of *Vannella* with a well-defined, thick plasma membrane, one nucleus (sometimes two or three), numerous mitochondria and a rich vacuolar system with large digestive vacuoles containing remnants of degraded bacteria, and some phagosomes containing individual bacteria (Figure S2).

To study LGP32 - *Vannella* sp. AP1411 interactions, grazing assays were performed on solid media (non-nutrient seawater agar). *Vibrio* strains (pathogenic or non-pathogenic to oysters) that constitutively express the Green Fluorescent Protein (GFP) were used to seed the plates and monitor the abundance of living vibrios by fluorescence quantification. In parallel, *Vannella* sp. AP1411 growth was monitored by counting the cell density at the surface of the grazing lawns. When the amoeba was cultured in the presence of the non-virulent *Vibrio tasmaniensis* LMG20012<sup>T</sup>, a rapid decay of GFP-expressing bacteria was observed 4 days after the beginning

of the experiment; whereas in the case of the pathogenic *V. tasmaniensis* LGP32, the amount of GFP-expressing bacteria remained stable for up to 7 days (Figure 3A). Amoeba growth (cells count) on the vibrio lawns counter mirrored the kinetics of the decay of GFP-expressing bacteria. Indeed amoeba density increased much faster in the presence of LMG20012<sup>T</sup> than in the presence of LGP32 (Figure 3B). These data indicate that *Vannella sp.* AP1411 do not prey efficiently on the virulent LGP32, whereas they can feed on the non-virulent LMG20012<sup>T</sup>.

To determine whether the amoebal growth defect observed in the presence of LGP32 was due to inefficient phagocytosis and/or resistance to intracellular degradation by LGP32, we next quantified the capacity of *Vannella sp.* AP1411 to phagocytose LGP32 and LMG20012<sup>T</sup>, using flow cytometry (Figure S3). The amount of amoebae carrying GFP-vibrios was quantified three and six days after grazing, at three days after the beginning of the interaction, 100% of amoebae were found to carry GFP-LMG20012<sup>T</sup> or GFP-LGP32 (Figure 3C). However at day 6, the percentage of amoebae carrying GFP-LMG20012<sup>T</sup> decreased down to 20% whereas it remained stable with GFP-LGP32 (Figure 3C) suggesting that LGP32 could resist intracellular degradation. Confocal microscopy investigations confirmed that the amoebae were able to phagocytose both vibrios (Figure 3D). These observations also confirmed that six days after the onset of the grazing assay, 80% of amoebae were devoid of any GFP-LMG20012<sup>T</sup> as quantified by flow cytometry, with only 20% of amoebae that were still carrying intracellular GFP-LMG20012<sup>T</sup>, whereas 100% of the amoebae were still carrying numerous GFP-LGP32 (Figure 3D). This suggests that GFP-LGP32 is able to resist intracellular degradation after engulfment in

contrary to the non-virulent control strain GFP-LMG20012<sup>T</sup> that seemed to be digested by the amoeba.

To investigate further the intracellular fate of the vibrios inside amoebae, transmission electron microscopy was performed three days after the onset of grazing (Figure 3E). Most LMG20012<sup>T</sup> cells were found clustered inside vacuoles and showed altered morphologies whereas LGP32 remained intact inside amoebae and were mostly found in individualized phagosomes confirming that LGP32 is resistant to intracellular degradation compared to LMG20012<sup>T</sup>. Altogether these results show that LGP32 is resistant to grazing by *Vannella sp.* AP1411 as opposed to the non-virulent LMG20012<sup>T</sup>, and that LGP32 is able to resist to intracellular degradation by the amoeba.

#### **The Vsm and CopA virulence factors participate in LGP32 resistance to grazing by marine amoeba.**

To investigate further the molecular basis of LGP32 resistance to predation by *Vannella sp.* AP1411 we studied the phenotype of mutant strains of LGP32 deleted for virulence factors previously identified to be involved in its interaction with oysters. On the one hand we examined stress resistance systems, i.e. the antioxidant SodA, and the copper resistance systems CopA and CusAB, involved in intraphagosomal resistance to antimicrobial activities of oyster immune cells (Vanhove *et al.*, 2016) and on the other hand, we examined secreted virulence factors, i.e. the metalloprotease Vsm and the serine protease Vsp that both play a role in oyster pathogenesis (Binesse *et al.*, 2008; Vanhove *et al.*, 2014); as well as the metalloprotease InhA (the ortholog of PrtV) that has been involved in multiple biotic interactions of vibrios (Vaitkevicius *et al.*, 2006),

and finally the T6SS1 secretion system required for cytotoxicity towards oyster immune cells (Rubio *et al.*, 2019). Among the height mutant strains of LGP32 tested here, only the  $\Delta copA$  and  $\Delta vsm$  deletion mutants were significantly less resistant to grazing by *Vannella sp.* AP1411 than the wild type strain, as shown by a faster clearance of GFP-vibrios on grazing lawns (Figure 4A and 4D; and Figure S4). Accordingly, on lawns of both  $\Delta copA$  and  $\Delta vsm$ , rapid bacterial clearance correlated with faster amoebal growth as shown by the higher amoeba density than on wild-type LGP32 lawns (Figures 4B and 4E). These results show that both the CopA resistance factor and the Vsm virulence factor play a role in the resistance of LGP32 to grazing by *Vannella sp.* AP1411 whereas the other tested virulence factors did not seem to be involved.

**The p-ATPase CopA confers resistance to intracellular killing and the extracellular protease Vsm inhibits amoebae motility during grazing.**

To investigate further the role of LGP32 virulence factors in resistance to amoeba grazing, we analyzed different steps of the grazing process, from phagocytosis to intracellular degradation. Flow cytometry analyses showed that the percentage of phagocytosis (Figures 4C and 4F) and the phagocytosis index (Figure S5) were not significantly different between the mutant strains  $\Delta copA$  and  $\Delta vsm$  and the wild-type LGP32, which suggested that the efficiency of the phagocytosis process was not disturbed by CopA or Vsm. Confocal and TEM microscopy, confirmed that both mutant strains were readily phagocytosed by the amoeba, and revealed that  $\Delta copA$  LGP32 vibrios were found mostly clustered in vacuoles and the bacteria appeared to have an altered morphology after three days of grazing. These observations suggest that  $\Delta copA$

LGP32 was more sensitive to intracellular degradation, indicating a role for *copA* in resistance of LGP32 to intracellular degradation. In contrast,  $\Delta vsm$  LGP32 appeared intact inside vacuoles and individual phagosomes in a similar manner than the wild type strain of LGP32, suggesting that Vsm is not involved in LGP32 resistance to intracellular degradation (Figure 5A and 5B).

In order to investigate further the role of Vsm in LGP32 resistance against grazing by *Vannella* sp. AP1411, amoeba grazing was recorded by time-lapse microscopy. Amoebae grazing behavior on vibrio lawns was analyzed using  $\Delta vsm$  LGP32 and  $\Delta copA$  LGP32 lawns and compared to wild type strain. Cell tracking was performed to measure amoeba migratory speed on the different lawns of vibrios (Figure 6) and the most striking difference that could be observed was that amoebae migratory speed was two fold faster ( $p < 0.001$ ) on lawns of  $\Delta vsm$  LGP32 compared to wild-type or  $\Delta copA$  LGP32 lawns (Supplementary Movies 1, 2 and 3), with mean velocities of 8.5, 3.9 and 3.8 microns per minute for  $\Delta vsm$  LGP32, wild-type LGP32 and  $\Delta copA$  LGP32, respectively (Figure 6D). These results indicate that the metalloprotease Vsm acts extracellularly by inhibiting amoeba motility, which likely disturbs grazing efficiency of amoeba, whereas CopA participate to the intracellular survival capacity of LGP32.

## DISCUSSION

In this study, we uncovered that the oyster pathogen *V. tasmaniensis* LGP32 is resistant to grazing and intracellular degradation by free-living marine amoebae of the *Vannella* genus. Here we used *Vannella* sp. AP1411 as a representative amoeba isolate, which belongs to genus of

amoeba that are ubiquitous in the oyster farming environment and that interact with vibrios in the environment. Additionally, we uncovered some of the molecular determinant that are involved in the resistance of LGP32 to *Vannella* sp. AP1411, the copper efflux p-ATPase CopA and the secreted metalloprotease Vsm. These results indicate that at least two different mechanisms that are contributing to *V. tasmaniensis* LGP32 pathogenicity in oysters are also involved in resistance to amoeba grazing.

Amoebae of the *Vannellidae* family were found to be ubiquitous in the Thau lagoon, a Mediterranean coastal lagoon used for oyster farming (20% of its total surface). Indeed, most of the isolated amoebae branched into the *Vannella* genus, independently of seasons. Together with the present study, the current literature suggests *Vannellidae* amoebae are frequent in aqueous environments, mostly in saltwater but are also frequently observed in freshwater (Smirnov *et al.*, 2007). By performing 16s barcoding sequencing and FISH analysis we were able to find that some *Vannella* isolates from oyster farms can be isolated together and remained associated for weeks *in vitro* suggesting that these amoebae can interact with vibrios in this environment. Although the ecological dynamics of these *Vannella*-Vibrio associations deserves to be investigated in greater details, our data suggest that the isolated *Vannella* in the present study are an ecologically relevant host to investigate the biotic interactions between *Vibrio* species and marine amoebae.

By studying the interactions between *Vannella* sp. AP1411 and the vibrio LGP32, we found that *V. tasmaniensis* LGP32 is able to resist predation by the marine amoeba. Vibrios belonging to

Accepted Article

the *tasmaniensis* species have been shown to resist hemocyte phagocytosis and to behave as facultative intracellular pathogen, a behavior that was shown to be central for its virulence in oysters (Duperthuy *et al.*, 2011; Vanhove *et al.*, 2016 ; Rubio *et al.*, 2019). The slow growth of amoebae on LGP32 was due, to some extent, to an inefficient intracellular killing of LGP32. Indeed, after phagocytosis most intracellular LGP32 were isolated in individual phagosomes inside the amoebae without any morphological alteration whereas the non-virulent control *V. tasmaniensis* LMG20012<sup>T</sup> were progressively clustered in vacuoles and degraded. This strongly suggests that LGP32 interferes with phagosome maturation and intracellular trafficking and this is reminiscent of its capacity to inhibit phagosome acidification in oyster phagocytes (Duperthuy *et al.*, 2011).

The copper efflux p-ATPase CopA was found here to play a major role in the capacity of LGP32 to resist intracellular degradation within the amoeba. As CopA is involved in the resistance of LGP32 to high concentration of copper within oyster hemocytes (Vanhove *et al.*, 2016), our results indicate that LGP32 resistance to grazing by the amoeba depends on its capacity to tolerate high concentrations of extracellular copper inside phagosomes and thus strongly suggest that *Vannella sp.* AP1411 of intra-phagosomal heavy metal poisoning in a similar manner than other professional phagocytes. Intra-phagosomal copper accumulation appear to be an ancestral highly conserved antimicrobial mechanism from free-living amoebae, social amoebae (Hao *et al.*, 2016), invertebrates (Vanhove *et al.*, 2016) to vertebrate professional phagocytes (Soldati and Neyrolles, 2012). As this antimicrobial mechanism is used by a variety of phagocytes we can

Accepted Article

hypothesize that, like other bacterial species resisting intracellular killing (Espariz *et al.*, 2007; Rowland and Niederweis, 2012), for some vibrios copper resistance confer to them a fitness advantage in diverse of host-vibrio interactions (Brooks *et al.*, 2014; Vanhove *et al.*, 2016).

Another determinant that was found here to be important for LGP32 resistance to grazing by *Vannella sp.* AP1411 was the secreted metalloprotease Vsm, a major secreted protease that plays a role as a virulence factor in oysters (Binesse *et al.*, 2008). In the present study, the Vsm deficient LGP32 strain showed an increased sensitivity to grazing by the amoeba, without losing its ability to resist intracellular degradation. This suggested that Vsm acts extracellularly rather than intracellularly, during LGP32 interaction with amoeba, which was confirmed by studying amoeba grazing behavior by time-lapse microscopy revealing an important increase of grazing migratory speed of amoebae on  $\Delta_{vsm}$  LGP32 bacteria lawns compared to lawns with the wild type strain. The involvement of Vsm metalloprotease in LGP32 resistance to grazing are reminiscent of the role played by other proteases secreted by vibrios, during their interaction with eukaryotic cells. In particular, the involvement of the secreted protease PrtV, that participates in the resistance of *V. cholerae* to the grazing by the bacteriovorous ciliate *Tetrahymena pyriformis* (Vaitkevicius *et al.*, 2006), and studies on the pathogenesis of LGP32 and *Vibrio aestuarianus*, another vibrio pathogenic for oysters, that showed that LGP32 Vsm can damage eucaryotic cells (Binesse *et al.*, 2008), and that its homolog Vam from *V. aestuarianus*, is a virulence factor inhibiting hemocyte phagocytosis (Labreuche *et al.*, 2010). Interestingly, recent work studying the interaction between clinical isolates of *V. cholerae* and the model

amoebae *Acanthamoeba castellanii* (Neff strain) reported that minor virulence factors like secreted enzyme, including the HapA metalloprotease and the hemolysin can be involved in fine tuning the fitness of *V. cholerae* during its interaction with grazers in the aquatic environment (Van der Henst *et al.*, 2018). Then the secretion of proteases by vibrios appear to be an important ancestral mechanism that may has been acquired to inhibit predation by grazers in aquatic environments but can also participate to their pathogenicity in animal hosts.

Hence copper resistance and secretion of proteases appear important virulence mechanisms for different vibrios and involved in interactions with diverse hosts, the presence of these virulence traits in vibrios against diverse hosts may represent evidences for the coincidental selection hypothesis (Adiba *et al.*, 2010).

As bacteriovorous protists exert a selective pressure on bacterial communities, different bacteria species are thought to have developed a diversity of survival strategies against predation through either pre-ingestional or post-ingestional adaptations (Matz and Kjelleberg, 2005). Pre-ingestional adaptations including high motility, filamentation, surface masking or toxin release are commonly found for extracellular pathogens; whereas post-ingestional adaptations including digestional resistance through vacuolar trafficking inhibition or vacuolar escaping, and toxin release are commonly found for intracellular pathogens and tend to favor bacterial growth (Matz and Kjelleberg, 2005). By establishing a new ecologically relevant cellular model to study the Vibrio-phagocytes interactions in marine environment, we found that, in the case of the facultative intracellular pathogen LGP32, a combination of pre-ingestional and post-ingestional

Accepted Article

adaptations could be involved. The involvement of the secreted Vsm represents a pre-ingestional adaptation, whereas the involvement of the CopA p-ATPase represents a post-ingestional adaptation. Thus, the predator-prey interactions between bacterial communities and bacterivorous protists may select for common of virulence factors allowing bacteria to resist to a diversity of phagocytic cells from amoeba to immune cells in animals, hence favoring the emergence of opportunistic pathogens able to colonize a diversity of hosts, like many vibrios do.

However not all virulence factors studied here were found to participate to LGP32 resistance to grazing by *Vannella sp.* AP1411. For instance, *V. tasmaniensis* T6SS1, Vsp, CusAB and SOD, which were found to play a role in oysters (Vanhove *et al.*, 2014, 2016; Rubio *et al.*, 2019), do not seem to play a role during interactions with *Vannella sp.* AP1411. These results strengthen the hypothesis that although some virulence factors could be involved in diverse biotic interactions, as observed here for CopA and Vsm, hence favoring opportunism, some other virulence factors could be more specifically involved in more specialized interactions. As for example, the metalloprotease InhA, the ortholog of PrtV in LGP32, does not appear to play a role during LGP32-Oyster interactions or LGP32-*Vannella sp.* AP1411 interaction, although it was shown to participate in the resistance of *V. cholerae* to the grazing by the bacterivorous ciliate *Tetrahymena pyriformis* (Vaitkevicius *et al.*, 2006). In a similar manner, T6SS of *Vibrio cholerae* which was shown to be involved during interaction with the social amoeba *D. discoideum* (Pukatzki *et al.*, 2006), did not appear to be involved in LGP32 resistance to *Vannella sp.* AP1411 although it was shown recently to be a major determinant of LGP32

Accepted Article

cytotoxic activity towards oyster hemocytes (Rubio *et al.*, 2019). Such host-specialization of for some vibrios virulence factors was also suggested recently for *V. cholerae*, as some of the virulence factors involved in the resistance to grazing by *A. castellanii* are considered to have a minor role in pathogenesis in human (Van der Henst *et al.*, 2018). These subtleties in the involvement of different virulence factors during different biotic interactions are reminiscent of the subtleties that are being uncovered for the intracellular pathogen *L. pneumophila*. These well studied intracellular pathogen harbors a broad range of virulence factors and can interact with a diversity of hosts, but this does not imply that all virulence factors have a similar importance in each interactions with these diverse hosts (Boamah *et al.*, 2017; Ghosh and O'Connor, 2017). Vibrios being mostly opportunistic pathogens, they probably have acquired, on the one hand, some virulence factors that are commonly involved in diverse host interactions that could be core virulence factors, and on the other hand, some virulence factors that are involved in more restricted host interactions that could be considered specialized virulence factors. Studying diverse host-vibrios interactions will help to decipher the diversity of their virulence factors and provide insights about opportunist pathogens co-evolutionary trajectories that are more complex than the host-parasite co-evolutionary arm race observed for host specialized pathogens (Brown *et al.*, 2012).

In conclusion, by developing a new ecologically relevant cellular model of interaction, we showed here that the oyster pathogen *V. tasmaniensis* LGP32 can resist phagocytosis by environmental grazers *Vannella spp.*, which are ubiquitous in oyster farming area and that appear

Accepted Article

to interact with vibrios in the oyster farming environment. By performing comparative cellular biology, we also demonstrated that some of LGP32 virulence factors can be involved in different types of interactions with multiple hosts from protozoans, such as environmental marine amoeba, to metazoans, e.g. oysters. This suggests that some virulence factors such as CopA and Vsm could play a central role in the interaction of *V. tasmaniensis* with diverse hosts whereas some other virulence factors appear to be involved in more specific host-vibrio interactions. Hence the present study implies that by studying more diverse host-vibrio interactions will help to gain further insights about the ecology of vibrios and to uncover the relative roles of biotic interactions in pathogen emergence.

## MATERIALS AND METHODS

### Bacterial strains and growth conditions

*E. coli* strain SBS363 was grown in Luria-Bertani (LB) or LB-agar (LBA) at 37°C. Marine amoebae *Vannella sp.* AP1411 and isolates 0517-TOEc-2, 0517-TOEc-3, 0517-TOEc-4, 0517-BWEc-1, 0517-BWEc-2, 0517-BWEc-3, 0517-BWEc-4 (isolated in this study) were grown in 70% sterile seawater (SSW) with *E. coli* wild-type strain SBS363 at 18°C, for 3 days prior experiments. Vibrios strains used in this study are *V. tasmaniensis* LMG20012<sup>T</sup>, *V. tasmaniensis* LGP32 and previously described LGP32 deletion mutants  $\Delta copA$ ,  $\Delta cusAB$ ;  $\Delta sodA$ ;  $\Delta vsm$ ,  $\Delta vsp$ ,  $\Delta inhA$  (Binesse *et al.*, 2008; Vanhove *et al.*, 2014, 2016) and insertion mutants  $\Delta vipA1$ ,  $\Delta vipA2$  (Rubio *et al.*, 2019). Vibrios strains carrying the pMRB-GFP or pD3-GFP-mut3 plasmids were

grown in LB + NaCl 0.5M supplemented with chloramphenicol (10 $\mu$ g mL<sup>-1</sup>) or kanamycin (20 $\mu$ g mL<sup>-1</sup>) at 20°C, for 24 hours prior experiments.

### **Isolation of environmental amoebae from the Thau Lagoon and Banuyls-sur-mer.**

Between 2014 and 2015, seawater, oysters and sediment from the Thau lagoon (South France) were sampled at least two times per season. Seawater was collected next to oyster tables at the Bouzigues station Ifremer-REPHY (GPS: N 43°26.058' E 03°39'.878') and filtered, on the boat, with a 180  $\mu$ m pore size nylon filter. In the lab, the water was re-filtered using a 8.0  $\mu$ m pore size MF-Millipore membrane. The 8.0  $\mu$ m pore size membrane was then cut in four pieces and each quarter was put upside down on a lawn of *E. coli* SBS363 seeded on SSW-agar. Oyster gills were cut in 4 pieces of roughly 0.5 cm<sup>2</sup> and each quarter was put upside down on a lawn of *E. coli* SBS363 seeded on non-nutrient SSW-agar. Sediment was collected under the oyster tables (depth of 9 meters) by core sampling. One gram of sediment was then deposited in the center of a lawn of *E. coli* SBS363 seeded on SSW-agar, in triplicate. After 1 to 3 weeks, depending of the sampling season, migrating amoebae were observed at the periphery of the plates. An agar square of 0.5x0.5 cm containing a single amoeba was then cut and deposited upside down on a fresh lawn of *E. coli* SBS363 seeded on SSW-agar. This was repeated several times to isolate and replicate amoebae. DNA was extracted using the High Pure PCR Template Preparation Kit (Roche), according to the manufacturers' protocol. In May 2017, sampling was performed from oyster gills from Thau lagoon at Bouzigues station Ifremer-REPHY (GPS: N 43°26.058' E 03°39'.878') and seawater from Banuyls-sur-mer at SOLA station (GPS: 42°29'300 N –

03°08'700 E). Samples were treated with the same protocol than previously and put on a lawn of *E. coli* SBS363 seeded on SSW-agar. After 2 weeks, amoebae were flushed by SSW and clonally isolated by dilutions within 96 well plates with statistically less than one amoeba per well. Amoebae clones were amplified by culturing in 6 well plates and DNA was extracted using the NucleoSpin Tissue kit (Macherey-Nagel), according to the manufacturer's protocol.

### **Identification of isolated marine amoebae from Thau Lagoon and Banuyls-sur-Mer and their associated bacteria.**

In order to infer the phylogenetic assignment of environmental amoebae, the v7 region of the 18S rDNA gene was sequenced using specific primers. The primers were designed using SILVA database dedicated tool (<http://www.arb-silva.de/search/testprime>). The selected primers, Amo\_AP\_1154\_F: 5' GAGRAAATTAGAGTGTTYAAAG 3' and Amo\_AP\_1470\_R: 5'TTATRGTTAAGACTA CGACGG 3', were used to amplified the hypervariable region V7 of 18S rDNA gene at an annealing temperature of 54°C. PCR amplicons were cloned using the TOPO TA Cloning Kit (Invitrogen), according to the manufacturers' protocol. Nucleotide sequences were determined by Sanger sequencing (GenSeq platform, Labex CEMEB, Montpellier, France). Sequence homologies were searched using BLASTn available at the NCBI web site (National Centre for Biotechnology Information (Wheeler *et al.*, 2007) (Table S1). For phylogenetic analyses, sequences were aligned in MOTHUR (Schloss, 2009) and imported into ARB software (Ludwig *et al.*, 2004) loaded with the silva database (<http://www.arb-silva.de>). A base frequency filter was applied to exclude highly variable positions before adding sequences to

the maximum parsimony backbone tree using the parsimony quick add marked tool implemented in ARB, thereby maintaining the overall tree topology provided by default. In order to verify that the Sanger sequencing strategy used here allowed us to identify the entire diversity of amoebae that could be isolated through culturing techniques. The total DNA from our different amoebae cultures was analyzed by performing barcoding on the v7 region of the 18S rDNA gene using generic primers that allowed amplifying more than 80% of all the eukaryote 18s rDNA sequences available in SILVA database (F-v7-1173: CCT GCG GCT TAA TTT GAC and R-v7-1438: CAT CAC AGA CCT GTT ATT GC). Sequencing was performed by Genome Quebec facility, Montreal, Canada.

Bacteria associated with Amoeba clonally isolated in May 2017 sampling were investigated using barcoding sequencing on the V3/V4 region of the 16S rDNA gene with Illumina generic primers according to the manufacturer's protocol (16Sv3v4-ILLAFWD: 5'-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG YRC CTA CGG GNG GCW GCAG and 16Sv3v4-ILLAREV: 5'-GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GYR GAC TAC HVG GGT ATC TAA TCC). Sequencing was performed by GenSeq platform, Labex CEMEB, Montpellier, France. OTUs taxonomic affiliation was performed using FROGS tool from galaxy software (Escudié *et al.*, 2018). To determine the taxonomic affiliation of clones isolated during the second sampling campaign, the sequence of the V4 hypervariable region of the 18S rDNA gene were analyzed using TAREuk454FWD1: 5'-CCAGCA(G/C)C(CT)GCGG-TAATTCC-3' and TAREukREV3: 5'-ACTTTCGTTCTTGAT(CT)(A/G)A-3' primers (Stoeck *et*

*al.*, 2010). Nucleotide sequences were determined by Sanger sequencing (GenSeq platform, Labex CEMEB, Montpellier, France) and aligned with MUSCLE using MEGA7 software (maximum-likelihood by bootstrap method, 1000 replications).

### **Grazing assay**

To prepare the co-culture of vibrios and amoebae, 1 mL of vibrio overnight culture ( $3.10^9$  bacteria.  $\text{mL}^{-1}$ ) was mixed with 100  $\mu\text{L}$  of three days old *Vannella sp.* AP1411 culture ( $5.10^5$  cells.  $\text{mL}^{-1}$ ) or with 100  $\mu\text{L}$  of 70% SSW for control condition. A volume of 50  $\mu\text{L}$  per well of the mixed culture were seeded on top of 500  $\mu\text{L}$  of 1% SSW-agar, in 24-well plates with transparent flat bottoms. Amoebae and bacteria lawns were carefully homogenized in the wells and let dry 4 hours at room temperature in a laminar flow cabinet and then incubated at 18 °C in a humidified atmosphere. GFP fluorescence intensity was measured every day over 7 days using a TECAN plate reader ( $\lambda_{\text{ex}}$  480 nm/ $\lambda_{\text{em}}$  520 nm). To estimate the effect of the amoebae grazing activity on the abundance of living vibrios expressing GFP, the fluorescence intensity of the wells containing amoebae was compared to the fluorescence of vibrios lawn without amoebae, and express as a ratio. Each condition was performed in technical triplicates and the results shown are the average of three independent experiments. Error bars represent the standard error of the mean ( $\pm$  SEM). Statistical analysis was performed using RM-ANOVA over the independent experiments.

### **Monitoring amoeba growth and association with GFP-vibrios.**

Accepted Article

To estimate the proliferation of *Vannella sp.* AP1411 at days 1, 3 and 6 in grazing assays, amoebae were directly imaged by phase contrast microscopy and enumerated in triplicate for each condition. Cells were flushed from the SSW-agar surface with 1 mL of 70% SSW and fixed for 30 minutes with 2% paraformaldehyde at room temperature. Pellets were washed and suspended in 500  $\mu$ L of PBS for flow cytometry analysis (BD FACS Canto). Data analyses were performed using Flowing Software by gating on amoeba cells using SSC/FSC parameters then quantifying the percentage of GFP fluorescence associated amoebae (as shown in Figure S2). In order to estimate the phagocytosis index, which corresponds to the average number of phagocytosed bacteria per cell, the mean intensity of GFP fluorescence per cell was normalized to the mean fluorescence per GFP + bacteria (as shown in Figure S3). The experiments were performed with three technical replicates per condition in each experiment, and depicted results are representative of at least two independent experiments. Error bars represent the standard error of the mean ( $\pm$  SEM). Statistical analysis was performed using two ways ANOVA with Holm-Sidak's multiple comparisons test.

#### **Fluorescence in situ hybridization.**

In situ hybridization for *Vibrionaceae* was performed on *Vannella sp.* AP141, 0517-TOEc-2, 0517-TOEc-3 and 0517-TOEc-4. After at least two weeks of culturing in flask in 70% SSW complemented with *E. coli* SBS363, amoebae were washed twice with 3 mL of 70% SSW and resuspended in 70% SSW using a cell scraper and fixed with 2% formaldehyde for 25 min at room temperature. After fixation, formaldehyde was removed by centrifugation of the cells by

centrifugation 10 min at 6000g, pellets were washed and resuspended in 500  $\mu$ L of PBS. Cell suspensions were cytopun on glass slides for 10 min at 500g. Glass slides were then labeled with the *Vibrionaceae* specific probe VIB572a at 10 ng mL<sup>-1</sup> (5'-ACCACCTGCATGCGCTTT-3') (Huggett et al., 2008) using hybridization buffer (900 mM NaCl, 20 mM Tris-HCl, 30% Formamide, 0.01% SDS, dH<sub>2</sub>O) for 90 minutes at 46 °C in the dark. The samples were then washed for 25 min in washing buffer (0.112 mM NaCl, 20 mM Tris-HCl, 0.01% SDS, 5 mM EDTA, dH<sub>2</sub>O) at 48 °C in the dark. Then, cells were counterstained using FITC at 0.5 ng mL<sup>-1</sup> for 30 min (Sigma). Confocal fluorescence imaging was performed using a 63X oil objective and images were captured using a Leica TCS SPE confocal scanning laser microscope, at 1 airy to ensure the focal plan.

### **Microscopy**

The different amoebae shapes were observed on living amoebae using phase contrast (on inverted wide-field microscope, Zeiss). The shape of the floating forms was recorded by scraping in the cell suspension followed by immediate observation. The cystic form was observed after amoebae were starved for a week. Intracellular organization of amoebae was observed using transmission electron microscopy. Briefly, amoebae cultured in flasks were washed twice using 70% SSW, and fixed in 2.5% glutaraldehyde for 2 hours in the dark at room temperature, followed by overnight at 4°C. After centrifugation, the supernatant was replaced by 3% low melting-point agarose (Sigma) and the pellet was resuspended. Cells entrapped in 3% agar were

immersed in a solution of 2.5% glutaraldehyde in PHEM buffer (1X, pH 7.4) overnight at 4°C. Then they were rinsed in PHEM buffer and post-fixed in a 0.5% osmic acid for 2 h in the dark at room temperature. After two rinses, samples were dehydrated in a graded series of ethanol solutions (30-100%). Samples were embedded in EmBed 812 using an Automated Microwave Tissue Processor for Electronic Microscopy, Leica EM AMW. Ultrathin sections (70nm; Leica-Reichert Ultracut E) were collected at different levels of each block, then counterstained with uranyl acetate 1.5% in 70% Ethanol and lead citrate and observed using a Tecnai F20 transmission electron microscope at 120KV in the CoMET MRI facilities, INM, Montpellier France.

Confocal fluorescence microscopy was performed on amoebae suspensions after PFA fixation to observe the localization of intra-amoeba vibrios. Cell suspensions were cytopun on glass slides for 10 min at 800 g. Glass slides were then stained with 0.25  $\mu\text{g mL}^{-1}$  DAPI (Sigma) and 0.1  $\mu\text{g mL}^{-1}$  Blue Evans (Sigma). Confocal fluorescence imaging was performed using a 63X oil objective and images were captured using a Leica TCS SPE confocal scanning laser microscope, at 1 airy to ensure the focal plan.

### **Time-lapse imaging and cell tracking**

Co-cultures of vibrio and amoebae were prepared following the same protocol as for grazing assays. Time-lapse imaging started 24 hours after the beginning of the grazing assays. Acquisitions were obtained using an inverted wide-field epifluorescence microscope (Zeiss) and B&W coolsnap camera. Time-lapses were done during 30 minutes with 1 frame every 30

seconds. TrackMate Tools software (ImageJ) was used for manual tracking to determine velocity and migration distance by amoebae on different vibrios lawns (Supplementary Movie 1, 2 and 3). Migration distances were measured for 18 amoebae at the same time (i.e. 20 frames per amoebae trace) for each condition. Each track depicted corresponds to the migration of an amoeba during 10 minutes. The migration path was determined by the x and y positions of amoebae tracked at each frame. Depicted results are representative of at least two independent experiments. Velocities were measured for 16 amoebae tracks, only tracks made of at least 5 frames were considered. Each presented result corresponds to the average velocity of one amoeba per track. Depicted results are representative of at least two independent experiments. Error bars represent the standard deviation ( $\pm$  SD). Statistical analysis was performed using Kruskal-Wallis test with Dunn's multiple comparisons test.

## **ACKNOWLEDGMENTS**

We are grateful to Thibaut Groult, Audrey Caro, and Marc Leroy for precious help in sample collection and preparation, to Eric Abadie for field trip coordination and to Michel Cantou from the University of Montpellier for scuba diving allowing sediment collection. We thank Marie Buysse and Olivier Duron for technical advices. This work, through the use of the GENSEQ platform (<http://www.labex-cemeb.org/fr/genomique-environnementale-2>) from the labEx CeMEB. The authors also thank the Montpellier RIO imaging platform (<https://www.mri.cnrs.fr>). The present study was supported by the EU funded project VIVALDI (H2020 program, n°678589), by the Ec2co-CNRS funded Intervibrio and VibrAm projects, by

Labex CEMEB Amibadapt project, and by Ifremer, University of Montpellier and University of Perpignan via Domitia.

Authors declare there are no competing financial interest in relation to the work described here.

## REFERENCES

- Adiba, S., Nizak, C., van Baalen, M., Denamur, E., and Depaulis, F. (2010) From grazing resistance to pathogenesis: The coincidental evolution of virulence factors. *PLoS One* **5**: 1–10.
- Austin, B. (2010) Vibrios as causal agents of zoonoses. *Vet Microbiol* **140**: 310–7.
- Binesse, J., Delsert, C., Saulnier, D., Champomier-Vergès, M.-C., Zagorec, M., Munier-Lehmann, H., et al. (2008) Metalloprotease vsm is the major determinant of toxicity for extracellular products of *Vibrio splendidus*. *Appl Environ Microbiol* **74**: 7108–17.
- Boamah, D.K., Zhou, G., Ensminger, A.W., and O’Connor, T.J. (2017) From Many Hosts, One Accidental Pathogen: The Diverse Protozoan Hosts of *Legionella*. *Front Cell Infect Microbiol* **7**..
- Boulais, J., Trost, M., Landry, C.R., Dieckmann, R., Levy, E.D., Soldati, T., et al. (2010) Molecular characterization of the evolution of phagosomes. *Mol Syst Biol* **6**: 423.
- Brooks, J.F., Gyllborg, M.C., Cronin, D.C., Quillin, S.J., Mallama, C.A., Foxall, R., et al. (2014)

Global discovery of colonization determinants in the squid symbiont *Vibrio fischeri*. *Proc Natl Acad Sci* **111**: 17284–17289.

Brown, S.P., Cornforth, D.M., and Mideo, N. (2012) Evolution of virulence in opportunistic pathogens: generalism, plasticity, and control. *Trends Microbiol* **20**: 336–342.

Bruto, M., James, A., Petton, B., Labreuche, Y., Chenivresse, S., Alunno-Bruscia, M., et al. (2017) *Vibrio crassostreae*, a benign oyster colonizer turned into a pathogen after plasmid acquisition. *ISME J* **11**: 1043–1052.

Constantin de Magny, G., Murtugudde, R., Sapiano, M.R.P., Nizam, A., Brown, C.W., Busalacchi, A.J., et al. (2008) Environmental signatures associated with cholera epidemics. *Proc Natl Acad Sci U S A* **105**: 17676–81.

Duperthuy, M., Schmitt, P., Garzón, E., Caro, A., Rosa, R.D., Le Roux, F., et al. (2011) Use of OmpU porins for attachment and invasion of *Crassostrea gigas* immune cells by the oyster pathogen *Vibrio splendidus*. *Proc Natl Acad Sci U S A* **108**: 2993–8.

Escudié, F., Auer, L., Bernard, M., Mariadassou, M., Cauquil, L., Vidal, K., et al. (2018) FROGS: Find, Rapidly, OTUs with Galaxy Solution. *Bioinformatics* **34**: 1287–1294.

Espariz, M., Checa, S.K., Audero, M.E.P., Pontel, L.B., and Soncini, F.C. (2007) Dissecting the *Salmonella* response to copper. *Microbiology* **153**: 2989–97.

Gay, M., Renault, T., Pons, A., and Le Roux, F. (2004) Two *Vibrio splendidus* related strains

collaborate to kill *Crassostrea gigas*: taxonomy and host alterations. *Dis Aquat Organ* **62**: 65–74.

Ghosh, S. and O'Connor, T.J. (2017) Beyond Paralogs: The Multiple Layers of Redundancy in Bacterial Pathogenesis. *Front Cell Infect Microbiol* **7**: 1–14.

Hao, X., Luthje, F., Ro, Nn, R., German, N.A., Li, X., et al. (2016) A role for copper in protozoan grazing - two billion years selecting for bacterial copper resistance. *Mol Microbiol* **00**: 1–14.

Van der Henst, C., Vanhove, A.S., Drebes Dörr, N.C., Stutzmann, S., Stoudmann, C., Clerc, S., et al. (2018) Molecular insights into *Vibrio cholerae*'s intra-amoebal host-pathogen interactions. *Nat Commun* **9**: 3460.

Huggett, M., Crocetti, G., Kjelleberg, S., and Steinberg, P. (2008) Recruitment of the sea urchin *Heliocidaris erythrogramma* and the distribution and abundance of inducing bacteria in the field. *Aquat Microb Ecol* **53**: 161–171.

Labreuche, Y., Le Roux, F., Henry, J., Zatylny, C., Huvet, A., Lambert, C., et al. (2010) *Vibrio aestuarianus* zinc metalloprotease causes lethality in the Pacific oyster *Crassostrea gigas* and impairs the host cellular immune defenses. *Fish Shellfish Immunol* **29**: 753–758.

Lemire, A., Goudenège, D., Versigny, T., Petton, B., Calteau, A., Labreuche, Y., and Le Roux, F. (2014) Populations, not clones, are the unit of vibrio pathogenesis in naturally infected

oysters. *ISME J* 1–9.

Lopez-Joven, C., Rolland, J.L., Haffner, P., Caro, A., Roques, C., Carré, C., et al. (2018) Oyster Farming, Temperature, and Plankton Influence the Dynamics of Pathogenic Vibrios in the Thau Lagoon. *Front Microbiol* **9**: 2530.

de Lorgeril, J., Lucasson, A., Petton, B., Toulza, E., Montagnani, C., Clerissi, C., et al. (2018) Immune-suppression by OsHV-1 viral infection causes fatal bacteraemia in Pacific oysters. *Nat Commun* **9**: 4215.

Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, et al. (2004) ARB: a software environment for sequence data. *Nucleic Acids Res* **32**: 1363–71.

Ma, A.T., McAuley, S., Pukatzki, S., and Mekalanos, J.J. (2009) Translocation of a *Vibrio cholerae* type VI secretion effector requires bacterial endocytosis by host cells. *Cell Host Microbe* **5**: 234–43.

Matz, C. and Kjelleberg, S. (2005) Off the hook – how bacteria survive protozoan grazing. *Trends Microbiol* **13**: 302–307.

Molmeret, M., Horn, M., Wagner, M., Santic, M., and Abu Kwaik, Y. (2005) Amoebae as training grounds for intracellular bacterial pathogens. *Appl Environ Microbiol* **71**: 20–8.

Pukatzki, S., Ma, A.T., Sturtevant, D., Krastins, B., Sarracino, D., Nelson, W.C., et al. (2006) Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the

Dictyostelium host model system. *Proc Natl Acad Sci U S A* **103**: 1528–33.

Ritchie, J.M., Rui, H., Zhou, X., Iida, T., Kodoma, T., Ito, S., et al. (2012) Inflammation and Disintegration of Intestinal Villi in an Experimental Model for *Vibrio parahaemolyticus*-Induced Diarrhea. *PLoS Pathog* **8**: e1002593.

Rosenberg, E. and Falkovitz, L. (2004) The *Vibrio shiloi*/*Oculina patagonica* model system of coral bleaching. *Annu Rev Microbiol* **58**: 143–59.

Le Roux, F. Le, Wegner, K.M., Baker-Austin, C., Vezzulli, L., Osorio, C.R., Amaro, C., et al. (2015) The emergence of *Vibrio* pathogens in Europe: ecology, evolution, and pathogenesis (Paris, 11–12th March 2015). *Front Microbiol* **6**: 830.

Rowland, J.L. and Niederweis, M. (2012) Resistance mechanisms of *Mycobacterium tuberculosis* against phagosomal copper overload. *Tuberculosis (Edinb)* **92**: 202–10.

Rubio, T., Oyanedel, D., Labreuche, Y., Toulza, E., Luo, X., Bruto, M., et al. (2019) Species-specific mechanisms of cytotoxicity toward immune cells determine the successful outcome of *Vibrio* infections. *Proc Natl Acad Sci U S A* **116**: 14238–14247.

Schloss, P.D. (2009) A High-Throughput DNA Sequence Aligner for Microbial Ecology Studies. *PLoS One* **4**: e8230.

Smirnov, A. V., Nasonova, E.S., Chao, E., and Cavalier-Smith, T. (2007) Phylogeny, Evolution, and Taxonomy of Vannellid Amoebae. *Protist* **158**: 295–324.

Soldati, T. and Neyrolles, O. (2012) Mycobacteria and the intraphagosomal environment: take it with a pinch of salt(s)! *Traffic* **13**: 1042–52.

de Souza Santos, M. and Orth, K. (2014) Intracellular *Vibrio parahaemolyticus* Escapes the Vacuole and Establishes a Replicative Niche in the Cytosol of Epithelial Cells. *MBio* **5**: e01506-14.

Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M.D.M., Breiner, H.W., and Richards, T.A. (2010) Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Mol Ecol* **19**: 21–31.

Takemura, A.F., Chien, D.M., and Polz, M.F. (2014) Associations and dynamics of Vibrionaceae in the environment, from the genus to the population level. *Front Microbiol* **5**: 38.

Vaitkevicius, K., Lindmark, B., Ou, G., Song, T., Toma, C., Iwanaga, M., et al. (2006) A *Vibrio cholerae* protease needed for killing of *Caenorhabditis elegans* has a role in protection from natural predator grazing. *Proc Natl Acad Sci* **103**: 9280–9285.

Vanhove, A.S.A.S., Dupertuy, M., Charrière, G.M.G.M., Le Roux, F., Goudenège, D., Gourbal, B., et al. (2014) Outer membrane vesicles are vehicles for the delivery of *Vibrio tasmaniensis* virulence factors to oyster immune cells. *Environ Microbiol* **32**.

Vanhove, A.S.A.S., Rubio, T.P.T.P., Nguyen, A.N.A.N., Lemire, A., Roche, D., Nicod, J., et al. (2016) Copper homeostasis at the host vibrio interface: Lessons from intracellular vibrio

transcriptomics. *Environ Microbiol* **18**: 875–888.

Vidal-Dupiol, J., Ladriere, O., Meistertzheim, A.-L., Foure, L., Adjeroud, M., and Mitta, G. (2011) Physiological responses of the scleractinian coral *Pocillopora damicornis* to bacterial stress from *Vibrio coralliilyticus*. *J Exp Biol* **214**: 1533–1545.

Wheeler, D.L., Barrett, T., Benson, D.A., Bryant, S.H., Canese, K., Chetvernin, V., et al. (2007) Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* **36**: D13–D21.

## FIGURE LEGENDS

**Fig. 1. Most of the isolated free-living amoebae from the Thau lagoon belong to *Vannellidae* family.**

Phylogenetic 18S rRNA-based trees of the Amoebozoa constructed with ARB software (Ludwig et al., 2004; <http://www.arb-home.de>) loaded with the silva database (<http://www.arb-silva.de>). A base frequency filter was applied using the parsimony quick add marked tool implemented in ARB. Our isolates are written in red. Most of the isolated amoebae during the environmental survey from water column (blue squares) and/or the sediment (brown squares), and/or oyster gills (light green squares) during winter (light blue circles) and/or spring (green circles) and/or summer (yellow circles) and/or autumn (orange circles) belong to the *Vannellidae* family.

**Fig. 2. Amoebae belonging to the *Vannella* genus can be found associated with *Vibrionaceae* bacteria in oyster farms.**

*Vannella* were clonally isolated from oyster gills from Thau lagoon (*Vannella* sp. AP1411, 0517-TOEc-2, 0517-TOEc-3, and 0517-TOEc-4) or from seawater in Banyuls-sur-Mer (0517-BWEc-1, 0517-BWEc-2, 0517-BWEc-3, and 0517-BWEc-4), then 16s barcoding sequencing was performed to identify potentially associated *Vibrionaceae*. (A) The 18S rRNA tree was constructed with maximum-likelihood by bootstrap method (1000 replications), isolates found associated with *Vibrionaceae* by 16s barcoding analyzes are indicated with red dots. (B) *Vibrionaceae* were revealed by FISH using the VIB572a *Vibrionaceae* specific probe (Red) and

amoebae cells were counterstained with FITC. Vibrios could be observed both outside and inside (arrows) amoebae by confocal microscopy. Scale bar: 5  $\mu$ m.

**Fig. 3. LGP32 is resistant to grazing by seawater amoebae *Vannella sp.* AP1411.**

Grazing assays were performed with *Vannella sp.* AP1411 with GFP expressing strains of LGP32 or of the avirulent LMG20012<sup>T</sup>. **(A)** Bacteria survival was assessed by monitoring the intensity of the GFP fluorescence. Each condition was performed in triplicate and the results represent the average of three independent experiments. Values are presented  $\pm$  SEM \*\*\*  $P < 0.001$  (RM-ANOVA). **(B)** Amoebae growth was monitored by manual counting under phase light microscopy. Each condition was performed in 3 technical replicates and depicted results are representative of 2 independent experiments  $\pm$  SEM. Data were analyzed by two-way ANOVA with Holm-Sidak's multiple comparisons test, \*\*\*  $P < 0.001$ . **(C)** The percentage of amoebae carrying GFP expressing bacteria was quantified by flow cytometry. Each condition was performed in 3 technical replicates and depicted results are representative of 2 independent experiments  $\pm$  SEM. Data were analyzed by two-way ANOVA with Holm-Sidak's multiple comparisons test, \*\*\*  $P < 0.001$ . **(D)** LMG20012<sup>T</sup> and LGP32 were observed inside amoebae by confocal microscopy 3 days and 6 days after contact. Nuclei were stained with DAPI (Blue) and amoeba proteins with Blue Evans (Red). Scale bar: 5  $\mu$ m. **(E)** Transmission electron microscopy was performed on amoebae, 3 days after grazing on LGP32 or LMG20012<sup>T</sup> lawns. Avirulent LMG20012<sup>T</sup> vibrios were found clustered in intracellular vacuoles with an altered shape and

appearance (arrows), compared to extracellular vibrios around the amoeba. Virulent LGP32 vibrios were found mostly in individualized phagosomes, and appeared intact (arrows). Scale bar: 1  $\mu\text{m}$  and 2  $\mu\text{m}$ .

**Fig. 4. Copper p-ATPase efflux pump CopA and metalloprotease Vsm are involved in LGP32 resistance to grazing by *Vannella sp.* 1411.**

(A-D) Bacterial resistance of  $\Delta copA$  and  $\Delta vsm$  strains of LGP32 was assessed by measurement the fluorescence of the GFP-expressing bacteria, after contact with amoebae. Each condition was performed in triplicate and the results shown are the average of three independent experiments. Values are presented  $\pm$  SEM, \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$  (RM-ANOVA). (B-E) Amoeba growth was monitored by manual counting under phase light microscopy. Each condition was counted in three technical replicates. The results shown are representative of two independent experiments  $\pm$  SEM. Data were analyzed by two-way ANOVA with Holm-Sidak's multiple comparisons test, \*\*\*  $P < 0.001$ . (C-F) In two independent experiments, percentage of amoebae associated with GFP bacteria was measured by flow cytometry. Each condition was performed in three technical replicates. The results shown are representative of two independent experiments  $\pm$  SEM. Data were analyzed by two-way ANOVA with Holm-Sidak's multiple comparisons test.

**Fig. 5. Copper p-ATPase efflux pump CopA is necessary to resist to intracellular degradation in amoeba whereas the metalloprotease Vsm is not.**

(A) Intracellular vibrios were observed 3 days and 6 days after infection, by confocal microscopy. Nuclei were stained with DAPI and amoeba proteins with Blue Evans. Scale bar: 5  $\mu\text{m}$ . (B) Transmission electron microscopy was performed on amoeba 3 days after grazing on  $\Delta\text{copA}$  or  $\Delta\text{vsm}$  mutant strains. Mutant  $\Delta\text{copA}$  vibrios were found clustered in intracellular vacuoles with an altered shape and appearance (arrows). Vibrios  $\Delta\text{vsm}$  LGP32 were found mostly in individualized phagosomes, and appeared intact (arrows), scale bar: 2  $\mu\text{m}$ .

**Fig. 6. Extracellular metalloprotease Vsm decreases the velocity of the amoebae *Vannella* sp. AP1411 and alters amoeba grazing efficiency.**

Amoebae migration on LGP32wt (A),  $\Delta\text{vsm}$  (B) or  $\Delta\text{copA}$  (C) mutant strains lawns was recorded by time-lapse phase light microscopy during 10 minutes. Each presented track corresponds to the migration of one amoeba. The results shown are representative of at least two independent experiments. (D) Average velocity of amoebae on LGP32wt,  $\Delta\text{copA}$  or  $\Delta\text{vsm}$  mutant strains. The results are representative of at least two independent experiments  $\pm$  SD. Data were analyzed by Kruskal-Wallis test with Dunn's multiple comparisons test, \*\*\*  $P < 0.001$ .

#### SUPPLEMENTARY FIGURE LEGENDS

**Fig. S1. Phylogeny of the diversity of amoebae OTUs found by barcoding of the v7 region of the 18s rDNA gene.** Phylogenetic 18S rRNA-based trees of the Amoebozoa constructed with ARB software (<http://www.arb-home.de> (Ludwig *et al.*, 2004)) loaded with the silva database

(<http://www.arb-silva.de>). A base frequency filter was applied using the parsimony quick add marked tool implemented in ARB. All 18S rDNA gene sequences assigned to amoeba belonged to the *Vannellidae* family (written in red).

**Fig. S2. *Vannella sp.* AP1411 isolated from Thau lagoon, France.**

(A) Different forms of *Vannella sp.* AP1411 under phase light microscopy. Trophozoite (left), pelagic (center) and cystic form (right). Scale bar : 10  $\mu\text{m}$ . (B) Transmission electron microscopy of trophozoite shape of the amoeba *Vannella sp.* AP1411. Whole cell overview showing nuclei (N) and several digestive vacuoles (DV) with or without *E. coli* SBS363 bacteria inside (B). Scale bar: 2  $\mu\text{m}$ .

**Fig. S3. Quantification of *Vannella sp.* AP1411 phagocytosis by cytometry.** (A) FSC/SSC Gate corresponding to amoebae (B) Determination of fluorescence background on amoebae alone. (C) Gate determination for GFP+ vibrios (Bacteria). (D) Measurement of the percentage of amoebae carrying GFP fluorescence after 3 days of co-culture with GFP expressing vibrios.

**Fig. S4. *CusAB*, *SodA*, *Vsp*, *InhA*, *VipA1* and *VipA2* are not involved in LGP32 resistance to grazing by *Vannella sp.* 1411.**

Bacterial resistance to grazing by *Vannella sp.* AP1411 of  $\Delta cusAB$ ,  $\Delta sodA$ ,  $\Delta vsp$ ,  $\Delta inhA$ ,  $\Delta vipA1$  and  $\Delta vipA2$  mutant strains of LGP32 was assessed by measuring the fluorescence of the GFP-

expressing bacteria, after contact with amoebae. Each condition was performed in triplicate and the results shown are the average of three independent experiments. Values are presented  $\pm$  SEM.

**Fig. S5. Vsm or CopA do not affect the phagocytosis rate of LGP32 by the amoeba *Vannella sp.* AP1411.** Grazing amoebae on GFP-vibrio LGP32wt, LGP32  $\Delta copA$  or of LGP32  $\Delta vsm$  were monitored by flow cytometry to estimate phagocytosis index per amoeba, which correspond to the average number of phagocytosed bacteria per cell. The results shown are representative of two independent experiments. Data were analyzed by one-way ANOVA with Friedman test.

#### **Description of Additional Supplementary Files**

File name: Supplementary Movie 1

Description: Time-lapse microscopy movie of *Vannella sp.* AP1411 and LGP32wt lawn in co-culture.


File name: Supplementary Movie 2


Description: Time-lapse microscopy of *Vannella sp.* AP1411 and LGP32  $\Delta copA$  lawn in co-culture.

File name: Supplementary Movie 3

Description: Time-lapse microscopy of *Vannella* sp. AP1411 and LGP32  $\Delta vsm$  lawn in co-culture.

Accepted Article



**A****B**