

Supplementary Material




FIGURE S1 | Larvae standard length (mm) measured at 5 dpf in zebrafish (*Danio rerio*) exposed to a concentration of 10 mg/L of microplastics (MP) and microplastics spiked with oxybenzone (MP + BP3), benzo[a]pyrene (MP + BaP), perfluorooctane sulfonate (MP + PFOS) as well as a mixture of the 3 pollutants (MP + 3POPs) and the control. Data shown as mean \pm SD, n=24 per condition. ANOVA $F(5, 133)=2.1581$, $p=0.062$.

Table S1. Sequences of primers used for qPCR experiment with associated amplicon size and efficiency (mean \pm SEM).

Gene	Forward primer (5' \rightarrow 3')	Reverse primer (5' \rightarrow 3')	Amplicon size	Efficiency %
<i>cyp1a</i>	GACAGGCGCTCCTAAAACAG	CTGAACGCCAGACTCTTTCC	93	98.6 \pm 2.7
<i>actb1</i>	CCCAGACATCAGGGAGTGAT	CACAATACCGTGCTCAATGG	123	95.3 \pm 2.3
<i>g6pd</i>	GTCCCGAAAGGCTCCACTC	CCTCCGCTTTCCTCTC	124	95.8 \pm 2.2
<i>rpl13a</i>	GGACTGTAAGAGGTATGCTTCCA	ACTTCCAGCCAACCTTCATGG	200	97.5 \pm 1.2

