

Environmental DNA for detecting *Bulinus truncatus*: A new environmental surveillance tool for schistosomiasis emergence risk assessment

Stephen Mulero¹  | Jérôme Boissier¹  | Jean-François Allienne¹ |
Yann Quilichini²  | Joséphine Foata² | Jean-Pierre Pointier³ | Olivier Rey¹ 

¹IHPE UMR 5244, CNRS, IFREMER, Univ. Montpellier, University of Perpignan, Perpignan, France

²UMR SPE 6134, CNRS - Università di Corsica Pascal Paoli, Corte, France

³USR3278 CRILOBE EPHE-CNRS-UPVD, PSL University, Perpignan, France

Correspondence

Olivier Rey, IHPE UMR 5244, CNRS, IFREMER, Univ. Montpellier, University of Perpignan, F-66860 Perpignan, France.
Email: olivier.rey@univ-perp.fr

Funding information

Agence Nationale de Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail, Grant/Award Number: PNREST 2019/1/059 Molrisk; Agence Nationale de la Recherche, Grant/Award Number: ANR-18-CE35-0001

Abstract

Under ongoing climate changes, the development of large-scale monitoring tools for assessing the risk of disease emergence constitutes an urging challenge. This is particularly the case for snail-borne diseases such as the urogenital bilharziasis that emerged in Corsica and threat European countries. The expansion of this tropical disease mainly relies on the local presence of competent snail hosts such as *Bulinus truncatus*. Unfortunately, very little is known about the actual repartition of freshwater snails worldwide which makes new emergences difficult to predict. In this study, we developed two ready-to-use environmental DNA-based methods for assessing the distribution of *B. truncatus* from water samples collected in the field. We used two approaches, a quantitative PCR (qPCR) and a droplet digital PCR (ddPCR) approach. We successfully detected *B. truncatus* in natural environments where the snail was previously visually reported. Our environmental DNA diagnostic methods showed a high sensitivity (≈ 60 DNA copy per mL of filtered water) and a high specificity to *B. truncatus*. Results obtained in qPCR and ddPCR were very similar. This study demonstrates that environmental DNA diagnostics tools enable a sensitive large-scale monitoring of snail-borne diseases hence allowing the delimitation of areas potentially threatened by urogenital schistosomiasis.

KEYWORDS

Bulinus truncatus, Corsica, ddPCR, environmental DNA, environmental monitoring, qPCR, schistosomiasis

1 | INTRODUCTION

Global changes, including climate changes (e.g., temperatures, precipitations), land use, chemical pollutants, and human-driven dispersion, greatly modify the spatial distribution of organisms at a global scale (Parmesan & Yohe, 2003; Pereira et al., 2010). While some

species become locally extinct due to climatic and/or anthropic pressures (Pereira et al., 2010), other species extend their geographical range with some populations sometimes reaching impressive abundance at the global scale (i.e., invasive species; Capinha, Essl, Seebens, Moser, & Pereira, 2015). Such rapid changes in both the distribution and the abundance also concern pathogens (Jones et al.,

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Environmental DNA* published by John Wiley & Sons Ltd

2008; Kim, Kabir, & Ara Jahan, 2014), and the resulting (re-)emerging diseases constitute a threat to animal (Cohen et al., 2017; Gozlan, St-Hilaire, Feist, Martin, & Kent, 2005) and to human health (Blum & Hotez, 2018; Morens & Fauci, 2013). Understanding and predicting the dynamics and epidemiology of emerging diseases in response to environmental changes have thus become an urging challenge in the last decades (Ellwanger, Kaminski, & Chies, 2019).

Among the emerging diseases documented so far, 17% are induced by pathogens transmitted by vectors and/or intermediate hosts (i.e., vector-borne diseases; VBD) (WHO, 2017). Predicting the emergence and spread of VBD is particularly challenging because it requires knowledge on the distribution and the spatio-temporal dynamics of pathogens and of their intermediate hosts and/or vectors that are also potentially influenced by current global change (Blum & Hotez, 2018; Richards, Lord, Pesko, & Tabachnick, 2009). In this regard, recent studies together with the development of international monitoring networks (e.g., VBORNET in Europe; ECDC, 2011) have made it possible to establish real-time risk maps of emergence and spread of some arthropod-borne-diseases such as malaria, chikungunya, and dengue (ECDC, 2014; Schaffner et al., 2013).

The monitoring of snail-borne diseases (SBD), and more particularly those transmitted via freshwater snail species, still remains in its infancy (Kincaid-Smith, Rey, Toulza, Berry, & Boissier, 2017). The neglected aspect of most SBD diseases contrast with their significant impact on human health. For instance, more than 250 million peoples are infected by *Schistosoma* parasites worldwide (Stensgaard, Vounatsou, Sengupta, & Utzinger, 2019). This infectious disease is also responsible for more than 200,000 deaths per year (WHO, 2002). The SBDs concern several vertebrates including humans, livestock, and wild animals. Consequently, these diseases also have a significant economic impact especially in the livestock industry (e.g., 2.5€ billion of annual losses linked to the fascioliasis worldwide; Jones et al., 2018). They are induced by some parasites that depend on a snail host to complete their lifecycle. Our current knowledge on the distribution and dynamics of SBD mainly relies on epidemiological studies based on the prevalence and intensities of adult parasites within their definitive hosts (either humans or animals), while the distribution of snail species involved in the transmission of SBD is generally poorly documented (Kincaid-Smith et al., 2017). This lack of knowledge can be explained by at least four main reasons: first, until very recently, freshwater ecosystems received less ecological attention than terrestrial systems (Collen et al., 2014; Puth & Post, 2005). Second, the distribution of snail populations is highly heterogeneous at both the spatial and temporal scale which makes their monitoring difficult (Lamy et al., 2012). Third, the available classical methods used to collect and identify snail species are laborious and time-consuming and important sampling efforts are generally necessary for assessing, with certainty, the presence/absence of a targeted species in the field (Lamy et al., 2012). Thus, it is currently inconceivable to monitor snails' populations at large spatial and temporal scales. Finally, once the snails collected from a given site, the taxonomic identification of specimens at the species level requires thorough malacological expertise

and generally these identifications still need to be completed by DNA analyses.

Methods based on environmental DNA (eDNA) constitute a promising approach for monitoring biodiversity (Bohmann et al., 2014). These methods were recently applied in a parasitological context to detect (cryptic) parasite species, their hosts and/or vectors (Bass et al., 2015). In this respect, eDNA has been used for detecting targeted parasites species (Huver, Koprivnikar, Johnson, & Whyard, 2015; Richey et al., 2018; Rusch et al., 2018), for characterizing parasite communities (Hartikainen et al., 2016), and to a lower extent for the detection of parasites' vectors (e.g., the invasive mosquito vectors *Aedes albopictus*; Schneider et al., 2016). Regarding SBDs, studies have mainly focused on the detection of free-living parasite infectious stages present in the water including *Opisthorchis viverrini* (Hashizume et al., 2017); *Fasciola hepatica* (Jones et al., 2018); *Schistosoma japonicum* (Worrell et al., 2011); and *Schistosoma mansoni* (Sengupta et al., 2019). This approach targeted toward the detection of parasites is particularly relevant to identify active transmission sites, that is, where the parasite is currently present and susceptible to infect definitive hosts such as humans or livestock species. Surprisingly, very few studies have developed eDNA-based methods to detect the presence of snail species that constitute intermediate hosts for the parasites and that are thus responsible for the transmission of the associated diseases. To the best of our knowledge, eDNA-based methods were developed only for the detection *Galba truncatula* and *Austropeplea tomentosa* (Jones et al., 2018; Rathinasamy et al., 2018), both species being involved in the transmission of the liver fluke (*Fasciola hepatica*), and more recently for the detection of *Oncomelania hupensis* involved in the transmission of *S. japonicum* (Calata et al., 2019). Such approach targeted toward the detection of snail hosts is of prime interest to identify sites where the associated disease could emerge (Kincaid-Smith et al., 2017).

In this study, we develop an eDNA-based method to detect the presence of *B. truncatus* from freshwater water samples. This snail species hosts two flatworm parasites *S. bovis*, a cattle parasite, and *S. haematobium*, the latter being responsible for the urogenital form of schistosomiasis. The urogenital schistosomiasis is one of the most severe and prevalent forms of schistosomiasis in Sub-Saharan Africa threatening 436 million individuals and affecting 112 million people throughout the world (WHO, 2018). Moreover, infection with *S. haematobium* also constitutes an extra risk factor for secondary infections with other infectious agents such as HIV, especially in women (WHO, 2002). The morbidity associated with urogenital schistosomiasis fortunately globally decreased in the last decades although its prevalence has increased throughout Africa (Herricks et al., 2017). Moreover, the future distribution of this disease in response to climate change is difficult to predict at the continental scale (Blum & Hotez, 2018) and outside the endemic area (Kincaid-Smith et al., 2017). In this regard, an outbreak of urogenital schistosomiasis has recently occurred in Southern Europe in Corsica (Boissier et al., 2016; Holtfreter, Mone, Muller-Stover, Mouahid, & Richter, 2014). This situation has called for an urgent need to develop environmental

diagnosis tools to monitor the distribution of compatible freshwater mollusks and hence predict possible new transmission sites in Europe (Kincaid-Smith et al., 2017).

Importantly, no vaccine exists against schistosomiasis and mass chemotherapy by the administration of praziquantel, an anti-helminth drug, remains the main strategy to control the disease in endemic areas. Today, the control of snail populations under a certain threshold to limit transmission is acknowledged as a complementary strategy necessary to eradicate schistosomiasis (Bergquist et al., 2017). Better characterizing the distribution of host snails is thus fundamental for targeting snail control strategies. So far, eDNA-based methods were developed to detect the presence of free-living stages (cercariae) of *S. japonicum* (Worrell et al., 2011) and *S. mansoni* (Sengupta et al., 2019) present in the water and susceptible to infect humans, or within their snail hosts (Amarir et al., 2014), but no methods were developed to detect host snail species except for *O. hupensis* (Calata et al., 2019).

We here propose a ready-to-use eDNA protocol to detect the presence of *B. truncatus* established in natural river streams from water samples. We also compared a classical optimized qPCR approach to a protocol using the droplet digital PCR technology (ddPCR), also called "third PCR generation" that provides interesting properties for its application in eDNA (Doi, Takahara, et al., 2015). First, ddPCR allows the absolute quantification of low levels of nucleic acids. Second, this technology is expected to be more robust and hence provide more repeatable results when amplifying DNA in the presence of PCR inhibitors (Hindson et al., 2011; Taylor, Laperriere, & Germain, 2017), the latter being an important technical hurdle in eDNA studies (Doi, Takahara, et al., 2015). So far, only 11 studies used ddPCR for eDNA detection (Baker et al., 2018; Coble et al., 2019; Doi, Takahara, et al., 2015; Doi, Uchii, et al., 2015; Hunter, Ferrante, Meigs-Friend, & Ulmer, 2019; Hunter, Meigs-Friend, Ferrante, Smith, & Hart, 2019; Jerde et al., 2016; Nathan, Simmons, Wegleitner, Jerde, & Mahon, 2014; Rusch et al., 2018; Uthicke, Lamare, & Doyle, 2018; Wacker et al., 2019). Based on our results, we compare and discuss the pros and cons of these two approaches in an eDNA context.

2 | MATERIALS AND METHODS

2.1 | Field sampling

2.1.1 | Sampling sites

We focused our sampling on five sites along the Cavu and the Solenzara Rivers (Corsica), two geographically isolated rivers where a urogenital schistosomiasis outbreak occurred very recently (Boissier et al., 2016; Ramalli et al., 2018) (Table 1). These sites in Corsica were chosen because the temporal dynamic of the local *B. truncatus* populations is monitored in these two rivers annually since 2014. In the Cavu River, we sampled water from three sites where populations of *B. truncatus* are well established and where

transmission occurred (Table 1): at the Mulinu Bridge, hereafter denoted "MB"; at the recreational A Tyroliana Park, hereafter named "AT," and at the "3 Pool" site, hereafter denoted "3P" these sites respectively correspond to site 5, 8, and 9 in Boissier et al. (2016). We sampled water from a fourth site upstream the Cavu River, hereafter denoted "WI" (i.e., Water Intake; corresponding to site 10 in Boissier et al., 2016) where no *B. truncatus* has ever been observed. This site was sampled to validate or invalidate the absence of *B. truncatus* at this site so far determined through visual inspection only. Along the Solenzara River, we sampled water from one site (the U Rosumarinu Snack, hereafter denoted "UR") where local infection with schistosomiasis occurred recently (Noel et al., 2017). Additionally to these sampling sites in Corsica, we also sampled water from an artificial channel in Perpignan city (Southern France) where *B. truncatus* never occurred. This site was considered as a field negative control. All samples were collected during the first week of September 2018. This period succeeds to the highest density period of *B. truncatus* in Corsican rivers (i.e., august; Kincaid-Smith et al., 2017) and before high-water period (i.e., October–November). At each site, we also measured environmental conditions: including water temperature, pH, flow at the water surface (measured in the river bed), and human activity (i.e., presence/absence of anthropic structures/individuals).

2.1.2 | Water sampling protocol

Our sampling protocol was designed to quantify possible effects of filtration volume and sampling location relative to the streambed (i.e., shore vs. streambed of the river) on our ability to detect *B. truncatus* DNA at a given site. To this aim, a total of 7 water samples were collected at each site. Specifically, three water volumes (1, 3, and 5 L) were filtered at two different locations within the river (i.e., shore vs. streambed) at each sampling site. Moreover, 500 ml of commercial spring water was filtered at each site and rigorously following the same protocol as a technical field negative control. Each water filtration was achieved using a filtration unit including a 0.45- μ m PES sterile membrane of 90 mm (VWR PES filter Unit 1 L, Model 514-0301) connected to a manual vacuum pump (Mityvac, Model MV8500). This mesh size constitutes a good compromise between membrane clogging rate and eDNA capture capacity for lotic environment (Hinlo, Gleeson, Lintermans, & Furlan, 2017). The resulting filtration membrane was then removed from its initial filtration unit, immediately stored in a 50 ml falcon filled with a Longmire buffer (Longmire, Maltbie, & Baker, 1997), and stored at room temperature until subsequent DNA extraction. We followed classical specific recommendations to avoid DNA contamination overall the sampling process (Taberlet, Coissac, Hajibabaei, & Rieseberg, 2012). In particular, scissors and forceps used to remove the membranes from filtration units were decontaminated before each sampling by successively placing instruments in a 10% bleach bath during 1 min, 1 min in a DNA AWAY™ solution (Thermo Scientific) followed by a flame sterilization step.

TABLE 1 Geographical coordinates (using Google Mercator projection), biotic, and abiotic parameters at each prospected sites

Site acronym	MB	AT	3P	WI	UR	Control
Geographical coordinates	41°42'16.82"N 9°20'5.23"E	41°43'12.22"N 9°18'6.27"E	41°43'53.14"N 9°17'36.87"E	41°44'29.30"N 9°17'39.58"E	41°50'37.86"N 9°20'36.87"E	42°40'55.60"N 2°54'4.14"E
Sampling date	26/09/2018 3:30 p.m.	26/09/2018 11:40 a.m.	26/09/2018 9:40 a.m.	26/09/2018 8:30 a.m.	27/09/2018 9 a.m.	04/10/2018 15 p.m.
pH	8.3	8.5	8.7	9	8.3	8.4
T (°C)	21.5	20	18.8	18	19.5	17.4
Human activity	Yes	Yes	Yes	Yes	Yes	No
Water flow	1 m/s	0.5 m/s	1 m/s	0.1 m/s	0.5 m/s	0.5 m/s
Snail density	10/m ²	50/m ²	>100/m ²	0/m ²	>100/m ²	0/m ²

Abbreviations: 3P, 3 Pool; AT, A Tyroliana Park; Control, Perpignan University channel; MB, Mulinu Bridge; UR, Solenzara snack; WI, Water intake.

Snail species	Origin	Sampling date	Collaborators
<i>Bulinus truncatus</i> ^a	Corsica, France	2018	
<i>Bythinia tentaculata</i>	Corsica, France	2018	
<i>Radix balthica</i> ^a	Corsica, France	2018	
<i>Ancylus fluviatilis</i> ^a	Corsica, France	2018	
<i>Gyraulus laevis</i>	Corsica, France	2018	
<i>Gyraulus sp.</i> ^a	Corsica, France	2018	
<i>Physa acuta</i> ^a	Corsica, France	2018	
<i>Potamopyrgus antipodarum</i> ^a	Corsica, France	2018	
<i>Pysidium sp.</i>	Corsica, France	2018	
<i>Theodoxus fluviatilis</i>	Corsica, France	2018	
<i>Biomphalaria glabrata</i>	Brazil	2018	
<i>Bulinus forskalii</i>	South-Africa	2017	Tine Huyse, KU Leuven, Belgium
<i>Bulinus globosus</i>	Senegal	2012	Tine Huyse, KU Leuven, Belgium
<i>Bulinus africanus</i>	South-Africa	2017	Tine Huyse, KU Leuven, Belgium
<i>Radix natalensis</i>	South-Africa	2017	Tine Huyse, KU Leuven, Belgium
<i>Galba truncatula</i>	Corsica, France	2018	

^aSpecies which occurrence was reported in the Cavu River and/or Solenzara River.

To evaluate the repeatability of our results, technical duplicates were achieved at two sampling sites (i.e., MB and AT) for which we respectively observed the lowest *B. truncatus* density (≈ 10 mollusks/m²) and an intermediate density of ≈ 50 mollusks/m². A total of 13 filtering membranes were thus obtained for each of these two sites ([3 filtrations volumes \times 2 river localities \times 2 duplicates] + 1 technical negative control).

Once all the water samples were collected, a visual inspection was achieved to qualify the local density of *B. truncatus* in the sampling area (low: 10–50 mollusks; medium: 50–100 mollusks; and high: more than 100 mollusks). This was achieved by inspecting rocks and the vegetation on a 1 m² surface on the shore of the river and following the same capture effort (30 min per site).

TABLE 2 Snails' species used for assessing the specificity of the developed qPCR and ddPCR primers

2.2 | Development of qPCR primers

2.2.1 | In silico development of qPCR primers

To design specific primers, we focused on the mitochondrial *cytochrome c oxidase I (cox1)* gene that have been first proposed for barcoding studies on animal DNA (Hebert, Cywinska, Ball, & deWaard, 2003) and classically (although not exclusively) used for the detection of animal eDNA (Tsuji, Takahara, Doi, Shibata, & Yamanaka, 2019). A total of 25 primer pairs were developed based on an alignment of 210 *cox1* sequences from *B. truncatus* from several countries available from the Barcode of Life Database BOLD (Ratnasingham & Hebert, 2007) to account for intraspecific variability. Primer pairs

were designed using Geneious® 8.1.4. (Kearse et al., 2012) and following classical criteria so as to optimize hybridization. The specificity of the resulting primer pairs to *B. truncatus* was first assessed in silico. To do so, a local database regrouping *cox1* sequences from 14,500 *Mollusca* species and available on BOLD and NCBI nucleotide databases was first created. Virtual PCRs using our designed primer sets were ran on the resulting database using ecoPCR 0.5.0. (Boyer et al., 2016) for detecting possible cross-amplification on related snail species. To check whether these primers could amplify a similar amplicon on other nonfreshwater mollusk species, we also ran a Primer-BLAST analysis against the nr database (Ye et al., 2012). The nr database contains nonredundant set of sequences (i.e., genomic DNA, transcripts, and proteins) from several databases hence allowing the use of BLAST analysis against DNA sequences from other species. This bioinformatic pipeline is summarized in Figure S1. All primer pairs showing a 100% specificity to *B. truncatus* ($N = 4$) were kept for subsequent qPCR specificity tests.

2.2.2 | Testing the specificity of the qPCR primers

The specificity of the resulting qPCR primers was assessed in vitro by qPCR on DNA templates from a total of 16 freshwater snail species (see Table 2), including three *B. truncatus* sister species (i.e., *B. forskalii*, *B. globosus*, and *B. africanus*). Moreover, qPCRs were also performed on an equimolar pool of DNA extracts from each of these 16 species (at 0.5 ng/μl each) and including or excluding *B. truncatus* DNA. Only the primer pairs amplifying exclusively *B. truncatus* DNA in individual DNA or pooled DNA were kept for the ensuing experiment.

2.3 | Molecular analyses

All molecular experiments were carried out under a sterile table top hood. Before and after each protocol (i.e., DNA extraction, PCR reaction preparation), the working surface was bleached, washed using a DNA AWAY™ solution (Thermo Scientific™), and then exposed to UV light during one hour.

The DNA positive controls were extracted from 5 *B. truncatus* individuals that were collected in the field in Corsica. Total genomic DNAs from these mollusks were extracted using the E.Z.N.A.® Tissue DNA kit (OMEGA Bio-Tek, Inc) following the “tissue protocol” for animal tissues, supplied in the kit handbook.

Total DNA from each membrane was extracted using the DNeasy® Blood & Tissue Kit (QIAGEN) which is commonly used in eDNA studies and which is supposed to yield an important amount of extracted DNA (Goldberg et al., 2016; Hinlo et al., 2017). We followed an adapted version of the “tissue” protocol from the kit handbook. Briefly, membranes were subdivided into 4 equal sections to allow the use of microcentrifuge tubes during subsequent extraction steps. Lysis was performed at 65°C during 1 hr using a final solution containing 567 μl of ATL buffers and 63 μl of proteinase K. We next

added 630 μl of AL buffers and 630 μl of 100% ethanol in the following steps (steps 2 and 3). The subsequent steps were performed following the manufacturer protocol, and the resulting genomic DNA was finally eluted in 100 μl of AE buffer. At the end of the extraction, we thus obtained four eluates of 100 μl for each membrane.

2.3.1 | Quantitative PCR approach

The four sections from each membrane were analyzed twice meaning that a total of 8 qPCR reactions were ran for each membrane. A total of 200 sections from 50 filter membranes (4 sections per membranes) were thus duplicated and processed in qPCR. Hence, 400 qPCR reactions were used for analyzing the 50 membranes collected. These 400 reactions were performed on two independent qPCR runs each of which containing the negative and positive qPCR controls. To limit possible biases due to inter-run C_t differences in subsequent statistical analyses, all samples that were included in our analyses were processed on the same first qPCR run.

The qPCRs reactions were performed using the Takyon™ No ROX SYBR® kit (Eurogentec) in a total volume of 10 μl, containing 2.5 μl of undiluted template DNA, 5 μl of MasterMix 2X (final concentration = 1X), and 2.5 μl of 10 μM primer mix (final concentration = 0.2 μM). The qPCR program was set as follow: 95°C for 3 min followed by 45 cycles of 95°C for 15 s and 60°C for 1 min, using a LightCycler®480 qPCR device. The resulting fluorescence signals were analyzed using the LightCycler®480 software v1.5.0. A signal was considered positive (i.e., with the presence of *B. truncatus* DNA) if the melting peak obtained for the sample displayed a melting temperature comprised between 79 and 82°C as defined by the classical melting curve obtained for the PCR positive control used in all qPCR runs (i.e., a 0.15 ng/μl DNA extract from a *B. truncatus* individual). Finally, we considered that eDNA from *B. truncatus* was present on a filtering membrane if two qPCR replicates from the same membrane section (i.e., at least 1 section fully positive) displayed a melting curve specific to *B. truncatus*.

Finally, the eDNA signal of *B. truncatus* obtained in qPCR was validated on a subset of 20 membranes section. Sequences from these qPCR products were obtained using ABI 3730xl sequencer at the GenoScreen platform (Montpellier, France).

2.3.2 | Droplet digital PCR approach

Because ddPCR is highly sensitive and because it is supposed to be robust even in the potential presence of PCR inhibitors, the ddPCR reactions were ran on pools of DNA extracts from the four sections of each membrane. Noteworthy, this pooling strategy was also tested in our qPCR analyses but failed to provide repeatable results. The ddPCRs were ran using the TaqMan chemistry on a QX200 AutoDG Droplet Digital System (Bio-Rad) at the I.A.G.E. platform (Montpellier, France). More specifically, we amplified a 165 bp region of the *cox1* gene using the Btco2F (5' ATTTTGACTTTTACCACCAT 3') and

Btco2R (5' GATATCCCAGCTAAATGAAG 3') primers and using the FAM-labeled Btco2P probe (5' TCGAAGGAGGGTTGGAACAGG-FAM 3') specific to *B. truncatus* in the ddPCR reactions. These primers and probe were generated on a consensus sequence of *B. truncatus* obtained from the same sequence file used as a template for the design of qPCR primers (i.e., alignment of 210 *B. truncatus* *cox1* sequences downloaded from the NCBI). Then, the best primers and probe combination were selected through melting temperature comparison and BLAST analyses against online databases and a local sequence file containing 108 *cox1* sequences of other freshwater snail species. The specificity of the developed primers and probe complex was next tested using the same mollusk DNA templates as those used for testing the specificity of the qPCR Btco1 primers, that is, individual ddPCR reactions were run on four equimolar pools, two of them containing pooled DNA of 3 *B. truncatus* sister species including and excluding *B. truncatus* DNA, the two others containing pooled DNA from endemic freshwater snail species of Corsica (Table 2) with and without *B. truncatus* DNA. According to the results from this test of specificity, only the primers and probe complex mentioned above showed a perfect specificity to *B. truncatus* and was selected for the ensuing experiments. Individual ddPCR reactions were performed on all membranes including negative controls as well as on the same positive control as for qPCR in a final volume of 20 μ l containing 9 μ l of template DNA. The resulting fluorescence signals were analyzed using QuantaSoft™ software v1.7. (BioRad). A signal was considered positive (i.e., with the presence of *B. truncatus* DNA) if (a) at least one positive droplet was detected and (b) the positive droplet displayed the same order of fluorescence magnitude as the positive droplets obtained for a ddPCR positive control. Finally, we considered that eDNA from *B. truncatus* was present on a filtering membrane if at least one positive droplet met the above criteria.

From the results obtained using the ddPCR approach, the number of *B. truncatus* eDNA copies per liter (c/L) of filtered water for each sample (i.e., each membrane) was estimated using the following equation (Agersnap et al., 2017; Rusch et al., 2018):

$$C_L = \frac{C_{rdd}}{0.9} * \frac{V_e}{V_r} \cdot \frac{V_w}{V_e}$$

With C_L : the number of eDNA c/L for the amplified sample; C_{rdd} : the copy number per reaction volume (c/20 μ l) and applying a 10% correction of losses during droplet generation; V_e : the total volume of eluted DNA after extraction; V_r : the volume of extracted DNA used for ddPCR reaction; and V_w : the total volume of filtered water (Agersnap et al., 2017).

2.4 | Data analysis

Several factors can influence the concentrations of eDNA isolated from environmental samples including the sampling protocols used (Dickie et al., 2018; Li, Lawson Handley, Read, & Hanfling, 2018) and the local abundance of the targeted species, the latter factor being

linked to the temporal demographic fluctuations of natural populations (Wilcox et al., 2018). With these considerations in mind, we statistically defined the most suitable conditions for optimizing the eDNA detection following our filtration protocol. To this end, we ran several generalized linear models (GLM) to infer the effect of the volume of filtered water (vol), the sampling position (pos), and the local density of *B. truncatus* (dens) on eDNA yielding, that is, cycle threshold values (C_t) obtained in qPCR and log-transformed c/L obtained in ddPCR. We also accounted for possible interactions between these variables in our models. The most parsimonious model for explaining c/L and C_t were selected based on the Akaike information criterion (AIC) (Akaike, 1974; Burnham & Anderson, 2003; Buxton, Groombridge, & Griffiths, 2017). When several models were equally supported (i.e., Δ AIC < 2; Burnham & Anderson, 2003), we retained the model containing the lowest number of parameters. All GLMs were generated using the function glm implemented in R 3.4.3 using Rstudio 1.1.383 interface (RStudio, 2016).

3 | RESULTS

3.1 | Sampling sites

The environmental conditions (e.g., temperature, pH) were very similar among the sampling sites. Likewise, all sites presented human activities (generally bathing) except the control site that is inaccessible to the public. The water flow varied from one site to another, ranging from 0.1 m/s for the site WI to 1 m/s for the site MB; these data are displayed in Table 1.

3.2 | Development of qPCR primers

Among all the primer pairs tested, Btco1F (5' TYGAAGGAG GGGTTGGAACA 3') coupled with Btco1R (5' RKTRATTCC TGGTGCYCGT 3') provided the best results regarding the qPCR specificity tests (i.e., only the DNA from *B. truncatus* or pooled DNA containing *B. truncatus* DNA were positive in qPCR) and were thus used for subsequent qPCR analyses. This primer pair resulted in the amplification of a 179 bp amplicon specific to *B. truncatus*.

3.3 | Molecular analyses

3.3.1 | Quantitative PCR approach

Based on the result obtained from the qPCR approach and according to our validation criteria, DNA from *B. truncatus* was detected from a total of 35 among the 44 field membranes analyzed (79.5%) at sites MB, AT, 3P, and UR. No *B. truncatus* DNA was detected in the water samples collected from the WI site which confirms our previous environmental diagnostic through visual inspections at this site. No *B. truncatus* DNA was detected in our negative control site

(i.e., University channel) or in the 6 field negative controls (i.e., spring water filtered at each sampling site). Noteworthy, two among the total of 48 field negative control replicates showed a measurable C_t value (Site UR: 37.39 and Site MB: 39.17). However, these two samples displayed unspecific melting curves compared to those obtained from positive controls. Likewise, none of our 6 qPCR negative controls showed a measurable C_t value. These results indicate that DNA contamination during the overall process from water sampling to qPCR reactions is very unlikely.

The water samples filtered on the river shore and in the streambed provided similar results, that is, 85.7% [18/21] of positive samples at the shore and 81% [17/21] at the streambed location (Figure 1). However, we observed a slight tendency in the C_t values to be higher for samples filtered in the river shore (35.67 ; $SD \pm 1.79 C_t$) than those filtered in the river streambed (37.71 ; $SD \pm 1.69 C_t$). The median C_t value for all eDNA samples was 35.2 ; $SD \pm 1.7$, ranging from 32.3 to 38.84 . Moreover, our sequencing validation step confirmed that the amplified DNA fragments during qPCR belonged to *B. truncatus* with a pairwise identity ranging from 97.52% to 100% and with a coverage of 97% to the matching sequences (NCBI accession number: MG407340.1 and MG407341.1).

3.3.2 | Droplet digital PCR approach

The ddPCR provided the same results to those obtained using the qPCR approach. DNA extracts from the filtered membranes obtained from sites MB, AT, 3P, and UR were positive to *B. truncatus* DNA (79.5%, 35/44 membranes) using ddPCR (i.e., at least one positive droplet presenting the same fluorescence amplitude as the DNA control was detected). Eighteen of the twenty-one filtration achieved on the river shore were positive to *B. truncatus* DNA (i.e., 85.7%), and 17/21 (i.e., 81%) filtration achieved on the streambed of the river were positive (Figure 2). No positive droplet was detected for the two ddPCR negative controls, for the six field negative controls, and for samples collected at site WI. The estimated number of DNA copies per ddPCR reaction volume ranged from $1.27 c/20 \mu l$ to $42.82 c/20 \mu l$ with a median value of $9.14 c/20 \mu l$; $SD \pm 9.61 c/20 \mu l$. The median number of DNA copies per liter of filtered water was $550.15 c/L$; $SD \pm 610.21 c/L$.

The lowest sensitivity threshold estimated from the ddPCR analyses (i.e., the lowest eDNA concentration detected in our samples) was $0.06 c/\mu l$ which represents approximately $60 c/ml$ of filtered water.

3.4 | Statistical analyses

Among all GLMs performed using the qPCR dataset, the best-supported model was the following: $C_t \sim \text{volume} + \text{position} + \text{density}$ (Table 3). According to this model, the snail density ($p < .001$) and the volume of filtered water ($p < .01$; see Table 3 and Figure 3) significantly affect the C_t values obtained by qPCR after molecular process.

Accordingly, a higher water volume and a higher mollusk density decrease the C_t value obtained which means that the concentration of *B. truncatus* eDNA is higher for these samples. A significant positive effect of the sampling position (shore) was also found ($p < .05$; Table 3), hence indicating that the C_t value obtained by qPCR tends to be higher and thus eDNA concentration is lower in water samples collected on the river shore compared to the streambed.

Concerning the ddPCR dataset, the best-supported model explaining c/L predicted a single positive effect of the *B. truncatus* densities ($p < .001$; Table 3). This result indicates that a higher mollusk density results in a higher eDNA concentration which is congruent with our expectations (Figure 4).

4 | DISCUSSION

Under the current global change context, it is essential to develop large-scale monitoring tools to understand and predict the dynamics and epidemiology of emerging diseases (Ellwanger et al., 2019). In this study, we present the first application of two eDNA-based methods, namely a ddPCR- and a qPCR-based protocols, to detect environmental DNA of *B. truncatus*, an intermediate host of *S. haematobium*, in natural rivers. Indeed, we were able to detect the presence of *B. truncatus* from water samples collected at four natural transmission sites previously reported in Corsica (Ramalli et al., 2018), while no *B. truncatus* eDNA was detected in sites where this species had never been visually reported before. The sensitivity of our two protocols reached 100% (assessed in silico and in vitro) when using 3 or 5 L of water during filtration but dropped to 91.7% when filtering 1 L of water using both qPCR and ddPCR. This result indicates that according to our results, at least 3 L of water is necessary to reach an optimal sensitivity whatever the density of mollusks at the sampling site.

More specifically, the sampling procedure that provides the best detection rate consists of a 5-liter filtration preferentially in the river streambed. Whatever the PCR technology used, we can argue that in this current version our protocols designed for water collection, sample preservation, and eDNA extraction are robust. The considerations of these three protocol aspects are critical for establishing a reliable eDNA-based approach (Goldberg et al., 2016) but are generally neglected in most publications. Only 5% of eDNA-based studies provided critical aspects of their sampling protocols (Dickie et al., 2018) and most eDNA studies suffer from a lack of reproducibility especially when their protocols are applied in natural environments for the first time (Tsuji et al., 2019). Finally, no standardized method for eDNA sampling is available so far (Tsuji et al., 2019). This technical lack raises the necessity of further studies benchmarking all existing sampling protocols in an attempt to define standardized sampling pipelines adapted to different environments and environmental samples (i.e., water, sediment, and air).

In this line, several aspects could be developed to improve our eDNA-based monitoring tools (Dickie et al., 2018). First, the robustness of our protocols should be validated in different freshwater




FIGURE 1 Membranes positiveness obtained from the quantitative PCR approach. The displayed percentage is calculated on the 8 qPCR replicates from the samples filtered on the river shore (a) and at the river streambed (b). Membranes are considered positive if at least two replicates from the same membrane section are positive






FIGURE 2 Membranes positiveness obtained from the droplet digital PCR approach. All positive membranes from the samples filtered on the river shore (a) and at the river streambed (b) are displayed in green. A membrane is considered positive if at least one positive PCR reaction compliant with the fluorescent signal from the DNA control is detected



environments with different hydrodynamic profiles. In the present study, all of the prospected sites, but the negative University channel site, presented a lotic hydrodynamism (i.e., flowing water) and whether the sampling strategy developed in this study could be applied in lentic environments (i.e., still freshwater) still needs to be assessed. Lentic environments are known to present some specific characteristics that can influence the eDNA decay (e.g.,


temperature, sunlight, and pH; Klobucar, Torrey, & Phaedra, 2017; Strickler, Fremier, & Goldberg, 2015). They also generally contain significantly more suspended solids (Takahara, Minamoto, Yamanaka, Doi, & Kawabata, 2012), which could increase the PCR inhibitors concentration in eDNA samples. Second, the ergonomic aspect and efficiency of our sampling tools could be further optimized. For instance, filtration units are voluminous and are probably

TABLE 3 Best fitted GLM models that explain (C_t) and (c/L)s obtained by qPCR and ddPCR based on the AIC criterion and number of parameters (nPars) (a) and the parameter estimates obtained from the chosen models

(a)				
GLM models	nPars	AIC	Δ AIC	Platform
$C_t \sim \text{vol} + \text{pos} + \text{dens}$	5	134.92	0	qPCR
$C_t \sim \text{vol} * \text{pos} + \text{dens}$	6	134.94	0.02	
$C_t \sim \text{dens}$	3	135.94	1	
$C_t \sim \text{vol} + \text{dens}$	4	136.07	1.13	
$c/L \sim \text{dens}$	3	106.11	0	ddPCR
$c/L \sim \text{pos} + \text{dens}$	4	108.08	1.97	
(b)				
Variable	Estimate	SE	z-value	p-value
qPCR				
Intercept	36.685959	0.446015	82.253	<2e-16
3 L	-0.764357	0.418130	-1.828	.06754
5 L	-1.181857	0.418130	-2.827	.00471
Shore	0.861609	0.338734	2.544	.01097
Density	-0.022992	0.004628	-4.968	6.76e-07
ddPCR				
Intercept	4.800520	0.303499	15.817	<2e-16
Density	0.017062	0.004619	3.694	.00022

inappropriate when field conditions are difficult (e.g., isolated prospecting sites; long fieldwork). The size-reduction of filtration devices could also be contemplated, for instance using smaller membrane holders coupled with a vacuum automated pump, although the latter could also be a limitation in some field conditions. Likewise, the use of silicate micro-beads (Wilcox et al., 2018) or dried membranes (Balint et al., 2018) could be more convenient than Longmire

solution for sample preservation although this would need to be validated. In terms of eDNA capture, the previous studies benchmarking different filtration membrane composition found that nitrate cellulose membranes (NC) coupled with DNA extraction with DNeasy® Blood & Tissue Kit provide higher eDNA yielding (Hinlo et al., 2017), these findings could also constitute relevant improvement to consider.

**FIGURE 3** Representation of the estimated qPCR cycle thresholds (residuals) as a function of the volume of filtered water and the *B. truncatus* densities at the sampling sites. ** and *** represent differences with significant levels below 0.01 and 0.001 respectively

Another major finding of this study is the potential of the ddPCR as an alternative quantification platform to the qPCR for environmental samples. In our study, qPCR and ddPCR gave strictly the same results regarding sensitivity since the same samples were found positive whatever the technology used. The only difference between these two PCR technologies is that ddPCR was able to provide reliable results on the pooled samples while qPCR did not. This result hence provides a small advantage of ddPCR compared to qPCR because it requires less replicates to give reliable results. Finally, because we did not conduct comparative analyses using qPCR and ddPCR on the same dilution standards, it is difficult to conclude on the most accurate technology regarding absolute quantification of environmental DNA and ultimately mollusk density in the field. Thus, based on our results and because ddPCR is generally argued as very sensitive and accurate even for the amplification of low quality DNA samples (e.g., high amount of inhibitors and/or untargeted DNA; Hindson et al., 2011; Taylor et al., 2017), we predict that the ddPCR will be increasingly used for detecting eDNA in the next few years.

The financial aspect of these protocols is also an important consideration for the development of large-scale monitoring tools. To this end, we investigated and compared the prices and time cost between the three approaches for monitoring *B. truncatus* presented in this paper (i.e., qPCR, ddPCR and visual prospecting), following the same method as in (Sengupta et al., 2019). The estimated prices for each site included the reusable materials (e.g., forceps, DNA primers, and probe), meaning that these prices are applicable for a first-time application of the protocol but should be devaluated in subsequent applications (Table S1). As expected, traditional visual survey remains the most economic approach (i.e., ≈20 EUR for all prospected sites; 12 hr/8 sites; Table S1). However, this approach is not adapted for large-scale monitoring as they require extensive workforce proportional to the number of prospected sites and huge malacological knowledge. Regarding the two eDNA-based methods developed in

this study, we estimated that the price between the two PCR technologies differed by 5 EUR (i.e., ≈60 EUR/sites; 6 hr/5 L × 8 sites for qPCR and ≈65 EUR/sites; 5h20/5 L × 8 sites for ddPCR; Table S1). Importantly, however, the estimation for ddPCR included service delivery. When estimating the price of ddPCR if realized in our laboratory facilities, prices are expected to be similar or even less expensive than qPCR analysis in a recurrent use framework. Expenditures related to ddPCR apparatus acquisition, which is much more expensive than those for qPCR, could be the only limiting aspect of this technology (Doi, Uchii, et al., 2015). However, this expense is compensated by the ability to obtain an absolute quantification from a single reaction compared with qPCR. Hence as denoted in the previous studies, ddPCR technology would be an interesting alternative to qPCR in quantifying eDNA from environmental samples as this enables absolute quantification from a single sample and due to the inhibitors dilutions during the sample singulation step (Doi, Uchii, et al., 2015).

In a schistosomiasis prevention perspective, these two eDNA-based approaches constitute promising tools for assessing the current distribution of *B. truncatus* from water samples collected in the field, and hence to identify new environments where schistosomiasis could emerge within and outside its endemic zone. In this regard, tropical schistosomes were recently found to be preadapted to temperate temperatures (Mulero, Rey, Arancibia, Mas-Coma, & Boissier, 2019), meaning that the spread of urogenital schistosomiasis northward mainly relies on the presence of locally adapted and compatible *B. truncatus* populations as those established in Corsica (Doby, Rault, Deblock, & Chabaud, 1966; Ramalli et al., 2018) in Spain and Portugal (Martínez-Ortí, Vilavella, Bargues, & Mas-Coma, 2019). In this context, we believe that these developed tools will help better assessing the risk of schistosomiasis emergence worldwide.

Importantly however, other factors need to be accounted to establish efficient risk maps of emergence and spread of schistosomiasis at large scale, which include human population movements,




FIGURE 4 Representation of the estimated *B. truncatus* DNA copy per liters of filtered water estimated using ddPCR as a function of *B. truncatus* densities at the sampling sites. *** represent differences with significant levels below 0.001

the spatial fluctuation of schistosomes' intermediate hosts under current climate changes, the invasion ability of African schistosomes species, the effect of hybridization between schistosome species on the invasion ability of the parasite, and finally the susceptibility of other European freshwater snails' species to *Schistosoma* parasites (Kincaid-Smith et al., 2017). To this end, the combination of eDNA-based tools with spatial modeling approaches such as ecological niche modeling (ENM) would improve our ability to generate "real-time" risk maps of schistosomiasis in Europe and endemic countries (Kincaid-Smith et al., 2017). ENM approaches were already used for spatial monitoring of freshwater snail species involved in SBD transmission such as fascioliasis and schistosomiasis (Cordellier & Pfenninger, 2009; Pedersen et al., 2014). Therefore, the use of ENM modeling approaches coupled with eDNA-diagnosis tools appears to be a next development step for SBD risk assessment endeavors.

ACKNOWLEDGMENTS

This study was funded by the French Agency for Food, Environmental and Occupational Health & Safety (PNREST 2019/1/059 Molrisk), the ANR (Program HySWARM ANR-18-CE35-0001), and the Occitania region (MOLRISK). This study is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR-10-LABX-41). Thank you to Franz Durandet and Maxime Juvin from Genome Editing engineering and analytics (I.A.G.E) who carried out the ddPCR experiments and subsequent first step analysis. We also wish to thank Sebastien Calvignac-Spencer and another anonymous referee for their help in the revision of previous versions of the manuscript. SM is also grateful to Didier Mulero for his help during field work and also the Occitania region for its involvement in this project.

CONFLICT OF INTEREST

None declared.

AUTHORS CONTRIBUTIONS

OR, JB, and SM conceived the original idea. SM, YQ, JF, and JFA carried out the field sampling. SM carried out the qPCR experiment, I.A.G.E. carried out the ddPCR experiment and first analyses. SM, OR, and JB analyzed and interpreted the data. JPP provided malacological expertise. SM and OR drafted the manuscript. OR, SM, JB, YQ, JF, JFA, and JPP revised the manuscript. All authors read and approved the final version of the manuscript.

ORCID

Stephen Mulero  <https://orcid.org/0000-0002-0853-0730>

Jérôme Boissier  <https://orcid.org/0000-0002-0793-3108>

Yann Quilichini  <https://orcid.org/0000-0003-0739-0155>

Olivier Rey  <https://orcid.org/0000-0003-4462-3712>

DATA AVAILABILITY STATEMENT

The dataset supporting the conclusions of this article is included within the article.

REFERENCES

- Agersnap, S., Larsen, W. B., Knudsen, S. W., Strand, D., Thomsen, P. F., Hesselsoe, M., ... Møller, P. R. (2017). Monitoring of noble, signal and narrow-clawed crayfish using environmental DNA from freshwater samples. *PLoS One*, *12*(6), e0179261. <https://doi.org/10.1371/journal.pone.0179261>
- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, *19*(6), 716–723. <https://doi.org/10.1109/tac.1974.1100705>
- Amarir, F., Sebti, F., Abbasi, I., Sadak, A., Fellah, H., Nhammi, H., ... Rhajaoui, M. (2014). *Schistosoma haematobium* detection in snails by Dral PCR and Sh110/Sm-SI PCR: Further evidence of the interruption of schistosomiasis transmission in Morocco. *Parasit Vectors*, *7*, 288–288. <https://doi.org/10.1186/1756-3305-7-288>
- Baker, C. S., Steel, D., Nieu Kirk, S., & Klinck, H. (2018). Environmental DNA (eDNA) from the wake of the whales: Droplet digital PCR for detection and species identification. *Frontiers in Marine Science*, *5*, 1–11. <https://doi.org/10.3389/fmars.2018.00133>
- Bálint, M., Nowak, C., Márton, O., Pauls, S. U., Wittwer, C., Aramayo, J. L., ... Jansen, M. (2018). Accuracy, limitations and cost efficiency of eDNA-based community survey in tropical frogs. *Molecular Ecology Resources*, *18*(6), 1415–1426. <https://doi.org/10.1111/1755-0998.12934>
- Bass, D., Stentiford, G. D., Littlewood, D. T. J., & Hartikainen, H. (2015). Diverse applications of environmental DNA methods in parasitology. *Trends in Parasitology*, *31*(10), 499–513. <https://doi.org/10.1016/j.pt.2015.06.013>
- Bergquist, R., Zhou, X. N., Rollinson, D., Reinhard-Rupp, J., & Klohe, K. (2017). Elimination of schistosomiasis: The tools required. *Infectious Diseases of Poverty*, *6*(1), 158. <https://doi.org/10.1186/s40249-017-0370-7>
- Blum, A. J., & Hotez, P. J. (2018). Global "worming": Climate change and its projected general impact on human helminth infections. *PLoS Neglected Tropical Diseases*, *12*(7), e0006370. <https://doi.org/10.1371/journal.pntd.0006370>
- Bohmann, K., Evans, A., Gilbert, M. T. P., Carvalho, G. R., Creer, S., Knapp, M., ... de Bruyn, M. (2014). Environmental DNA for wildlife biology and biodiversity monitoring. *Trends in Ecology & Evolution*, *29*(6), 358–367. <https://doi.org/10.1016/j.tree.2014.04.003>
- Boissier, J., Grech-Angelini, S., Webster, B. L., Allienne, J.-F., Huysse, T., Mas-Coma, S., ... Mita, G. (2016). Outbreak of urogenital schistosomiasis in Corsica (France): An epidemiological case study. *The Lancet Infectious Diseases*, *16*(8), 971–979. [https://doi.org/10.1016/S1473-3099\(16\)00175-4](https://doi.org/10.1016/S1473-3099(16)00175-4)
- Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., & Coissac, E. (2016). obitools: A unix-inspired software package for DNA metabarcoding. *Molecular Ecology Resources*, *16*(1), 176–182. <https://doi.org/10.1111/1755-0998.12428>
- Burnham, K., & Anderson, D. (2003). *Model selection and multimodel inference: A practical information-theoretic approach* (2nd ed., Vol. 1, p. 488). New-York, NY: Springer Science & Business Media.
- Buxton, A. S., Groombridge, J. J., & Griffiths, R. A. (2017). Is the detection of aquatic environmental DNA influenced by substrate type? *PLoS One*, *12*(8), e0183371. <https://doi.org/10.1371/journal.pone.0183371>
- Calata, F. I. C., Caranguian, C. Z., Mendoza, J. E. M., Fornillos, R. J. C., Tabios, I. K. B., Fontanilla, I. K. C., ... Sato, M. O. (2019). Analysis of environmental DNA and edaphic factors for the detection of the snail intermediate host *Oncomelania hupensis* quadrasi. *Pathogens*, *8*(4), 1–24. <https://doi.org/10.3390/pathogens8040160>
- Capinha, C., Essl, F., Seebens, H., Moser, D., & Pereira, H. M. (2015). The dispersal of alien species redefines biogeography in the Anthropocene. *Science*, *348*(6240), 1248–1251. <https://doi.org/10.1126/science.aaa8913>

- Coble, A. A., Flinders, C. A., Homyack, J. A., Penaluna, B. E., Cronn, R. C., & Weitemier, K. (2019). eDNA as a tool for identifying freshwater species in sustainable forestry: A critical review and potential future applications. *Science of the Total Environment*, *649*, 1157–1170. <https://doi.org/10.1016/j.scitotenv.2018.08.370>
- Cohen, A. J., Brauer, M., Burnett, R., Anderson, H. R., Frostad, J., Estep, K., ... Forouzanfar, M. H. (2017). Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: An analysis of data from the Global Burden of Diseases Study 2015. *The Lancet*, *389*(10082), 1907–1918. [https://doi.org/10.1016/s0140-6736\(17\)30505-6](https://doi.org/10.1016/s0140-6736(17)30505-6)
- Collen, B., Whitton, F., Dyer, E. E., Baillie, J. E. M., Cumberlidge, N., Darwall, W. R. T., ... Böhm, M. (2014). Global patterns of freshwater species diversity, threat and endemism. *Global Ecology and Biogeography*, *23*(1), 40–51. <https://doi.org/10.1111/geb.12096>
- Cordellier, M., & Pfenninger, M. (2009). Inferring the past to predict the future: Climate modelling predictions and phylogeography for the freshwater gastropod *Radix balthica* (Pulmonata, Basommatophora). *Molecular Ecology*, *18*(3), 534–544. <https://doi.org/10.1111/j.1365-294X.2008.04042.x>
- Dickie, I. A., Boyer, S., Buckley, H. L., Duncan, R. P., Gardner, P. P., Hogg, I. D., ... Weaver, L. (2018). Towards robust and repeatable sampling methods in eDNA-based studies. *Molecular Ecology Resources*, *18*(5), 940–952. <https://doi.org/10.1111/1755-0998.12907>
- Doby, J. M., Rault, B., Deblock, S., & Chabaud, A. (1966). Snails and bilharziasis in Corsica. Distribution, frequency and biology of "*Bulinus truncatus*". *Annales De Parasitologie Humaine Et Comparée*, *41*(4), 337–349.
- Doi, H., Takahara, T., Minamoto, T., Matsuhashi, S., Uchii, K., & Yamanaka, H. (2015). Droplet digital polymerase chain reaction (PCR) outperforms real-time PCR in the detection of environmental DNA from an invasive fish species. *Environmental Science & Technology*, *49*(9), 5601–5608. <https://doi.org/10.1021/acs.est.5b00253>
- Doi, H., Uchii, K., Takahara, T., Matsuhashi, S., Yamanaka, H., & Minamoto, T. (2015). Use of droplet digital PCR for estimation of fish abundance and biomass in environmental DNA surveys. *PLoS One*, *10*(3), e0122763. <https://doi.org/10.1371/journal.pone.0122763>
- ECDC (2011). VBORNET, European Network for Arthropod Vector Surveillance for Human Public Health. VBORNET. Retrieved from <http://www.vbornet.eu/>
- ECDC (2014). Guidelines for the surveillance of native mosquitoes in Europe. ECDC. <https://doi.org/10.2900/37227>
- Ellwanger, J. H., Kaminski, V. L., & Chies, J. A. B. (2019). Emerging infectious disease prevention: Where should we invest our resources and efforts? *Journal of Infection and Public Health*, *12*(3), 313–316. <https://doi.org/10.1016/j.jiph.2019.03.010>
- Goldberg, C. S., Turner, C. R., Deiner, K., Klymus, K. E., Thomsen, P. F., Murphy, M. A., ... Taberlet, P. (2016). Critical considerations for the application of environmental DNA methods to detect aquatic species. *Methods in Ecology and Evolution*, *7*(11), 1299–1307. <https://doi.org/10.1111/2041-210x.12595>
- Gozlan, R. E., St-Hilaire, S., Feist, S. W., Martin, P., & Kent, M. L. (2005). Biodiversity: Disease threat to European fish. *Nature*, *435*(7045), 1046. <https://doi.org/10.1038/4351046a>
- Hartikainen, H., Bass, D., Briscoe, A. G., Knipe, H., Green, A. J., & Okamura, B. (2016). Assessing myxozoan presence and diversity using environmental DNA. *International Journal for Parasitology*, *46*(12), 781–792. <https://doi.org/10.1016/j.ijpara.2016.07.006>
- Hashizume, H., Sato, M., Sato, M. O., Ikeda, S., Yoonuan, T., Sanguankiat, S., ... Minamoto, T. (2017). Application of environmental DNA analysis for the detection of *Opisthorchis viverrini* DNA in water samples. *Acta Tropica*, *169*, 1–7. <https://doi.org/10.1016/j.actatropica.2017.01.008>
- Hebert, P. D., Cywinska, A., Ball, S. L., & deWaard, J. R. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *270*(1512), 313–321. <https://doi.org/10.1098/rspb.2002.2218>
- Herricks, J. R., Hotez, P. J., Wanga, V., Coffeng, L. E., Haagsma, J. A., Basañez, M.-G., ... Murray, C. J. L. (2017). The global burden of disease study 2013: What does it mean for the NTDs? *PLoS Neglected Tropical Diseases*, *11*(8), e0005424. <https://doi.org/10.1371/journal.pntd.0005424>
- Hindson, B. J., Ness, K. D., Masquelier, D. A., Belgrader, P., Heredia, N. J., Makarewicz, A. J., ... Colston, B. W. (2011). High-throughput droplet digital PCR system for absolute quantitation of DNA copy number. *Analytical Chemistry*, *83*(22), 8604–8610. <https://doi.org/10.1021/ac202028g>
- Hinlo, R., Gleeson, D., Lintermans, M., & Furlan, E. (2017). Methods to maximise recovery of environmental DNA from water samples. *PLoS One*, *12*(6), e0179251. <https://doi.org/10.1371/journal.pone.0179251>
- Holtfreter, M. C., Mone, H., Muller-Stover, I., Mouahid, G., & Richter, J. (2014). Schistosoma haematobium infections acquired in Corsica, France, August 2013. *Eurosurveillance*, *19*(22), 2–4. <https://doi.org/10.2807/1560-7917.es2014.19.22.20821>
- Hunter, M. E., Ferrante, J. A., Meigs-Friend, G., & Ulmer, A. (2019). Improving eDNA yield and inhibitor reduction through increased water volumes and multi-filter isolation techniques. *Scientific Reports*, *9*(1), 5259. <https://doi.org/10.1038/s41598-019-40977-w>
- Hunter, M. E., Meigs-Friend, G., Ferrante, J. A., Smith, B. J., & Hart, K. M. (2019). Efficacy of eDNA as an early detection indicator for Burmese pythons in the ARM Loxahatchee National Wildlife Refuge in the greater Everglades ecosystem. *Ecological Indicators*, *102*, 617–622. <https://doi.org/10.1016/j.ecolind.2019.02.058>
- Huver, J. R., Koprivnikar, J., Johnson, P. T., & Whyard, S. (2015). Development and application of an eDNA method to detect and quantify a pathogenic parasite in aquatic ecosystems. *Ecological Applications*, *25*(4), 991–1002. <https://doi.org/10.1890/14-1530.1>
- Jerde, C. L., Olds, B. P., Shogren, A. J., Andruszkiewicz, E. A., Mahon, A. R., Bolster, D., & Tank, J. L. (2016). The influence of stream bottom substrate on retention and transport of vertebrate environmental DNA. *Environmental Science & Technology*, *50*(16), 8770–8779. <https://doi.org/10.1021/acs.est.6b01761>
- Jones, K. E., Patel, N. G., Levy, M. A., Storeygard, A., Balk, D., Gittleman, J. L., & Daszak, P. (2008). Global trends in emerging infectious diseases. *Nature*, *451*(7181), 990–993. <https://doi.org/10.1038/nature06536>
- Jones, R. A., Brophy, P. M., Davis, C. N., Davies, T. E., Emberson, H., Rees Stevens, P., & Williams, H. W. (2018). Detection of *Galba truncatula*, *Fasciola hepatica* and *Calicophoron daubneyi* environmental DNA within water sources on pasture land, a future tool for fluke control? *Parasit Vectors*, *11*(1), 342. <https://doi.org/10.1186/s13071-018-2928-z>
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., ... Drummond, A. (2012). Geneious basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, *28*(12), 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Kim, K. H., Kabir, E., & Ara Jahan, S. (2014). A review of the consequences of global climate change on human health. *Journal of Environmental Science and Health. Part C, Environmental Carcinogenesis & Ecotoxicology Reviews*, *32*(3), 299–318. <https://doi.org/10.1080/10590501.2014.941279>
- Kincaid-Smith, J., Rey, O., Toulza, E., Berry, A., & Boissier, J. (2017). Emerging schistosomiasis in Europe: A need to quantify the risks. *Trends Parasitol*, *33*(8), 600–609. <https://doi.org/10.1016/j.pt.2017.04.009>
- Klobucar, S. L., Torrey, R. W., & Phaedra, B. (2017). At the forefront: Evidence of the applicability of using environmental DNA to quantify the abundance of fish populations in natural lentic waters with additional sampling considerations. *Canadian Journal of Fisheries and Aquatic Sciences*, *74*(12), 2030–2034. <https://doi.org/10.1139/cjfas-2017-0114>

- Lamy, T., Pointier, J. P., Jarne, P., & David, P. (2012). Testing meta-population dynamics using genetic, demographic and ecological data. *Molecular Ecology*, 21(6), 1394–1410. <https://doi.org/10.1111/j.1365-294X.2012.05478.x>
- Li, J., Lawson Handley, L. J., Read, D. S., & Hanfling, B. (2018). The effect of filtration method on the efficiency of environmental DNA capture and quantification via metabarcoding. *Molecular Ecology Resources*, 18(5), 1102–1114. <https://doi.org/10.1111/1755-0998.12899>
- Longmire, J. L., Maltbie, M., & Baker, R. J. (1997). *Use of "lysis buffer" in DNA isolation and its implication for museum collections* (Vol. 163, pp. 1–3). Museum of Texas Tech University. Occasional Papers.
- Martínez-Ortí, A., Vilavella, D., Bargues, M. D., & Mas-Coma, S. (2019). Risk map of transmission of urogenital schistosomiasis by *Bulinus truncatus* (Audouin, 1827) (Mollusca: Gastropoda, Bulinidae) in Spain and Portugal. *Animal Biodiversity and Conservation*, 42(2), 257–266. <https://doi.org/10.32800/abc.2019.42.0257>
- Morens, D. M., & Fauci, A. S. (2013). Emerging infectious diseases: Threats to human health and global stability. *PLoS Pathogens*, 9(7), e1003467. <https://doi.org/10.1371/journal.ppat.1003467>
- Mulero, S., Rey, O., Arancibia, N., Mas-Coma, S., & Boissier, J. (2019). Persistent establishment of a tropical disease in Europe: The pre-adaptation of schistosomes to overwinter. *Parasites & Vectors*, 12(1), 379. <https://doi.org/10.1186/s13071-019-3635-0>
- Nathan, L. M., Simmons, M., Wegleitner, B. J., Jerde, C. L., & Mahon, A. R. (2014). Quantifying environmental DNA signals for aquatic invasive species across multiple detection platforms. *Environmental Science & Technology*, 48(21), 12800–12806. <https://doi.org/10.1021/es5034052>
- Noël, H., Ruello, M., Maccary, A., Pelat, C., Sommen, C., Boissier, J., ... Berry, A. (2017). Large outbreak of urogenital schistosomiasis acquired in Southern Corsica, France: Monitoring early signs of endemicization? *Clinical Microbiology and Infection*, 24(3), 295–300. <https://doi.org/10.1016/j.cmi.2017.06.026>
- Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421(6918), 37–42. <https://doi.org/10.1038/nature01286>
- Pedersen, U. B., Stendel, M., Midzi, N., Mduluzi, T., Soko, W., Stensgaard, A.-S., ... Kristensen, T. K. (2014). Modelling climate change impact on the spatial distribution of fresh water snails hosting trematodes in Zimbabwe. *Parasit Vectors*, 7, 536. <https://doi.org/10.1186/s13071-014-0536-0>
- Pereira, H. M., Leadley, P. W., Proença, V., Alkemade, R., Scharlemann, J. P. W., Fernandez-Manjarrés, J. F., ... Walpole, M. (2010). Scenarios for global biodiversity in the 21st century. *Science*, 330(6010), 1496–1501. <https://doi.org/10.1126/science.1196624>
- Puth, L. M., & Post, D. M. (2005). Studying invasion: Have we missed the boat? *Ecology Letters*, 8(7), 715–721. <https://doi.org/10.1111/j.1461-0248.2005.00774.x>
- Ramalli, L., Mulero, S., Noel, H., Chiappini, J. D., Vincent, J., Barre-Cardi, H., ... Berry, A. (2018). Persistence of schistosomal transmission linked to the Cavu river in southern Corsica since 2013. *Eurosurveillance*, 23(4), 2-5. <https://doi.org/10.2807/1560-7917.ES.2018.23.4.18-00017>
- Rathinasamy, V., Hosking, C., Tran, L., Kelley, J., Williamson, G., Swan, J., ... Spithill, T. W. (2018). Development of a multiplex quantitative PCR assay for detection and quantification of DNA from *Fasciola hepatica* and the intermediate snail host, *Austropeplea tomentosa*, in water samples. *Veterinary Parasitology*, 259, 17–24. <https://doi.org/10.1016/j.vetpar.2018.06.018>
- Ratnasingham, S., & Hebert, P. D. (2007). bold: The barcode of life data system (<http://www.barcodinglife.org>). *Molecular Ecology Notes*, 7(3), 355–364. <https://doi.org/10.1111/j.1471-8286.2007.01678.x>
- Richards, S. L., Lord, C. C., Pesko, K., & Tabachnick, W. J. (2009). Environmental and biological factors influencing *Culex pipiens quinquefasciatus* Say (Diptera: Culicidae) vector competence for Saint Louis encephalitis virus. *The American Journal of Tropical Medicine and Hygiene*, 81(2), 264–272. <https://doi.org/10.4269/ajtmh.2009.81.264>
- Richey, C. A., Kenelty, K. V., Van Stone Hopkins, K., Stevens, B. N., Martínez-Lopez, B., Barnum, S. M., ... Soto, E. (2018). Distribution and prevalence of *Myxobolus cerebralis* in postfire areas of Plumas National Forest: Utility of environmental DNA sampling. *J Aquat Anim Health*, 30(2), 130–143. <https://doi.org/10.1002/aah.10014>
- RStudio Team (2016). *RStudio: Integrated development for R*. RStudio, Inc. Retrieved from <http://www.rstudio.com>
- Rusch, J. C., Hansen, H., Strand, D. A., Markussen, T., Hytterod, S., & Vralstad, T. (2018). Catching the fish with the worm: A case study on eDNA detection of the monogenean parasite *Gyrodactylus salaris* and two of its hosts, Atlantic salmon (*Salmo salar*) and rainbow trout (*Oncorhynchus mykiss*). *Parasit Vectors*, 11(1), 333. <https://doi.org/10.1186/s13071-018-2916-3>
- Schaffner, F., Bellini, R., Petric, D., Scholte, E. J., Zeller, H., & Rakotoarivony, L. M. (2013). Development of guidelines for the surveillance of invasive mosquitoes in Europe. *Parasit Vectors*, 6, 209. <https://doi.org/10.1186/1756-3305-6-209>
- Schneider, J., Valentini, A., Dejean, T., Montarsi, F., Taberlet, P., Glaiot, O., & Fumagalli, L. (2016). Detection of invasive mosquito vectors using environmental DNA (eDNA) from water samples. *PLoS One*, 11(9), e0162493. <https://doi.org/10.1371/journal.pone.0162493>
- Sengupta, M. E., Hellström, M., Kariuki, H. C., Olsen, A., Thomsen, P. F., Mejer, H., ... Vennervald, B. J. (2019). Environmental DNA for improved detection and environmental surveillance of schistosomiasis. *Proceedings of the National Academy of Sciences of the United States of America*, 116(18), 8931–8940. <https://doi.org/10.1073/pnas.1815046116>
- Stensgaard, A. S., Vounatsou, P., Sengupta, M. E., & Utzinger, J. (2019). Schistosomes, snails and climate change: Current trends and future expectations. *Acta Tropica*, 190, 257–268. <https://doi.org/10.1016/j.actatropica.2018.09.013>
- Strickler, K. M., Fremier, A. K., & Goldberg, C. S. (2015). Quantifying effects of UV-B, temperature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation*, 183, 85–92. <https://doi.org/10.1016/j.biocon.2014.11.038>
- Taberlet, P., Coissac, E., Hajibabaei, M., & Rieseberg, L. H. (2012). Environmental DNA. *Molecular Ecology*, 21(8), 1789–1793. <https://doi.org/10.1111/j.1365-294X.2012.05542.x>
- Takahara, T., Minamoto, T., Yamanaka, H., Doi, H., & Kawabata, Z. (2012). Estimation of fish biomass using environmental DNA. *PLoS One*, 7(4), e35868. <https://doi.org/10.1371/journal.pone.0035868>
- Taylor, S. C., Laperriere, G., & Germain, H. (2017). Droplet Digital PCR versus qPCR for gene expression analysis with low abundant targets: From variable nonsense to publication quality data. *Sci Rep*, 7(1), 2409. <https://doi.org/10.1038/s41598-017-02217-x>
- Tsuji, S., Takahara, T., Doi, H., Shibata, N., & Yamanaka, H. (2019). The detection of aquatic macroorganisms using environmental DNA analysis—A review of methods for collection, extraction, and detection. *Environmental DNA*, 1(2), 1–10. <https://doi.org/10.1002/edn3.21>
- Uthicke, S., Lamare, M., & Doyle, J. R. (2018). eDNA detection of corallivorous seastar (*Acanthaster cf. solaris*) outbreaks on the Great Barrier Reef using digital droplet PCR. *Coral Reefs*, 37(4), 1229–1239. <https://doi.org/10.1007/s00338-018-1734-6>
- Wacker, S., Fossøy, F., Larsen, B. M., Brandsegg, H., Sivertsgård, R., & Karlsson, S. (2019). Downstream transport and seasonal variation in freshwater pearl mussel (*Margaritifera margaritifera*) eDNA concentration. *Environmental DNA*, 1(1), 64–73. <https://doi.org/10.1002/edn3.10>
- WHO (2002). *The partnership for parasite control, second meeting, Rome*. 20 February 2018. WHO. Retrieved from <http://www.who.int/news-room/fact-sheets/detail/schistosomiasis>
- WHO (2017). *Vector-borne diseases*. 31 October 2017. WHO, 1. Retrieved from <https://www.who.int/news-room/fact-sheets/detail/vector-borne-diseases>

- WHO (2018). *Schistosomiasis*. 27 December 2018. WHO. Retrieved from <https://www.who.int/schistosomiasis/en/>
- Wilcox, T. M., Zarn, K. E., Piggott, M. P., Young, M. K., McKelvey, K. S., & Schwartz, M. K. (2018). Capture enrichment of aquatic environmental DNA: A first proof of concept. *Molecular Ecology Resources*, 18(6), 1392–1401. <https://doi.org/10.1111/1755-0998.12928>
- Worrell, C., Xiao, N., Vidal, J. E., Chen, L., Zhong, B., & Remais, J. (2011). Field detection of *Schistosoma japonicum* cercariae in environmental water samples by quantitative PCR. *Applied and Environmental Microbiology*, 77(6), 2192–2195. <https://doi.org/10.1128/AEM.01561-10>
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., & Madden, T. L. (2012). Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*, 13, 134. <https://doi.org/10.1186/1471-2105-13-134>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Mulero S, Boissier J, Allienne J-F, et al. Environmental DNA for detecting *Bulinus truncatus*: A new environmental surveillance tool for schistosomiasis emergence risk assessment. *Environmental DNA*. 2020;2: 161–174. <https://doi.org/10.1002/edn3.53>