

Observation of a mesoscale eddy dipole on the northern Madagascar ridge: Consequences for the circulation and hydrography in the vicinity of a seamount

Vianello Patrick ¹, Herbette Steven ^{2,*}, Ternon Jean-Francois ³, Demarcq Herve ³, Roberts Michael J. ^{1,4}

¹ SA-UK Bilateral Chair: Ocean Science and Marine Food Security, Nelson Mandela University, Port Elizabeth, South Africa

² Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, Univ. Brest - CNRS - IRD - Ifremer, Brest, France

³ MARBEC, IRD, Univ Montpellier, CNRS, Ifremer, Sète, France

⁴ National Oceanography Centre, Southampton, United Kingdom

* Corresponding author : Steven Herbette, email address : steven.herbette@univ-brest.fr

Abstract :

Based on satellite and in situ data, the dynamic characteristics and vertical structure of a surface intensified mesoscale dipole recently expelled from the South East Madagascar Current (SEMC) is described for the first time. The dipole was surveyed 250 nautical miles south of Madagascar between 14 and 23 November 2016, during west-east and south-north transects carried out over the northern Madagascar Ridge. The dipole consisted of two counter-rotating vortices of similar size (100 km) and intensity (0.7 f), and an intense southwestward jet (150 cm s⁻¹) in the frontal region between the two eddies. The cyclonic eddy was lying on the western side of the anticyclonic eddy. With azimuthal velocities reaching 100 cm s⁻¹ at the surface and decreasing slowly with depth (40 cm s⁻¹ at -600 m), this MAD-Ridge dipole was defined as a highly non-linear ($Ro \sim 0.7$) isolated eddy-type structure ($c\beta \sim 11$ cm s⁻¹ and $U/c\beta \sim 0.7$) capable of trapping and advecting water masses over large distances. The enhanced concentration of chlorophyll-a found in the cyclone relative to the anticyclone could be tracked back to the spin-up phase of the two eddies and attributed to eddy-pumping. The eddy cores were located above the pycnocline (1026.4 kg m⁻³), within the upper 600 m, and consisted of varieties of Subtropical Underwater (STUW) found within the SEMC. The STUW found in the anticyclone was more saline and oxygenated than in the cyclone, highlighting mixing with the inshore shelf waters from the southeastern coastal upwelling cell off Madagascar. Observations suggest that the dipole interacted strongly with the chaotic bathymetry of the region, characterized by a group of five seamounts lying between -240 m and -1200 m. The bathymetry blocked its westward advection, trapping it in the vicinity of the seamount for more than 4 weeks, so enhancing the role of the eddy-induced velocities in stirring the surrounding water masses. Squeezed between the southern Madagascar shelf and the northern flank of the anticyclone, two filament-like dynamic features with very different water-mass properties could be observed on the south-north transect: i) one filament highly concentrated in chlorophyll-a demonstrating the capacity of the eddy to export shelf water offshore; ii) intrusions of a more southern-type of STUW generally found south of the South Indian Counter Current (SICC) recirculating on the external flanks of the anticyclone. Although the observed circulation and hydrography were largely constrained by the presence of the mesoscale eddy

dipole, unmistakable fine-scale dynamics were also observed in the vicinity of the MAD-Ridge seamount, superimposed onto the mesoscale eddy flow.

49 **1. Introduction**

50 *1.1 Context*

51 Seamounds are ubiquitous in the World's Oceans. In recent years, advances in satellite
52 altimetry have allowed >14 000 seamounds with a vertical extension exceeding 1000 m to be
53 spotted (Kitchingman et al., 2007). According to Lavelle and Mohn (2010), >100 000 tall
54 seamounds still remain uncharted as a consequence of limitations in the resolution of satellite
55 altimetry. The proportion of seamounds that have had their environment monitored is also
56 extremely low, even though seamounds are known to play crucial roles in structuring the
57 ecology of the oceans, and more recently, for their vulnerability to human exploitation (Clark
58 et al., 2010; Schlacher et al., 2010). Seamounds are often seen as key habitats for marine life,
59 even more so in oligotrophic waters where they are considered as hotspots for life and
60 biodiversity (Genin and Boehlert, 1985; Dower et al., 1992; Rogers, 1994; Mouriño et al.,
61 2001). In addition, many are located in Areas Beyond National Jurisdiction (ABNJ) where
62 there is little regulation, leaving them targeted by industrial fisheries (Marsac et al., 2020) and
63 sometimes resulting in the total collapse of the fishery (Koslow, 1997; Clark, 2001; Pitcher et
64 al., 2010).

65 There seems to be general consensus in the literature that the associated with seamounds
66 are linked tightly to the dynamics of oceanic circulation. At least two of today's general
67 concepts arose in the literature of the 1980s and 1990s, when interest in seamount biology
68 started to rise. The first states that there is increased primary productivity and chlorophyll-*a*

69 (hereafter chl-*a*) around seamounts because of the enhanced vertical flux of nutrients towards
70 the euphotic layer. The second is that currents around seamounts favour the retention of
71 organic matter and organisms, which contributes to the specificity of the ecosystems, isolated
72 from the surrounding environment, and sheltering a restricted and unique biodiversity (Genin
73 and Boehlert, 1985; Dower et al., 1992; Boehlert and Mundy, 1993; Comeau et al., 1995;
74 Mouriño et al., 2001; Genin, 2004). However, there has been little tangible evidence to sustain
75 these concepts (Genin and Dower, 2007; Rowden et al., 2010).

76

77 *1.2 Influence of seamounts on the ocean circulation*

78 Theoretical and idealized modelling has helped in understanding the physical processes at
79 play when a tidal or non-tidal flow encounters an isolated seamount. For instance, Garrett
80 (2003) found that steep seamounts, located in areas of strong tidal flow, act as stirring rods for
81 the ocean where the energy from lunar and solar barotropic tides is converted into an internal
82 wave field, commonly referred to as the internal (baroclinic) tide. Internal waves then
83 propagate into the ocean interior inducing motions of the density surfaces (isopycnals).
84 Whether this internal wave field breaks locally or far away, such dissipation generates sites of
85 intense vertical turbulent mixing that contribute to the local stratification and nutrient
86 enrichment of surface layers. Non-tidal flows impinging on a seamount may also generate
87 internal waves, commonly referred to as Lee waves. The latter will also either dissipate
88 locally or radiate away depending on the characteristics of the flow and the topography
89 (Nikurashin and Ferrari, 2010). When the non-tidal flow is characterized by low Rossby
90 numbers, it will deviate anticyclonically around the seamount. In some conditions, this
91 anticyclonic flow may even remain trapped over an obstacle, constituting a feature known as a
92 Taylor cap (or Taylor column; Huppert, 1975). The formation of a Taylor column is also
93 accompanied by the detachment of a cyclonic eddy that may remain trapped in the vicinity of
94 the obstacle or advected away (Huppert and Bryan, 1976; Royer, 1978; Verron and Le
95 Provost, 1985; Herbette et al., 2003). Oceanic currents induced by intense mesoscale eddies
96 enter this category of non-tidal flows. Taylor caps can produce large uplift of the interior
97 isopycnals (Dower and Mackas, 1996), which could enhance phytoplankton blooms. They can
98 also be long-life features facilitating the retention of particles and biota near the seamount
99 summit (Mullineaux and Mills, 1997).

100

101 *1.3 The northern Madagascar Ridge*

102 In the South West Indian Ocean (SWIO), the 1500-km long Madagascar Ridge is an
103 elongated aseismic plateau that extends from the tip of the southern Madagascar shelf all the
104 way down to 35°S (Sinha et al., 1981). It separates the Mozambique and Madagascar basins,
105 two ocean basins of mid- to late Cretaceous age, and typically rises from abyssal depths (-
106 5000 m) to between 1500 and 2500 m of the sea surface (Fig. 1). On its southern portion (at
107 33°12'S), the Walters Shoal, a seamount almost reaching the sea surface, is its most
108 prominent feature. On its northern portion, just south of the Madagascan shelf, the ridge
109 widens and becomes a rough plateau composed of at least five seamounts, which have never
110 been monitored.

111 The northern Madagascar Ridge is a productive region of highly complex and turbulent
112 dynamics. It is influenced by cold filaments, highly concentrated in chl-*a*, detaching from the
113 adjacent southern Madagascar coastal upwelling cells (Lutjeharms and Machu, 2000; Quartly,
114 2006; Quartly et al., 2006; Ramanantsoa et al., 2018; Demarcq et al., 2020). It is also located
115 in the very energetic retroflection region of the South East Madagascar Current (SEMC)
116 (Pous et al., 2014; Vianello et al., 2020), which flows south along the east coast of
117 Madagascar, transporting around 35 Sv¹ of warm, saline water from the subtropical South
118 Indian Ocean (Siedler et al., 2009). It originates from the bifurcation, at 20°S, of the Indian
119 Ocean South Equatorial Current (SEC) (DiMarco et al., 2002) and forms the northern part of
120 the western boundary current of the South Indian Ocean subtropical gyre. At the southern tip
121 of Madagascar, the dynamics of the SEMC becomes highly complex with three possible
122 modes (Quartly et al., 2006; Ramanantsoa et al., 2020): i) an early retroflection mode in
123 which the SEMC veers eastwards several hundreds of kilometres north of the southern tip of
124 Madagascar (~23°S); ii) a canonical retroflection mode in which the SEMC overshoots the
125 southern tip of Madagascar, flowing south, before finally veering east; iii) a third mode in
126 which the SEMC continues to flow west following the southern Madagascan shelf edge. The
127 last two modes contribute to the formation of intense mesoscale eddies or dipoles that
128 propagate westwards over the Madagascar Ridge and towards the Agulhas Current (De
129 Ruijter et al., 2004; Nauw et al., 2008; Siedler et al., 2009; Halo et al. 2014). Ridderinkhof et
130 al. (2013) showed that, south of Madagascar, these mesoscale eddies often take the form of
131 large dipoles, among which some may remain strong enough to subsequently trigger an early
132 retroflection of the Agulhas Current.

133

¹ Sverdrup (Sv) = 10⁶ m³ s⁻¹

134 *1.4 Previous cruises on the northern Madagascar Ridge*

135 A few research cruises had previously taken place over the northern Madagascar Ridge. In
136 2001, the ACSEX hydrographic cruise (Agulhas Current Sources Experiment) consisted of
137 four transects perpendicular to the Madagascar shelf that captured the SEMC as well as an
138 anticyclonic and cyclonic eddy dipole (De Ruijter et al., 2004; Nauw et al., 2008). In 2005,
139 the Madagascar Experiment (MadEx) highlighted the presence of intensified currents at all
140 depths of the water column (Quartly, 2006; Quartly et al., 2006). In September 2008, eight
141 across-shore transects carried out on board the RV *Dr Fridtjof Nansen* along the eastern
142 Madagascan coast, emphasized the complex dynamics of the northern and southern branches
143 of the East Madagascar Current (Voldsund et al., 2017). One year later, the same vessel
144 returned to the area and surveyed the south and west coasts of Madagascar within the
145 framework of the Agulhas Somali Large Marine Ecosystem (ASCLME) programme (Pripp et
146 al., 2014). Evidence of coastal upwelling was found along the southeast coast of Madagascar
147 and at two sites on its west coast, Cap Saint André and Nosy Be Island (Alvheim et al., 2009).
148 In November/December 2009, a multidisciplinary cruise was conducted over a group of
149 seamounts of the SWIO, whose summits lay at depths situated between -100 m and -1250 m.
150 The objective of that programme was to gain knowledge on the pelagic ecosystems around the
151 seamounts and to determine the dominant physical processes at play (Read and Pollard,
152 2017). Five of the seamounts were located over the South West Indian Ridge and one over the
153 southern Madagascar Ridge, close to the Walters Shoal (Rogers, 2016). Results showed that
154 oceanic currents around the seamounts were linked to the internal wave field originating from
155 the tidal flow, or to the presence of mesoscale eddies. Although Taylor caps were detected at
156 a few locations during that survey (Pollard and Read, 2017), their existence was intermittent
157 and did not influence the observed circulation and hydrography.

158

159 *1.5 The MADRidge project*

160 In 2016/2017, an international programme (the MADRidge project) was designed to
161 monitor the ecosystems in the vicinity of shallow seamounts in the SWIO (Fig. 1). The
162 International Union for Conservation of Nature (IUCN) and the French Institut de Recherche
163 pour le Développement (IRD), together with partners in France, South Africa and
164 Madagascar, carried out three multidisciplinary research cruises that surveyed three shallow
165 seamounts lying in three very different dynamic environments (Roberts et al., 2020). The
166 study describes the physical (currents and hydrography) and biogeochemical (oxygen and
167 fluorescence) *in situ* data collected during the MAD-Ridge Leg 1 expedition that focused only

168 on one of the three seamounts: an unnamed seamount, thereafter named MAD-Ridge, located
169 on the northern Madagascar Ridge. The work here aims at providing knowledge of the
170 underlying dynamics within this very turbulent environment, focusing specifically on the role
171 of the bathymetry in constraining the circulation and hydrography. The *in situ* survey offered
172 an unique opportunity to characterize in detail a surface-intensified mesoscale eddy dipole
173 that had been freshly expelled from the SEMC and stayed trapped in the vicinity of the MAD-
174 Ridge seamount during the whole cruise. Our objectives are to describe in detail the synoptic
175 conditions in place during the MAD-Ridge Leg 1 cruise and to analyse how the conditions
176 may have influenced the nature of the flow-topography interactions and the environmental
177 response in term of chl-*a* concentration.

178

179 *1.6 Outline*

180 Section 2 below describes the MAD-Ridge Leg 1 cruise, focusing on the vertical profiling
181 of the physical (currents and hydrography) and biogeochemical (oxygen and fluorescence) *in*
182 *situ* data, and their subsequent validation and calibration. The satellite data (sea surface height
183 and chl-*a*) and the methods used to track mesoscale eddies in the region are also briefly
184 explained. Section 3 then highlights the presence during the survey of a surface-intensified
185 coherent mesoscale cyclonic/anticyclonic dipole expelled from the SEMC. The vertical
186 structure of the dipole is characterized in terms of velocities, water mass properties and its
187 impact on the vertical distribution of chl-*a* and nutrients. Fine-scale turbulent dynamic
188 features such as filaments, which superimpose onto the dominant flow induced by the
189 mesoscale eddy dipole, are described in Section 4. The observations are discussed in
190 Section 5 in the light of theoretical work on eddy-seamount interactions and recent progress in
191 the understanding of the variability of the SEMC. Finally, Section 6 summarizes our
192 observations and discusses the important role played by the northern Madagascar Ridge in
193 Global Ocean circulation by governing aspects of connectivity between the SEMC and the
194 Agulhas Current.

195

196 **2. Data and Methods**

197 *2.1 The MAD-Ridge cruise*

198 The MAD-Ridge Leg 1 cruise took place between 8 and 25 November 2016 (doi:
199 10.17600/16004800) on board the RV *Antea*. It focused on the MAD-Ridge seamount at
200 27°29'S, 46°16'E. The cruise consisted of two perpendicular transects of ~150 nautical miles
201 that crossed the summit of the seamount (Fig. 2). The west-east transect was carried out

202 during the period 14-18 November, and the south-north transect from 19 to 23 November.
203 Each station along the transects consisted of conductivity-temperature-depth (CTD) and
204 lowered acoustic Doppler current profiler (L-ADCP) vertical profiling down to -1000 m.
205 Stations were every 15 nautical miles outwards from the seamount, and at intervals reduced to
206 5 nautical miles over the slopes and summit of the seamount. The west-east transect (45°-
207 47°30'E, at 27°30'S) had 15 stations, the south-north transect (28°17'S - 25°40'S, at 46°15'E)
208 16 stations. The northernmost station of the meridional transect was located on the outer edge
209 of the southern Madagascan continental shelf on the -840 m isobath. Ship acoustic Doppler
210 current profiler (S-ADCP) measurements were collected along the whole cruise track.

211

212 *2.2 CTD and nutrients data*

213 *In situ* vertical profiles of temperature, salinity, dissolved oxygen and fluorescence were
214 collected using a Seabird SBE 911+ CTD-O₂ equipped with a Wetlabs ECO FL fluorometer.
215 The CTD-O₂ probe had two sensors for temperature, salinity and dissolved oxygen. The
216 vertical profiles were made from the surface to 1000 m. Seawater samples were collected at
217 different depths (up to 11 samples per cast) to calibrate the salinity (measured on board using
218 a Portasal salinometer and OSIL normal seawater), oxygen (measured on board using the
219 Winkler method) and fluorescence (filtration on board and phytoplankton pigment analysis at
220 the laboratory using High Pressure Liquid Chromatography) sensors. Nutrients (NO₂, NO₃,
221 PO₄ and Si(OH)₄) were determined by the classical colorimetric method (Oudot et al., 1998)
222 on samples collected at each station. CTD-O₂ calibration was performed using the
223 CADYHAC software from IFREMER (Kermabon et al., 2015). Conservative temperature and
224 absolute salinity were calculated according to the TEOS-10 equations, and the vertical
225 stretching term of the potential vorticity (PV) was derived as $|f|N^2/g$, with f the Coriolis
226 parameter, N the Brunt-Väisälä frequency and g the constant for gravity (Talley et al., 2011).

227

228 *2.3 In situ current measurements*

229 The RV *Antea* has a 75 kHz RDI Ocean Surveyor hull mounted S-ADCP, which allows for
230 continuous vertical profiling of the ocean currents along the ship's track. Velocity
231 components were time-averaged over 2 min. The vertical resolution (bin size) was set to 16
232 m, with a maximum measurement depth down to -600 m. The S-ADCP data were processed
233 using the CASCADE software from IFREMER (Le Bot et al., 2011). A tidal correction was
234 applied using the TPX08-atlas (Egbert et al., 2002). The final S-ADCP product consisted of a

235 2-km horizontal resolution set of vertical profiles for the zonal u and meridional v components
 236 of the velocity vector.

237 Two 300 kHz RDI Workhorse (upward and downward orientated) L-ADCPs were attached
 238 to the CTD frame to measure the zonal and meridional components of the velocity through the
 239 water column at each CTD station, with an 8 m vertical bin-size resolution. L-ADCP data
 240 were processed on board, then calibrated after the cruise using a software developed by IFM-
 241 GEOMAR/LDEO (Thurnherr, 2014). The L-ADCP failed at stations 2 and 3. Despite the
 242 coarser horizontal resolution of the L-ADCP sampling (~ 25 km for the L-ADCP vs. 2 km for
 243 the S-ADCP), the vertical structure of the velocity fields given by the S-ADCP and L-ADCP
 244 were similar (Supplementary material Fig. S1). Hence, the S-ADCP data are used in the
 245 analysis.

246

247 *2.4 Altimetry data*

248 Daily interpolated merged delayed time altimetry data gridded at $1/4^\circ$ resolution, produced
 249 by Ssalto/Duacs and distributed by the Copernicus Marine Environment Monitoring Service
 250 (CMEMS, <http://marine.copernicus.eu/>) were used to describe the surface mesoscale synoptic
 251 conditions over the northern Madagascar ridge. Mean Eddy Kinetic Energy (EKE) was
 252 derived from Sea Level Anomaly (SLA) data over a large portion the SWIO (Fig. 1) as
 253 follows:

$$\overline{EKE} = \frac{1}{2} (\overline{u'_{gs}{}^2} + \overline{v'_{gs}{}^2}), \quad (1)$$

254 where u'_{gs} and v'_{gs} are the zonal and meridional components of the surface geostrophic
 255 current anomaly, and the $\bar{\cdot}$ stand for a linear time average operator from 1995 to 2015. Further,
 256 Absolute Dynamic Topography (MADT) data were used to compute the absolute surface
 257 geostrophic currents, relative vorticity and the Okubo-Weiss quantity² (Okubo, 1970; Weiss,
 258 1991).

259

260 *2.5 Ocean surface colour and chl-a satellite data*

261 Daily 4-km resolution MODIS ocean colour data provided by NASA
 262 (<https://oceancolor.gsfc.nasa.gov/>) were processed to produce composite 3-day images of the
 263 chl- a surface distribution.

² The Okubo-Weiss quantity λ_{ok} measures the local influences of the shear/strain rate against the relative vorticity. It is calculated by subtracting the relative vorticity $\zeta = (\partial_x v_{gs} - \partial_y u_{gs})^2$ from the deformation rate $\sigma = (\partial_x u_{gs} - \partial_y v_{gs})^2 + (\partial_x v_{gs} + \partial_y u_{gs})^2$: $\lambda_{ok} = \sigma - \zeta$.

264

265 *2.6 Eddy tracking and dipole occurrence*

266 Seven-day SLA products at $\frac{1}{4}^\circ$ resolution were used to monitor the long-term eddy activity
 267 in the region over the period 1993-2016 and to infer statistics on the presence of surface-
 268 intensified mesoscale eddy dipoles in the region. Eddies were tracked using the algorithm
 269 developed by Chelton et al. (2007). The method consists of finding SLA extrema sitting
 270 inside closed SLA contours (Chelton et al., 2007; Mason et al., 2011; Halo et al., 2014). Once
 271 an eddy is identified, the eddy centre coordinates are recorded. The method was further
 272 adapted to: i) discard weak eddies that have SLA extrema <10 cm in amplitude; ii) only retain
 273 eddies potentially interacting with the seamount – the typical eddy radius in the area being 90
 274 km [Halo et al., 2014], eddies found farther from the seamount summit were discarded; iii)
 275 distinguish single eddies from dipoles. Dipoles were diagnosed when a cyclone and an
 276 anticyclone could both be observed during the same 7-day period, <180 km from the
 277 seamount summit, and when the maximum velocity in the frontal region between the two
 278 eddies was at least $1.5\times$ the velocity found on the eddy periphery. For each dipole detected, a
 279 “dipole strength” (DS) was computed as an estimate of the gradient of SLA in the frontal
 280 region, subtracting the minimum SLA found within the cyclone (SLA_{\min}) from the maximum
 281 SLA found within the anticyclone (SLA_{\max}) and dividing the difference by the distance
 282 between the two eddy centres ($d_{c/ac}$): $DS = (SLA_{\max} - SLA_{\min}) / (d_{c/ac})$.

283

284 *2.7 Bathymetry*

285 The bathymetry of the MAD-Ridge seamount was surveyed on board using the two single-
 286 beam echo-sounders (12 and 38 kHz) mounted on the RV *Antea*. The echo-sounder
 287 measurements differed significantly from the ETOPO 1³ and GEBCO 30⁴ products based on
 288 satellite altimetry. The MAD-Ridge seamount summit was indeed found 6 km farther south
 289 than expected. In addition, although the seamount reached the sea surface in ETOPO 1 and
 290 GEBCO 30, it was found at -240 m during the cruise. The SRTM⁵ (Shuttle Radar Topography
 291 Mission) bathymetry product which showed the seamount at the correct position just 150 m
 292 below the sea surface, is used in the following for displaying the bathymetry of the area.

293 According to the *in situ* data, the seamount summit consists of a 20-km wide oval plateau,
 294 slightly elongated along a south-north axis, that plunges steeply from -240 m to the seafloor at

³ ETOPO 1 : doi:10.7289/V5C8276M

⁴ GEBCO 30: doi:10.5285/a29c5465-b138-234d-e053-6c86abc040b9)

⁵ SRTM: https://topex.ucsd.edu/WWW_html/srtm30_plus.html

295 -1600 m (Fig. 2). It should be stressed that the MAD-Ridge seamount is not an isolated
 296 structure; it is surrounded by four deeper summits situated between -600 m and -1200 m. The
 297 detailed topography of these neighbouring seamounts was not monitored during the cruise.

298

299 2.8 In situ geostrophic velocities

300 The components of the geostrophic velocity perpendicular to the west-east and south-north
 301 transects were calculated integrating vertically the thermal wind equation:

$$\frac{\partial v_{\perp}}{\partial p} = \frac{-1}{f} \frac{1}{\Delta x_{//}} \Delta \left(\frac{1}{\rho} \right). \quad (3)$$

302 v_{\perp} is the component of the geostrophic velocity perpendicular to each segment separating two
 303 CTD vertical profiles, p the pressure, f the Coriolis parameter, $\Delta x_{//}$ the segment length and
 304 $\Delta \left(\frac{1}{\rho} \right)$ the variation of specific volume over the segment. The right side of Eq. (3) was
 305 computed from the TEOS-10 Gibbs equation of state using conservative temperature and
 306 absolute salinity. Eq. (3) was then integrated vertically from a pressure of reference. This
 307 pressure of reference was calculated for each segment as the pressure at which the vertical
 308 shear of the S-ADCP velocity component perpendicular to the segment balanced the right-
 309 side term of Eq. (3). A horizontal low-pass Lanczos filter was applied to both the S-ADCP
 310 and temperature and salinity data prior to the integration, to remove spurious signal associated
 311 with non-geostrophic dynamics. The cut-off wave number was set to $1/20 \text{ km}^{-1}$, a value ~ 3
 312 times less than the Rossby radius of deformation found in the region (Chelton et al., 1998).
 313 Ageostrophic velocities were calculated by subtracting the calculated geostrophic velocities
 314 from the non-filtered S-ADCP velocity data.

315

316 3. Characteristics of a strong surface intensified mesoscale eddy dipole

317 A map of surface EKE, a proxy for mesoscale turbulence in the ocean, provides robust
 318 evidence that the MAD-Ridge seamount is located in a region characterized by a high level of
 319 turbulent mesoscale activity, with EKE values ranging between 630 and $800 \text{ cm}^2 \text{ s}^{-2}$ (Fig. 1).
 320 Although such levels of EKE are about $3\times$ less than those found in the most energetic western
 321 boundary current systems (Pilo et al., 2015), they are higher than in most parts of the ocean
 322 and suggest the presence of highly variable synoptic conditions.

323

324 3.1 Surface signature and coherence of the mesoscale eddy dipole

325 The cruise took place when a surface-intensified anticyclonic/cyclonic eddy dipole was
326 present over the northern Madagascar Ridge (Fig. 3). The anticyclone was centred over the
327 seamount, with its cyclonic counterpart lying on its western flank. Both eddies were roughly
328 the same size, with a radius of approximately 100 km, and of similar amplitude. Within the
329 anticyclone, the maximum values of SLA and surface relative vorticity were >35 cm (Fig. 3)
330 and of the order of $-0.7f$ ($f \sim -6.6 \times 10^{-5} \text{ s}^{-1}$) (Fig. 4). Similar but opposite values were found
331 within the cyclone, with minimum values of SLA below -35 cm and surface relative vorticity
332 of the order of $+0.7f$. The frontal region between the two eddies was characterized by intense
333 southwestward geostrophic jet-sustaining velocities $>150 \text{ cm s}^{-1}$ (Fig. 3), which highlights the
334 extreme intensity of the mesoscale dipole. Using an eddy-tracking algorithm, the vortices
335 could be traced back to 26 October 2016, coinciding with the time when the SEMC started to
336 subdivide into two branches as seen in the surface relative vorticity maps (Fig. 4). The west-
337 flowing branch followed the shelf edge, whereas the southwest-flowing branch detached itself
338 from the coast at 25°S , 47°E . The southward flow was observed until 27°S , where the SEMC
339 started to veer westwards. Cyclonic vorticity developed on the inshore side of the current, and
340 anticyclonic vorticity strengthened on the offshore side. On 2 November 2016, the dipole was
341 fully formed, although it was still embedded within the SEMC. While strengthening, it
342 detached itself from the SEMC and propagated southwest towards the MAD-Ridge seamount.
343 From 9 to 23 November 2016, the southwestward propagation of the dipole slowed, and the
344 dipole stayed in the vicinity of the seamount for two full weeks. By 30 November, the
345 cyclonic eddy had moved slightly southwest and the anticyclone had elongated notably in a
346 northeast-southwest direction. Another cyclonic eddy could be observed on the eastern flank
347 of the anticyclone. On 7 December 2016, the anticyclone split into two eddies. One stayed
348 trapped over the seamount, but the most intense one continued to form a dipole with the
349 original cyclone. The dipole then accelerated its southwestward propagation. It was tracked
350 until 24 December 2016 (not shown) when both eddies finally dissipated, and another similar
351 dipole began to interact with the MAD-Ridge seamount. Hence, the west-east and south-north
352 transects of Leg 1 provided a unique opportunity to survey a strong mesoscale eddy dipole
353 freshly expelled from the SEMC and interacting with the northern Madagascar Ridge.

354

355 *3.2 Vertical structure of the mesoscale eddy dipole: focus on azimuthal velocities*

356 The ship-mounted S-ADCP measurements provide additional information on the vertical
357 structure of the currents within the dipole. The west-east transect crossed the entire
358 anticyclonic eddy through its centre and captured the southeastern portion of the cyclone (Fig.

359 3). The agreement between the low-pass spatially filtered S-ADCP currents and the surface
360 geostrophic currents derived from altimetry shows that the dipole is accurately located by the
361 altimetry (Fig. 3), and lends confidence in the ability of the low-pass Lanczos filter to retrieve
362 the geostrophic part of the currents from the S-ADCP data. The vertical structure of the flow
363 confirms that the dipole was surface-intensified (Fig. 5). Down to -400 m, the mesoscale
364 circulation was in total accord with the existence of the mesoscale eddy dipole: clockwise and
365 anticlockwise circulations were observed within the cyclone and anticyclone, respectively.
366 The highest velocities were found in the southwestward jet that lay within the frontal region
367 between the two eddies (stations 5 and 6), with values above 150 cm s^{-1} at the surface and still
368 as high as 70 cm s^{-1} at -600 m. On the flanks of the dipole, velocities were slightly slower, but
369 still as high as 100 cm s^{-1} in the upper 100 m of the water column, and of the order of 70 cm s^{-1}
370 at -400 m (Fig. 5a, c, e).

371 Between the two transects, the anticyclone moved above the seamount while being
372 stretched along a southeast-northwest axis (Supplementary Fig. S2). Hence, the south-north
373 transect only captured one arc of the anticyclonic eddy in which the flow was mostly to the
374 southwest, with velocities of the order of $80\text{--}100 \text{ cm s}^{-1}$. Nevertheless, south of 27°S , the flow
375 veered anticlockwise to the east, confirming the anticyclonic rotation. Subsurface velocities
376 were also weaker, not exceeding 40 cm s^{-1} below -400 m. Geostrophic velocities computed
377 from the vertical profiling of density confirm this overall circulation pattern (Fig. 6).

378

379 *3.3 Vertical structure of the mesoscale eddy dipole: focus on the hydrography*

380 Additional characteristics on the vertical structure of the two eddies is provided by the *in*
381 *situ* temperature, salinity, oxygen and chl-*a* data collected during the two transects (Fig. 7 and
382 8). The vertical stretching term of the potential vorticity highlights the squeezing and
383 stretching of the isopycnals and provides extra information on the process of formation of a
384 water mass (Talley et al., 2011).

385 The doming of the isopycnals (black contours) in Fig. 7 and 8 allows accurate location of
386 the core of the anticyclone (X1) and the southern flank of the cyclone (X2). Considering the
387 average vertical density profile found in the MAD-Ridge region, the 1026.4 kg m^{-3} isopycnal
388 was hereafter selected as the pycnocline that separates the surface stratified waters from the
389 slightly deeper non-stratified waters ($N^2 < 10^{-4} \text{ s}^{-2}$). On the west-east transect, the curvature of
390 this pycnocline clearly showed the presence of a surface-intensified mesoscale eddy dipole.
391 This isopycnal depth is found at -400 m within the anticyclone, and at -300 m within the
392 cyclone (Fig. 7). The south-north transect only intersected the southwestern portion of the

393 anticyclone. At stations 17, 18 and 19 (south of the seamount), the depth of the pycnocline
394 was similar to that (-400 m) observed within the anticyclone on the west-east transect
395 (stations 11–15). The rise of the pycnocline north of the seamount and adjacent to the
396 southern Madagascan shelf edge is in accord with the intensified westward flow observed on
397 the northern flank of the anticyclone (Fig. 8).

398 The cores of these two eddies were located above the pycnocline. At such depths,
399 azimuthal velocities are at their highest (Fig. 5 and 6), and the water mass properties differed
400 substantially whether they belonged to the anticyclone (X1), cyclone (X2) or the frontal zone
401 between the two eddies (Fig. 7). Surface waters found within the cyclonic eddy were on
402 average 0.6°C cooler and 0.1 g kg^{-1} more saline than those within the anticyclonic eddy. This
403 difference is even more visible when considering the salinity maximum centred on the
404 1026.0 kg m^{-3} isopycnal: the salinity was 0.3 g kg^{-1} higher in the anticyclone (36 g kg^{-1} at
405 stations 13, 14 and 24) than in the cyclone.

406

407 *3.4 Water mass properties within the mesoscale eddy dipole*

408 Water mass properties can be investigated by plotting the CTD vertical profiles on two
409 diagrams, conservative temperature (CT) vs. absolute salinity (SA) (Fig. 9a, b) and
410 conservative temperature vs. dissolved oxygen (O_2) (Fig. 9c). Profiles are grouped into three
411 classes, depending on whether they were collected within the cyclonic eddy, within the
412 anticyclonic eddy or within the frontal zone in between the two eddies. This classification was
413 made using altimetry data.

414

415 *3.4.1 Below the pycnocline, within the depth range of Antarctic Intermediate Water (AAIW):*

416 Between -800 m and -1000 m, for waters heavier than 1027.0 kg m^{-3} , the signature of
417 AAIW is clearly visible (Fig. 9a, b), with a minimum in salinity falling below 34.6 g kg^{-1} and
418 a minimum in temperature $<10^{\circ}\text{C}$ (Emery and Meincke, 1986). These properties match
419 observations carried out within the Agulhas Current, confirming the widespread nature of this
420 water mass in the SWIO (Beal et al., 2006).

421 Within this depth range too, the isopycnals were still deflected, mirroring the surface-
422 intensified dipole. However, the fact that all datapoints reported on the CT/SA and CT/ O_2
423 diagrams for that depth range are superimposed, independent of their location in the dipole
424 (Fig. 9), is an indication that the isopycnal variations of temperature, salinity and oxygen were
425 weak. The low values of geostrophic velocities ($\sim 10\text{ cm s}^{-1}$) at those depths (Fig. 6) confirm

426 the belief that these water masses did not belong to the core of the eddies forming the
427 mesoscale eddy dipole, but rather were being entrained by the surface eddy cores.

428

429 *3.4.2 Below the pycnocline, within the depth range of South Indian Central Water (SICW):*

430 On both transects, just below the eddy core, between the 1026.4 kg m^{-3} and 1026.7 kg m^{-3}
431 isopycnals, South-East Indian Sub-Antarctic Mode water (SEISAMW) was identified on all
432 the vertical CT/SA-profiles (Hanawa and Talley, 2001). Within that depth range, we observed
433 no spatial variation of temperature, salinity or oxygen on any given isopycnal (Fig 9). The
434 characteristics in temperature ($10\text{--}15^\circ\text{C}$) and salinity ($34.7\text{--}35.3 \text{ g kg}^{-1}$) are similar to the
435 heavier range of central waters commonly found within the subtropical gyre of the SWIO
436 (Emery and Meincke, 1986; Sprintall and Tomczak, 1992; Beal et al., 2006). In addition, we
437 observed high oxygen concentration of $>200 \mu\text{mol kg}^{-1}$ (Fig. 9c) and low potential vorticity
438 values (Fig. 7d and 8d). This helps to identify more accurately this water mass as the
439 SEISAMW, a heavy variety of Sub-Antarctic Mode Water (SAMW), formed within the deep-
440 winter mixed layer of the South East Indian Ocean (Hanawa and Talley, 2001). New et al.
441 (2007) have identified SEISAMW over the Mascarene plateau on the southern side of the
442 SEC. These observations suggest that the SEMC transported this water mass, ensuring a
443 connection between the Mascarene plateau and the northern Madagascar Ridge.

444

445 *3.4.3 Above the pycnocline, within the depths range of South Tropical Underwater (STUW)*
446 *and Tropical Surface Water (TSW)*

447 The STUW, characterized by salinity $>35.5 \text{ g kg}^{-1}$ and a high potential vorticity of
448 $\sim 150 \times 10^{-11} \text{ s}^{-1}$ (Hanawa and Talley 2001; Nauw et al., 2006), can be seen in all the MAD-
449 Ridge CTD profiles above the pycnocline between the 1026.4 and 1024.8 kg m^{-3} isopycnals
450 (Fig. 7 and 8) and on the CT/SA diagram (Fig. 9). That water mass constituted the core of
451 both eddies forming the mesoscale eddy dipole. Nevertheless, there was some indication that
452 the anticyclonic eddy core (X1 in Fig. 7 and 8) contained less-altered STUW than anywhere
453 else. Indeed, extremely high values of salinity ($>36 \text{ g kg}^{-1}$) were observed on the west-east
454 transect at station 13, on 18 November, east of the seamount (X1 in Fig. 7), and then a few
455 days later on 21 November, at stations 23-24 just north of the seamount when the eddy had
456 moved onto the seamount summit (X1 in Fig 8).

457 We now attempt to backtrack these properties to the formation of the dipole within the
458 SEMC. As already mentioned, maps of surface relative vorticity suggest that the cyclone was
459 generated inshore of the SEMC, whereas the anticyclone was formed on its offshore side (Fig.

460 4). The similarity between the CT/SA profiles inside the cyclonic eddy and those observed on
461 the southern continental shelf of Madagascar during the ASCEX cruise (de Ruijter et al.,
462 2004) adds weight to this assumption. The observed difference in salinity between the two
463 eddies was attributable to an existing cross-shore gradient of salinity within the SEMC itself
464 that can be linked to the water mass properties of the SEC. The latter transports a mixture of
465 Tropical Surface Water (TSW) and Sub-Tropical Surface Water (STSW) west, right across
466 the Indian Ocean, the STSW being much more saline than the former, and on the southern
467 side of the SEC (New et al., 2005, 2007). The densest part of the STSW subducts under the
468 Tropical Front to form some kind of intra-thermocline waters commonly referred as Sub-
469 Tropical Underwater (STUW; O'Connor et al., 2002). This water mass is reported to be about
470 0.2 g kg^{-1} more saline than the TSW. When the SEC flows over the Mascarene plateau, water
471 masses are partially mixed, which smooths out the difference in salinity (New et al., 2007).
472 Nonetheless, water masses on the southern edge of the SEC remain more saline than that on
473 the northern edge. When the SEC splits into two branches, as it approaches Madagascar, its
474 southern part forms the southern branch of the EMC that flows south along the Madagascar
475 coast, known as the SEMC. It is made up of STSW and STUW, but the offshore waters are
476 more saline than the inshore ones. As the SEMC flows south along the eastern Madagascar
477 coast, the offshore entrainment of fresh Madagascar shelf water into the SEMC and its
478 subsequent mixing with the waters within the current reinforce the cross-shore salinity
479 gradient, agreeing with the water masses observed within the eddy cores.

480

481 *3.5 Impact of the mesoscale eddy dipole on chl-a and nutrient distribution*

482 Three-day composite maps of satellite sea surface chl-*a* concentration show enhanced
483 phytoplankton concentration within the cyclone (Fig. 10). This fits with the widely spread
484 paradigm that the uplift of isopycnals within cyclones brings more nutrients into the euphotic
485 layer, enhancing primary production (McGillicuddy et al., 1998; Oschlies and Garçon, 1998;
486 Lévy, 2008). The vertical distribution of chl-*a* along the west-east transect confirms this
487 enhancement in the surface layer, accompanied by an uplift of the Deep Chlorophyll-*a*
488 Maximum (DCM), following the upward doming of the isopycnals induced by the cyclonic
489 eddy (X2 on Fig. 7). The DCM within the cyclone reached 0.40 mg m^{-3} at -55 m at station 3,
490 but only 0.20 mg m^{-3} was measured at -125 m at station 14 in the anticyclone.

491 The daily evolution of satellite chl-*a* concentration within both eddies was calculated over
492 their lifetime, from 27 October to 24 December 2016 (Fig. 11). The corresponding chl-*a*
493 concentration was extracted from the centre of both eddies and smoothed with a 3-day-

494 window moving average to account for missing data caused by cloud cover. The
495 concentration of chl-*a* within the cyclone clearly increased during the spin-up phase of the
496 eddy when the eddy pumping mechanism that uplifts nutrients towards the euphotic zone is
497 meant to be at its maximum (Lévy, 2008). The concentration then decreased, but still
498 remained higher than within the anticyclone by at least 0.05 mg m^{-3} until mid-December
499 2016.

500 Although linking the distributions of nitrates and chl-*a* is beyond the scope of this paper, it
501 is worth mentioning that the vertical distribution of nitrate along the two transects is also
502 clearly constrained by the presence of the dipole (Fig. 12). The 1024.0 kg m^{-3} isopycnal
503 separates the nutrient-depleted surface layers from the nutrient-rich subsurface waters, while
504 following a remarkable, classic eddy shape.

505

506 **4. Evidence of small-scale turbulence**

507 In addition to the presence of a strong mesoscale eddy dipole, the analysis of the MAD-
508 Ridge Leg 1 dataset reveals a series of indications also of fine-scale turbulent dynamics in the
509 region during the cruise.

510

511 *4.1 Fine scale undulations of the isopycnals*

512 Fine-scale structures, smoothed out when considering the balanced geostrophic flow (Fig.
513 6a, b) are clearly visible on the west-east and south-north 2-km horizontal resolution S-ADCP
514 transects (Fig. 5). The most striking example was in the vicinity of the seamount, on its
515 western side during the west-east transect, where a series of upward (stations 5 and 7) and
516 downward (stations 4, 6 and 8) undulations of isopycnal depth can be seen (e.g. X3 in Fig. 7).
517 Deviations are of 30 m magnitude and are greatest at the depth of the seamount (-240 m) for
518 the 1025.5 kg m^{-3} isopycnal. These perturbations have a strong signature ($>40 \text{ cm s}^{-1}$) in the
519 non-geostrophic velocity field (Fig. 6c), reinforcing the southward velocity of the flow.

520

521 *4.2 Sharp horizontal density front within the frontal zone of the dipole*

522 The frontal region that separates the two eddies of the dipole was characterized by sharp
523 horizontal gradients of temperature and salinity in the 150-m-thick surface layer (Fig. 7). On
524 the west-east transect, between stations 4 and 6 and separated by just 35 km, the vessel
525 thermosalinograph, which samples water 2 m below the sea surface, reported a 1°C increase

526 in temperature⁶ and a 0.15 g kg^{-1} decrease in salinity⁷ over 6 h (not shown). Such variations
527 cannot be explained by the net local surface heat and freshwater fluxes and must therefore be
528 linked to the intrinsic properties of the two eddies, i.e. the presence of warmer, more saline
529 water within the anticyclone than in the cyclone. Theoretical studies predict that non-linear
530 processes associated with a turbulent mesoscale eddy field can lead to the enhancement of a
531 pre-existing horizontal density gradient within the surface mixed layer, and in turn generate
532 sub-mesoscale ageostrophic instabilities and strong vertical velocities (McWilliams, 2016).
533 The coarse resolution of the CTD casts during the two transects does not allow any diagnosis
534 of frontogenesis (Capet et al., 2008) nor vertical velocities though inversion of the ω -equation
535 (Pollard and Regier, 1992; Legal et al., 2007; Rousselet et al., 2019). However, the frontal
536 region between the two eddies showed high positive values of the Okubo-Weiss quantity, of
537 the order of $1.7 \times 10^{-10} \text{ s}^{-2}$, a marker for areas characterized by growth of horizontal tracer
538 gradient (Okubo, 1970; Weiss, 1991).

539

540 *4.3 Vertical tilting of the anticyclonic eddy*

541 The anticyclonic eddy was not made of a homogeneous positive vorticity core when the
542 west-east transect (14–18 November 2016) was sampled (Fig. 4). On 16 November 2016,
543 several poles of positive vorticity were seen within the +20 cm SLA, used here to identify the
544 boundary of the anticyclone. According to altimetry, the second part of the west-east transect
545 crossed two of these poles (Fig. 4). One was centred on the seamount summit at stations 7, 8
546 and 9 on 16 November whereas the other one coincided with a maximum of salinity noted
547 farther west at stations 13, 14 and 15 on 18 November. The downward doming of the
548 pycnocline (1026.4 kg m^{-3}) observed at those stations confirms this picture (Fig. 7). A close
549 look at the vertical structure of the isopycnals along this west-east transect reveals that the
550 anticyclone was slightly tilted vertically towards the west, with deeper isopycnals below the
551 eastern pole at station 13 than below the western pole at station 8.

552

553 *4.4 Entrainment of southern STUW waters*

554 The CT/SA and CT/O₂ diagrams (Fig. 9) show that, at station 28 (black dots), the
555 properties of the subsurface water corresponding to the isotherms 17–22°C (between -250 m
556 and -100 m) differed significantly relative to any of the other stations sampled. These
557 subsurface water masses were 0.2 g kg^{-1} more saline and $40 \text{ } \mu\text{mol kg}^{-1}$ more oxygenated than

⁶ The temperature increases from 23.8°C to 24.8°C.

⁷ The salinity decreases from 35.35 g kg^{-1} to 35.2 g kg^{-1} .

558 the other stations. The DCM (Fig. 8) was also weaker and deeper (0.30 mg m^{-3} at -133 m)
559 than at the two neighbouring stations on either side, just 15 miles away. Station 29 to the
560 north had a DCM of 0.80 mg m^{-3} at -50 m, whereas station 27 to the south had a DCM of 0.35
561 mg m^{-3} at -80 m. The World Ocean Atlas annual climatology (WOA18) shows that such type
562 of more saline and more oxygenated STUW is found south of the South Indian Counter
563 Current (SICC) between 30°S and 35°S . This more southern type of STUW was also
564 observed near the MAD-Ridge area, in a cross-shore transect carried out off the eastern
565 Madagascan shelf at 25°N in 2008 (Voldsund et al., 2017, their Fig. 8 and 9). Its presence was
566 identified 200 km offshore, beyond the SEMC, within a northward flow of southern waters.
567 The location of station 28 beyond the northern edge of the anticyclone but south of the
568 southern Madagascan slope, in a narrow region of strong westward velocity (Fig. 3), suggests
569 that a filament of this southern type STUW was entrained there by the anticyclonic flow.

570

571 4.5 Detachment of coastal filaments with high surface chl-a content

572 The 3-day composite image of chl-a for 20–22 November 2016 (Fig. 10) shows that a
573 patch of water highly concentrated in chl-a was sampled at stations 29, 30 and 31 during the
574 south-north transect (red line). The elongated, filament-like shape of this patch, along with the
575 evolution of the absolute surface geostrophic velocities in the area (Fig. S2), suggest that it
576 was torn off from the enriched coastal shelf waters of the South-East Madagascar coastal
577 upwelling cell (Ramanantsoa et al., 2018), then advected onto the northern Madagascar
578 Ridge. *In situ* data show indeed that the DCM was stronger and shallower than at any other
579 station of the survey (0.74 mg m^{-3} at -73 m at station 31; Fig. 8).

580

581 5. Discussion

582 Based on satellite and *in situ* data, we have described for the first time the dynamic
583 characteristics and vertical structure of a surface-intensified mesoscale dipole recently
584 expelled from the SEMC (Fig. 1 and 2). The dipole consisted of two counter-rotating vortices
585 of similar size (100 km) and intensity (0.7 f), and an intense southwestward jet (150 cm s^{-1})
586 lying in the frontal region between the two eddies (Fig. 3 and 4). CTD and S-ADCP vertical
587 profiling revealed that the cores of the two eddies forming the dipole were located above the
588 1026.4 kg m^{-3} isopycnal, within the upper 600 m (Fig. 5, 6, 7 and 8). Observations also
589 provide evidence that, close to the seamount, fine-scale dynamics superimpose onto the
590 mesoscale eddy field.

591

592 *5.1 Overall circulation and hydrography: the dominant role of the mesoscale eddy*

593 A non-linear isolated eddy-type structure has the strength to remain coherent over an
 594 extended life, trapping water masses within its core and advecting them over long distances
 595 (McWilliams and Flierl, 1979; Chelton et al., 2007). Eddies associated with high Rossby
 596 number⁸ ($Ro > 0.1$) are generally considered non-linear. They may also be defined as isolated
 597 when their azimuthal velocities decrease faster than $1/r$ (r being the distance to the eddy
 598 centre) away from their core (Morel and McWilliams, 1997). Obtaining a reliably accurate
 599 estimate of the azimuthal velocity according to r is usually difficult because of coarse
 600 resolution in the observations, background noise, and the fact that eddies are rarely observed
 601 as purely axisymmetric features, but rather elongated deformed shapes. A cruder but more
 602 reliable estimation of the capacity of an eddy to trap water masses was proposed by Chelton et
 603 al. (2007) and relies on its ability to resist dispersion into planetary Rossby waves. This ability
 604 may be measured by the ratio of the maximum azimuthal velocity (U) over the eddy
 605 propagation speed (c_β). The eddy propagation speed is here estimated as the zonal phase
 606 speed of planetary Rossby waves $c_\beta = \beta R_d^2$, with $R_d = NH/|f|$ the Rossby deformation radius, N
 607 the Brünt-Väisälä frequency, and β the meridional gradient of the Coriolis parameter f
 608 (Sutyrin and Morel, 1997). With maximum relative vorticity values of the order of $0.7 |f|$,
 609 azimuthal velocities $> 70 \text{ cm s}^{-1}$ within the upper 600 m layer and a stratification of the order
 610 of $N^2 \sim 7 \times 10^{-5} \text{ s}^{-2}$, the MAD-Ridge dipole classifies itself as a highly non-linear isolated eddy-
 611 type structure ($c_\beta \sim 11 \text{ cm s}^{-1}$ and $U/c_\beta \sim 0.7$).

612 Hence, during the MAD-Ridge Leg 1 cruise, the circulation, hydrography and primary
 613 production over the northern Madagascar Ridge were largely dominated by the signature of a
 614 surface-intensified mesoscale eddy feature. The water masses found within the cyclonic and
 615 anticyclonic eddy cores corresponded to the water masses at the formation site, i.e. a mixture
 616 of coastal upwelled waters from the southeastern Madagascar upwelling cell and STUW
 617 found within the SEMC. In addition, the distribution of chl-*a* within the dipole was originally
 618 generated during the spin-up phase of the two eddies. Upward eddy pumping within the
 619 cyclonic eddy led to enhanced primary production, which was then advected by the dipole
 620 onto the ridge (Fig. 10).

621 However, our study shows that these dipoles were more than just intense and long-life
 622 coherent structures. The strong induced velocities also entrained and stirred the surrounding

⁸ The Rossby number $Ro = U/(|f|L)$ is a non-dimensional parameter computed as the ratio of the non-linear terms of the momentum equations over the Coriolis terms. For an eddy-like structure, U and L correspond to the eddy radius and the maximum azimuthal velocity, respectively. The ratio U/L is sometimes replaced by the maximum relative vorticity in the eddy core.

623 wates masses. Chl-*a* patches were torn off from the South East Madagascar coastal upwelling
624 cell onto the northern Madagascar Ridge, along with intrusions of nearby southern
625 Madagascan shelf waters originating from south of the SICC.

626

627 *5.2 Influence of the bathymetry on the eddy flow*

628 The mesoscale eddy-dipole was observed in the vicinity of a tall and shallow seamount,
629 whose summit lies 240 m below the sea surface within the isopycnal layer where the cores of
630 the eddies resided. Therefore, the dynamics and evolution of the dipole would be expected to
631 be strongly influenced by the seamount, and more generally by the chaotic bathymetry of the
632 northern Madagascar Ridge, itself made up of several seamounts lying between -1200 m and -
633 240 m (Fig. 2). A series of observations, described below, support this hypothesis.

634 Surface-intensified mesoscale eddies typically self-propagate westwards at the zonal phase
635 speed of the planetary Rossby waves, and dipoles can even propagate faster because of their
636 mutual advecting effect (Hogg and Stommel, 1985). In the absence of bathymetry, the
637 mesoscale eddy dipole should have therefore been moving west at a speed $>10 \text{ km day}^{-1}$.
638 However, it remained trapped in the vicinity of the seamount for more than 4 weeks. Only an
639 eastward barotropic flow or some topography-induced effect could have in theory inhibited
640 the westward propagation of the eddy (Morel, 1995; Vandermeirsch et al., 2001). Hence, in
641 the absence of the former, the dynamic influence of the topography must be responsible for
642 the trapping of the eddy above the seamount. A seamount may in fact slow down the
643 propagation of an eddy (Herbette et al., 2003). In the presence of chaotic topography, eddies
644 can even remain trapped in the area for several weeks (Richardson and Tychensky, 1998;
645 Herbette, 2003; Sutyryn et al., 2011).

646 The interaction of a mesoscale eddy with a seamount favours its erosion through
647 filamentation and may lead to its vertical or horizontal splitting (Herbette et al., 2003, 2005).
648 Erosion is always accompanied by the deformation of the vortex, and results from an external
649 shear induced by the formation of two extra vortices, a cyclone that detaches from the
650 seamount and an anticyclone that forms over the seamount as a Taylor cap (Herbette et al.,
651 2003). Maps of surface relative vorticity show that the shape of the mesoscale eddy dipole
652 kept evolving during the cruise. The anticyclonic eddy was notably deformed between 9 and
653 23 November (Fig. 4), which may have resulted into the multiple poles of positive vorticity
654 observed within the +20 cm SLA closed contour. Differences in the vertical structure of the
655 flow between the two transects also highlighted the evolution of the eddy. The dipole intensity
656 was weaker at depth on the south-north transect than on the west-east transect about 4 days

657 earlier (Fig. 6). In addition, there was vertical tilting of the dipole vertical structure. These
 658 observed deformations are similar to results obtained from idealized simulations of an eddy
 659 encountering a seamount (Herbette et al., 2003, 2005; Sutyrin et al., 2011) and tend to support
 660 our hypothesis that the dipole studied here was, at a fine scale, influenced by the seamount.

661

662 *5.3 The northern Madagascar Ridge: a region characterized by highly complex circulation*

663 *5.3.1 Mesoscale variability and retroflexion of the SEMC*

664 The 1993-2016 time-series of daily SLA allowed us to track the presence of cyclones,
 665 anticyclones and dipoles over the northern Madagascar Ridge (Section 2.6 above). Dipoles
 666 were found in the area >38% of the time, single cyclones and anticyclones about 25% and
 667 30%, respectively (not shown). Although the strength of the MAD-Ridge dipole ($DS_{MAD-Ridge} = 0.32 \text{ cm km}^{-1}$)
 668 was among the strongest of the time-series ($\overline{DS} = 0.20 \pm 0.06 \text{ cm km}^{-1}$
 669 and $DS \in [0.07, 0.50] \text{ cm km}^{-1}$), the analysis demonstrates that such surface-intensified
 670 dipolar eddies are not exceptional in the area. The northern Madagascar Ridge is in fact
 671 characterized by high values of mean EKE. Previous work attributed this intense variability to
 672 the passage of intense mesoscale eddies travelling from east to west, coming either from the
 673 nearby SEMC or from the SWIO (Quartly et al., 2006; Ridderinkhof et al., 2013; Halo et al.,
 674 2014). A recent study based on SLA data showed that much of the variability in circulation in
 675 the region was related to three retroflexion regimes of the SEMC (Ponsoni et al., 2016;
 676 Ramanantsoa et al., 2020). A preliminary analysis of a 2-year current-meter time-series from
 677 two moorings deployed on the eastern and western flanks of the MAD-Ridge seamount
 678 confirmed that the variability of the circulation over the northern Madagascar Ridge is largely
 679 dominated by the retroflexion modes of the SEMC (unpublished data).

680 When the retroflexion is in a canonical mode, dipoles similar to that surveyed during the
 681 MAD-Ridge cruise are expelled from the SEMC. Surface relative vorticity showed that the
 682 dipole surveyed during the MAD-Ridge cruise resulted from the coupling between a large
 683 patch of cyclonic vorticity that was formed on the southeastern tip of Madagascar, forcing the
 684 SEMC to flow south. This cyclonic patch later detached from the current after forming a
 685 dipole with an anticyclonic vorticity patch of the SEMC (Fig. 4).

686

687 *5.3.2 Influence of sub-mesoscale dynamics:*

688 Our results have shown that sub-mesoscale dynamics may superimpose the dominant
 689 mesoscale eddy-driven flow. Some fine-scale undulations of the isopycnals were also evident

690 on the eastern side of the seamount along the west-east transect (X3 in Fig 7). Although there
691 is evidence that they were induced by eddy-topography interactions, they could also be the
692 signature of: i) sub-mesoscale features generated in the frontal region between the two eddies;
693 ii) internal tidal/Lee waves radiating away from the seamount after being generated by
694 tidal/geostrophic flow impinging over the seamount (Nikurashin and Ferrari, 2010). There is
695 evidence too that the northern Madagascar Ridge could be an area of intense internal tide
696 generation (Arbic et al., 2010; A. Koch-Larouy, pers. comm.) and that the steepness of the
697 MAD-Ridge seamount could make it a candidate for internal tide dissipation (Hosegood et al.,
698 2019). Nonetheless, no direct influence of the seamount on the vertical distribution of chl-*a*
699 was observed along the two transects (Fig. 6 and 7). The DCM was even found slightly
700 deeper over the summit (-150 m at station 8) than on the slopes of the seamount. Even if these
701 undulations corresponded to internal waves, the resolution of the CTD vertical profiles along
702 the two transects was too coarse to capture the patchiness of vertical mixing events which by
703 essence act at very local and small scales. The search for local overturning cells through the
704 determination of the Thorpe scale in the CTD vertical profiles (Dillon, 1982; Finnigan et al.,
705 2002) might have provided evidence of vertical mixing, but was beyond the scope of this
706 work.

707

708 *5.3.3 The ghost Taylor column effect*

709 The presence of Taylor columns above seamounts seems to be a deeply anchored
710 theoretical concept for biologists looking for an impact of the seamount on the distribution of
711 the lower trophic components of pelagic ecosystems. Taylor columns may indeed be
712 generated on top of a seamount by mesoscale eddies. However, one still queries their
713 effectiveness in impacting primary production and facilitating retention of organisms in the
714 context of a rapidly changing environment.

715 The time-scale of this biological response vs. the time-scale of ocean circulation variability
716 is an essential aspect of the problem. Although it is generally admitted that phytoplankton
717 responds within a day or two to the presence of nutrients within the euphotic layer, the
718 response of zooplankton is delayed by several weeks (Genin and Boehlert, 1985; Genin and
719 Dower, 2007). The 1993-2016 time-series of surface geostrophic velocity computed from
720 altimetry at the MAD-Ridge seamount was used to estimate the probability of Taylor column
721 occurrences using velocities $<30 \text{ cm s}^{-1}$ as a proxy (see Appendix). Results show that this
722 threshold was met only 27% of the time over the period 1993-2016 (not shown). In addition,
723 the time-scale (~ 10 days) of eddy variability in the region (de Ruijter et al., 2004; Nauw et al.,

2008; Halo et al., 2014; Ramanantsoa et al., 2020) seems too short to trigger a biological response of the zooplankton at the seamount (Annasawmy et al., 2020; Noyon et al., 2020). Only 17 incidences of low velocity events lasted >25 days within the 23-year time-series.

727

728 **6 Summary and conclusions**

729 The dipole surveyed during the MAD-Ridge cruise originated from the SEMC when the
730 latter was in a canonical retroflexion mode. In such a mode, intense long-life coherent dipoles
731 are expelled from the SEMC. The cruise highlighted the fact that these dipoles interacted
732 strongly with the complex bathymetry of the northern Madagascar Ridge. By blocking eddy
733 propagation and favouring its erosion, the topography contributes to the stirring of the
734 surrounding water masses by the strong eddy-induced velocities, which themselves contribute
735 indirectly to the mixing of water masses in the region and their subsequent westward
736 advection by non-linear isolated eddies. As such eddies will continue their journey towards
737 the Agulhas Current (Siedler et al., 2009), the northern Madagascar Ridge is concluded to
738 play a key role in World Ocean circulation. In particular, because fresh, cool upwelled water
739 is usually found at the southeastern tip of Madagascar when the SEMC overshoots southwards
740 (Dilmahamod et al., 2019; Ramanantsoa et al., 2020), more of this water mass is expected to
741 be exported within the Agulhas Current (Beal et al., 2006, 2011).

742 The mesoscale variability in the region is largely constrained by the variability of the
743 SEMC (Ramanantsoa et al., 2020). Eddies are therefore expected to encounter the northern
744 Madagascar Ridge when the SEMC is in the canonical retroflexion mode (34% of the time)
745 or early retroflexion mode (13% of the time). When the SEMC continues westwards (no
746 retroflexion, 53% of the time), the northern Madagascar Ridge sits between the SEMC and
747 the SICC, with limited mesoscale eddies. Current-topography interactions may only then
748 determine the circulation and hydrography of the region.

749 We stress that a biological signature resulting from a Taylor column effect is unlikely at
750 the MAD-Ridge seamount because of the intense mesoscale variability there. These results
751 are consistent with observations reported by Read and Pollard (2017), who described the
752 circulation and hydrography around six seamounts located over the South West Indian Ridge,
753 in the vicinity of the Agulhas Return Current, an area also characterized by the frequent
754 passage of strong mesoscale eddies.

755 Further, using satellite-derived chl-*a*, Demarcq et al. (2020) could not identify any
756 phytoplankton signature over the MAD-Ridge seamount. Those authors showed that chl-*a*
757 variability in the region was dominated by filaments torn off from the coastal upwelling cells

758 and advected in the vicinity of the seamount by the mesoscale and sub-mesoscale dynamics.
759 Therefore, in the vicinity of seamounts where the circulation is dominated by large mesoscale
760 variability, the distribution of chl-*a* is expected to be governed by the mesoscale eddy flow.

761

762 **Acknowledgements**

763 We thank C. Bachelier and P. Rousselot (IRD, Brest) for their S-ADCP (CB) and CTD and
764 L-ADCP (PR) calibration, and F. Baurand and S. Hillion (IRD, Brest) for the chemical
765 analysis (nutrients and phytoplankton pigments), and the officers, crew and scientists of the
766 RV *Antea* for assistance during the cruise. The work was supported financially and
767 logistically by the Institut de Recherche pour le Développement (IRD) and the Flotte
768 Océanographique Française. Additional funding was received from the Fonds Français pour
769 l'Environnement Mondial (FFEM) on the Areas Beyond National Jurisdiction (ABNJ) in the
770 SWIO and LABEX Mer. The research was carried out under the auspices of the Western
771 Indian Ocean Upwelling Research Initiative (WIOURI), part of the 2016–2020 International
772 Indian Ocean Expedition (IIOE-2) programme.

773

774 **Appendix**

775 **A Taylor cap at the MAD-Ridge seamount**

776 When a geophysical flow encounters a seamount, a closed isolated anticyclonic circulation
777 can grow and remain trapped above the seamount summit. This feature is commonly referred
778 as a Taylor cap or a Taylor column (Huppert, 1975; Huppert and Bryan, 1976). In a situation
779 of moderate stratification, like that found over the northern Madagascar ridge during leg 1, the
780 conditions for a Taylor cap to grow resume to $H_T/(H R_o) > 2$ and $R_o < 0.15$, where H_T is the
781 height of the seamount, H the bottom depth, $R_o = U/(f L)$ the Rossby number, U the flow
782 velocity, f the Coriolis parameter, L the seamount radius (White et al., 2007; Chapman and
783 Haidvogel, 1992; Sutyurin et al., 2011). Considering the characteristics of the MAD-Ridge
784 seamount ($L = 27.5$ km and $H_T = 1400$ m, $H = 1600$ m) and its latitude ($27^{\circ}30'S$), one finds
785 that the most constraining condition relates to the smallness of the Rossby number, so
786 requiring the velocity of the flow to be < 30 cm s⁻¹.

787

788 **References**

789 Alvheim, O., Torstensen, E., Fennessy, S., MacKay, F., Zaera, D., Bemiasa, J., 2009. Cruise
790 reports “Dr Fridtjof Nansen”, West Madagascar: Pelagic Ecosystem Survey.

- 791 Annasawmy, P., Ternon, J-F., Lebourges-Dhaussy, A., Roudaut, G., Cotel, P., Herbette, S.,
792 Ménard, F., Marsac, F., 2020a. Micronekton distribution as influenced by mesoscale
793 eddies, Madagascar shelf and shallow seamounts in the south-western Indian Ocean: an
794 acoustic approach. *Deep Sea Res. II* (this issue).
- 795 Arbic, B. K., Wallcraft, A.J., Metzger, E.J., 2010. Concurrent simulation of the eddying
796 general circulation and tides in a global ocean model, *Ocean Modell.* 32, 175–187.
- 797 Beal, L.M., Chereskin, T.K., Lenn, Y.D., Elipot, S., 2006. The sources and mixing
798 characteristics of the Agulhas Current. *J. Phys. Oceanogr.* 36, 2060–2074.
- 799 Beal, L. M., De Ruijter, W. P. M., Biastoch, A., Zahn, R., 2011. On the role of the Agulhas
800 system in ocean circulation and climate. *Nature*, 472(7344), 429–436.
801 doi:10.1038/nature09983
- 802 Boehlert, G.W., Mundy, B.C., 1993. Ichthyoplankton assemblages at seamounts and oceanic
803 islands. *Bull. Mar. Sci.* 53, 336–361.
- 804 Capet, X., McWilliams, J.C., Molemaker, M.J., Shchepetkin, A.F., 2008. Mesoscale to
805 submesoscale transition in the California current system. II. Frontal processes. *J. Phys.*
806 *Oceanogr.* 38, 44–64.
- 807 Chapman, D.D., Haidvogel, D.B., 1992. Formation of Taylor caps over a tall isolated
808 seamount in a stratified ocean. *Geophys. Astrophys. Fluid Dyn.* 64, 31–65.
- 809 Chelton, D. B., deSzoeke, R.A., Schlax, M.G., El Naggar, K., Siwertz, N., 1998.
810 Geographical variability of the first baroclinic Rossby radius of deformation. *J. Phys.*
811 *Oceanogr.* 28, 433–460.
- 812 Chelton, D.B., Schlax, M.G., Samelson, R.M., de Szoeke, R.A., 2007. Global observations of
813 large oceanic eddies. *Geophys. Res. Lett.* 34(15). doi.org/10.1029/2007GL030812
- 814 Clark, M.R., 2001. Are deepwater fisheries sustainable? The example of orange roughy
815 (*Hoplostethus atlanticus*) in New Zealand. *Fish. Res.* 51, 123–135.
- 816 Clark, M.R., Rowden, A.A., Schlacher, T., Williams, A., Consalvey, M., Stocks, K.I., Rogers,
817 A.D., O’Hara, T.D., White, M., Shank, T.M., Hall-Spencer, J.M., 2010. The ecology of
818 seamounts: structure, function and human impacts. *A. Rev. Mar. Sci.* 2, 253–278.
- 819 Comeau, L.A., Vézina, A.F., Bourgeois, M., Juniper, S.K., 1995. Relationship between
820 phytoplankton production and the physical structure of the water column near Cobb
821 Seamount, northeast Pacific. *Deep-Sea Res. I.* 42, 993–1005.
- 822 Demarcq, H., Noyon, M., Roberts, M.J., 2020. Satellite observations of phytoplankton
823 enrichments around seamounts in the South West Indian Ocean, with a special focus on
824 the Walters Shoal. *Deep-Sea Res. II* (this issue).
- 825 De Ruijter, W.P.M., van Aken, H.M., Beier, E.J., Lutjeharms, J.R.E., Matano, R.P., Schouten,
826 M.W., 2004. Eddies and dipoles around South Madagascar: formation, pathways and
827 large-scale impact. *Deep-Sea Res. I.* 51, 383–400.
- 828 Dilmahamod, A.F., Penven, P., Aguiar-González, B., Reason, C.J.C., Hermes, J.C., 2019. A
829 new definition of the South-East Madagascar bloom and analysis of its variability. *J.*
830 *Geophys. Res. Oceans*, 124. doi.org/10.1029/2018JC014582.
- 831 Dillon, T.M., 1982. Vertical overturns: a comparison of Thorpe and Ozmidov lengthscales. *J.*
832 *Geophys. Res.*, 87, 9601–9613.
- 833 Di Marco, S.F., Chapman, P., Nowlin, W.D., Hacker, P., Donohue, K., Luther, M., Johnson,
834 G.C., Toole, J., 2002. Volume transport and property distributions of the Mozambique
835 Channel. *Deep-Sea Res. II.* 49, 1481–1511.
- 836 Dower, J., Freeland, H., Juniper, K., 1992. A strong biological response to oceanic flow past
837 Cobb Seamount. *Deep-Sea Res. I.* 39, 1139–1145.
- 838 Dower, J., Mackas, D. L., 1996. “Seamount effects” in the zooplankton community near Cobb
839 Seamount. *Deep Sea Res. I.* 43, 837–858.

- 840 Emery, W.J., Meincke, J., 1986. Global water masses: summary and review. *Oceanologica*
841 *Acta* 9(4).
- 842 Egbert, G.D., Erofeeva, S.Y., 2002. Efficient inverse modelling of barotropic ocean tides. *J.*
843 *Atmos. Oceanic Technol.* 19, 183–204.
- 844 Finnigan, T., Luther, D., Lukas, R., 2002. Observations of enhanced diapycnal mixing near
845 the Hawaiian Ridge. *J. Phys. Oceanogr.* 32, 2988–3002.
- 846 Garcia, H.E., Weathers, K., Paver, C.R., Smolyar, I., Boyer, T.P., Locarnini, R.A., Zweng,
847 M.M., Mishonov, A.V., Baranova, O.K., Seidov, D., Reagan, J.R., 2018. World Ocean
848 Atlas 2018, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen
849 Saturation. A. Mishonov (Technical Ed.); NOAA Atlas NESDIS 83, 38 pp.
- 850 Garrett, C., 2003. Internal tides and ocean mixing. *Science* 301, 1858–1859.
- 851 Genin, A., 2004. Bio-physical coupling in the formation of zooplankton and fish aggregations
852 over abrupt topographies. *J. Mar. Syst.* 50, 3–20.
- 853 Genin, A., Boehlert, G.W., 1985. Dynamics of temperature and chlorophyll structures above a
854 seamount: an oceanic experiment. *J. Mar. Res.* 43, 907–924.
- 855 Genin, A., Dower, J.F., 2007. Seamount plankton dynamics. In *Seamounts: Ecology,*
856 *Fisheries and Conservation.* Pitcher T.J., Morato, T., Hart, P.J.B., Clark, M.R., Haggan,
857 N., Santos, R.S., (Eds) Blackwell Publishing, Oxford, UK. 84–100.
- 858 Halo, I., Penven, P., Backeberg, B., Ansorge, I., Shillington, F., Roman, R., 2014. Mesoscale
859 eddy variability in the southern extension of the East Madagascar Current: seasonal
860 cycle, energy conversion terms, and eddy mean properties. *J. Geophys. Res.* 119, 7324–
861 7356.
- 862 Hanawa, K., Talley, L.D., 2001. Mode waters. In *Ocean Circulation and Climate: Observing*
863 *and Modelling the Global Ocean.* Siedler, G., Church, J., Gould, J. (Eds). Elsevier, New
864 York. 373–392.
- 865 Herbette, S., Morel, Y., Arhan, M., 2003. Erosion of a surface vortex by a seamount. *J. Phys.*
866 *Oceanogr.* 33, 1664–1679.
- 867 Herbette, S. 2003. Erosion des tourbillons océaniques : influence de la topographie et d'un
868 front isopycnal – application aux anneaux des aiguilles. PhD thesis, University of Brest.
- 869 Herbette, S., Morel, Y., Arhan, M., 2005. Erosion of a surface vortex by a seamount on the
870 beta plane, *J. Phys. Oceanogr.* 35, 2012–2030.
- 871 Hogg, N.G., Stommel, H.M., 1985. The heton, an elementary interaction between discrete
872 baroclinic geostrophic vortices, and its implication concerning eddy heat-flow. *Proc. R.*
873 *Soc. Lond.* 397, 1–20. doi.org/10.1098/rspa.1985.0001.
- 874 Hosegood, P.J. Nimmo-Smith, W.A.M., Proud, R., Adams, K., Brierley, A.S., 2019. Internal
875 lee waves and baroclinic bores over a tropical seamount shark 'hot-spot', *Prog.*
876 *Oceanogr.* 172, 34–50. doi.org/10.1016/j.dsr2.2020.104744.
- 877 Huppert, H., 1975. Some remarks on the initiation of inertial Taylor columns. *J. Fluid Mech.*
878 67, 397–412.
- 879 Huppert, H., Bryan, K., 1976. Topographically generated eddies. *Deep-Sea Res.* 23, 655–679.
- 880 Kermabon, C., Le Bot, P., Thierry, V., Lherminier, P., Branellec, P., 2015. CADYHAC:
881 Chaîne d'Ajustage des Données d'HYdrologie Après Campagne – Documentation
882 utilisateur (V1.1). R.INT. ODE/LPO/15-01.
- 883 Kitchingman, A., Lai, S., Morato, T., Pauly, D., 2007. How many seamounts are there and
884 where are they located. In *Pitcher, T.J., Morato, T., Hart, P.J.B., Clark, M.R., Haggan,*
885 *N., Santos, R.S. (Eds). Seamounts: Ecology, Fisheries and Conservation.* Blackwell
886 Publishing, Oxford, UK. 26–40.
- 887 Koslow, J.A., 1997. Seamounts and the ecology of deep-sea fisheries. *Am. Scient.* 85, 168–
888 176.

- 889 Lavelle, J.W., Mohn, C., 2010. Motion, commotion, and biophysical connections at deep
890 ocean seamounts. *Oceanogr.* 23, 90–103.
- 891 Le Bot, P., Kermabon, C., Lherminier, P., Gaillard, F., 2011. CASCADE V6.1 : Logiciel de
892 validation et de visualisation des mesures ADCP de coque. OPS/LPO 11-01.
893 <https://archimer.ifremer.fr/doc/00342/45285/>
- 894 Legal, C., Klein, P., Treguier, A-M., Paillet, J., 2007. Diagnosis of the vertical motion in a
895 mesoscale stirring region. *J. Phys. Oceanogr.* 37, 1413–1424, doi: 10.1175/JPO3053.1
- 896 Lévy, M., 2008. The modulation of biological production by oceanic mesoscale turbulence.
897 *Lect. Notes Physics* 744, 219–261.
- 898 Locarnini, R.A., Mishonov, A.V., Antonov, J.I., Boyer, T.P., Garcia, H.E., Baranova, O.K.,
899 Zweng, M.M., Paver, C.R., Reagan, R., Johnson, D.R., Hamilton, M., Seidov, D., 2013.
900 *World Ocean Atlas 2013. Vol. 1: Temperature.*, S. Levitus, 3028 Ed.; A. Mishonov,
901 Technical Ed.; NOAA Atlas NESDIS, 73 (September), 40, doi:10.3029 1182/blood-
902 2011-06-357442.
- 903 Lutjeharms, J., Machu, E., 2000. An upwelling cell inshore of the East Madagascar Current.
904 *Deep-Sea Res. I.* 47, 2405–2411. [http://dx.doi.org/10.1016/S0967-0637\(00\)00026-1](http://dx.doi.org/10.1016/S0967-0637(00)00026-1).
- 905 Marsac, F., Galletti, F., Ternon, J-F., Romanov, E.V., Demarcq, H., Corbari, L., Bouchet, P.,
906 Roest, W.R., Jorry, S.J., Olu, K., Loncke, L., Roberts, M.J., Ménard, F., 2020.
907 Seamounts, plateaus and governance issues in the southwestern Indian Ocean, with
908 emphasis on fisheries management and marine conservation, using the Walters Shoal as
909 a case study for implementing a protection framework. *Deep-Sea Res. II* (this issue).
- 910 Mason, E., Colas, F., Molemaker, J., Shchepetkin, F., Troupin, C., McWilliams, J.C., Sangra,
911 P., 2011. Seasonal variability of the Canary Current: a numerical study. *J. Geophys. Res.*
912 116, doi:10.1029/2010/JC006665.
- 913 McGillicuddy, D.J., Robinson, A.R., Siegel, D.A., Jannasch, H.F., Johnson, R., Dickey, T. D.,
914 McNeil, J., Michaels, A.F., Knap, A.H., 1998. Influence of mesoscale eddies on new
915 production in the Sargasso Sea. *Nature* 394, 263–266.
- 916 McWilliams, J.C., 2016. Submesoscale currents in the ocean. *Proc. R. Soc. A* 472: 20160117.
917 <http://dx.doi.org/10.1098/rspa.2016.0117>
- 918 McWilliams, J., Flierl, G., 1979. On the evolution of isolated nonlinear vortices. *J. Phys.*
919 *Oceanogr.* 9, 1155–1181.
- 920 Morel, Y., 1995. The influence of an upper thermocline current on intrathermocline eddies. *J.*
921 *Phys. Oceanogr.* 25, 3247–3252.
- 922 Morel, Y., McWilliams, J.C., 1997. Evolution of isolated interior vortices in the ocean., *J.*
923 *Phys. Oceanogr.*, 27,727–748.
- 924 Mouriño, B., Fernández, E., Serret, P., Harbour, D., Sinha, B., Pingree, R., 2001. Variability
925 and seasonality of physical and biological fields at the Great Meteor Tablemount
926 (subtropical NE Atlantic). *Oceanologica Acta* 24, 167–185.
- 927 Mullineaux, L.S., Mills, S., 1997. A test of the larval retention hypothesis in seamount-
928 generated flows. *Deep-Sea Res.* 44, 745–770.
- 929 Nauw, J.J., van Aken, H.M., Lutjeharms, J.R.E., de Ruijter, W.P.M., 2006. Intrathermocline
930 eddies in the southern Indian Ocean. *J. Geophys. Res.* 111(C3).
931 <doi.org/10.1029/2005JC002917>
- 932 Nauw, J.J., van Aken, H.M., Webb, A., Lutjeharms, J.R.E., de Ruijter, W.P.M., 2008.
933 Observations of the southern East Madagascar Current and undercurrent and
934 countercurrent system. *J. Geophys. Res.* 113(C8). <doi.org/10.1029/2007JC004639>
- 935 New, A.L., Alderson, S.G., Smeed, D.A., Stansfield, K.L., 2007. On the circulation of water
936 masses across the Mascarene Plateau in the South Indian Ocean. *Deep-Sea Res. I.* 54,
937 42–74.

- 938 New, A.L., Stansfield, K., Smythe-Wright, D., Smeed, D.A., Evans, A.J., Alderson, S.G.,
939 2005. Physical and biochemical aspects of the flow across the Mascarene Plateau in the
940 Indian Ocean. *Phil. Trans. R. Soc. Lond. A.* 363, 151–168.
- 941 Nikurashin, M., Ferrari, R., 2010. Radiation and dissipation of internal waves generated by
942 geostrophic motions impinging on small-scale topography: theory. *J. Phys. Oceanogr.*
943 40, 1055–1074. doi: 10.1175/2009JPO4199.1
- 944 Noyon, M., Rasoloarijao, Z., Huggett, J., Terson, J-F., Roberts, M., 2020. Comparison of
945 mesozooplankton communities at three shallow seamounts in the South West Indian
946 Ocean. *Deep Sea Res. II* (this issue).
- 947 O'Connor, B.M., Fine, R.A., Maillet, K.A., Olson D.B., 2002. Formation rates of subtropical-
948 underwater in the Pacific Ocean. *Deep-Sea Res. I.* 49, 1571–1590.
- 949 Okubo, A., 1970. Horizontal dispersion of floatable particles in the vicinity of velocity
950 singularities such as convergences. *Deep-Sea Res.* 17, 445–454.
- 951 Oschlies, A., Garçon, V., 1998. Eddy-induced enhancement of primary production in a model
952 of the North Atlantic Ocean. *Nature* 394, 266–269.
- 953 Oudot, C., Morin, P., Baurand, F., Wafar, M., Le Corre, P., 1998. Northern and southern
954 water masses in the Equatorial Atlantic: distribution of nutrients on the WOCE A6 and
955 A7 lines. *Deep-Sea Res. I.* 45, 873–902.
- 956 Pilo, G.S., Mata, M.M., Azevedo, L.L., 2015. Eddy surface properties and propagation at
957 Southern Hemisphere western boundary current systems. *Ocean Sci.* 11, 629–641.
- 958 Pitcher, T.J., Clark, M.R., Morato, T., Watson, R., 2010. Seamount fisheries: do they have a
959 future? *Oceanography* 23, 1. 134–144.
- 960 Pollard, R.T., Regier, L.A., 1992. Vorticity and vertical circulation at an ocean front. *J. Phys.*
961 *Oceanogr.* 22, 609–625.
- 962 Pollard, R., Read, J., 2017. Circulation, stratification and seamounts in the Southwest Indian
963 Ocean. *Deep-Sea Res. II.* 136, 36–43. doi:/10.1016/j.dsr2.2015.02.018
- 964 Ponsoni, L., Aguiar-Gonzalez, Ridderinkhof, H., Maas, L.R., 2016. The East Madagascar
965 Current; volume transport and variability based on long-term observations., *J. Phys.*
966 *Oceanogr.* 46, 1045–1065, doi:10.1175/JPO-D-15-0154.1
- 967 Pous, S., Lazure, P., André, G., Dumas, F., Halo, I., Penven, P., 2014. Circulation around La
968 Réunion and Mauritius islands in the south-western Indian Ocean: a modeling
969 perspective. *J. Geophys. Res. Oceans* 119, 1957–1976. doi:10.1002/2013JC009704
- 970 Pripp, T., Gammelsrod, T., Krakstad, J., 2014. Physical influence on biological production
971 along the western shelf of Madagascar. *Deep-Sea Res. II.* 100, 174–183.
- 972 Quartly, G.D. (2006) RRS Discovery Cruise 288, 26 January - 21 February 2005. Madagascar
973 Experiment (MadEx). National Oceanography Centre, Southampton, Cruise Report 8.
974 Southampton, UK, National Oceanography Centre Southampton, 105 pp.
- 975 Quartly, G.D., Buck, J.J.H., Srokosz, M.A., Coward, A.C., 2006. Eddies around Madagascar
976 – the retroflexion re-considered. *J. Mar. Syst.* 63, 115–129.
- 977 Ramanantsoa, J.D., Krug, M., Penven, P., Rouault, M., Gula, J., 2018. Coastal upwelling
978 south of Madagascar: temporal and spatial variability. *J. Mar. Syst.* 178, 29–37.
- 979 Ramanantsoa, J.D., P. Penven, P., Raj, R.P., Renault, L., Ponsoni, L., Ostrowski, M.,
980 Dilahamod, A.F., Rouault, M., 2020. Where and how the East Madagascar Current
981 retroflexion generates? (submitted to *J. Geophys. Res.*)
- 982 Read, J., Pollard, R.T., 2017. An introduction to the physical oceanography of six seamounts
983 in the southwest Indian Ocean. *Deep-Sea Res. II.* 136, 44–58.
984 doi:/10.1016/j.dsr2.2015.06.022
- 985 Richardson, P.L., Tychensky, A., 1998. Meddy trajectories in the Canary Basin measured
986 during the SEMAPHORE experiment, 1993–1995. *J. Geophys. Res.* 103(C11), 25029–
987 25045.

- 988 Ridderinkhof, W., Le Bars, D., von der Heydt, A.S., De Ruijter, W.P.M., 2013. Dipoles of the
 989 South East Madagascar Current. *Geophys. Res. Lett.* 40(3), 558–562.
- 990 Roberts, M.J., Ternon J-F., Marsac, F., Noyon, M., Payne, A.I.L., 2020. The MADRidge
 991 Project: Bio-Physical coupling around a shallow seamount on the northern Madagascar
 992 Ridge, South West Indian Ocean. *Deep-Sea Res. II.* (this issue).
- 993 Rogers, A. D., 1994. The biology of seamounts. *Adv. Mar. Biol.* 30, 305–350.
- 994 Rogers, A.D., 2016. Pelagic ecology of the South West Indian Ocean Ridge seamounts:
 995 introduction and overview. *Deep-Sea Res. II.* 136, 1–4. doi:/10.1016/j.dsr2.2016.12.009
- 996 Rousselet, L., Doglioli, A. M., de Verneil, A., Pietri, A., Della Penna, A., Berline, L., Marrec,
 997 P., Grégori, G., Thyssen, M., Carlotti, F., Barrillon, S., Simon-Bot, F., Bonal, M.,
 998 d’Ovidio, F., Potrenko, A., 2019. Vertical motions and their effects on a biogeochemical
 999 tracer in a cyclonic structure finely observed in the Ligurian Sea. *J. Geophys. Res.*
 1000 *Oceans* 124, 3561–3574. <https://doi.org/10.1029/2018JC014392>.
- 1001 Rowden, A.A., Dower, J.F., Schlacher, T.A., Consalvey, M., Clark, M.R., 2010. Paradigms in
 1002 seamount ecology: fact, fiction and future. *Mar. Ecol.* 31, 226–241.
 1003 <https://doi.org/10.1111/j.1439-0485.2010.00400.x>
- 1004 Royer, T.C.1978. Ocean eddies generated by seamounts in the North Pacific. *Science* 199,
 1005 1063–1064.
- 1006 Schlacher, T.A., Rowden, A.A., Dower, J.J., Consalvey, M., 2010. Seamount science scales
 1007 undersea mountains: new research and outlook. *Mar. Ecol.* 31(Suppl. 1), 1–13.
- 1008 Siedler, G., Rouault, M., Biastoch, A., Backeberg, B., Reason, C.J.C., Lutjeharms, J.R.E.,
 1009 2009. Modes of the southern extension of the East Madagascar Current. *J. Geophys.*
 1010 *Res.* 114(C01005). doi:10.1029/2008JC004921
- 1011 Sinha, M.C., Loudon, K.E., Barry Parsons, B., 1981. The crustal structure of the Madagascar
 1012 Ridge. *Geophys. J. R. Astron. Soc.* 66, 351–377.
- 1013 Sprintall, J., Tomczak, M., 1992. On the formation of central water and thermocline
 1014 ventilation in the southern hemisphere. *Deep-Sea Res. I.* 40, 827–848.
- 1015 Sutyryn, G., Herbette, S., Carton, X., 2011. Deformation and splitting of baroclinic eddies
 1016 encountering a tall seamount. *Geophys. Astrophys. Fluid Dyn.* 105, 478–505. doi:
 1017 10.1080/03091929.2011.566566.
- 1018 Sutyryn, G., Morel, Y., 1997. Intense vortex motion in a stratified fluid on the beta-plane: an
 1019 analytical theory and its validation. *J. Fluid Mech.* 336, 203–220.
 1020 doi:10.1017/S0022112096004685.
- 1021 Talley, L.D., Pickard, G.L., Emery, W.J., Swift, J.H., 2011. *Descriptive Physical*
 1022 *Oceanography*, 6th edn. Academic Press, New York.
- 1023 Thurnherr, A.M., 2014. How to process LADCP data with the LDEO Software (Versions IX.7
 1024 – IX.10).
- 1025 Vandermeersch, F., Morel, Y., Sutyryn, G., 2001. The net advective effect of a vertically
 1026 sheared current on a coherent vortex. *J. Phys. Oceanogr.* 31, 2210–2225. doi:
 1027 10.1175/1520-0485(2001)031<2210:TNAEOA>2.0.CO;2.
- 1028 Vianello, P., Ternon, J-F., Demarcq, H., Herbette, S., Roberts, M.J., 2020. Ocean currents and
 1029 gradients of surface layer properties in the vicinity of the Madagascar Ridge, South
 1030 West Indian Ocean. *Deep-Sea Res. II* (this issue).
- 1031 Verron, J., Le Provost, C., 1985. A numerical study of quasigeostrophic flow over
 1032 topography. *J. Fluid Mech.* 154, 231–252.
- 1033 Voldund, A., Aguiar-Gonzalez, B., Grammelsrod, T., Krakstad, J-O., Ullgren, J., 2017.
 1034 Observations of the East Madagascar Current system: dynamics and volume transports.
 1035 *J. Mar. Res.* 75, 531–555.
- 1036 Weiss, J., 1991. The dynamics of enstrophy transfer in two-dimensional hydrodynamics.
 1037 *Physica D* 48, 273–294.

- 1038 White, M., Bashmachnikov, I., Arístegui, J., Martins, A., 2007. Physical processes and
 1039 seamount productivity. In Pitcher, T.J., Morato, T., Hart, P.J.B., Clark, M.R., Haggan,
 1040 N., Santos, R.S. (Eds), Seamounts: Ecology, Fisheries and Conservation. Blackwell
 1041 Publishing, Oxford, UK. 65–84.
- 1042 WOA18. World Ocean Atlas 2018. Temperature: Locarni et al., 2013; Salinity: Zweng et al.,
 1043 2018; Oxygen: Garcia et al., 2018.
- 1044 Zweng, M.M., Reagan, J.R., Seidov, D., Boyer, T.P., Locarnini, R.A., Garcia, H.E.,
 1045 Mishonov, A.V., Baranova, O.K., Weathers, K., Paver, C.R., Smolyar, I., 2018. World
 1046 Ocean Atlas 2018, Volume 2: Salinity. A. Mishonov (Technical Ed.). NOAA Atlas
 1047 NESDIS 82, 50 pp.

1048 Figure Legends

1049 **Figure Legends**

1050

1051 Fig. 1. Mean (1995–2015) surface eddy kinetic energy (EKE) of the western Indian Ocean
 1052 with contours (1000 and 3000 m) of the satellite (SRTM) bathymetry superimposed (solid
 1053 grey). The black box indicates the area in which mesoscale eddies was tracked. The three
 1054 seamounts surveyed during the broader MADRidge project are represented: Walters Shoal,
 1055 south of Madagascar, La Pérouse, north of Réunion Island (yellow circles), MAD-Ridge
 1056 seamount, northern Madagascar Ridge (red circle). Black arrows schematize the major
 1057 features of the oceanic circulation in the region: Agulhas Current (AC); Mozambique Channel
 1058 Anticyclonic Eddies (MCAE); South Equatorial Current (SEC); North East Madagascar
 1059 Current (NEMC); South East Madagascar Current (SEMC); South Indian Counter Current
 1060 (SICC).

1061

1062 Fig. 2. (a) Satellite (SRTM) bathymetry with the location of the east–west and south–north
 1063 transects surveyed during the MAD-Ridge Leg 1 cruise. The two transects intersect at the
 1064 MAD-Ridge seamount. Positions of the CTD and fluorometer vertical profiles (stations) are
 1065 superimposed (black dots). An index is given to each cast (yellow boxes). (b) Same as (a),
 1066 zooming in over the seamount summit. The SRTM bathymetry has been replaced by one
 1067 resulting from optimal interpolation of echo-sounder bathymetry data collected on board the
 1068 RV *Antea* during the cruise. Casts 8, 21 and 22 are located over the summit (depth ~240 m),
 1069 whereas casts 7, 9, 20 and 23 are located over the slopes of the seamount (depth ~650 m).

1070

1071 Fig. 3. (Top) Weekly average sea level anomaly (SLA) describing the mesoscale eddy field in
 1072 place during the MAD-Ridge Leg 1 cruise, with geostrophic currents (vectors) calculated
 1073 from satellite ADT superimposed: (top left) 16 November 2016; (top right) 20 November
 1074 2016. The location of the two transects (black solid lines) is superimposed on the altimetry
 1075 maps. The trajectory of the cyclone and anticyclone forming a mesoscale eddy dipole (thin
 1076 black lines with dots) is superimposed from 29 October to 24 December 2016, with positions
 1077 of the eddy centres reported every 7 days (dots). (Bottom) Low-pass filtered S-ADCP surface
 1078 current along the west–east (bottom left) and south–north (bottom right) transects. The west–
 1079 east and south–north transects were undertaken between 14 and 18 November and 19 and 23
 1080 November 2016, respectively.

1081

1082 Fig. 4. Maps (from 16 October to 14 December 2016) of surface geostrophic relative vorticity
 1083 (s^{-1}) over the northern Madagascar Ridge calculated from weekly satellite absolute dynamic
 1084 topography (ADT). The ± 20 cm SLA contours delimiting the cores of the anticyclonic and
 1085 cyclonic eddies forming the mesoscale eddy dipole are superimposed (solid white).

1086

1087 Fig. 5. Vertical sections of S-ADCP data along the west–east (14–18 November, left) and
 1088 south–north (19–23 November, right) transects, with iso-density (kg m^{-3}) contours
 1089 superimposed (solid black). Vertical sections include the current magnitude (a, b), and its
 1090 zonal u-component (c, d) and meridional v-component (e, f). Iso-contours of current
 1091 magnitude (solid white) are superimposed every 50 cm s^{-1} from -1 m s^{-1} to 1 m s^{-1} . Indices of
 1092 the CTD-fluorometer profiles (stations) are reported on the top x-axis and superimposed as
 1093 black dashed vertical lines. The seamount is also superimposed (black filled).

1094
 1095 Fig. 6. Vertical sections of geostrophic (a, b) and ageostrophic (c, d) current components
 1096 along the west-east (14–18 November) (left) and south–north (19–23 November) (right)
 1097 transects, with iso-density (kg m^{-3}) contours superimposed (solid black). The meridional v-
 1098 component/zonal u-component is shown for the east–west and south–north transects.

1099
 1100 Fig. 7. Vertical sections of conservative temperature ($^{\circ}\text{C}$), absolute salinity (g kg^{-1}), dissolved
 1101 oxygen ($\mu\text{mol kg}^{-1}$), potential vorticity ($\times 10^{-11} \text{ s}^{-1}$) and chl-*a* (mg m^{-3}), along the west–east
 1102 transect (stations 1–15), with iso-density (kg m^{-3}) contours superimposed (solid black). The
 1103 blue triangle at 46.25°E refers to the seamount. Vertical dashed lines indicate the position of
 1104 the CTD vertical profiles. Station indices are reported on the top x-axis. Note the reduced
 1105 vertical scale (0–500 m) used for potential vorticity and chl-*a*. Important features described in
 1106 the text are reported: X1: anticyclone (STUW + high salinity + high O_2); X2: cyclone (STUW
 1107 + high chl-*a*); X3: vertical deviations of the isopycnals; X4: oxygen hotspot within the SICW.

1108
 1109 Fig. 8. Same as Fig. 7, for the north–south transect. X1: Anticyclone (STUW + high salinity +
 1110 high O_2); X5: subsurface oxygen hotspot; X6: high chl-*a* (Madagascar Shelf-enriched waters);
 1111 X7: vertical undulations of isopycnal depth.

1112
 1113 Fig. 9. (a, b) CT-SA diagram, with iso-density (kg m^{-3}) contours superimposed (solid black)
 1114 for all CTD casts of the MAD-Ridge Leg 1 cruise. In (a), the dots' colour indicates whether
 1115 the CTD cast was within the cyclonic eddy (blue, stations 2–4), the anticyclonic eddy (red,
 1116 stations 8–13 and 16–26), the frontal region in between the two eddies (green, stations 5–7),
 1117 or a non-classified region (grey). Station 28 is highlighted in black dots. In (b), the colour
 1118 scale represents the depth of measurement. Water masses are identified: TSW = Tropical
 1119 Surface Water, STUW = Subtropical Underwater, SAMW = Sub Antarctic Mode Water,
 1120 SEISAMW = South East Indian Sub Antarctic Mode Water, AAIW = Antarctic Intermediate
 1121 Water. (c) Same as (a) for a CT- O_2 diagram.

1122
 1123 Fig. 10. 3-day composite (20–22 November 2016) map of satellite chl-*a* with geostrophic
 1124 current vectors superimposed (black arrows). The positions of the anticyclonic (AC) and
 1125 cyclonic (C) eddy centres are also superimposed, as well as the two transects surveyed during
 1126 the MAD-Ridge Leg 1 cruise (solid black and red). The red portion of the south–north
 1127 transect corresponds to the *in situ* fluorometer profiles that showed high chl-*a* concentrations
 1128 when integrated vertically.

1129
 1130 Fig. 11. Comparative evolution of sea level anomaly (SLA) at the centre of the cyclonic (blue)
 1131 and anticyclonic (red) eddies, and their respective maximum satellite chl-*a* concentrations
 1132 (green dotted/solid for the cyclone/anticyclone, respectively) from 25 November to 25
 1133 December 2016, including the MAD-Ridge cruise period. The increase of chl-*a* concentration
 1134 during the growing phase of the cyclone suggests a phytoplankton response to eddy pumping.

1135

1136 Fig. 12. Vertical sections of nitrate concentration ($\mu\text{mol kg}^{-1}$) along the (a) west–east and (b)
1137 south–north transects, with iso-density (kg m^{-3}) contours superimposed (solid black). Dashed
1138 vertical lines indicate the position of the CTD vertical profiles and black dots show the
1139 sampling depths. Station indices are reported on the top x-axis.

1140

1141 **Supplementary material**

1142

1143 Fig. S1. (a, b) L-ADCP and (c, d) S-ADCP current magnitude for the west–east (left) and
1144 south–north (right) transects, with iso-density (kg m^{-3}) contours superimposed (solid black).
1145 The 50 cm s^{-1} and 100 cm s^{-1} contours are overlaid (solid white). (e, f) Magnitude of the
1146 difference between the L-ADCP and S-ADCP current vectors. Vertical dashed lines indicate
1147 the position of the CTD profiles. Station indices are reported on the top x-axis. White-shaded
1148 areas indicate missing data. Note that there were no L-ADCP profiles at stations 2 or 3.

1149

1150 Fig. S2. Daily maps of absolute surface geostrophic current magnitude (colour scale), with
1151 current vectors superimposed. (a) 29 October 2016, (b) 12 November 2016, (c) 19 November
1152 2016, (d) 26 November 2016, (e) 3 December 2016, (f) 10 December 2016. The west–east
1153 and south–north transects are superimposed (solid black).








Figure 1













Figure 3






Fig. 5









Fig. 6:








Fig. 9

Author's Declaration of Interest

Regarding the re-submission of the research manuscript :

DSR2_2019_147

Observation of a mesoscale eddy dipole on the northern Madagascar Ridge: consequences for the circulation and hydrography in the vicinity of a seamount

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfied the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us.

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

We confirm that the work described has not been published previously, that it is not under consideration for publication elsewhere, that its publication is approved by all authors and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder.

We further confirm that any aspect of the work covered in this manuscript that has involved experimental animals has been conducted with the ethical approval of all relevant bodies.


We understand that the Corresponding Author is the sole contact for the Editorial process (including Editorial Manager and direct communications with the office). She is responsible for communicating with the other authors about progress, submissions of revisions and final approval of proofs. We confirm that we have provided a current, correct email address which is accessible by the Corresponding Author and which has been configured to accept email from steven.herbette@univ-brest.fr Signed by all authors as follows:

LIST OF AUTHORS

Patrick Vianello

Date: 05/06/2020

Signature:



Steven Herbette (corresponding author)

Date: 05/06/2020 Signature :



Jean-Fran çois Ternon

Date : 05/06/2020

Signature :



Herv éDemarcq

Date 05/06/2020



Signature

Michael J. Roberts

Date 05/06/2020

Signature

Journal Pre-proof