

Global importance of oxic molybdenum sinks prior to 2.6 Ga revealed by the Mo isotope composition of Precambrian carbonates

Marie Thoby¹, Kurt O. Konhauser², Philip W. Fralick³, Wladyslaw Altermann⁴, Pieter T. Visscher⁵, and Stefan V. Lalonde¹

¹CNRS-UMR6538 Laboratoire Géosciences Océan, Institut Universitaire Européen de la Mer, Université de Bretagne Occidentale, 29280 Plouzané, France

²Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, Alberta T6G 2E3, Canada

³Department of Geology, Lakehead University, Thunder Bay, Ontario P7B 5E1, Canada

⁴Department of Geology, University of Pretoria, Private Bag X20, Hatfield, Pretoria 0028, South Africa

⁵Department of Marine Sciences, University of Connecticut, Groton, Connecticut 06340, USA

ABSTRACT

Sedimentary molybdenum (Mo) isotope compositions are a promising paleoredox indicator because the Mo isotope composition of seawater reflects the balance between anoxic and oxic sinks. Most available data are from shales; however, the Mo isotope composition of carbonates also reflects the composition of ancient seawater. Here, we provide an expanded data set of carbonate Mo isotope compositions, including the first data for carbonates older than 2.64 Ga, which we evaluate against a compilation of published data for carbonates, shales, and iron formations spanning geological time. Archean carbonate samples reveal maximum $\delta^{98}\text{Mo}$ values that are generally above 1‰. These heavy values indicate that Mn(IV)-oxide or Fe(III)-oxide sinks were sufficiently important to influence the Mo isotope composition of seawater as far back as 2.93 Ga. Comparison of Mo isotope and rare earth element data, as well as residence time considerations, indicates that this metal-oxide influence was likely global. Available Mo isotope data for shales over the same time period generally show crustal values, which we attribute to negligible authigenic enrichment of Mo from seawater due to low ambient concentrations and a paucity of euxinic conditions. Our work demonstrates that the carbonate record provides important new insights into marine paleoredox conditions, especially when shale records are absent or unsuitable, and reinforces the emerging paradigm that oxic Mo sinks were important in the marine realm prior to 2.7 Ga.

INTRODUCTION

Molybdenum (Mo) stable isotopes have emerged as a powerful proxy for marine redox evolution that is based on the redox-sensitive nature of Mo sources and sinks. Under modern oxic conditions, seawater shows a homogeneous value of 2.36‰ (Siebert et al., 2003; Fig. DR1 in the GSA Data Repository¹). Modern Mo seawater sources are principally derived from the oxidative weathering of sulfide minerals in continental crust (~90%; mean $\delta^{98}\text{Mo}$ value of 0.20‰; Siebert et al., 2003; Voegelin et al., 2012) variably modified during transport to values as high as 2.39‰ (e.g., Voegelin et al., 2012). Modern Mo sinks are dominated by Mn(IV)- and Fe(III)-oxide minerals that sequester Mo by adsorp-

tion and coprecipitation. With an equilibrium Mo fractionation of 3.3‰ onto Mn(IV) oxides (Siebert et al., 2003; Barling and Anbar, 2004) and from 0.83‰ to 2.19‰ onto Fe(III)-oxides (Goldberg et al., 2009), they constitute the isotopically lightest sink and are responsible for the heavy Mo isotope enrichment of seawater. The second most important sink is sedimentary Mo scavenging under suboxic, anoxic, and euxinic conditions, which also favors the lighter isotope but to a lesser degree (see Fig. DR1). As sulfide increases, sediment isotope compositions approach that of seawater as Mo scavenging becomes near-quantitative (e.g., Neubert et al., 2008). This explains why black shales are used to track marine paleoredox using Mo isotopes.

The Mo isotope proxy has been used to provide critical insights into the relative importance of oxic and anoxic marine waters through geological time. Arnold et al. (2004) reported $\delta^{98}\text{Mo}$ values of ~0.69‰ for mid-Proterozoic euxinic sediments of the Velkerri (1.40 Ga) and Wollongorang (1.70 Ga) Formations, Australia, implying significant areas of seafloor anoxia at that time. Heavy Mo isotope signatures in shales extend back to 2.60 Ga. Conversely, prior to 2.60 Ga, shales generally show crustal Mo isotope values (Siebert et al., 2005; Wille et al., 2007), suggesting that either oxic Mo sinks were minimal prior to the Great Oxygenation Event (GOE) ca. 2.45 Ga, or Mo in shales is a poor proxy for Archean oxygenation. Interestingly, Planavsky et al. (2014) demonstrated that iron formations (IFs) as old as 2.95 Ga show highly fractionated $\delta^{98}\text{Mo}$ values ranging from -0.71‰ to 2.32‰, which vary as a function of Fe/Mn ratio. The lightest values are best explained by sydepositional adsorption of Mo to Mn(IV)-oxides, implying that sufficient O₂ for Mn(II) oxidation, and thus the evolution of oxidative photosynthesis, had already occurred by 2.95 Ga.

Here, we turn to another sedimentary proxy capable of recording ancient Mo cycling: the molybdenum isotope composition of Mo hosted in carbonates ($\delta^{98}\text{Mo}_{\text{carb}}$; Voegelin et al., 2009, 2010; Wen et al., 2011; Eroglu et al., 2015; Romaniello et al., 2016). Although carbonates are a negligible Mo sink, they may record the $\delta^{98}\text{Mo}$ signature of seawater from which they were formed (Voegelin et al., 2009; Romaniello et al., 2016). Our study provides an expanded

¹GSA Data Repository item 2019205, supplemental discussion on marine Mo cycling and postdepositional modification, description of samples and their rare earth element spectra, detailed methods and compilation of data sources, and data tables, is available online at <http://www.geosociety.org/datarepository/2019/>, or on request from editing@geosociety.org.

CITATION: Thoby, M., et al., 2019, Global importance of oxic molybdenum sinks prior to 2.6 Ga revealed by the Mo isotope composition of Precambrian carbonates: *Geology*, v. 47, p. 559–562, <https://doi.org/10.1130/G45706.1>

data set for modern and Archean carbonate Mo isotopic compositions, which we compare to published data from carbonate, shale, and IF records to shed new light on marine Mo cycling in the Archean.

SAMPLES AND METHODS

The newly acquired Mo isotopic data presented here include modern thrombolites and stromatolites from the Bahamas, 2.52 Ga stromatolites from the Gamohaana Formation (Ghaap Group, South Africa), crystal fans and stromatolites from the 2.80 Ga Mosher Carbonate (Steep Rock Group, Canada) and 2.93 Ga Ball Assemblage (Red Lake Greenstone Belt, Canada), and 2.97 Ga stromatolitic and nonstromatolitic carbonates from the Chobeni Formation (Nsuzze Group, South Africa). See Figure 1 and the Data Repository for descriptions and locations of each unit and sample. Major-element concentrations were measured on a HORIBA Ultima 2 inductively coupled plasma–atomic emission spectrometer (ICP-AES) after HF-HNO₃ total digestion, trace elements were measured on carbonate leaches (5% acetic acid) using a ThermoFisher Scientific Element2 high-resolution–inductively coupled plasma–mass spectrometer (HR-ICP-MS), and Mo isotopic compositions were measured on a ThermoFisher Scientific Neptune multicollector ICP-MS (MC-ICP-MS) after 6 N HCl digestion, double spike (DS; ⁹⁷Mo/¹⁰⁰Mo) addition, and chromatographic purification of Mo. Mo concentrations were calculated by isotope dilution. All analyses were performed at Le Pôle Spectrométrie Océan, Brest, France. For all data (new and compiled), $\delta^{98}\text{Mo}$ values are reported relative to the NIST3134 (U.S. National Institute of Standards and Technology) standard set to 0.25‰ (Nägler et al., 2014; see Table DR1 for details), uncertainties are reported as two standard deviations (2SD), and data sets are tabulated in Tables DR2–DR6. See the Data Repository for extended methods.

RESULTS AND DISCUSSION

Phanerozoic Mo Isotope Record

The modern seawater $\delta^{98}\text{Mo}$ value of 2.36‰ ($\pm 0.1\%$; Siebert et al., 2003) is reflected in the maximal $\delta^{98}\text{Mo}$ values of both modern black shales and carbonates (see Fig. 2A; Tables DR3 and DR4). With a maximum value of 1.76‰, our new data from modern Bahamian microbialites fall within the range observed for contemporaneous carbonates, but due to detrital impurities, or local production of sulfide, they do not reach the modern seawater value (see sample description in the Data Repository; Fig. 2A; Table DR2). For all available carbonate data, $\delta^{98}\text{Mo}$ values are never heavier than modern seawater and are thus useful for constraining the minimum value of seawater at any given time (Fig. 2A).

During the Phanerozoic, carbonates recorded a range of $\delta^{98}\text{Mo}$ values, from 2.42‰ to -0.91%




Figure 1. Maps showing Canadian (A) and South African (B) Archean carbonate occurrences analyzed in this study. See the Data Repository (see footnote 1) for additional details and GPS coordinates.

(see Fig. 2A), with maximum values universally attaining the modern seawater value of $2.36\% \pm 0.1\%$ (Fig. 2A). Shales recorded a range of $\delta^{98}\text{Mo}$ values similar to carbonates (from 2.44‰ to -1%), but they also included a few outliers lower than -1% . Only about one fifth of compiled Phanerozoic shale values fall within 1‰ of the modern seawater value, in part due to nonquantitative Mo removal. Carbonates are also limited in their ability to record seawater signatures, notably by their low Mo contents, which make them susceptible to diagenetic resetting or dilution by detrital materials, by the presence of organic matter or pore-water sulfide during diagenesis, which may alter whole-rock values, and/or their pervasive recrystallization at even low metamorphic grades. Nonetheless, carbonates provide complementary insight into Mo cycling during periods when shale records fall short.

Mo Cycling in the Precambrian

For the Proterozoic, no Mo isotope data are available for carbonate rocks. Maximum values for shales are highly variable, ranging from

-0.09% to 1.98‰ and averaging $0.81\% \pm 1.31\%$; Arnold et al. (2004) inferred a Proterozoic $\delta^{98}\text{Mo}_{\text{sw}}$ value of 1.08‰. Available IF data during this period are limited, but universally heavy in $\delta^{98}\text{Mo}$, with maximum values for each deposit ranging from 1.03‰ to 2.08‰ ($n = 3$). Archean Mo isotope data are more abundant in literature for all three lithologies, and a significant contrast in Mo isotope compositions among the three lithologies appears to occur prior to ca. 2.6 Ga. Maximum shale values range from 1.72‰ to 1.76‰ and average $1.73\% \pm 0.05\%$ from 2.60 Ga to 2.50 Ga, yet they are universally low (ranging from -0.46% to 0.89‰, averaging $0.31\% \pm 0.52\%$) before 2.64 Ga. In contrast, and with the exception of the Chobeni carbonates, maximum values for carbonates and IF are consistently positive from 2.97 Ga to 2.50 Ga (0.82‰ to 1.97‰, with an average of $1.31\% \pm 0.72\%$, for carbonates; 0.97‰ to 1.87‰, with an average of $1.52\% \pm 0.79\%$, for IF; see Fig. 2B). As discussed below, this contrast likely relates to the chemical conditions that enabled the capture of seawater molybdenum in shale records.




Figure 2. New and compiled $\delta^{98}\text{Mo}$ data for shales, iron formations, and carbonates through geological time. A: Data from 3.5 Ga to present. B: A focus on the period from 3.0 Ga to 2.5 Ga. See the Data Repository (see footnote 1) for values and data sources. Dark-gray area represents modern continental crust values (± 2 standard deviations [2SD]; Siebert et al., 2003), and light-gray area represents value of modern seawater (± 2 SD; Siebert et al., 2003). C: New and compiled Mo concentration data for shales, iron formations, and carbonates through geological time.

From 2.60 to 2.50 Ga, maximum $\delta^{98}\text{Mo}$ values in shales and carbonates varied from 1.40‰ to 1.97‰, and their average places a lower limit on global seawater Mo isotopic composition of $1.63\text{‰} \pm 0.48\text{‰}$ during this period (Fig. 2B). IF deposits analyzed during this period present a maximum value between 0.97‰ and 1.76‰. New data from 2.52 Ga carbonate samples from the Gamohaana Formation, Griqualand West basin, show a maximum $\delta^{98}\text{Mo}$ of 1.11‰, in line with elevated values observed for carbonates of the adjacent Transvaal basin (up to 1.97‰; Eroglu et al., 2015; Fig. 2B). The general agreement in maximum $\delta^{98}\text{Mo}$ values for available shale, carbonate, and IF records during this period is remarkable. All three archives appear to record seawater that is isotopically heavy due to sequestration of light Mo isotopes by Mn(IV)- and Fe(III)-oxide exit channels, consistent with evidence for increasing surface redox potential in the runup to the GOE. Enhanced supply of sulfate (and likely Mo) to the oceans at this time and the development of euxinic conditions (e.g., Reinhard et al., 2009) would have promoted the authigenic enrichment of seawater Mo into organic-rich shales.

In several carbonate samples measured by others (Voegelin et al., 2010; Eroglu et al., 2015), $\delta^{98}\text{Mo}$ compositions are isotopically light (as low as -0.82‰), indicating the former presence of Mn(IV)- or Fe(III)-oxides in the samples. As chemical sediments that often contain little detrital material, Precambrian carbonates are generally characterized by low Mo concentrations (<200 ppb), with average isotope-dilution whole-rock Mo concentrations of 140 ± 470 ppb for the period 2.97–2.50 Ga (Voegelin et al., 2010; Eroglu et al., 2015; this study). By contrast, shales generally reflect crustal Mo abundances (1.1 ppm; Rudnick and Gao, 2003), unless authigenic enrichment from seawater has occurred. Accordingly, detritus-poor carbonate rocks should be more sensitive to the addition of small amounts of isotopically light Mo associated with Mn(IV)- or Fe(III)-oxide sedimentary inputs.

Our data set includes the first available Mo isotope data for carbonates older than 2.64 Ga. Microbialitic calcite and aragonitic crystal fans (now replaced by calcite) from the 2.80 Ga Mosher Formation at Steep Rock (Canada) show a maximum $\delta^{98}\text{Mo}$ value of $1.22\text{‰} \pm 0.04\text{‰}$, comparable to dolomitic stromatolites and dolomitized crystal fans from the 2.93 Ga Ball Assemblage at Red Lake (Canada), which display a maximum $\delta^{98}\text{Mo}$ value of $1.03\text{‰} \pm 0.03\text{‰}$ (Fig. 2B). Both of these values are remarkably heavy. While the low Mo contents of these samples might make their $\delta^{98}\text{Mo}$ values more susceptible to diagenetic alteration, multiple lines of evidence indicate that this was not responsible for the heavy isotope enrichments prior to 2.64 Ga. Briefly, there is no relationship between $\delta^{98}\text{Mo}$ and $\delta^{18}\text{O}_{\text{carb}}$, and $\delta^{98}\text{Mo}$ data tend toward crustal values as a function of degree of


silicification (see the discussion and Figs. DR4 and DR5 in the Data Repository). Furthermore, burial of light Mo with organic carbon at concentrations typical of Archean shale is unlikely to have affected the global marine Mo cycle (see Data Repository). Instead, these heavy values suggest that Mn(IV)- and Fe(III)-oxide exit channels for Mo were sufficiently important to impart a heavy Mo isotope signature on Mesoproterozoic seawater. This conclusion is similar to that of Planavsky et al. (2014) and Ossa Ossa et al. (2018), who also concluded that an Mn(IV)-oxide exit channel was in operation ca. 2.95 Ga. Crucially, the isotopically light data from these studies imply some unknown degree of Mo removal by an oxide-based exit channel, while the isotopically heavy data in our new data set (also present in the IFs examined by Planavsky et al., 2014) reflect the composition of residual seawater.

If both carbonates (this study) and IFs (Planavsky et al., 2014) are characterized by Mo isotope compositions reflecting an important role for metal oxide-based Mo sinks prior to 2.7 Ga, then with the exception of recently analyzed shales from the Pongola Supergroup (Ossa Ossa et al., 2018), why has this not been detected in the larger shale record? Several possible explanations for this discrepancy exist, notably the possibility of spatial heterogeneity in seawater $\delta^{98}\text{Mo}$ (see below and supplementary discussion). However, considering that nearly all available data for shales older than 2.64 Ga show $\delta^{98}\text{Mo}$ values in the crustal range, we suggest that the most parsimonious explanation is that most shales simply did not experience authigenic Mo enrichment prior to 2.64 Ga. Indeed, shales older than ca. 2.64 Ga rarely show signs of enrichment in Mo above crustal values (Fig. 2B) and generally reflect detrital Mo sources. This can be explained by the fact that euxinic conditions that strongly promoted Mo enrichment in shales were generally absent, whereas after 2.63 Ga, when the first euxinic conditions were established (e.g., Reinhard et al., 2009),

carbonates and shales show comparable $\delta^{98}\text{Mo}$ records. Moreover, shales deposited under anoxic but noneuxinic conditions today still become authigenically enriched in Mo; it would appear that lower marine Mo concentrations rendered this process less effective in the Archean. In this context, carbonates that contain a smaller Mo contribution from detrital components, and do not depend on local euxinia or a large marine Mo reservoir for authigenic enrichment, provide unique insight into the upper limit of $\delta^{98}\text{Mo}$ evolution of seawater prior to 2.63 Ga that is otherwise cryptic in the shale sedimentary record. The rare cases where authigenic signals are recorded in shales (e.g., Ossa Ossa et al., 2018) point to the intriguing possibility of localized Mo isotope responses to local redox conditions.

The new carbonate $\delta^{98}\text{Mo}$ data presented here show no relation with Mn content or Mn/Fe ratio, and, considered in light of other redox indicators such as the presence of negative Ce anomalies (represented here as $\text{Pr}/\text{Pr}^* > 1$; see the Data Repository), it becomes apparent that interpretation of carbonate $\delta^{98}\text{Mo}$ data may not always be straightforward. Simple mixing calculations reveal that the Mo isotope composition of carbonates is more resistant to dilution by detrital material than rare earth element (REE)-based signatures (Fig. 3). However, paired $\delta^{98}\text{Mo}$ -Ce anomaly data from the 2.97 Ga Chobeni Formation (Fig. 3) reveal the presence of significant negative Ce anomalies, as reported elsewhere (Siahi et al., 2018), yet they show no sign of Mo isotope fractionation outside of the crustal range. This apparent contradiction may be reconciled by the significantly shorter residence times of light REEs (<300 yr; see the Data Repository) compared to Mo (likely greater than 17 k.y. during the Archean; Asael et al., 2015), making REE signals a local proxy, and as a result, the two proxies are necessarily recording redox at different spatial and temporal scales. Thus, one interpretation of the paired Chobeni $\delta^{98}\text{Mo}$ -Ce anomaly data set is that the

Figure 3. $\delta^{98}\text{Mo}$ data for carbonate samples analyzed in this study plotted against Post-Archean Australian Shale (PAAS)-normalized Pr anomalies as a proxy for true Ce anomalies (see footnote 1). Gray areas represent crustal ranges for both proxies, with darker-gray box indicating their intersection. Black lines represent mixing trajectories between crust ($\delta^{98}\text{Mo} \approx 0\text{‰}$ and $\text{Pr}/\text{Pr}^* = 1$) and different authigenic end members, including modern seawater ($\delta^{98}\text{Mo} = 2.36\text{‰}$ and $\text{Pr}/\text{Pr}^* = 1.4$) and different hypothetical $\delta^{98}\text{Mo}$ compositions of ancient seawater and Mn- and Fe-oxides ($\delta^{98}\text{Mo}$ ranging from 1.5‰ to 0‰; Pr/Pr^* fixed at 1.2). Gray arrows show trajectory of admixing of Mo associated with Mn- and Fe-oxide sources.



basin experienced oxidative Ce cycling at a local scale, while oxic sinks for Mo remained minor in seawater at a global scale. The reverse is true for Red Lake samples, while the Mosher Carbonate shows evidence for both local Mn(IV)-oxide precipitation (positive Pr/Pr*; consistent with Riding et al., 2014; Fralick and Riding, 2015) as well as global Mn(IV)-oxide precipitation (positive $\delta^{98}\text{Mo}$). For the Chobeni data, another possibility is that mixing occurred between authigenic Mo from seawater and oxide sources, with the resulting mixture falling within the crustal $\delta^{98}\text{Mo}$ range. Some points fall below the crustal $\delta^{98}\text{Mo}$ range, supporting this possibility. While caution is warranted because late oxidative alteration of surface samples may be difficult to detect (Albut et al., 2018), our Chobeni data set is consistent with at least some O_2 present locally in the basin, similar to recent findings based on REE data (Siahi et al., 2018) and Fe, Mo, and S isotope data (Eickmann et al., 2018; Ossa Ossa et al., 2018).

CONCLUSION

This study presents the first $\delta^{98}\text{Mo}$ data for carbonates older than 2.64 Ga, with values ranging from -1.54‰ to 1.22‰ . By contrast, Mo isotope compositions of shales of similar age generally show crustal values. We suggest that this discrepancy results from low concentrations of Mo in seawater as well as a general absence of euxinic conditions. This, in turn, prevented widespread authigenic Mo enrichment in shales prior to the onset of oxidative continental weathering, an increased Mo reservoir, and the first establishment of localized euxinia at ca. 2.7–2.6 Ga (e.g., Reinhard et al., 2009). Carbonate rocks are not subject to these constraints, and along with IFs, provide additional perspective on the $\delta^{98}\text{Mo}$ composition of Mesoproterozoic seawater. Our data set further indicates minimum global seawater $\delta^{98}\text{Mo}$ values of 1.0‰ and 1.2‰ at 2.93 Ga and 2.80, respectively, consistent with the idea that Mn(IV)- and Fe(III)-oxide sinks for Mo were globally important as far back as 2.95 Ga (Planavsky et al., 2014; Ossa Ossa et al., 2018). Despite possible limitations stemming from detrital contamination and mixing of different authigenic components in the same sample, the $\delta^{98}\text{Mo}$ composition of carbonates is a promising proxy for understanding marine molybdenum cycling and paleoredox evolution before conditions permitted widespread authigenic Mo enrichment in shales.

ACKNOWLEDGMENTS

We thank Dawn Sumner for providing samples and Olivier Rouxel for providing Mo standards and double spike. We also thank Emmanuel Ponzevera, Alexis de Prunel , Yoan Germain, Bleuenn Guegen, Marie-Laure Rouget, and C line Liorzou for analytical assistance, and Matthieu Plasman for geographic information system assistance. This work was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant to Konhauser

(RGPIN-165831), an NSERC Discovery Grant to Fralick (RGPIN 16400), and a U.S. National Science Foundation grant to Visscher (EAR 1561173). Altermann acknowledges financial support by the Exxaro Chair at the University of Pretoria (South Africa) and by National Research Foundation Incentive Funding. Lalonde acknowledges support from the European Union's Horizon 2020 research and innovation program (grant agreement no. 716515). Ronny Sch nberg and two anonymous reviewers are thanked for their constructive and insightful reviews.

REFERENCES CITED

- Albut, G., Babechuk, M.G., Kleinhans, I.C., Benger, M., Beukes, N.J., Steinhilber, B., Smith, A.J.B., Kruger, S.J., and Schoenberg, R., 2018, Modern rather than Mesoproterozoic oxidative weathering responsible for the heavy stable Cr isotopic signatures of the 2.95 Ga old Ijzermijn iron formation (South Africa): *Geochimica et Cosmochimica Acta*, v. 228, p. 157–189, <https://doi.org/10.1016/j.gca.2018.02.034>.
- Arnold, G.L., Anbar, A., Barling, J., and Lyons, T., 2004, Molybdenum isotope evidence for widespread anoxia in mid-Proterozoic oceans: *Science*, v. 304, p. 87–90, <https://doi.org/10.1126/science.1091785>.
- Asael, D., Rouxel, O., Bekker, A., and Scott, C., 2015, Dissolved Mo in the Archean oceans—A case study from the 2.63 Ga Jeerinah Formation, Australia: *Goldschmidt Abstracts*, 2015, p. 121.
- Barling, J., and Anbar, A.D., 2004, Molybdenum isotope fractionation during adsorption by manganese oxides: *Earth and Planetary Science Letters*, v. 217, p. 315–329, [https://doi.org/10.1016/S0012-821X\(03\)00608-3](https://doi.org/10.1016/S0012-821X(03)00608-3).
- Eickmann, B., Hofmann, A., Wille, M., Bui, T.H., Wing, B.A., and Schoenberg, R., 2018, Isotopic evidence for oxygenated Mesoproterozoic shallow oceans: *Nature Geoscience*, v. 11, p. 133–138, <https://doi.org/10.1038/s41561-017-0036-x>.
- Eroglu, S., Schoenberg, R., Wille, M., Beukes, N., and Taubald, H., 2015, Geochemical stratigraphy, sedimentology, and Mo isotope systematics of the ca. 2.58–2.50 Ga-old Transvaal Supergroup carbonate platform, South Africa: *Precambrian Research*, v. 266, p. 27–46, <https://doi.org/10.1016/j.precamres.2015.04.014>.
- Fralick, P.W., and Riding, R., 2015, Steep Rock Lake: Sedimentology and geochemistry of an Archean carbonate platform: *Earth-Science Reviews*, v. 151, p. 132–175, <https://doi.org/10.1016/j.earscirev.2015.10.006>.
- Goldberg, T., Archer, C., Vance, D., and Poulton, S.W., 2009, Mo isotope fractionation during adsorption to Fe(oxyhydr) oxides: *Geochimica et Cosmochimica Acta*, v. 73, p. 6502–6516, <https://doi.org/10.1016/j.gca.2009.08.004>.
- N gler, T.F., Anbar, A.D., Archer, C., Goldberg, T., Gordon, G.W., Greber, N.D., Siebert, C., Sohrin, Y., and Vance, D., 2014, Proposal for an international molybdenum isotope measurement standard and data representation: *Geostandards and Geoanalytical Research*, v. 38, p. 149–151, <https://doi.org/10.1111/j.1751-908X.2013.00275.x>.
- Neubert, N., N gler, T.F., and B ttcher, M.E., 2008, Sulfidity controls molybdenum isotope fractionation into euxinic sediments: Evidence from the modern Black Sea: *Geology*, v. 36, p. 775–778, <https://doi.org/10.1130/G24959A.1>.
- Ossa Ossa, F., Hofmann, A., Wille, M., Spangenberg, J.E., Bekker, A., Poulton, S.W., Eickmann, B., and Schoenberg, R., 2018, Aerobic iron and manganese cycling in a redox-stratified Mesoproterozoic epicontinental sea: *Earth and Planetary Science Letters*, v. 500, p. 28–40, <https://doi.org/10.1016/j.epsl.2018.07.044>.

- Planavsky, N.J., et al., 2014, Evidence for oxygenic photosynthesis half a billion years before the Great Oxidation Event: *Nature Geoscience*, v. 7, p. 283–286, <https://doi.org/10.1038/ngeo2122>.
- Reinhard, C.T., Raiswell, R., Scott, C., Anbar, A.D., and Lyons, T.W., 2009, A late Archean sulfidic sea stimulated by early oxidative weathering of the continents: *Science*, v. 326, p. 713–716, <https://doi.org/10.1126/science.1176711>.
- Riding, R., Fralick, P., and Liang, L., 2014, Identification of an Archean marine oxygen oasis: *Precambrian Research*, v. 251, p. 232–237, <https://doi.org/10.1016/j.precamres.2014.06.017>.
- Romaniello, S.J., Herrmann, A.D., and Anbar, A.D., 2016, Syndiagenetic diagenetic control of molybdenum isotope variations in carbonate sediments from the Bahamas: *Chemical Geology*, v. 438, p. 84–90, <https://doi.org/10.1016/j.chemgeo.2016.05.019>.
- Rudnick, R.L., and Gao, S., 2003, *Composition of the Continental Crust: Treatise on Geochemistry Volume 3*: Oxford, UK: Elsevier, 659 p., <https://doi.org/10.1016/B0-08-043751-6/03016-4>.
- Siahi, M., Hofmann, A., Master, S., Wilson, A., and Mayr, C., 2018, Trace element and stable (C, O) and radiogenic (Sr) isotope geochemistry of stromatolitic carbonate rocks of the Mesoproterozoic Pongola Supergroup: Implications for seawater composition: *Chemical Geology*, v. 476, p. 389–406, <https://doi.org/10.1016/j.chemgeo.2017.11.036>.
- Siebert, C., N gler, T.F., von Blanckenburg, F., and Kramers, J.D., 2003, Molybdenum isotope records as a potential new proxy for paleoceanography: *Earth and Planetary Science Letters*, v. 211, p. 159–171, [https://doi.org/10.1016/S0012-821X\(03\)00189-4](https://doi.org/10.1016/S0012-821X(03)00189-4).
- Siebert, C., Kramers, J., Meisel, T., Morel, P., and N gler, T.F., 2005, PGE, Re-Os, and Mo isotope systematics in Archean and early Proterozoic sedimentary systems as proxies for redox conditions of the early Earth: *Geochimica et Cosmochimica Acta*, v. 69, p. 1787–1801, <https://doi.org/10.1016/j.gca.2004.10.006>.
- Voegelin, A.R., N gler, T.F., Samankassou, E., and Villa, I.M., 2009, Molybdenum isotopic composition of modern and carboniferous carbonates: *Chemical Geology*, v. 265, p. 488–498, <https://doi.org/10.1016/j.chemgeo.2009.05.015>.
- Voegelin, A.R., N gler, T.F., Beukes, N.J., and Lacassie, J.P., 2010, Molybdenum isotopes in late Archean carbonate rocks: Implications for early Earth oxygenation: *Precambrian Research*, v. 182, p. 70–82, <https://doi.org/10.1016/j.precamres.2010.07.001>.
- Voegelin, A.R., N gler, T.F., Pettke, T., Neubert, N., Steinmann, M., Pourret, O., and Villa, I.M., 2012, The impact of igneous bedrock weathering on the Mo isotopic composition of stream waters: Natural samples and laboratory experiments: *Geochimica et Cosmochimica Acta*, v. 86, p. 150–165, <https://doi.org/10.1016/j.gca.2012.02.029>.
- Wen, H., Carignan, J., Zhang, Y., Fan, H., Cloquet, C., and Liu, S., 2011, Molybdenum isotopic records across the Precambrian-Cambrian boundary: *Geology*, v. 39, p. 775–778, <https://doi.org/10.1130/G32055.1>.
- Wille, M., Kramers, J.D., N gler, T.F., Beukes, N., Schr der, S., Meisel, T., Lacassie, J., and Voegelin, A., 2007, Evidence for a gradual rise of oxygen between 2.6 and 2.5 Ga from Mo isotopes and Re-PGE signatures in shales: *Geochimica et Cosmochimica Acta*, v. 71, p. 2417–2435, <https://doi.org/10.1016/j.gca.2007.02.019>.

Printed in USA