Retreat rates, modalities and agents responsible for erosion along the coastal chalk cliffs of Upper Normandy: The contribution of terrestrial laser scanning

Letortu Pauline ^{1, 2, *}, Costa Stephane ², Maquaire Olivier ², Delacourt Christophe ³, Augereau Emmanuel ³, Davidson Robert ², Suanez Serge ¹, Nabucet Jean ⁴

¹ Univ Bretagne Occidentale, IUEM, Lab LETG Brest Geomer, F-29280 Plouzane, France.

² Univ Caen Basse Normandie, Lab LETG Caen Geophen, Esplanade Paix, F-14000 Caen, France.

³ Univ Bretagne Occidentale, IUEM, Lab Domaines Ocean, F-29280 Plouzane, France.

⁴ Univ Rennes 2, Lab LETG Rennes Costel, F-35000 Rennes, France.

* Corresponding author : Pauline Letortu, email address : pauline.letortu@univ-brest.fr

Abstract :

In order to follow all the changes affecting the coastal chalk cliff face in Upper Normandy and improve knowledge about cliff erosion, repeated terrestrial laser scanning (TLS) surveys were carried out frequently between 2010 and 2013 (every 4-5 months). They were conducted at two sites with similar lithostratigraphic characteristics but different exposures to marine actions (the former being an abandoned cliff and the latter an active cliff). They provide a quantification of the production of debris with centimeter precision (from +/- 0.01 to 0.04 m). These surveys provided three major outcomes: 1) cliff retreat rates were measured at high spatial resolution with retreat values, unsurprisingly, 3-4 times higher for an active cliff than for an abandoned cliff. This result highlights that marine actions should be seen as not only a transport agent but also a particularly effective erosion agent; 2) a significant proportion of debris fall production (about 25%) in the total active cliff retreat was identified; and 3) one of the modalities of retreat was visualized as the creation of a basal notch, which propagates instability towards the upper part of the cliff face. Later, this instability generates rock falls coming from the whole cliff face.

Highlights

► Two sites with similar lithology but different exposures to marine actions are studied. ► Cliff face changes over time and by location are examined using TLS. ► Marine actions are an effective agent of erosion. ► The proportion of debris fall production is about 25% in the active cliff retreat. ► A basal notch propagates instability towards the upper part of the cliff face.

Keywords : Coastal chalk cliff, Retreat rates, Retreat modalities, Agents for erosion, Terrestrial laser scanning, Upper Normandy

1. Introduction

45	To understand the regressive dynamics of coastal cliffs, the knowledge of retreat rates at fine
46	scale and the study of agents leading to erosion are major challenges (Trenhaile, 1987;
47	Sunamura, 1992; Griggs and Trenhaile, 1994; Stephenson et al., 2013). Erosion is
48	traditionally quantified by studying the rates of retreat of a spatial object tracked over time,
49	often the cliff top (Bird, 2008). However, these average retreat rates are incomplete
50	information because they do not reflect the sudden nature of the hazard. This results in rock
51	falls that threaten populations located at the cliff top and at the cliff foot. Thus, retreat occurs
52	in "jerks", generated by the interaction of both internal factors (e.g., rock strength and
53	structure) and external factors (e.g., rainfall, temperature variations, and wave action). The
54	contribution of the latter to triggering rock falls is difficult to determine (Letortu et al., in
55	press). "Rock fall" is used in this paper to describe movements of coherent rock (Varnes,
56	1978). From Varnes' typology, two types of movement can be distinguished according to the

fallen volume: 1) debris falls describe the small-scale release of tiny blocks or flakes (up to 57 58 decimeter) from across the cliff face; and 2) rock falls describe large-scale movements from all or part of the cliff face. The former are common on rocky slopes but are often omitted in 59 60 quantification due to their fine scale or unsuitable point of view. Far from negligible, their participation is estimated at about 10% of the total retreat (May and Heeps, 1985; Hénaff et 61 al., 2002) and may be an early sign of instability. Monitoring these jerks requires a high 62 spatial and temporal resolution and a horizontal point of view that enables all changes to be 63 monitored (Young et al., 2009). 64

Due to unstable and subvertical cliff faces, quantification is difficult and sometimes 65 dangerous. For these reasons, remote sensing technologies are mainly used. Aerial images 66 can be used but the data accuracy is, at best, pluridecimeter and the point of view is 67 68 inappropriate (vertical or oblique). Recent advances in remote sensing technology, in 69 particular the improvement in the spatial resolution, may provide effective measurement tools. They may also offer the opportunity of conducting surveys at higher temporal 70 71 frequency. Satellite data are interesting but their spatial resolution still appears insufficient for 72 monitoring debris falls. Photogrammetry is not used because its reliability is affected by the 73 height of the cliffs (Lim et al., 2005) but new developments are promising. Terrestrial Laser 74 Scanning (TLS) is particularly convenient because the temporal and spatial resolution, as 75 well as the point of view, can be defined by the user.

The TLS technique, described as "one of the most promising surveying techniques for

77 rockslope characterization and monitoring" (Abellán et al., 2009), has been widely used in

78 the study of mass transfers to:

1) identify structural and geomorphological characteristics of landslides (Oppikofer et al.,

80 2009; Sturzenegger and Stead, 2009; Rothmund et al., 2013), landslide mapping (Rowlands

et al., 2003) and displacement tracking (Delacourt et al., 2007; Oppikofer et al., 2008, 2012;

82 Teza et al., 2008; Travelletti et al., 2008, 2013; Abellán et al., 2009, 2010);

2) analyze rock falls including those affecting cliffs (Lim et al., 2005, 2010; Rosser et al.,

2005, 2013; Quinn et al., 2010; Dewez et al., 2013; Abellán et al., 2014; Kuhn and Prüfer,
2014);

3) analyze warning movements of falls (Rosser et al., 2007; Abellán et al., 2009, 2010;

87 Royán et al., 2014).

In our study, TLS is used for repeated surveys in order to observe the evolution of the cliff face at two sites with a similar lithological context but subject to different marine forcing (abandoned and active cliffs). The aim is to quantify erosion at fine scale, to visualize the modalities of retreat and to contribute to the debate about the agents responsible for erosion of the Upper Normandy chalk cliffs. This paper first explains the study area and details the material and the TLS survey methodology. Then, the results of the cliff face monitoring are described and discussed.

95

96 2. Study area

97 2.1. Regional setting

Upper Normandy is located in the northwestern part of France, on both sides of the 50th 98 northern parallel, along the English Channel (epicontinental sea, 86 m deep on average). 99 100 The environment is macrotidal with a tidal range of 8 m. Swell is limited but the wind sea can 101 reach a significant wave height of 4 m in Dieppe (annual return period). Upper Normandy has 102 a marine west coast climate. From Météo-France data (1971–2000), average winter 103 temperatures are positive but an average of 26 daily freeze/thaw cycles is recorded per year 104 (minimal temperature can reach -15° C). Rainfall is distributed over the year (\approx 800 mm) although fall and winter are the wettest seasons (51 mm in August and 94 mm in November). 105 106 Daily rainfall can exceed 77 mm in October. The Upper Normandy cliffs, 60–70 m high on average, extend 120 km from Le Havre (SW) to Le Tréport (NE) (Seine-Maritime) (Fig. 1). At 107 their foot, there is a marine erosion platform (from 150 to 350 m wide), hidden, at the upper 108 part of the beach, by a thin gravel beach. The cliffs are intersected by numerous drained or 109

dry valleys perpendicular to the coast, protected at their outlet by often wide (from 30 to 100
m) gravel beaches due to the presence of groins or harbor jetties.

At the northwestern end of the sedimentary Paris Basin and in contact with the English 112 113 Channel, the Upper Normandy cliffs between Cap d'Antifer and Le Tréport are made of chalk with flints, dated as Upper Cretaceous (Pomerol et al., 1987; Mortimore and Duperret, 2004). 114 On this highly karstified frame (Rodet, 1991), there are residual flint formations and 115 Quaternary loess (Lautridou, 1985). The main tectonic deformations (NW-SE) have led to 116 117 the outcrop of various Upper Cretaceous geological strata (Fig. 1). The different stages of chalk have slight variations in facies and fine sedimentary discontinuities, which generate 118 119 some subtle resistance contrasts (Juignet and Breton, 1992; Laignel, 1997; Lasseur, 2007). From the oldest to the newest, Cenomanian chalk outcrops a few meters at the base of the 120 121 cliffs of Cap d'Antifer and Etretat, and east of Fécamp. It is heterogeneous, sometimes rich in detrital components (clay and guartz), and may be glauconitic or nodular. Turonian chalk is 122 the overlying strata. It is composed of clayish, grayish to whitish chalk, with frequent nodular 123 124 beds and little or no flint. This stage outcrops from Cap d'Antifer to Etretat, from Fécamp to Eletot, and from Puys (east of Dieppe) to Le Tréport. Turonian chalk is covered with 125 Coniacian chalk, outcropping between Cap d'Antifer and Saint-Valéry-en-Caux, and between 126 127 Dieppe and Le Tréport. The overlying stage, Santonian chalk, appears in the central part of the study area between Saint-Valéry-en-Caux and Puys. Finally, Campanian chalk, the most 128 recent one, is only found between Quiberville and Pourville-sur-Mer (Hautot-sur-Mer). In 129 130 relation to geotechnical studies, Santonian and Campanian chalk seem more favorable to weathering than other stages of the Upper Cretaceous (Laignel, 1997). Over these chalk 131 strata, a bed of clay and sand sediments about 10 to 30 m thick of Paleogene age (Bignot, 132 133 1962) replaces the usual residual flint formation (Laignel, 1997; Costa et al., 2006a), especially in Sainte-Marguerite-sur-Mer, Varengeville-sur-Mer (these two towns are located 134 along Cap d'Ailly) and Sotteville-sur-Mer (Fig. 1). 135







139

140 The regressive dynamics of the Upper Normandy cliffs are expressed by instantaneous rock falls affecting all or part of the cliff. At the foot of the slopes, a fine gravel beach develops but 141 the deposits are only a few meters thick. The tracking and distribution of these beaches are 142 affected by rock falls and man-made structures (harbor jetties and groins), which are 143 obstacles to longshore drift. The absence of a beach or a reduced amount of beach 144 sediments at the cliff foot may alter the effectiveness of marine actions, and therefore the 145 intensity and modalities of erosion (Costa et al., 2006b). 146 The regressive dynamics followed by the cliff top between 1966 and 2008 (photogrammetry 147 and orthophotography) reveal a retreat rate of 0.15 m y⁻¹ with high spatial variability in Upper 148 Normandy (Costa et al., 2004; Letortu, 2013; Letortu et al., 2014). However, these average 149

150 retreat rates provide only fragmentary information because they do not describe the sudden

nature of the departure of material. For example, on 19 December 2012 in Dieppe, a rock fall of approximately 100,000 m³ resulted in a cliff top retreat of 40 m in a few seconds while the average annual retreat rate on this site is 0.21 m y⁻¹ (Letortu et al., 2014; Michoud et al., 2014).

155

156 2.2. Site selection

157 To decipher the relative contribution of marine and subaerial agents in the erosion rates, two 158 neighboring sites with similar lithological contexts (cliff made up of Coniacian and Santonian chalk) but subject to different marine forcing were selected (Figs. 1 and 2). One site is an 159 abandoned cliff only affected by subaerial agents; the other site is an active cliff, which is 160 evolving under the influence of both subaerial and marine agents. The abandoned cliff face is 161 located in Dieppe. This site was divided into two areas with different cliff face orientations: 162 one corresponding to an exposure of 280°N (named Dieppe 1 W, 45 m long, 30 m high); and 163 the other corresponding to an exposure of 010°N (named Dieppe 2 N, 80 m long, 30 m high). 164 165 The site characterized by an active cliff face is along Cap d'Ailly. More precisely, it lies on either side of the Petit Ailly dry valley in Varengeville-sur-Mer (exposure of 010°N, 250 m 166 linear, maximum height of 40 m) (Fig. 2). 167



- 170 Fig. 2: Location, lithostratigraphy and panorama of Petit Ailly *and* Dieppe (1 W and 2 N)
- 171

- 172 3. Material and methods
- 173 3.1. TLS surveys
- 174 Terrestrial laser scanner is a measuring instrument that uses pulse laser technology to
- determine the distance from the device to the point to be measured. This distance is based
- 176 on a non-contact and reflectorless acquisition of a point cloud and is measured by the time
- between the transmission of an infrared laser pulse and the return of the reflected pulse. The

instrument used in this study is a RIEGL LMS-Z390i emitting a wavelength of 1,550 nm,
which records a single time-of-flight without any access to the complete signal form (full
waveform). The theoretical range of this device is 400 m for an 80% reflective surface and
140 m for a 10% reflective surface (RIEGL, 2007).

182 This technique is subject to use restrictions of two types: geometric and radiometric.

183 Regarding the geometric limitations, Lichti (2007) demonstrated the influence of the

incidence angle on the measurement accuracy. This method is also affected by radiometric

limitations including environmental ones. The accuracy and resolution of the scan depend on

the reflection of the scanned material and the atmospheric water content. During our

187 surveys, meteorological conditions were often good: no rainfalls, clear visibility and low wind.

A discussion on the principles of TLS and its performance is provided by Teza et al. (2007).

189 The main parameter that controls the spatial resolution is the distance between the

190 measured point and the TLS device. The area near the TLS device has a high spatial

resolution (millimeter). The distance to the scanned object is dependent on the length of the

192 coast being monitored and its height (the vertical swath of TLS is only 80°).

During repeated surveys at these two sites, 19 point clouds were acquired over 28 months
(between 10/07/2010 and 02/12/2013), every 4–5 months (Fig. 3).

195



Fig. 3: Temporal distribution of point clouds between 10/07/2010 and 02/12/2013 (gray: point
clouds affected by a high error margin; dotted line: unusable point cloud)

199

200 The TLS station is 33 m from the abandoned cliff of Dieppe 1 W and 37 m from Dieppe 2 N. 201 For Petit Ailly, the TLS station is positioned at 80 m. For repeated surveys of high accuracy, 202 the process of data acquisition requires additional equipment (Fig. 2): targets, a total station 203 (TPS400 Leica TC410C) and a DGPS (Trimble with a base station 5,700 and a mobile 204 receiver 5,800). The total station, where the point of setup is previously known by DGPS 205 measurements, can replace the point cloud acquired in a relative coordinate system in an absolute coordinate system (Lambert 93 and associated RGF93 and IGN69). Although the 206 207 TLS instrumental precision is ± 0.003 m to 50 m, the main source of error in data processing 208 comes from the georeferencing of the point cloud in an absolute coordinate system via the targets (Jaud, 2011). The use of a large number of targets (16 at Dieppe 1 W, 13 at Dieppe 2 209 N, 17 at Petit Ailly) and the maximizing of their distance from the TLS, as long as they all 210 211 remain visible, reduce the alignment error of the point cloud.

212

213 3.2. Data processing

The main steps in the data processing are 1) georeferencing and point cloud assembly; 2) point cloud cleaning; 3) meshing and interpolation to produce a Digital Elevation Model (DEM); and 4) creation of a DEM of Difference (DoD) (Jaud, 2011). Retreat is calculated by dividing the volume of change on the cliff face by the monitored area. The volume of change does not include debris material after failure because this is generally quickly removed (in a few days or weeks) at Petit Ailly, and quickly covered by vegetation at Dieppe. The retreat is then divided by years to obtain the annual retreat rate.

Of the 19 point clouds acquired, one was unusable (12/18/2011 at Dieppe 2 N) (Fig. 3,
dotted line). Observation of the point cloud highlighted artifacts including the rain, which

absorbed the laser pulse. Three other point clouds did not have good georeferencing (error

from 0.04 to 0.08 m). Two point clouds (07/05/2011 at Dieppe 1 W and 02/12/2013 at Dieppe

225 2 N) were certainly disturbed during acquisition as a docking cross-Channel ferry created significant temporary vibrations. The other one (02/24/2011 at Petit Ailly) was probably 226 227 disturbed by depression of the TLS during acquisition on a wet sandy foreshore. These three 228 point clouds were not used to quantify the rate of retreat, but only to study the spatial distribution of material departures (Fig. 3, dates in gray). In February 2013, human activities 229 (clearing work and setup of safety nets) disturbed measurements in Dieppe. All human 230 231 impacts were removed in the calculations and tables to monitor the "natural" evolution of the 232 chalk cliff in Dieppe.

233

234 3.3. Error margin

To quantify the error between the DEM, fixed areas were selected, such as urban furniture in 235 Dieppe or groin at Petit Ailly in each original point cloud (just after georeferencing, step 1 236 previously described). These fixed areas are equivalent to approximately 7,000 points. These 237 data were processed with the same methodology previously mentioned in order to obtain a 238 239 DEM about fixed areas at each site. The final DoD revealed the margin of error of measurement. The lowest margin of error was ± 0.01 m in Dieppe 2 N. At Petit Ailly, the 240 accuracy was ± 0.03 m. In Dieppe 1 W, it was ± 0.04 m. These differences were mainly due 241 242 to the quality of topometric data. The greatest difference was located in Dieppe 1 W because 243 topometric measurements were temporarily disturbed by a docking cross-Channel ferry and 244 renovation of the passenger terminal (building visible in Fig. 2).

245

246 **4. Results**

247 4.1. Retreat rates and spatial distribution of erosion

During the repeated surveys, in Petit Ailly, the total retreat of the cliff face was 0.57 m (\pm 0.03 m), corresponding to an annual retreat rate of 0.24 m y⁻¹. Dieppe 1 W receded by 0.08 m y⁻¹ while Dieppe 2 N receded by 0.06 m y⁻¹ (Fig. 4). Between both orientations of Dieppe, the retreat rate differences are within the margin of error. These first results highlight that the site

of Petit Ailly had a more intense retreat than Dieppe (0.24 m y⁻¹ against 0.06–0.08 m y⁻¹),

marked by sudden accelerations (up to 1.16 m y⁻¹ for the DoD F, Fig. 4). Unsurprisingly, 253 abandoned cliffs receded more slowly than active cliffs (3-4 times slower). This difference 254 255 confirms the strong influence of marine actions on the regressive dynamics of the chalk cliffs. For both sites, the retreat rates were higher for the period including all or part of winter than 256 other seasons. For example, in Petit Ailly during winter 2011/2012, the retreat rate was 0.17 257 m y^{-1} while during spring and summer 2012, it was 0.06 m y^{-1} . 258 259 TLS surveys provide a fine location of erosion - the cliff foot corresponds to the first third of the cliff height and the cliff top is the rest. The cliff top receded faster than the cliff foot, at 260 both cliffs, the active (Petit Ailly) and the abandoned ones (Dieppe): 1) 0.25 m y⁻¹ at the cliff 261

top against 0.21 m y⁻¹ at the cliff foot at Petit Ailly; and 2) 0.07–0.10 m y⁻¹ at the cliff top

against 0.02–0.03 m y^{-1} at the cliff foot in Dieppe (Fig. 4).

Previous DEM	Next DEM	Time gap (month)	Name of DoD	Retreat rate (m y ⁻¹)	
10/08/2010	02/24/2011	4.5	A	0.00	
02/24/2011	09/27/2011	7.0	В	0.09	
09/27/2011	12/18/2011	3.0	С	0.00	
12/18/2011	03/21/2012	3.0	D	0.17	
03/21/2012	09/18/2012	6.0	E	0.06	
09/18/2012	02/11/2013	4.5	F	1.16	
10/08/2010	02/11/2013	28.0	Total	0.24	.0

Petit Ailly in the context of an active cliff (error margin: ± 0.03 m)

Dieppe 1 W in the context of an abandoned cliff (error margin: ± 0.04 m)

Previous DEM	Next DEM	Time gap (month)	Name of DoD	Retreat rate (m y ⁻¹)	
10/07/2010	02/23/2011	4.5	А	0.07	
02/23/2011	07/05/2011	4.5	В	0.06]
07/05/2011	12/18/2011	5.5	С	0.03]
12/18/2011	03/22/2012	3.0	D	0.24	Setup of safety net
03/22/2012	02/12/2013	10.5	E	-	 disturbing measurements
10/07/2010	03/22/2012	17.5	Total	0.08	Cliff top erosion: 0.10 m y ⁻¹ Cliff foot erosion: 0.03 m y ⁻¹

Dieppe 2 N in the context of an abandoned cliff (error margin: ± 0.01 m)

Previous DEM	Next DEM	Time gap (month)	Name of DoD	Retreat rate (m y ⁻¹)	
10/07/2010	02/23/2011	4.5	A	0.10	
02/23/2011	07/05/2011	4.5	В	0.00	
07/05/2010	12/18/2011	6.5	С		
07/05/2011	03/22/2012	8.5	D	0.04	Setup of safety net
03/22/2012	02/12/2013	10.5	E	-	disturbing measurements
10/07/2010	03/22/2012	17.5	Total	0.06	Cliff top erosion: 0.07 m y^{-1}

265

Fig. 4: Successive retreat rates as a function of DoD (gray DEM: affected by a high error

267 margin)

268

269 4.2. Proportion of rock falls and debris falls in the retreat

270 Understanding the evolution of cliff faces depends on the frequency of surveys and the

erosive dynamics of cliffs. Their frequency (every 4–5 months) does not ensure that a

starting area that appears homogeneous in the DoD is the result of a single rock fall.

Cliff top erosion: 0.25 m y⁻¹ Cliff foot erosion: 0.21 m y⁻¹

Successive rock falls may be located at the same place. However, these different dynamics
could be determined thanks to an inventory of falls collected weekly by the non-profit
organization ESTRAN on the coastline studied (Letortu et al., 2014, in press). Due to the
centimeter accuracy of the data and the high spatial resolution of the TLS, it is possible to
separate debris falls (Fig. 5, photographs a and b) and rock falls (Fig. 5, photographs D and
F).

279



- Fig. 5: Examples of debris falls (circles a and b) and rock falls (stars D and F) at Petit Ailly identified by TLS in Fig. 8 (ESTRAN organization, 2012)
- 283
- Over the period studied (October 2010–February 2013), the abandoned cliff was only
- affected by debris falls (blue in electronic version of Figs. 6 and 7). The active cliff was
- affected by debris falls (dark blue in electronic version of Fig. 8) and rock falls (from light blue

- to red in electronic version of Fig. 8). Rock falls (10 cases were identified from 10 to 1,636
 m³) generated local significant retreats, especially on the eastern cliff face (up to 4.71 m
 deep). However, the largest ones (1,636 m³ for rock fall F and 194 m³ for rock fall G) were
 observed on the DoD F between 09/18/2012 and 02/11/2013 (Fig. 8).
- 291



292

Fig. 6: DoDs of the cliff face at Dieppe 1 W between 10/07/2010 and 02/12/2013 (gray title:

- 294 DoD affected by a high error margin; F: fall, W: winter, SP: spring, SU: summer; *: with
- 295 human actions)



Fig. 7: DoDs of the cliff face at Dieppe 2 N between 10/07/2010 and 02/12/2013 (gray title:

299 DoD affected by a high error margin; F: fall, W: winter, SP: spring, SU: summer; *: with

300 human actions)



³⁰¹

- Fig. 8: DoDs of the cliff face at Petit Ailly between 10/08/2010 and 02/11/2013 (gray title:
- 303 DoD affected by a high error margin; F: fall, W: winter, SP: spring, SU: summer)

Thus, on the active cliff, debris falls represented 25% of the total retreat over the period studied (Fig. 9). Debris falls are responsible for a retreat of 0.06 m y⁻¹, against 0.18 m y⁻¹ for rock falls over 28 months under a total retreat rate of 0.24 m y⁻¹. The rate related to debris falls is consistent with the retreat rate for the abandoned cliffs (0.06–0.08 m y⁻¹) where only debris falls were observed (Fig. 9).



311 Fig. 9: Retreat rates as a function of departure area and type of movement



The DoD B in Fig. 8 shows departures of rocky material at the active cliff foot (east face). 314 These departures occur by two rock falls, which have various depths. Rock fall A generates a 315 316 maximal basal notch of 2.26 m (59 m³) while rock fall B creates a maximal notch of 1.61 m 317 (89 m³). Later, in the DoD D (Fig. 8), debris fall b (0.61 m maximum deep, 2 m³) and rock fall D (2.05 m maximum deep, 86 m³) occur near these initial notches. Notches seem to 318 propagate instability gradually at their periphery and often towards the top of the cliff. A few 319 320 months later (DoD F), rock falls of the whole cliff face are observed at or next to these locations (rock falls F and G with a volume of 1,636 m³ and 194 m³, respectively). This study 321 highlights one of the retreat modalities: the creation of a basal notch that, by overhanging, 322 will gradually destabilize the top of the cliff to generate a rock fall of massive volume from the 323 324 whole cliff face.

325

326 **5. Discussion**

Due to its centimeter precision and high temporal frequency, monitoring by TLS has provided a new understanding of the retreat of chalk cliff faces at the study sites. Some of the results generate discussion, which is organized in four points: 1) the spatial distribution of the retreat over the short and long terms; 2) the proportion of debris falls in retreat; 3) the modality of cliff retreat; and 4) the determination of possible agents and processes responsible for erosion.

333 Over 28 months of repeated surveys, the cliff top retreated faster than the cliff foot for both 334 active and abandoned cliffs. At the abandoned cliff, this result seems to be a normal evolution because, since the polder construction in the early 1980s, the slope has evolved 335 exclusively under subaerial agents. These agents are particularly efficient at the cliff top. 336 337 Over a longer timescale, the retreat rates should not converge. The final result should be a terrestrial slope. In contrast, this situation may appear surprising at the active cliff of Petit 338 Ailly. The foot of the active cliff should be more eroded due to a combination of marine and 339 subaerial agents. This result can be explained by the lithology and the timescale of our study. 340 This active cliff has a specific morphostructural context: the chalk has overlying Tertiary 341

342 strata. In these strata, there are water tables from where runoff may affect the chalk cliff top.

343 This process may create a temporarily high retreat of the chalk cliff top. Over a longer

- timescale, the retreat difference between the foot and the top of the cliff should change.
- Indeed, there is a "disconnect in the timescale of our monitoring and the cliff morphology"
- 346 (Rosser et al., 2013). On the active cliff, these two rates should evolve to fit with the sub-
- vertical profile of the current cliff (Fig. 10). Otherwise, the form of the cliff should change.



- Fig. 10: Sub-vertical cliff profiles and basal notches in Sainte-Marguerite-sur-Mer (Capd'Ailly)
- 351

348

Our study also highlights that debris falls correspond to 25% of the total retreat of the active 352 353 cliff. They are responsible for a retreat rate of 0.06 m y^{-1} , while rock falls are responsible for a retreat rate of 0.18 m y⁻¹ during the 28 months. This share is much greater than those 354 reported by May and Heeps (1985) and Hénaff et al. (2002) (10% and 11%, respectively). 355 Nevertheless, as suggested by Costa (1997), this proportion of debris falls is highly variable 356 357 in time and may be guickly removed. Costa (1997) measured a retreat of 0.02 m for a single event (a rapid thaw on 31 December 1995) along the same coastline. Fortunately, these 358 debris falls were observed but were removed at high tide. Our repeated surveys by TLS and 359 the resultant visualization of erosion in the centimeter scale better quantify these debris falls. 360 These surveys over 28 months also enable one of the modalities of cliff retreat to be 361 identified. On the east side of the dry valley at Petit Ailly, the upward propagation of failure 362

from a destabilizing notch at the cliff foot is observed. This recalls the usual scenario 363 364 (sometimes considered simplistic) of a basal notch (Sunamura, 1988, 1992; Trenhaile, 1987; Stephenson et al., 2013; Trenhaile et al., 2013) leading to larger rock falls named 365 366 overhanging movements (Hantz et al., 2003; Andriani and Walsh, 2007). Our work reveals, like Rosser et al. (2013) along the cliffs of the North York Moors National Park (UK), that 367 departure areas evolve over time around their periphery with a dominant upward direction 368 (vertical and subvertical). Thus, the authors stated that the distribution of rock falls may not 369 370 be random: "there is clear spatial clustering indicative of propagation when rock falls are considered as cumulative through time". These notches (created by debris falls or rock falls) 371 372 are visible in many places along the Upper Normandy coast (Costa, 1997), especially along 373 Cap d'Ailly (Fig. 10). They may be a warning sign of more massive failure (the date is still 374 difficult to predict) which can cover the whole cliff face (Young et al., 2009). As Terzaghi 375 (1950) suggested, slow surface movements precede catastrophic landslides. He added, "if a landslide comes as a surprise to the eyewitnesses, it would be more accurate to say that the 376 377 observers failed to detect the phenomena which preceded the slide...". Further investigations 378 are needed to understand better the link between preparatory phenomena and rock falls. 379 The combination of knowledge about the location, the time of cliff retreat, and the type of 380 movement (debris falls and rock falls) may provide hypotheses about the agents (marine or 381 subaerial) and processes responsible for erosion. Debris falls affect both sites (Dieppe and 382 Petit Ailly) and are present on the whole cliff faces. A large production of debris falls was 383 observed at the foot of the Petit Ailly active cliff (green circles in electronic version of Fig. 8). The location of erosion is specific: it occurs only at the first 10 m of the cliff face. This erosion 384 is not found at the base of the abandoned cliffs in Dieppe. These features raise questions 385 386 about the agents and processes responsible for erosion. This basal erosion of the active cliff 387 may be induced by the water table, which saturates the chalk with water, prone to hydroclasty and cryoclasty. However, the abandoned cliffs where there is the water table and 388 389 other subaerial agents should also have this basal erosion. Thus, marine actions may lead to basal erosion by debris falls. The first 10 m of erosion at the cliff foot are consistent with the 390

height of the waves that can reach it locally. Erosion due to marine actions may be quantified 391 by deduction. At the foot of the abandoned cliffs, the annual retreat rate related to debris 392 393 falls, and exclusively due to subaerial agents, is 0.03 m y⁻¹. Along the active cliff, the retreat rate related to debris falls is 0.14 m y^{-1} (Fig. 9) but includes subaerial and marine agents. 394 Marine actions may be responsible for a retreat rate by debris falls of 0.11 m y⁻¹ of the 395 observed 0.14 m y^{-1} . Thus, at the active cliff foot, marine actions may generate a retreat by 396 397 debris falls four times higher than subaerial actions. This simple (perhaps simplistic) 398 deduction may provide the first elements of quantification of marine actions. Again, marine actions may be an efficient agent of erosion. 399

The marine actions may not be uniform at the active cliff foot (Fig. 8). The sides of the dry 400 valley evolve differently. The west face has higher debris fall retreat at the cliff foot than the 401 402 east face. This difference at fine scale may be due to the presence of a man-made structure, a groin, located in the axis of the dry valley. Indeed, this structure leads to an accumulation 403 of gravels at the west face. As Robinson (1977) observed an increase in cliff erosion by a 404 405 factor of 18 when beach sediments were present compared to when they were not, we suggest these gravels may be used as projectiles by the swell and may explain the high 406 407 abrasion by debris falls at the cliff foot on the west face (Fig. 11). On the east face, waves 408 breaking without projectiles may be less effective. These observations converge towards 409 model results (Sunamura, 1982, 1992; Limber and Murray, 2011; Kline et al., 2014). 410 Sunamura (1982, 1992) showed in laboratory experiments that cliffs and beaches exhibit 411 internal feedbacks. The amount and configuration of beach material can bring positive feedback with higher wave erosive efficacy by providing abrasive agents (Sunamura, 1992). 412 Kline et al. (2014) studied the role of mechanical abrasion by beach sediments in 413 414 cliff/platform/beach evolution over a long timescale (up to 1,000 years). They suggest that "mechanical abrasion is a feasible mechanism of cliff and platform evolution". However, on 415 the whole cliff face scale at Petit Ailly, the east face has a higher retreat rate than the west 416 face $(0.39 \text{ m y}^{-1} \text{ against } 0.12 \text{ m y}^{-1})$. This is due to numerous rock falls that occur on the east 417 face but rarely on the west face. Dewez et al. (2013) also observed rock falls at the cliff foot 418

at the downdrift side of the groin of Criel-sur-Mer (20 km north of Dieppe). They mentioned 419 the caving effect of waves by hydraulic forces including water hammer and air compression, 420 421 particularly effective without a beach (gravel transit is blocked by the 120 m-long groin). At 422 the Petit Ailly, this effect is possible due to the lack of beach on the downdrift side of the groin. There is no protecting gravel beach. When a rock fall occurs, most fallen rocks are 423 guickly removed. Therefore, water hammer and air compression may be more intense and 424 425 may generate rock falls. This suggestion underlines the role of marine actions as an erosion 426 agent and the dual role of a gravel beach, which can protect the cliff foot or attack it (Costa et al., 2006b). Future TLS surveys here will be valuable to provide complementary information 427 over a longer timescale, leading to a better understanding of the roles of marine actions and 428 429 gravel beaches.

430



432 Fig. 11: Mid-tide (coefficient 106) at Petit Ailly and position of gravel beach

433

The departure of debris falls at the active cliff top, and sometimes reaching the mid-cliff, is visible on the DoDs B and E (purple circles in electronic version of Fig. 8) during off-winter periods, thus without frost. In the context of an abandoned cliff, the location of debris falls also highlights linear shapes (Figs. 6 and 7). The often linear shapes of the departure areas and their concentration in dihedrons seem to be runoff zones, and thus subaerial actions. Subaerial actions may not be negligible in the context of an active cliff. In this paper, erosion agents have been discussed – marine actions with mechanical

abrasion, water hammer, air compression, and subaerial agents with runoff. These

hypotheses open up research perspectives about the quantification of their action. Beyond
increasing the temporal representativeness of monitoring, reducing the time interval between
each survey and conducting surveys before/during/after specific weather and sea conditions
including freeze/thaw cycles and storms, supplementary measurements will be carried out to
1) monitor the behavior of water tables that could affect the stability of the cliff; 2) obtain a
fine acquisition of rainfall, thermal and swell data; and 3) quantify the role of marine actions
in the destabilization of the chalk massif.

It is necessary to focus not only on the external agents of erosion but also on internal factors such as chalk fracturing, which alters material resistance and often predetermines the extent of rock falls. To deal with the problem of spatial representativeness and check if our results are replicable, other lithologies of cliff faces are currently under study.

453

454 6. Conclusions

455 In Upper Normandy, the repeated surveys over 28 months on two sites with similar 456 lithostratigraphic characteristics but different exposures to marine actions, highlight that cliff retreat is highly variable in time, location, its modalities, and the agents/processes involved. 457 458 The active cliff of Petit Ailly (Varengeville-sur-Mer) has regressive dynamics 3-4 times 459 greater than the abandoned cliffs of Dieppe (0.24 m y⁻¹ against 0.06–0.08 m y⁻¹). This 460 difference highlights the importance of marine actions in erosion dynamics. Whereas the 461 abandoned cliff recedes only with debris falls, the active cliff has a combination of debris falls 462 (25% of the retreat) and rock falls (75%). On both sites, a spatial variation in retreat rates can be observed over this short timescale: the cliff top recedes faster than the cliff foot. 463 Moreover, due to the fine location of the retreat obtained by our methodology, one of the 464 465 modalities of retreat is detected. Precursory movements creating a basal notch later generate massive rock falls on the whole cliff face. These results indicate that TLS can be 466 very useful for the spatial prediction of rock falls and for contributing to the debate about the 467 468 agents and processes responsible for erosion.

470 Acknowledgements

We thank the reviewers for their detailed reviews of the manuscript and their helpfulcomments.

473

474 **References**

- Abellán, A., Jaboyedoff, M., Oppikofer, T., Vilaplana, J.M., 2009. Detection of millimetric
- deformation using a terrestrial laser scanner: experiment and application to a rock fall event.
- 477 Natural Hazard and Earth System Sciences 9(2), 365-372.
- 478 Abellán, A., Calvet, J., Vilaplana, J.M., Blanchard, J., 2010. Detection and spatial prediction
- of rock falls by means of terrestrial laser scanner monitoring. Geomorphology 119(3), 162-
- 480 171.
- Abellán, A., Oppikofer, T., Jaboyedoff, M., Rosser, N.J., Lim, M., Lato, M.J., 2014. Terrestrial
- 482 laser scanning of rock slope instabilities. Earth Surface Processes and Landforms 39(1), 80-
- 483 97.
- 484 Andriani, G.F., Walsh, N., 2007. Rocky coast geomorphology and erosional processes: A
- case study along the Murgia coastline South of Bari, Apulia SE Italy. Geomorphology 87(3),
 224-238.
- Bignot, G., 1962. Étude sédimentologique et micropaléontologique de l'Éocène du Cap
- d'Ailly (près de Dieppe, Seine-Maritime). Ph.D. Thesis, University of Paris, France.
- Bird, E., 2008. Coastal Geomorphology. An introduction (Second Edition). Wiley, New York.
- 490 Costa, S., 1997. Dynamique littorale et risques naturels: L'impact des aménagements, des
- 491 variations du niveau marin et des modifications climatiques entre la baie de Seine et la baie
- de Somme. Ph.D. Thesis, University of Paris I, France.
- 493 Costa, S., Delahaye, D., Freiré-Diaz, S., Di Nocera, L., Davidson, R., Plessis, E., 2004.
- 494 Quantification of the Normandy and Picardy chalk cliff retreat by photogrammetric analysis.
- In: Mortimore, R.N., Duperret, A. (Eds.), Coastal chalk cliff instability. Engineering Geology
- 496 Special Publications, London, pp. 139-148.

- 497 Costa, S., Laignel, B., Hauchard, E., Delahaye, D., 2006a. Facteurs de répartition des
- 498 entonnoirs de dissolution dans les craies du littoral du Nord-Ouest du Bassin de Paris.
- 499 Zeitschrift für Geomorphologie 50, 95-116.
- 500 Costa, S., Hénaff, A., Lageat, Y., 2006b. The gravel beaches of north-west France and their
- 501 contribution to the dynamic of the coastal cliff-shore platform system. Zeitschrift für
- 502 Geomorphologie, Suppl.-Vol 144, 199-214.
- 503 Delacourt, C., Allemand, P., Berthier, E., Raucoules, D., Casson, B., Grandjean, P.,
- Pambrun, C., Varel, E., 2007. Remote-sensing techniques for analysing landslide kinematics:
- a review. Bulletin de la Société Géologique de France 178(2), 89-100.
- 506 Dewez, T.J.B., Rohmer, J., Regard, V., Cnudde, C., 2013. Probabilistic coastal cliff collapse
- 507 hazard from repeated terrestrial laser surveys: case study from Mesnil Val (Normandy,
- northern France). Journal of Coastal Research, Special Issue 65, 702-707.
- 509 Griggs, G.B., Trenhaile, A.S., 1994. Coastal cliff and platform. In: Carter, R.W.G., Woodroffe,
- 510 C.D. (Eds.), Coastal Evolution. Cambridge Univ. Press, Cambridge, pp. 425-450.
- Hantz, D., Vengeon, J.M., Dussauge-Peisser, C., 2003. An historical, geomechanical and
- 512 probabilistic approach to rock-fall hazard assessment. Natural Hazards and Earth System
- 513 Sciences 3(6), 693-701.
- 514 Hénaff, A., Lageat, Y., Costa, S., Plessis, E., 2002. Le recul des falaises crayeuses du Pays
- 515 de Caux: détermination des processus d'érosion et quantification des rythmes d'évolution.
- 516 Géomorphologie: Relief, Processus, Environnement 8(2), 107-118.
- Jaud, M., 2011. Techniques d'observation et de mesure haute résolution des transferts
- 518 sédimentaires dans la frange littorale. Ph.D. Thesis, University of Bretagne Occidentale,
- 519 France.
- Juignet, P., Breton, G., 1992. Mid-Cretaceous sequence stratigraphy and sedimentary
- 521 cyclicity in the western Paris Basin. Palaeogeography, Palaeoclimatology, Palaeoecology
 522 91(3), 197-218.
- 523 Kline, S.W., Adams, P.N., Limber, P.W., 2014. The unsteady nature of sea cliff retreat due to
- mechanical abrasion, failure and comminution feedbacks. Geomorphology 219, 53-67.

- 525 Kuhn, D., Prüfer, S., 2014. Coastal cliff monitoring and analysis of mass wasting processes
- with the application of terrestrial laser scanning: A case study of Rügen, Germany.

527 Geomorphology 213, 153-165.

- Laignel, B., 1997. Les altérites à silex de l'ouest du Bassin de Paris: caractérisation
- 529 lithologique, genèse et utilisation potentielle comme granulats. Ph.D. Thesis, University of
- 530 Rouen, France.
- 531 Lasseur, E., 2007. La Craie du Bassin de Paris (Cénomanien-Campanien, Crétacé
- 532 supérieur). Sédimentologie de faciès, stratigraphie séquentielle et géométrie 3D. Ph.D.
- 533 Thesis, University of Rennes 1, France.
- Lautridou, J.P., 1985. Le cycle périglaciaire pléistocène en Europe du Nord-Ouest et plus
- particulièrement en Normandie. Ph.D. Thesis, University of Caen Basse-Normandie, France.
- Letortu, P., 2013. Le recul des falaises crayeuses haut-normandes et les inondations par la
- 537 mer en Manche centrale et orientale: de la quantification de l'aléa à la caractérisation des

risques induits. Ph.D. Thesis, University of Caen Basse-Normandie, France.

- 539 <u>https://halshs.archives-ouvertes.fr/file/index/docid/1018719/filename/These_Letortu.pdf</u>
- Letortu, P., Costa, S., Bensaid, A., Cador, J.M., Quénol, H., 2014. Vitesses et rythmes de
- 541 recul des falaises crayeuses de Haute-Normandie (France): méthodologie et variabilité du
- recul. Géomorphologie, Relief, Processus et Environnement 2, 133-144.
- 543 Letortu, P., Costa, S., Cador, J.M., Coinaud, C., in press. Statistical and empirical analyses
- of the triggers of coastal chalk cliff failure. Earth Surface Processes and Landforms.
- Lichti, D.D., 2007. Error modelling, calibration and analysis of an AM-CW terrestrial laser
- scanner system. ISPRS Journal of Photogrammetry and Remote Sensing 61(5), 307-324.
- Lim, N., Petley, D.N., Rosser, N.J., Allison, R.J., Long, A.J., Pybus, D., 2005. Combined
- 548 digital photogrammetry and time-of-flight laser scanning for monitoring cliff evolution. The
- 549 Photogrammetric Record 20(110), 109-129.
- Lim, M., Rosser, N.J., Allison, R.J., Petley, D.N., 2010. Erosional processes in the hard rock
- coastal cliffs at Staithes, North Yorkshire. Geomorphology 114(1), 12-21.

- Limber, P.W., Murray, A.B., 2011. Beach and sea-cliff dynamics as a driver of long-term
- rocky coastline evolution and stability. Geology 39(12), 1147-1150.
- May, V., Heeps, C., 1985. The nature and rates of changes on chalk coastlines. Zeitschrift
 für Geomorphologie, Suppl.-Vol 57, 81-94.
- 556 Michoud, C., Carrea, D., Costa, S., Derron, M.H., Jaboyedoff, M., Davidson, R., Delacourt,
- 557 C., Letortu, P., Maquaire, O., 2014. Landslide detection and monitoring capability of boat-
- based mobile laser scanning along Dieppe coastal cliffs, Normandy. Landslides, 1-16. DOI
 10.1007/s10346-014-0542-5.
- 560 Mortimore, R.N., Duperret, A., 2004. Coastal chalk cliff instability. Engineering Geology
- 561 Special Publications, London.
- 562 Oppikofer, T., Jaboyedoff, M., Keusen, H.R., 2008. Collapse at the eastern Eiger flank in the 563 Swiss Alps. Nature Geoscience 1(8), 531-535.
- 564 Oppikofer, T., Jaboyedoff, M., Blikra, L., Derron, M.H., Metzger, R., 2009. Characterization
- and monitoring of the Åknes rockslide using terrestrial laser scanning. Natural Hazards and
 Earth System Sciences 9(3), 1003-1019.
- 567 Oppikofer, T., Bunkholt, H., Fischer, L., Saintot, A., Hermanns, R.L., Carrea, D., Longchamp,
- 568 C., Derron, M.H., Michoud, C., Jaboyedoff, M., 2012. Investigation and monitoring of rock
- slope instabilities in Norway by terrestrial laser scanning. Proc. of the 11th International & 2nd
- 570 North American Symposium on Landslides, Banff, Canada, pp. 3-8.
- 571 Pomerol, B., Bailey, H.W., Monciardini, C., Mortimore, R.N., 1987. Lithostratigraphy and
- 572 biostratigraphy of the Lewes and Seaford chalks: a link across the Anglo-Paris basin at the
- 573 Turonian-Senonian boundary. Cretaceous Research 8(4), 289-304.
- Quinn, J.D., Rosser, N.J., Murphy, W., Lawrence, J.A., 2010. Identifying the behavioural
- 575 characteristics of clay cliffs using intensive monitoring and geotechnical numerical modeling.
- 576 Geomorphology 120(3), 107-122.
- 577 RIEGL, 2007. Documentation technique du TLS (LMS-Z390i).
- 578 Robinson, L.A., 1977. Marine erosive processes at the cliff foot. Marine Geology 23(3), 257-
- 579 271.

- Rodet, J., 1991. Les karsts de la craie: étude comparative. Ph.D. Thesis, University of Paris,
 France.
- 582 Rosser, N.J., Petley, D.N., Lim, M., Dunning, S.A., Allison, R.J., 2005. Terrestrial laser
- scanning for monitoring the process of hard rock coastal cliff erosion. Quarterly Journal of
- 584 Engineering Geology and Hydrogeology 38(4), 363-375.
- Rosser, N.J., Lim, M., Petley, D.N., Dunning, S.A., Allison, R.J., 2007. Patterns of precursory
- rockfall prior to slope failure. Journal of Geophysical Research: Earth Surface 112(F4).
- 587 Rosser, N.J., Brain, M.J., Petley, D.N., Lim, M., Norman, E.C., 2013. Coastline retreat via
- progressive failure of rocky coastal cliffs. Geology 41(8), 939-942.
- 589 Rothmund, S., Niethammer, U., Malet, J.P., Joswig, M., 2013. Landslide surface monitoring
- 590 based on UAV- and ground-based images and terrestrial laser scanning: accuracy analysis
- and morphological interpretation. First Break 31(8).
- Rowlands, K.A., Jones, L.D., Whitworth, M., 2003. Landslide laser scanning: a new look at
- an old problem. Quarterly Journal of Engineering Geology and Hydrogeology 36(2), 155-157.
- Royán, M.J., Abellán, A., Jaboyedoff, M., Vilaplana, J.M., Calvet, J., 2014. Spatio-temporal
- analysis of rock fall pre-failure deformation using Terrestrial LiDAR. Landslides 11(4), 697-709.
- 597 Stephenson, W.J., Dickson, M.E., Trenhaile, A.S., 2013. Rock Coasts. In: Shroder, J.F.
- 598 (Ed.), Treatise on Geomorphology. Academic Press, San Diego, pp. 289-307.
- 599 Sturzenegger, M., Stead, D., 2009. Quantifying discontinuity orientation and persistence on
- 600 high mountain rock slopes and large landslides using terrestrial remote sensing techniques.
- Natural Hazards and Earth Systems Science 9(2), 267-287.
- 602 Sunamura, T., 1982. A wave tank experiment on the erosional mechanism at a cliff base.
- Earth Surface Processes and Landforms 7(4), 333-343.
- Sunamura, T., 1988. Beach morphologies and their change. In: Horikawa, K. (Ed.),
- Nearshore dynamics and coastal processes. University of Tokyo Press, Tokyo, pp. 136-161.
- Sunamura, T., 1992. Geomorphology of Rocky Coasts. John Wiley & Son, London.

- Terzaghi, K., 1950. Mechanism of landslides. In: Paige, S. (Ed.), Application of Geology to
 Engineering Practice (Berkeley Volume). Geological Society of America, Washington, pp. 82123.
- Teza, G., Galgaro, A., Zaltron, N., Genevois, R., 2007. Terrestrial laser scanner to detect
- landslide displacement fields: a new approach. International Journal of Remote Sensing
 28(16), 3425-3446.
- Teza, G., Pesci, A., Genevois, R., Galgaro, A., 2008. Characterization of landslide ground
 surface kinematics from terrestrial laser scanning and strain field computation.
- 615 Geomorphology 97(3), 424-437.
- Travelletti, J., Oppikofer, T., Delacourt, C., Malet, J.P., Jaboyedoff, M., 2008. Monitoring
- 617 landslide displacements during a controlled rain experiment using a long-range terrestrial
- 618 laser scanning (TLS). International Archives of the Photogrammetry, Remote Sensing and
- 619 Spatial Information Sciences 37(B5), 485-490.
- Travelletti, J., Delacourt, C., Malet, J.P., Allemand, P., Schmittbuhl, J., Toussaint, R., 2013.
- 621 Performance of Image Correlation Techniques for Landslide Displacement Monitoring. In:
- Margottini, C., Canuti, P., Sassa, K. (Eds.), Landslide Science and Practice. Springer, Berlin
 Heidelberg, pp. 217-226.
- Trenhaile, A.S., 1987. The geomorphology of rock coasts. Oxford University Press, Oxford.
- Trenhaile, A.S., Prestanski, K.J., Porter, N.I., Gagnon, J., 2013. Rock coast erosional forms
- (notches and shore platforms) on the La Paz Peninsula, Southern Baja, Mexico. Geo-Temas14, 1-4.
- Varnes, D.J., 1978. Slope movement types and processes. In: Schuster, R.L., Krizek, R.J.
- 629 (Eds.), Landslides: Analysis and Control. Transportation and Road Research Board, National
- Academy of Science, Washington, pp. 11-33.
- 631 Young, A.P., Flick, R.E., Gutierrez, R., Guza, R.T., 2009. Comparison of short-term seacliff
- retreat measurement methods in Del Mar, California. Geomorphology 112(3), 318-323.