

Changes in productivity and intermediate circulation in the northern Indian Ocean since the last deglaciation: new insights from benthic foraminiferal Cd/Ca records and benthic assemblage analyses

Ruifang Ma^{1,2*}, Sophie Sépulcre¹, Laetitia Licari³, Frédéric Haurine¹, Franck Bassinot⁴, Zhaojie Yu⁵,
Christophe Colin¹

¹Université Paris-Saclay, CNRS, GEOPS, 91405, Orsay, France.

²State Key Laboratory of Cryospheric Science, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, 730000, Lanzhou, China.

³CEREGE, Aix-Marseille Université-Europole de l'Arbois-BP80, 13545, Aix-en-Provence, France.

⁴LSCE/IPSIL, CEA CNRS UVSQ, UMR 8212, F-91190 Gif Sur Yvette, France.

⁵Key Laboratory of Marine Geology and Environment, Institute of Oceanology, Chinese Academy of Sciences, 266071, Qingdao, China.

Contents of this file

Figures S1-S4

Additional Supporting Information (Files uploaded separately)

Captions for Tables S1 to S2

Introduction

Figure S1. a) – b) The Net primary productivity distribution in the Northern Indian Ocean during January and July, respectively. Maps based on MODIS chlorophyll-a, SST, PAR satellite data, using the standard vertically Generalized Production Model (VGPM) (Behrenfeld and Falkowski, 1997) as the standard algorithm.

Fig. S2. Down core variations of PC1 and 2 scores, together with the percentages of major species in: (a-c) benthic assemblages 1, 2 and 3, respectively. The color-shaded intervals and abbreviations are the same as in Figure 2.

Fig. S3. Compilation of benthic $\delta^{13}\text{C}$ records obtained from MD77-191 (1254 m) (Ma et al., 2020), MD77-176 (1325 m) (Ma et al., 2019), RC12-344 (2140 m, Naqvi et al., 1994),

MD97-2120 (1210 m, Pahnke and Zahn, 2005), and Core 905 (1580 m, Jung et al., 2009). The color-shaded intervals and abbreviations are the same as in Figure 2.

Fig. S4. Compilation of benthic $\delta^{13}\text{C}$, PC 1 scores and Cd_w obtained from core MD77-176 (BoB). The color-shaded intervals and abbreviations are the same as in Figure 2.

Table S1. Sub-bottom depth, Cd/Ca values ($\mu\text{mol/mol}$) of benthic foraminifera species from core MD77-191 (*Hoeglundina elegans*, *Cibicidoides pachyderma* and *Uvigerina peregrina*, *Globobulimina* spp.) and core MD77-176 (*Hoeglundina elegans*).

Table S2. Down-core depth and quantities of the major benthic foraminiferal species at core MD77-191.




Fig. S1

79 a)




Fig. S2

89 b)




Fig. S2




Fig. S2




Fig. S3




Fig. S4