

***Pleurolocina* from the western Atlantic and eastern Pacific Oceans: a new intertidal species from Curaçao with unusual shell microstructure (Mollusca, Bivalvia, Lucinidae)**

Emily A. Glover¹, John D. Taylor¹

¹ Department of Life Sciences, The Natural History Museum, London SW7 5BD, UK

Corresponding author: John D. Taylor (j.taylor@nhm.ac.uk)

Academic editor: Richard Willan | Received 15 June 2016 | Accepted 5 August 2016 | Published 29 September 2016

<http://zoobank.org/4203049F-C7DE-4BCE-AFDA-05A2EBDF6434>

Citation: Glover EA, Taylor JD (2016) *Pleurolocina* from the western Atlantic and eastern Pacific Oceans: a new intertidal species from Curaçao with unusual shell microstructure (Mollusca, Bivalvia, Lucinidae). ZooKeys 620: 1–19. doi: 10.3897/zookeys.620.9569

Abstract

A new shallow water species of the lucinid bivalve *Pleurolocina* is described from Curaçao in the southern Caribbean Sea and compared with known species of the genus from the western Atlantic and eastern Pacific Oceans. Although confused with the Floridian species *P. leucocyma*, it is most similar to the eastern Pacific *P. undata*. As in all studied lucinids, the new species possesses symbiotic bacteria housed in the ctenidia. The shell microstructure is unusual with repeated and intercalated conchiolin layers that have sublayers of ‘tulip-shaped’ calcareous spherules. Predatory drillings by naticid gastropods frequently terminate at the conchiolin layers.

Keywords

Bacterial symbionts, Caribbean, conchiolin layers, defensive adaptation, Lucinidae, *Pleurolocina*

Introduction

The tropical and subtropical western Atlantic is one of the major centres of marine molluscan diversity and bivalves in the speciose family Lucinidae, with an estimated 46 species in this ocean, have been the focus of many studies since the discovery of their chemosymbiosis with sulphide-oxidising bacteria (e.g. Giere 1985, Fisher and Hand 1984, Frenkiel and Mouëza 1995, Frenkiel et al. 1996, Gros et al. 1998, 1998, 2012). Nonetheless, new species from both shallow and deep water are still discovered and new genera identified (Taylor and Glover 2009, Taylor et al. 2013). Additionally, within the area there are several cryptic species with narrower ranges nestled among supposedly widespread species (Huber 2015, Taylor and Glover submitted). Distributional data for western Atlantic lucinids indicates that although some are widespread, others have more restricted ranges. A recurring pattern is of congeneric pairs, one largely restricted to the Gulf of Mexico and Florida and the other with a more southerly Caribbean range as exemplified by *Lucinisca nassula* and *L. muricata* (Taylor and Glover submitted). This dual distribution is similar to that proposed by Petuch (1982) as a relict of the Caloosahatchee-Gatunian pattern dating from the Pliocene but possibly inherited by present day taxa. Additionally, in the eastern Pacific, there are lucinids closely similar morphologically and genetically to those of the western Atlantic and presumably separated by the rise of the Central American Isthmus around 3.5 mya. Examples of these are the pair *Radiolucina amianta* (Atlantic) and *R. cancellaris* (Pacific) (see Garfinkle 2012), and the pair *Ctena imbricatula* (Atlantic) and *Ctena mexicana* (Pacific) (Taylor et al. 2011).

Pleurolucina (Dall, 1901) is a genus of small lucinids characterised by broad radial ribs. The type species, *Lucina leucocyma* Dall, 1886, first described from off the Florida Keys, is documented as having a distribution from North Carolina to Colombia including Yucatan Peninsula (Britton 1970, Vokes and Vokes 1983, Huber 2015). Two other species, *P. hendersoni* Britton, 1972 and *P. sombreroensis* (Dall, 1886), are known from the western Atlantic (Britton 1972), while three further species are recorded from the Eastern Pacific (Coan and Valentich-Scott 2012). During field sampling in shallow seagrass around Curaçao in May 2015 we collected a *Pleurolucina* that we recognised as similar to, but likely distinct from, *P. leucocyma*. Further research showed this to be an undescribed species more widely distributed in the southern Caribbean and confounded with *P. leucocyma*. An apparent high incidence of failed naticid drill holes focused attention on the shell microstructure revealing intercalated organic layers. Thought to be related to *Lucina* or *Cavilinga* (Britton 1972, Bretsky 1976) and included by Taylor et al. (2011) in the subfamily Lucininae, no *Pleurolucina* species has previously been included in molecular analyses.

We describe this new *Pleurolucina* from Curaçao in comparison with other western Atlantic and Eastern Pacific species, detail its phylogenetic position and illustrate its unusual shell microstructure with calcified conchiolin layers.

Material and methods

Samples of the new species were collected in southern Curaçao – location below. Details of ctenidia and sperm were studied using critical point dried glutaraldehyde-fixed specimens. Shells, microstructure and anatomy were imaged using a Quanta FEI 650 FEG scanning electron microscope. Comparative shell material was studied in USNM and NHMUK.

Institutional abbreviations

FMNH	Field Museum of Natural History, Chicago, USA
MCZ	Museum of Comparative Zoology, Harvard University, USA
MNHN	Muséum national d'Histoire Naturelle, Paris, France
RMNH	Rijksmuseum van Natuurlijke Historie, Leiden, Netherlands
NHMUK	The Natural History Museum, London, UK
SBMNH	Santa Barbara Museum of Natural History, USA
USNM	United States National Museum of Natural History, USA

Other abbreviations

H	shell height
L	shell length
LV	left valve
PI	protoconch I length
PII	protoconch II length
RV	right valve
SEM	scanning electron microscopy
T	tumidity single valve

Systematics

Family Lucinidae Fleming, 1828

Subfamily Lucininae Fleming, 1828

Pleurolocina Dall, 1901

Dallucina Olsson & Harbison, 1953. Type species, by original designation, *Lucina* (*Here*) *amabilis* Dall, 1898. Pliocene, Florida. Gender feminine.

Type species. *Lucina leucocyma* Dall, 1886, by original designation. Recent, western Atlantic Ocean. Gender feminine.

Diagnosis. Shell small, L to 27 mm (*P. sombreroensis* usually less than 10 mm), subcircular to ovate, generally higher than long, inflated to highly inflated. Sculpture of 4–6 broad radial ribs separated by broad sulci, sometimes absent in adult shells, crossed by closely-spaced, often terraced, commarginal lamellae. Lunule deeply excavated to shallow. Ventral margin finely beaded. Hinge: RV with two cardinal teeth, posterior-most sometimes bifid, anterior and posterior lateral teeth present; LV with two cardinal teeth, anterior smaller, with anterior and posterior lateral teeth. Anterior adductor muscle scar relatively short, broad, separate from pallial line for about $\frac{1}{2}$ to $\frac{2}{3}$ of length, pallial line entire.

Included species. Western Atlantic: *P. leucocyma* (Dall, 1886), *P. hendersoni* Britton, 1972, *P. sombreroensis* (Dall, 1886). Eastern Pacific: *P. leucocymoides* (Lowe, 1935), *P. taylori* Coan & Valentich-Scott, 2012, *P. undata* (Carpenter, 1865).

Distribution. Western Atlantic: northern Florida to Brazil (*P. sombreroensis* Espirito Santo, Rios 1994). East Pacific: Baja California Mexico to Ecuador, Galapagos Islands (Coan and Valentich-Scott 2012).

Geological range. Early Oligocene to Recent. *Pleurolocina amabilis* (Dall, 1898) is a distinctive, laterally compressed species from the Late Pliocene to mid-Pleistocene of Florida. It was made type species of the new genus *Dallucina* by Olsson and Harbison (1954) but other than the lateral compression it is similar in most characters to *P. leucocyma*. From Miocene deposits of Ecuador Olsson (1964) described *Paslucina* with *Lucina (Paslucina) follis* Olsson, 1964 as type species. This has the shape and radial folds typical of *Pleurolocina* species and may be an antecedent.

Pleurolocina quadricostata (Dall, 1903) from the Pliocene Bowden Formation of Jamaica (Woodring 1925: 121, pl. 16, figs 4–6) resembles the living *P. leucocyma*. From the same deposit, *Phacoides (Linga) tithonis* (Dall, 1903) (Woodring 1925: 120, pl. 16, figs 2, 3) is similar to *P. sombreroensis*. A species described as *Lucina (Cavilinga) triloba* (Dockery 1982, pl. 19, fig 4) from the Early Oligocene, Vicksburg Group, Mississippi, USA, has characters of *Pleurolocina* but with only two radial folds. From the same deposits, *Lucina (Cavilinga) imbricolamella* Dockery (1982 pl. 20, figs 11–12) resembles the Recent *Pleurolocina sombreroensis*.

Relationships. From morphological characters of the shells, *Pleurolocina* species are usually regarded as being related to *Lucina* s.s. or *Cavilinga* (Britton 1972, Bretsky 1976). *Pleurolocina harperae* below is the only member of the genus yet to be included in molecular analyses and results (Taylor et al. submitted) show that it groups within the Lucininae, close to *Cavilinga blanda*, in a subclade of *Lucina* and *Divalinga* species.

Remarks. In the absence of molecular evidence, other than for *P. harperae*, our concept of *Pleurolocina* embraces a range of shell morphologies from species like *P. leucocyma*, *P. undata* and *P. taylori* that have prominent radial ribs, through the less ribbed *P. hendersoni* and *P. leucocymoides*, to the small *P. sombreroensis* that has a rounded shell lacking radial ribs. Nevertheless, they are all rather inflated with similar dentition, anterior adductor muscle scars and beaded inner margins.

***Pleurolucina harperae* n. sp.**

<http://zoobank.org/D9916BAC-D208-4A5B-8499-6FE1B5ADC3BB>

Figs 1–5

Lucina leucocyma: Daccarett and Bossio 2011: 177, fig. 1243.

Pleurolucina leucocyma: Huber 2015: 433, fig. p. 85.

Type material. *Holotype*: 1 whole shell L 8.8, H 8.5 T 3.2 mm (NHMUK 20160338), southwestern Curaçao, channel into Spaanse Water, opposite Hyatt Resort, 12°03'57" N 68°51'13" W. BivAToL stn Cur-5-15-009, 22 May 2015.

Paratypes: 92 valves (NHMUK 20160339), 2 paired valves (RMNH 5003991–50003992), 3 paired valves (FMNH344698), 2 paired valves (USNM 1411553). Same locality as holotype.

Other material. 19 ethanol preserved specimens (NHMUK), same locality as holotype.

Description. Shell subovate, slightly anteriorly extended, L to 9.6 mm, H to 9.7 mm, H/L 0.99, moderately inflated, sculpture of flat, closely spaced commarginal lamellae, with four prominent, broad ribs with interspaces variable in width, but always narrower than ribs themselves; microsculpture of tight rows of shallow pits (Fig. 1 P). Umbones low, situated on midline. Anterior dorsal area arcuate. Protoconch: PI 217 µm, PI + PII 228 µm, PII a narrow rim with fine increments (Fig. 1 O). Lunule short, semicircular, slightly impressed. Ligament short, set in shallow resilifer. Hinge teeth: LV with two cardinal teeth; a robust anterior lateral tooth and smaller posterior lateral. RV with a single large cardinal tooth and anterior and posterior lateral teeth. Anterior adductor muscle scar short, broad, widely divergent from pallial line (60–70 µm) for about half of length (Fig. 2 A), posterior scar ovate; pallial line entire, pallial blood vessel scar sometimes visible. Shell margin finely beaded, sinuate with anterior sinus deeper. Shell within pallial line often patchily eroded to expose inner shell layers. Colour grey-white.

Anatomy. General anatomy resembles most other described lucinids (Fig. 3). Mantle fusion ventral to the posterior apertures is very short. Foot short and broad when retracted but can be vermiform when extended (Fig. 3 A) with a small heel. Visceral pouches absent. Distinct mantle gills are absent but the inner mantle ventral to the anterior adductor muscle is thickened (Fig. 3 C) and may be a respiratory area with blood space as seen in other lucinids (Taylor and Glover 2000). Labial palps are very short. In common with all other studied Lucinidae, *P. harperae* has ctenidia comprising inner demibranchs only; these were pink in life, large, thick and occupying much of the mantle cavity (Fig. 3 B). Ctenidial filaments are approx. 40 µm thick and 380 µm deep with a narrow 45 µm ciliated zone and a deep bacteriocyte zone (Fig 3 D). Bacteriocytes were packed with 'potato-shaped' bacteria 3–5 µm long and 1.5–2.0 µm wide (Figs 3 G, H). The surface of the microvilli-covered bacteriocytes and intercalary cells were colonised by abundant spirochaetes 2.5 µm long and 0.2 µm wide (Fig. 3 F) similar to those re-




Figure 1. *Pleurolocina harperae* sp. n. **A–C** Holotype NHMUK 20160338 exterior of right and interior of right and left valves L 8.8 mm. **D–P** Paratypes. NHMUK 20160339 dorsal view L 7.6 mm. **E** Exterior of left valve L 7.7 mm. **F** Interior of right valve L 6.3 mm. **G** Exterior of right valve L 7.9 mm. **H** Hinge area of right valve L 8.6 mm. **I** Exterior of left valve L 63 mm. **J, K** Interiors of right and left valves L 5.0 mm. **L** Dorsal view showing lunule. Scale bar = 0.5 mm. **M, N** Details of hinge teeth of J, K. **O** Protoconch. Scale bar = 100 μ m. **P** Detail of microsculpture. Scale bar = 20 μ m.




Figure 2. Outline drawings of shell interiors of **A** *P. harperae* and **B** *P. leucocyma*.

ported by Ball et al. (2009) from *Euanodontia ovum* (Reeve, 1850). In comparison the symbiotic bacteria of *Clathrolucina costata* collected at the same time and same habitat were longer and rod shaped, 8–10 μm in length and approx. 1 μm wide.

The sperm of *P. harperae* were 9 μm long and 1.2 μm wide at the base, tapering and curved distally (Figs 3 J, K). From the same locality, sperm of *Clathrolucina costata* were shorter, 4.8–5 μm and 1–1.2 μm wide with blunt tips. Oocytes of *P. harperae* were approx. 200 μm in diameter (Fig. 3 I). Comparative sperm data is available for a few other western Atlantic lucinids (Bigatti et al. 2004); sperm of *Codakia orbicularis* were 14–15 μm long, tapering with a width of 0.8 μm ; *Ctena orbiculata* were cylindrical, slightly curved, 7.5 μm long and 1–1.2 μm wide at base and *Lucina pensylvanica* were 15.5 μm long, with curved tapering heads and 1.1 μm wide at the posterior.

Shell microstructure. Within a very thin (ca 1 μm) periostracum, *Pleurolocina harperae* has a basic four layered shell (Figs 4 A,B); an outer composite prismatic layer, followed inwards by a thin crossed-lamellar layer, then a thicker layer of irregular spherulitic prisms and within the pallial line a complex crossed-lamellar layer with sublayers of irregular prisms. The shell layers are interrupted by sheets of conchiolin around 20–90 μm in thickness, each with repeated sublayers of small discrete ‘tulip-shaped’ calcified spherulites approx. 5 μm in diameter (Figs 4 D, F). Each spherulite is joined to those of the layer below with a narrow (0.5 μm) semicalcified channel through the conchiolin (Figs 4 E, F). At the shell surface, the conchiolin sheets correspond to major depositional halts (Fig. 4 A) visible as notches in the shell with the conchiolin appearing contiguous with the invaginated periostracum. In each shell there may be between 1–5 of such sheets.

Drill holes in *Pleurolocina harperae* produced by predatory naticid gastropods were observed with full penetration in 14 out of 114 single valves, but with 12 records of




Figure 3. *Pleurolucina harperae*, general anatomy, ctenidia, bacteria, oocytes and sperm. **A** Right side, with mantle removed, right demibranch and extended foot stained with methylene blue L 7 mm **B** Left demibranch and foot, critical point dried preparation. Scale bar = 1 mm **C** Cut section to show general anatomy, stained with methylene blue L 8 mm **D** Transverse section through single ctenidial demibranch. Scale bar = 100 μm **E** Surface of bacteriocytes and intercalary cells on lateral view of a ctenidial filament. Scale bar = 15 μm **F** Spirochaete bacteria on surface of bacteriocytes. Scale bar = 2 μm **G, H** Symbiotic bacteria contained in bacteriocyte. Scale bar = 5 μm **I** Developing oocytes. Scale bar = 500 μm **J, K** Sperm. Scale bars = 5, 2 μm respectively. **aa** anterior adductor muscle **bz** bacteriocyte zone **cz** ciliated zone **dg** digestive gland **f** foot **lp** labial palps **me** mantle edge **ov** ovary with oocytes **pa** posterior adductor **r** rectum **rd** right demibranch **st** stomach **tm** thickened mantle ventral to anterior adductor muscle.




Figure 4. Shell microstructure of *Pleurolocina harperae*. **A** Fractured section of shell margin showing major notch growth halt and conchiolin layer. Scale bar = 400 μm **B** Fractured section showing succession of shell layers. Shell exterior at top. Scale bar = 100 μm **C** Conchiolin layer with regular bands of spherulites. Scale bar = 40 μm **D** Individual spherulite. Scale bar = 2 μm **E** Adjacent spherulites embedded in conchiolin with narrow channels between layers. Scale bar = 5 μm **F** Single spherulites with channels below and above. Scale bar = 5 μm . **cl** crossed lamellar layer **co** conchiolin layer **cp** composite prismatic layer **ip** irregular prismatic layer **p** periostracum **sp** spherulitic prismatic layer.




Figure 5. Failed and multiple drill holes in shells of *Pleurolucina harperae*. **A** L = 6.8 mm **B** L = 9.8 mm **C** L = 7.8 mm **D** SEM of failed drill hole terminating at conchiolin layer. Scale bar = 1.0 mm.

incomplete drill holes that terminated at an internal conchiolin layer (Fig. 5). In one shell there were three failed drills and in another two failures before successful penetration. Incidences of apparent multiple completed drill holes in dead shells may have resulted from post-mortem degradation of organic layers in failed drill holes.

Similar conchiolin calcified sheets were identified in *Pleurolucina hendersoni* (Figs 6 A, B) and *P. undata* (Figs 6 C–E) but not in *P. leucocyma* (2 shells examined) or *P. sombreroensis* (2 shells examined). Also conchiolin sheets with multiple layers of calcareous spherules were observed in *Lucina pensylvanica* from the Florida Keys (Figs F,G), apparently confined to the inner shell layer within the pallial line. This is distinct from the calcified periostracum of this species (Fig. 6 H) as described by Taylor et al. (2004). No conchiolin sheets were observed in a single *Cavilinga blanda* examined. For comparison, the repeated conchiolin sheets reported in *Cardiolumina* species by Ishikawa and Kase (2007) were studied in *C. quadrata* from the Philippines. These sheets were approx. 10–15 μm thick and only lightly calcified with sporadic spherulitic crystal aggregations (Figs 6 I–K) with no multiple sub-layers.




Figure 6. Shell microstructure of other species *Pleurolocina hendersoni*, *P. undata*, *Lucina pensylvanica* and *Cardiolucina quadrata*. **A** *Pleurolocina hendersoni* Guadeloupe, fractured section with prominent calcified conchiolin layer, periostracum at base. Scale bar = 20 μm **B** *P. hendersoni*, detail of conchiolin layer with lines of calcareous spherulites. Scale bar = 20 μm **C** *Pleurolocina undata* Baja California, fractured section with thin conchiolin layer. Scale bar = 200 μm **D** *P. undata*, detail of conchiolin layer with spherulites. Scale bar = 20 μm **E** *P. undata*, single spherulites embedded in conchiolin. Scale bar = 3 μm **F** *Lucina pensylvanica* Florida Keys, calcified conchiolin layer. Scale bar = 20 μm **G** *L. pensylvanica*, single spherulite. Scale bar = 2 μm **H** *L. pensylvanica*, section of periostracum with calcareous granules. Shell interior to top. Scale bar = 20 μm **I** *Cardiolucina quadrata* Philippines, fractured section with conchiolin layer. Scale bar = 200 μm **J** *C. quadrata* detail of conchiolin layer with calcareous aggregates. Scale bar = 50 μm **K** *C. quadrata* detail of calcareous aggregate. Scale bar = 10 μm .




Figure 7. *Pleurolucina leucocyma*. **A–C** *Lucina leucocyma* Dall, 1881 lectotype MCZ 7986, exterior, interior and dorsal view of right valve, L 5.7 mm, H 6.6 mm **D, E** *Lucina leucocyma* paralectotype USNM 83140, exterior of left valve and interior of right valve, L 4.8 mm, H 5.5 mm **F–K**, *Pleurolucina leucocyma* USNM 446563 Eolis Station 368, off Ajax Reef, Florida **F** Exterior of left valve, L 5.1 mm **G** Left valve of juvenile shell, L 3.1 mm **H** Lateral view of left valve, L 5.1 mm **I** Interior of left valve, L 5.5 mm **J** Interior of right valve, L 5.5 mm **K** Protoconch, scale bar = 100 µm.

Habitat. *Pleurolucina harperae* is an intertidal to shallow subtidal species collected from sand amongst seagrass rhizomes (largely *Thalassia testudinum*, *Halodule* sp.) in contrast to *P. leucocyma* that is usually recorded from deeper water, for example 30–180 m around the Florida Keys (Britton 1970). Records of *P. harperae* from Atlantic Panama (USNM below) are also from shallow water seagrass habitats. At Curaçao it co-occurred with several other lucinid species: *Clathrolucina costata* (d’Orbigny, 1845), *Ctena imbricatula* (C.B. Adams, 1845), *Anodontia alba* Link, 1807, *Codakia orbicularis* (Linnaeus, 1758), *Lucina roquesana* J. & W. Gibson-Smith, 1982 and *Divalinga quadrisulcata* (d’Orbigny, 1845).




Figure 8. Bivariate height/length plots comparing *P. harperae* with *P. leucocyma*, and *P. undata*. Length and height in millimetres.

Distribution. Southern Caribbean: Panama (USNM 759784; 620716, 759825) Colombia -Taganga (Daccarett and Bossio 2011), Curaçao. The distribution of *Pleurolucina harperae* in the southern Caribbean is uncertain but it may be restricted to the southwestern area. There have been no records from the Antilles and intensive sampling of molluscs around Guadeloupe by Muséum national d'Histoire Naturelle (KARUBENTHOS 2012, 2015) recorded only *P. hendersoni* and *P. sombreroensis* (Taylor and Glover submitted). Similarly, only *P. sombreroensis* was recorded from a recent survey of the marine molluscan fauna of French Guiana (MNHN - GUYANE 2014).

Etymology. Named for Elizabeth (Liz) Harper, University of Cambridge, bivalve researcher, colleague and friend, who helped collect the new species.

Comparison with other species. *Pleurolucina leucocyma* (Fig. 7) was thought to be widespread across the tropical Western Atlantic but we now consider it to be restricted to Florida and the Gulf of Mexico with the southern Caribbean records representing *Pleurolucina harperae*. The new species differs from *P. leucocyma* (mean L 6.2 mm, H 7.4 mm, H/L 1.13) in being larger, less inflated and usually longer than high in the adult (Fig. 8). The radial folds are usually lower and the anterior adductor muscle scar is shorter and more divergent from the pallial line (Fig. 2 B). In shape and sculpture, it is most similar to the somewhat larger *Pleurolucina undata* (Figs 9 E-G) (mean L 15.1 mm, H 15 mm, H/L 0.95) from the eastern Pacific, Gulf of California, intertidal zone to 60 m (Coan and Valentich-Scott 2012).

Other less similar species are: *P. hendersoni* (Figs 9 A, B) an offshore to deep water species (to 300 m) from the southern Caribbean (Cuba, Lesser Antilles) that reaches




Figure 9. Other *Pleurolucina* species. **A, B** *Pleurolucina hendersoni* Britton, 1972, exterior and interior of left valve Guadeloupe station GD 69 (MNHN), L 9.1 mm **C, D** *Pleurolucina sombrerensis* (Dall, 1886) exterior of left valve (L 4.9 mm) and interior of right valve (L 5.2 mm), USNM 446178, Eolis stn 48, off Miami, Florida, 110 m **E–G** *Pleurolucina undata* (Carpenter, 1865) exterior of left valve and interiors of right and left valves, NUMUK 1915.15.273 ‘California’, L 11.0 mm **H–J** *Pleurolucina leucocymoides* (Lowe, 1935), exterior of right valve and interiors of right and left valves SBMNH 141511, Baja California, NE of Isla Danzante, Mexico, L 11 mm **K–M** *Pleurolucina taylora* Coan & Valentich-Scott, 2012, holotype, exterior of left valve and interior of left and right valves, SBMNH 149647, Baja California, Los Frailes, Mexico, L 9.5 mm.

about 12 mm in length and resembles the eastern Pacific *P. leucocymoides*. Compared with other *Pleurolucina*, the sculpture of broad radial folds is less pronounced and the commarginal lamellae are widely spaced and prominent. *Pleurolucina sombrerensis* (Figs 9 C, D) lives in deeper water to 200 m from the Florida Keys to Brazil. The shell reaches about 6–7 mm in length and is rounded in outline, with a shallow radial anterior sulcus and prominent close commarginal lamellae, sometimes separated by deep interspaces. It does not closely resemble other *Pleurolucina* but shares some shell features including dentition and adductor scar shape. The larger *P. leucocymoides* (Figs 9 H–J) is known from shallow water to 150 m and ranges from Baja California to Ec-

cuador and Galapagos Islands. The sculpture of broad prominent commarginal lamellae and absence of prominent radial folds distinguish it from other *Pleurolocina*. Lastly, *P. taylori* (Figs 9 K–M) is known from the intertidal zone to 183 m in the Gulf of California; it is distinguished by the highly inflated shell and closely spaced, low commarginal lamellae with four to five radial folds and resembles the extinct late Pliocene – mid-Pleistocene Floridian species *P. amabilis*.

Discussion

Pleurolocina is a genus of seven living species from the tropical to subtropical western Atlantic and eastern Pacific with none recognised from the eastern Atlantic or Indo-West Pacific. In that respect, it is similar to *Radiolucina* (Garfinkle 2012) and *Lucinisca* that share similar distributions. In the western Atlantic, the most similar species to the southern Caribbean *Pleurolocina harperae* is *P. leucocyma* from Gulf of Mexico and Florida. This distributional pattern of northern and southern species pairs is seen in *Ctena* (*C. orbiculata* and *C. imbricatula*), *Lucinisca* (*L. nassula* and *L. muricata*) and *Lucina* (*L. pennsylvanica* and *L. roquesana*) (see Taylor and Glover submitted). Cognate pairs of bivalves have been recognised from morphology and/or molecules on either side of the central American Isthmus (Marko 2002, Marko and Moran 2009). Although molecular confirmation is lacking, *Pleurolocina harperae* is similar in shell form to *P. undata*, *P. hendersoni* resembles *P. leucocymoides* and perhaps *P. leucocyma* is a sister species to *P. taylori*.

An interesting and unusual feature of *Pleurolocina harperae* is the repeated conchiolin sheets that are calcified with layers of embedded spherules. A model of conchiolin sheet formation in another lucinid genus, *Cardiolucina*, was proposed by Ishikawa and Kase (2007 fig. 7). Periodically, normal shell secretion of outer, middle and inner shell layers stops and a conchiolin sheet is secreted across the inside of the shell from the margin and extending within the pallial line. This break in normal calcification is marked by a distinct notch at the shell surface. Calcification then resumes with secretion of normal shell layers. Conchiolin layer formation in *Pleurolocina* is essentially similar but each layer is thicker with repeated sublayers of aragonitic spherules. The narrow channels linking successive spherule layers suggest some sort of original tissue connection to the cells of the mantle surface.

Conchiolin layers within the shell have been recorded in several bivalve families but those in the Corbulidae have attracted most attention because of the supposed resistance to predation by drilling gastropods evidenced by the high incidence of failed borings that terminate at the organic layers (e.g. Lewy and Samtleben 1979, Harper 1994). Alternatively, organic layers may enhance resistance to shell dissolution, endolithic organisms or shell breakage (Anderson 1992, Harper 1994, Kardon 1998). In contrast to *Pleurolocina* where the conchiolin layers are secreted episodically, the layers in Corbulidae are secreted continuously as a sublayer of normal shell formation. In *Corbula gibba* the conchiolin layer is calcified with cone-shaped spher-

ules approx. 8 μm in diameter (Lewy and Samtleben 1979 figs 5A–F). The organic layers of *Pleurolocina harperae* are similar in position and mode of formation to those recorded for species of *Cardiolumina* (Ishikawa and Kase 2007), but are much more highly calcified. *Cardiolumina* spp also show a high incidence of multiple drill holes with many terminating at the organic layers (Ishikawa and Kase 2007). *Pleurolocina* and *Cardiolumina* are not closely related among the Lucininae and the occurrence of conchiolin layers in other lucinids seems to be sporadic and certainly absent in many genera although no comprehensive study has been made. Nonetheless, calcified conchiolin layers do occur in some individuals of *Lucina pensylvanica* that is more closely related to *Pleurolocina*. It is tempting to regard the conchiolin layers as an adaptation conferring some resistance to shell drilling predation but, as argued in the case of *Corbula* (e.g. Kardon 1998), the layers may be an exaptation having first developed with some other function such as resistance to shell dissolution or enhancement of mechanical strength.

Acknowledgements

The field workshop in Curaçao was organized by Rüdiger Bieler as part of the Bivalve Assembling the Tree-of-Life project (bivatol.org) and supported by the U.S. National Science Foundation (NSF) Assembling the Tree of Life (AToL) program (award DEB-0732854 to RB and colleagues). We thank our colleagues at the workshop for logistic support especially Liz Harper and André Sartori for help with digging and sieving. Ellen Strong (USNM) kindly hosted our visit to the collections and provided images of the paralectotype of *L. leucocyma* and loan of other specimens. We thank Gonzalo Giribet and Adam Baldinger (MCZ) for images of the lectotype of *P. leucocyma* and Paul Valentich-Scott (Santa Barbara Museum of Natural History) generously allowed us to use images of *P. leucocymoides* and *P. taylori*. We are also grateful to Chiho Ikebe (NHMUK) for help with molecular sequencing and some micro-images. We appreciate the careful attention of Editor Richard Willan and reviewer André Sartori.

References

- Anderson LC (1992) Naticid predation on corbulid bivalves: effects of physical factors, morphological features, and statistical artifacts. *Palaios* 7: 602–620. doi: 10.2307/3514872
- Ball AD, Purdy K, Glover EA, Taylor JD (2009) Ctenidial structure and multiple bacterial symbionts in *Anodontia* (*Euanodontia*) *ovum* (Reeve, 1850) from the Great Barrier Reef, Australia (Bivalvia: Lucinidae). *Journal of Molluscan Studies* 75: 175–185. doi: 10.1093/mollus/eyp009
- Bigatti G, Peharda M, Taylor JD (2004) Size at first maturity, oocyte envelopes and external morphology of sperm in three species of Lucinidae (Mollusca: Bivalvia) from Florida Keys, USA. *Malacologia* 46: 417–426.

- Bretsky SS (1976) Evolution and classification of the Lucinidae (Mollusca; Bivalvia). *Palaeontographica Americana* 8(50): 219–337.
- Britton JC (1970) The Lucinidae (Mollusca: Bivalvia) of the Western Atlantic Ocean. Unpublished PhD dissertation, George Washington University. University Microfilms 71–12, 288, 566 pp.
- Britton JC (1972) Two new species and a new subgenus of Lucinidae (Mollusca: Bivalvia), with notes on certain aspects of lucinid phylogeny. *Smithsonian Contributions to Zoology* 129: 1–19. doi: 10.5479/si.00810282.129
- Carpenter PP (1865) Diagnoses of new species of mollusks, from the west tropical region of North America, principally collected by the Rev. J. Rowell, of San Francisco. *Proceedings of the Zoological Society of London for the year 1865*: 278–282.
- Coan EV, Valentich-Scott P (2012) Bivalve seashells of tropical west America. Marine bivalve mollusks from Baja California to northern Peru. Santa Barbara Museum of Natural History, Santa Barbara, Part 1, 598 pp.
- Daccarett EY, Bossio VS (2011) Colombian Seashells from the Caribbean Sea. *Informatore Piceno*, Ancona, Italy, 384 pp.
- Dall WH (1886) Reports on the results of dredging, under the supervision of Alexander Agassiz, in the Gulf of Mexico and in the Caribbean Sea, 1877–79, by the United States Coast Survey steamer “Blake,” Lieutenant-Commander G.D. Sigsbee, U.S.N. and Commander J.R. Bartlett, U.S.N., commanding. XXIX. Report on the Mollusca. Part 1, Brachiopoda and Pelecypoda. *Bulletin of the Museum of Comparative Zoology* 12: 171–318.
- Dall WH (1889) A preliminary catalogue of the shell-bearing marine mollusks and brachiopods of the southeastern coast of the United States, with illustrations of many of the species. *Bulletin of the US National Museum* 37: 1–221.
- Dall WH (1898–1903) Contribution to the Tertiary fauna of Florida with especial reference to the Miocene Silex beds of Tampa and the Pliocene beds of the Caloosahatchee River. *Wagner Free Institute of Science of Philadelphia Transactions* 3(4): 571–947 [1898]; 3(6): 1219–1654 [1903].
- Dall WH (1901) Synopsis of the Lucinacea and of the American species. *Proceedings of the United States National Museum* 23: 779–833. doi: 10.5479/si.00963801.23-1237.779
- Dockery DT (1982) Lower Oligocene Bivalvia of the Vicksburg Group in Mississippi. *Mississippi Department of Natural Resources Bureau of Geology, Bulletin* 123: 1–261.
- Fisher MR, Hand SC (1984) Chemautotrophic symbionts in the bivalve *Lucina floridana* from seagrass beds. *Biological Bulletin, Woods Hole* 167: 445–459. doi: 10.2307/1541289
- Fleming J (1828) A history of British animals, exhibiting the descriptive characters and systematic arrangement of the genera and species of quadrupeds, birds, reptiles, fishes, Mollusca and Radiata of the United Kingdom; including the indigenous, extirpated and extinct kinds; together with periodical and occasional visitants. Bell & Bradfute, Edinburgh, 565 pp.
- Frenkiel L, Mouëza M (1995) Gill ultrastructure and symbiotic bacteria in *Codakia orbicularis* (Bivalvia, Lucinidae). *Zoomorphology* 115: 51–61. doi: 10.1007/BF00397934
- Frenkiel L, Gros O, Mouëza M (1996) Gill structure in *Lucina pectinata* (Bivalvia: Lucinidae) with reference to hemoglobin in bivalves with symbiotic sulphur-oxidising bacteria. *Marine Biology* 125: 511–524.

- Garfinkle EAR (2012) A review of North American Recent *Radiolucina* (Bivalvia, Lucinidae) with the description of a new species. *ZooKeys* 205: 19–31. doi: 10.3897/zookeys.205.3120
- Giere O (1985) Structure and position of bacterial endosymbionts in the gill filaments of Lucinidae from Bermuda (Mollusca, Bivalvia). *Zoomorphology* 105: 296–301. doi: 10.1007/BF00312060
- Gros O, Frenkiel L, Mouëza M (1998) Gill filament differentiation and experimental colonization by symbiotic bacteria in aposymbiotic juveniles of *Codakia orbicularis* (Bivalvia: Lucinidae). *Invertebrate Reproduction and Development* 34: 219–231. doi: 10.1080/07924259.1998.9652656
- Gros O, Elisabeth NH, Gustave SDD, Caro A, Dubilier N (2012) Plasticity of symbiont acquisition throughout the life cycle of the shallow-water tropical lucinid *Codakia orbiculata* (Mollusca: Bivalvia). *Environmental Microbiology* 14: 1584–1595. doi: 10.1111/j.1462-2920.2012.02748.x
- Harper EM (1994) Are conchiolin sheets in corbulid bivalves primarily defensive? *Palaeontology* 37: 551–578.
- Huber M (2015) *Compendium of bivalves 2*. ConchBooks, Harxheim, Germany, 907 pp.
- Ishikawa M, Kase T (2007) Multiple predatory drill; holes in *Cardiolucina* (Bivalvia: Lucinidae): effect of conchiolin sheets in predation. *Palaeogeography Palaeoclimatology Palaeoecology* 254: 508–522. doi: 10.1016/j.palaeo.2007.07.004
- Kardon G (1998) Evidence from the fossil record of an antipredatory exaptation: conchiolin layers in corbulid bivalves. *Evolution* 5: 68–79. doi: 10.2307/2410921
- Lewy Z, Samtleben C (1979) Functional morphology and paleontological significance of the conchiolin layers in corbulid pelecypods. *Lethaia* 12: 341–351. doi: 10.1111/j.1502-3931.1979.tb01019.x
- Lowe HN (1935) New marine Mollusca from west Mexico, together with a list of shells collected at Punta Penasco, Mexico. *San Diego Society of Natural History, Transactions* 8: 15–34.
- Marko PB (2002) Fossil calibration of molecular clocks and the divergence times of geminate species pairs separated by the Isthmus of Panama. *Molecular Biology and Evolution* 19: 2005–2021. doi: 10.1093/oxfordjournals.molbev.a004024
- Marko PB, Moran AL (2009) Out of sight, out of mind: high cryptic diversity obscures the identities and histories of geminate species in the marine bivalve subgenus *Acar*. *Journal of Biogeography* 36: 1861–1880. doi: 10.1111/j.1365-2699.2009.02114.x
- Mikkelsen PM, Bieler R (2007) *Seashells of Southern Florida. Living marine molluscs of the Florida Keys and adjacent regions. Bivalves*. Princeton University Press, Princeton, 503 pp.
- Olsson AA (1964) Neogene mollusks from northwestern Ecuador. *Paleontological Research Institute, Ithaca, New York*, 256 pp.
- Olsson AA, Harbison A (1953) Pliocene Mollusca of southern Florida. *The Academy of Natural Sciences of Philadelphia Monograph* 8: 1–457, 65 pls.
- Olsson AA, McGinty TL (1958) Recent marine mollusks from the Caribbean coast of Panama with the description of some new genera and species. *Bulletin of American Paleontology* 39: 5–58.
- Petuch EJ (1982) Paraprovincialism: remnants of paleoprovincial boundaries in Recent molluscan provinces. *Proceedings of the Biological Society of Washington* 95: 774–780.

- Redfern C (2013) Bahamian seashells: 1161 species from Abaco, Bahamas. Bahamianseashells.com, Inc., Boca Raton, Florida, 501 pp.
- Rios EC (1994) Seashells of Brazil (2nd edn). Fundação Universidade do Rio Grande, Museu Oceanográfico, Rio Grande, 368 pp.
- Taylor JD, Glover EA (2000) Functional anatomy, chemosymbiosis and evolution of the Lucinidae. In: The Evolutionary Biology of the Bivalvia. Special Publication of the Geological Society of London 177: 207–225. doi: 10.1144/GSL.SP.2000.177.01.12
- Taylor JD, Glover EA (2009) New lucinid bivalves from hydrocarbon seeps of the Western Atlantic (Mollusca: Bivalvia: Lucinidae). *Steenstrupia* 30: 127–140.
- Taylor JD, Glover EA (submitted) Lucinid bivalves of Guadeloupe: diversity and systematics, with a critical assessment of other western Atlantic species (Mollusca: Bivalvia: Lucinidae). *Zootaxa*.
- Taylor JD, Glover EA, Peharda M, Bigatti G, Ball A (2004) Extraordinary flexible shell sculpture; the structure and formation of calcified periostracal lamellae in *Lucina pensylvanica* (Bivalvia: Lucinidae). *Malacologia* 46: 277–294.
- Taylor JD, Glover EA, Smith L, Dyal P, Williams ST (2011) Molecular phylogeny and classification of the chemosymbiotic bivalve family Lucinidae (Mollusca: Bivalvia). *Zoological Journal of the Linnean Society* 163: 15–49. doi: 10.1111/j.1096-3642.2011.00700.x
- Taylor JD, Glover EA, Smith L, Ikebe C, Williams ST (submitted) New molecular phylogeny of Lucinidae: increased taxon base with focus on tropical western Atlantic species (Mollusca: Bivalvia). *Zootaxa*.
- Taylor JD, Glover EA, Williams ST (2013) Taxonomy and phylogeny of western Atlantic Lucinidae: new genus for *Lucina costata* d'Orbigny, 1846, a new species of *Ferrocina* and neotype designation for *Venus orbiculata* (Montagu, 1808). *The Nautilus* 127: 131–146.
- Vokes HE, Vokes EH (1984) Distribution of shallow-water marine Mollusca, Yucatan Peninsula, Mexico. Mesoamerican Ecology Institute Monograph 1. Middle American Research Institute Publication 54: 183 pp.
- Woodring WP (1925) Contributions to the geology and palaeontology of the West Indies. Miocene molluscs from Bowden, Jamaica. Pelecypods and scaphopods. Carnegie Institute Washington Publication 366, 222 pp.