

S1 Table List of the species recorded for each of the three major fish families caught by recreational fishermen around the Cerbère-Banyuls natural marine reserve (see Fig 1). Trophic level and maximum length (in cm) from www.fishbase.org.

Family	Sp. latin name	Sp. common name	Trophic level	Max. length
Sparidae	<i>Boops boops</i>	bogue	2.8	40
	<i>Dentex dentex</i>	common dentex	4.5	100
	<i>Diplodus annularis</i>	annular seabream	3.6	28
	<i>Diplodus cervinus</i>	zebra seabream	3.0	55
	<i>Diplodus puntazzo</i>	sheephead bream	3.2	60
	<i>Diplodus sargus</i>	white seabream	3.4	45
	<i>Diplodus vulgaris</i>	two-banded seabream	3.5	45
	<i>Lithognathus mormyrus</i>	sand steenbras	3.4	55
	<i>Oblada melanura</i>	saddled seabream	3.4	37
	<i>Pagellus acarne</i>	axillary seabream	3.8	36
	<i>Pagellus bogaraveo</i>	blackspot seabream	4.2	70
	<i>Pagellus erythrinus</i>	common pandora	3.5	60
	<i>Pagrus pagrus</i>	common seabream	3.9	91
	<i>Sarpa salpa</i>	cow bream	2.0	51
	<i>Sparus aurata</i>	gilt-head bream	3.7	70
<i>Spondyliosoma cantharus</i>	black seabream	3.3	60	
Serranidae	<i>Anthias anthias</i>	swallowtail seaperch	3.8	27
	<i>Serranus cabrilla</i>	comber	3.4	40
	<i>Serranus scriba</i>	painted comber	3.8	36
Labridae	<i>Coris julis</i>	Mediterranean rainbow wrasse	3.4	30
	<i>Labrus merula</i>	brown wrasse	3.6	45
	<i>Labrus mixtus</i>	cuckoo wrasse	3.9	40
	<i>Labrus viridis</i>	green wrasse	3.9	47
	<i>Symphodus ocellatus</i>	ocellated wrasse	3.5	12
	<i>Symphodus tinca</i>	East Atlantic peacock wrasse	3.3	44