Spatial Variability of Organic Matter and Phosphorus Cycling in Rhône River Prodelta Sediments (NW Mediterranean Sea, France): a Model-Data Approach

Ait Ballagh Fatima Ezzahra ^{1,2}, Rabouille Christophe ^{1,*}, Andrieux-Loyer Françoise ³,
Soetaert Karline ⁴, Lansard Bruno ¹, Bombled Bruno ¹, Monvoisin Gael ⁵, Elkalay Khalid ²,
Khalil Karima ²

¹ Violaine Laboratoire des Sciences du Climat et de l'Environnement, Laboratoire Mixte CEA-CNRS-UVSQ, IPSL et Université Paris-Saclay, CEA-Orme des Merisiers, 91191, Gif sur Yvette, France

² Laboratory of Applied Sciences for the Environment and Sustainable Development, Essaouira School of Technology, Cadi Ayyad University, Km 9, Route d'Agadir, BP. 383, Essaouira Aljadida, Morocco

³ Ifremer, DYNECO PELAGOS, ZI Pointe du Diable, 29280, Plouzané, France

⁴ Department of Estuarine and Delta Systems, Royal Netherlands Institute for Sea Research (NIOZ) and Utrecht University, P.O. Box 140, 4400 AC, Yerseke, The Netherlands

⁵ GEOPS, Université Paris-Saclay et CNRS, Campus d'Orsay, Orsay, France

* Corresponding author : Christophe Rabouille, email address : rabouille@lsce.ipsl.fr

Abstract :

The Mediterranean Sea (MS) is a large oligotrophic sea whose productivity is sensitive to riverine nutrient inputs. More specifically, phosphorus (P) river supply is crucial for the MS, with an important role of the estuarine/deltaic filter especially for the storage and recycling in sediments. A benthic dataset from the Rhône River prodelta was used to derive P budgets, by means of an early diagenetic model including the benthic P cycle. The model was fitted to pore water profiles of oxygen, nitrate, sulfate, dissolved inorganic carbon, ammonium, oxygen demand units, dissolved inorganic phosphorus (DIP) and solid data (organic carbon (OC), Fe-bound P, Ca-bound P and organic P). Results indicated that the intensity of biogeochemical processes occurring below the sediment–water interface decreased from the river mouth to the adjacent continental shelf with decreasing integrated rates of OC mineralization (160–10 mmol m⁻² day⁻¹). The organic P mineralization was intense near the river mouth and decreased offshore (1196–80 μmol m⁻² day⁻¹). Its contribution to DIP release was large (> 90%). Fe-bound P had a key role in transferring P to deeper layers. These deltaic sediments played an important role as a source of regenerated DIP. A significant part of DIP was recycled to the overlying waters (72–94%), representing 25% of the riverine DIP discharge. Simultaneously, 6–28% of DIP produced in sediments was buried as Ca-bound P. Overall, this study highlighted the importance of deltaic sediments as an additional source of DIP to the coastal sea, and a minor but permanent sink of phosphorus as solid P burial.

Keywords : Organic matter, Phosphorus, Mediterranean Sea, Rhône River prodelta, Sediment, Modelling

19 **Acknowledgements**

20 We would like to thank the captain and the crew members of N.O. Tethys II (INSU/CNRS) for
21 their assistance during sampling at sea. We are grateful to Laurie Brethous, and Anouk
22 Villedieu who provided help at sea during sampling and in the laboratory for measurements.

23 Florian Caradec and Emilie Rabiller are acknowledged for their help during P-speciation
24 measurements at IFREMER's laboratory. We thank the two anonymous reviewers whose
25 comments greatly improved the quality of this manuscript. This work was supported by the
26 INSU/EC2CO-MissRhodia project, the French State program "Investissement d'avenir" run by
27 the National Research Agency (AMORAD project ANR-11-RSNR-0002) and the CEDoc
28 cooperation research project between Morocco and France. This is LSCE publication number
29 XXXX.

2. Introduction

Phosphorus is a key macronutrient for all life forms (Ruttenberg 1992; Tyrrell 1999). It is a typical limiting element of primary production in marine areas (Benitez-Nelson 2000), such as the Mediterranean Sea, with a prominent role in the Eastern part (Krom et al. 1991). Overall, particulate P is partitioned as P associated to organic matter, Fe-oxyhydroxides (henceforth called Fe-oxides), aluminum oxides, vivianite and apatite (Lebo 1991). Meybeck (1982) showed on a global scale that rivers deliver globally 21×10^{12} g P yr⁻¹ to the sea. However, only 10% of this flux is available to the marine biota, because more than 90% of the P load is bound to particles and, therefore, settle rapidly to the seafloor in the coastal zones.

Although eutrophication has existed throughout the geological times (Nixon 1995; Cloern 2001), the second half of the 20th century has witnessed the dramatic increase of anthropogenic activities (i.e. agricultural practice, industrial and urban waste, dam building and perturbation of hydrological cycles and sediment loads) which degraded deltaic and coastal ecosystems and altered biogeochemical cycles of carbon (C), nitrogen (N) and phosphorus (P) in these vulnerable areas (Howarth et al. 1996; Vitousek et al. 1997; Jackson et al. 2001; Diaz and Rosenberg 2008; Wilkinson 2017). These anthropogenic changes led also to large organic matter inputs (McKee et al. 2004; Canfield et al. 2005; Bianchi and Allison 2009; Syvitski et al. 2009) and intense nutrient (N and P) loads to coastal margins compared to the middle of the 20th century (Downing et al. 1999; Cloern 2001; Ludwig et al. 2009). The nutrient enrichment

stimulates primary production and phytoplankton growth in adjacent coastal waters (Milliman and Meade 1983; Cloern 2001), which leads to phytoplankton proliferation, alteration in species compositions, hypoxia and deterioration of water quality (Smith et al. 1999).

River-dominated Ocean Margins (RiOMar) are important interfaces, connecting approximately 87% of land surfaces to the oceans (Ludwig and Probst 1998; McKee et al. 2004). They support a large human population and provide ecological, economical and societal development due to their richness (Batker et al. 2014). They are important burial sites and preserve more than 40% of organic matter in the world ocean (Muller-Karger et al. 2005; Burdige 2007). Their sediments are characterized by high spatio-temporal variability of riverine discharge to the seafloor and strong biogeochemical gradients offshore (McKee et al. 2004; Lansard et al. 2009; Rassmann et al. 2016).

The Mediterranean Sea is a quasi-enclosed marginal sea (0.8% of the world ocean surface; Powley et al. 2017), often characterized as a replicate of an open ocean (Antoine and Morel 1995). This regional sea is oligotrophic, with lower nutrient concentrations (N and P) compared to oceanic areas (Dugdale and Wilkerson 1988), especially for dissolved inorganic phosphorus. A west-east gradient of decreasing P concentrations (Moutin et al. 2012) leads to higher primary production in the western Mediterranean Basin (Van Den Broeck and Moutin 2002; Moutin and Raimbault 2002). Overall, it has been shown that the P availability plays a key role as a limiting nutrient in the entire Mediterranean Sea (Krom et al. 1991; Diaz et al. 2001).

Rivers discharge large amounts of sediment and freshwater in the Mediterranean Sea (Ludwig et al. 2009). After the Nile damming, the Rhône River became the most important source of terrestrial organic matter, nutrients and freshwater to the Mediterranean Sea (Antonelli et al. 2008; Ludwig et al. 2009). The average particulate discharge from the Rhône River to the Mediterranean was estimated to 7.4 Mt y^{-1} (Pont et al. 2002), with large seasonal

and interannual variability. Overall, riverine fluxes at the Mediterranean Sea's boundaries and the controlling processes at the river-sea connection are key for understanding P cycling and its control in this oligotrophic sea, especially sediments where particulates from the river rapidly settle and are biogeochemically transformed.

Reactive P reaching the sediment-water interface is mainly associated to organic matter with a molar C/P ratio that is determined by its origin. Additional inputs of P are bound to Fe-oxides and as carbonate fluorapatite (CFA; Emsley 1980; Sundby et al. 1992; Ruttenberg and Berner 1993; Slomp et al. 1996; Delaney 1998; Andrieux-Loyer and Aminot 2001; Slomp et al. 2004; Ruttenberg 2014). As organic matter is progressively buried below the sediment-water interface, the non-refractory fraction is degraded via a sequence of biogeochemical processes, releasing DIP to the pore waters (Krom and Berner 1981; Sundby et al. 1992; Joshi et al. 2015). In the oxic zone of sediments, this dissolved phase may diffuse to the overlying waters or be adsorbed onto sediment Fe-oxide surfaces reversibly and irreversibly (Fe-bound P; Syers et al. 1973; Froelich 1988; Fox 1990; Balls 1994; Filippelli and Delaney 1996; Anschutz et al. 1998; Slomp et al. 1998; Cade-Menun et al. 2019). In the anoxic layers, DIP concentrations generally show a large increase due to organic P mineralization and P release from Fe-oxides dissolution, as the latter are used as a terminal electron acceptor in the mineralization process (Krom and Berner 1981). Therefore, iron reduction can have a significant role in phosphorus cycle as it has been shown for sulfur cycle (Thamdrup et al., 1994). DIP can ultimately be taken up by an authigenic mineral such as carbonate fluorapatite (CFA) through precipitation with Ca^{2+} (Ruttenberg and Berner 1993) or by vivianite (Emerson and Widmer 1978; Egger et al. 2015; Dijkstra et al. 2016; Lenstra et al. 2018). This “sink-switching” represents the main long-term sink of reactive P, as long as the authigenic mineral acts as an insoluble phase in deep sediments without being reprocessed during the diagenetic processes (Ruttenberg and Berner 1993; Slomp et al. 1998; Kraal et al. 2015). The P regeneration and sink-switching needs to be better studied

in subaqueous deltaic sediments subjected to high nutrient loads. Considering the large P inputs from the Rhône River, it is necessary to determine the biogeochemical P dynamics and the fate of P input to the deltaic sediments: release vs burial with authigenic minerals.

Biogeochemical modelling by reactive-transport models has been largely used to the early diagenesis studies in the aquatic sediments (Berner 1980; Rabouille and Gaillard 1991; Tromp et al. 1995; Boudreau 1996; Soetaert et al. 1996; Wang and Van Cappellen 1996; Boudreau 1997; Wijsman et al. 2002; Berg et al. 2003; Canavan et al. 2006), but the benthic P cycle is rarely incorporated as a function of organic matter mineralization (Canavan and Slomp 2006; Gypens et al. 2008; Lenstra et al. 2018) especially in a highly dynamic system. Here, we analyzed spatial variability in the sediment composition, the organic matter transformation and nutrient retention in the Rhône prodelta sediments, in order to quantify the intensity of biogeochemical processes and its effect on the benthic P cycle.

The present paper aims at quantifying the biogeochemical processes that control phosphorus recycling and trapping in these sediments and at determining the spatial impact of riverine organic matter from the river mouth to the adjacent continental shelf. To this end, we used an early diagenetic model (OMEXDIA, Soetaert et al. 1996), with P diagenesis (Ait Ballagh et al. 2020), and applied it to a dataset of sediment and porewater composition in the biogeochemical gradient of the Rhône River prodelta and shelf sediments, including particulate P speciation.

3. Materials and methods

3.1. Study site

The Rhône River (RR; Fig. 1) is the largest river in the Mediterranean Sea in terms of sediment loads, nutrient and freshwater fluxes (Antonelli et al. 2008; Ludwig et al. 2009). The

RR is 812 km long, characterized by a catchment area of 97800 km² originating in the Alps and ending in a delta of 1455 km² (Pont et al. 2002; Antonelli et al. 2004). Fifty kilometers upstream from the river mouth, the Rhône River splits into two unequal branches : the eastern channel (Grand Rhône, 50 km long, also called Rhône), which discharges 80-90% of the freshwater in the Gulf of Lions, and the western channel (Petit Rhône, 70 km long) that carries 10-20% of total discharge (Ibañez et al. 1997; Sabatier et al. 2006; Provansal et al. 2012). During flood events, 80% to 90% of annual particulate loads are discharged into the Rhône River delta, from the catchment area (Arnau et al. 2004; Ollivier et al. 2010; Eyrolle et al. 2012). The remaining loads originate from the soil erosion in the catchment area, in stream production or resuspension of river sediments from the upstream during lower freshwater discharges (Eyrolle et al. 2012). Overall, the mean water discharge at the Rhône River mouth is 1700 m³ s⁻¹, with a peak discharge of 13000 m³ s⁻¹ in December 2003 (Lochet and Leveau 1990; Ibañez et al. 1997; Pont et al. 2002; Arnau et al. 2004).

Generally, the Rhône River's plume extends towards the southwest (SW), as it is oriented by wind regimes close to the Rhône River mouth and by the River flow (Estournel et al. 1997; Naudin et al. 1997; Gatti et al. 2006). Consequently, the delivered loads of particulate material to the delta are mostly transported in the SW direction and some are ultimately driven to the adjacent continental shelf. The largest share of the riverine inputs is deposited at the seafloor near the river mouth, leading to the formation of a deposition fan called prodelta (Got and Aloisi 1990; Lansard et al. 2007). Consequently, the water depth increases from 10 m at the river mouth to 75-90 m on the adjacent continental shelf.

The decrease of sediment loads with distance from the Rhône River mouth is accompanied by a progressive aging of organic particles (Cathalot et al. 2013). The decreasing sediment deposition offshore gives rise to a large biogeochemical gradient from the Rhône River mouth to the continental shelf (Cathalot et al. 2010; Pastor et al. 2011; Rassmann et al.

2016). Concerning the DIP inputs from the Rhône River, Ludwig et al. (2009) showed an increase of DIP loads from 1970 to around 1985, followed by a decrease that may be related to the mitigation of pollution (i.e., the ban of P detergents and/or the upgrading of the waste water treatment plants).

The Rhône prodelta (named because of its prograding properties) is characterized by two regions: the proximal zone located near the river mouth at a water depth of around 20 meters and a sedimentation rate of 20-35 cm yr⁻¹ depending on the period investigated and the tracer used (Charmasson et al. 1998; Radakovitch et al. 1999; Miralles et al. 2005; Lansard et al. 2007) whereas the prodelta zone is located further at 30-65 meters depth and displays sedimentation rates of 1-2 cm yr⁻¹. The adjacent continental shelf extends further away from the prodelta and has sedimentation rates of less than 0.1 cm yr⁻¹.

3.2. Measurements

3.2.1. Sampling stations

The sampling cruise MissRhoDia was carried out aboard the *RV Tethys II* in May 2018. Nine stations were sampled between 2 and 18 km from the Rhône River mouth, in the proximal (stations A, Z and AZ), prodelta (stations AK, K, B and L) and distal (stations C and E) domains over a southwest transect, corresponding to the main direction of the river plume. The locations of sampling stations are shown in Figure 1 and the main characteristics are in Table 1. Sediment cores were collected using an UWITEC[®] single corer (9.5 cm of inner diameter, 60 cm of length) for porewater and solid phases analysis. Bottom water samples were taken with a 12-L Niskin[®] bottle as close as possible to the seafloor (approximately 1 m). Salinity and temperature (°C) were measured using a salinometer with a precision of ± 0.1 and a thermometer with a precision of ± 0.1°C, respectively (Table 1).

3.2.2. In situ microelectrode measurements

The distribution of O₂ and pH at the sediment-water interface were measured using *in situ* microelectrode with a benthic microprofiler (Unisense®) (Rabouille et al. 2003; Lansard et al. 2008; Cathalot et al. 2010; Rassmann et al. 2016; 2020). The measurements of 4–5 vertical depth O₂ profiles per station and 2 pH profiles were achieved with a vertical resolution of 200 µm near the sediment-water interface. The dissolved O₂ concentrations were measured by polarographic oxygen microelectrodes that were provided with a built-in reference and internal guard cathode (Revsbech 1989). The microelectrodes were calibrated with a two-point calibration technique using the bottom water oxygen concentration determined by Winkler titration (Grasshof et al. 1983) and the anoxic pore waters of the sediment (Cai and Sayles 1996). The pH microelectrodes were calibrated by NBS pH buffers (pH 4.01, 7.00 and 9.00) which provided the slope of the sensor response to pH and which estimated the pH variation. The measurement of seawater pH in bottom waters was carried out by spectrophotometry with unpurified m-cresol purple (Clayton and Byrne 1993; Dickson et al. 2007) and is reported on the total proton scale (pH_T). The precision of the dissolved O₂ measurement was of ± 0.5 µM, whereas the uncertainty of measured pH was ± 0.01 pH units.

3.2.3. Ex situ measurements of solid and pore water profiles

In order to determine the sediment porosity, a core was sliced with a resolution of 2 mm for 0–10 mm depth, 5 mm for 10–60 mm, and 10 mm until the bottom of the core. The sediment porosity was calculated using water contents (the difference between the wet and dried weight after one week at 60 °C) and an averaged sediment density of 2.65 g cm⁻³. The sediment porosity was then corrected by the bottom water salinity.

A second sediment core was processed within 1–2 hours after sampling for the extraction of pore waters by its introduction in a glove bag flushed with N₂ atmosphere in order to avoid oxidation of reduced compounds. The extraction of pore waters was conducted by rhizons with

a mean pore size of 0.1 μm (Seeberg-Elverfeldt et al. 2005). After collection, the pore water Fe^{2+} and the total dissolved Fe were measured using the ferrozine method (Stookey 1970). Pore waters were poisoned using HgCl_2 for total alkalinity (TA) and dissolved inorganic carbon (DIC), acidified with HCl for SO_4^{2-} measurements, and stored at 4 °C until laboratory analysis. Subsamples for NH_4^+ , NO_3^- and DIP measurements were frozen on board. The pore water NH_4^+ concentration was diluted and quantified by the indophenol blue method (Grasshof et al. 1983), while the pore water NO_3^- concentration was measured using the Vanadium chloride reductant after dilution, followed by the addition of NEDD/sulphanilamide and the colorimetric method (Hansen and Koroleff 1999; Schnetger and Lehnert 2014). The uncertainties of the methods for NH_4^+ and NO_3^- were 5% and 2%, respectively.

The pore water DIC concentrations were quantified with a DIC analyzer (Apollo SciTech[®]), by the use of 1 ml of the sample volume with 4-7 replicates as described previously by Rassmann et al. (2016). The DIC method was calibrated using certified reference materials (oceanic CO_2 measurements, Scripps Institution of Oceanography-batch n°136) which provided an uncertainty of 0.3%. The pore water SO_4^{2-} concentrations were diluted by a factor of 100 and measured using ion chromatography on an ICS 1000 chromatograph with an AS14 column and an AG14 guard column, both in 4mm diameter (Dionex) (Rassmann et al. 2020). The uncertainty of this method was of $\pm 1.6 \%$.

For the pore water DIP, the measurements were performed spectrophotometrically (Murphy and Riley 1962) at 880 nm on 3 ml of each sample, using a mixed reagent prepared daily from sulfuric acid (H_2SO_4), potassium antimony (III) oxytartrate hemihydrate ($\text{K}(\text{SbO})\text{C}_4\text{H}_4\text{O}_6 \cdot 1/2\text{H}_2\text{O}$), ammonium molybdate ($(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$) and ascorbic acid ($\text{C}_6\text{H}_8\text{O}_6$). The precision of the method was 1%.

3.2.4. Sequential extraction of P forms

Another sediment core was sampled and freeze-dried for 48 hours, and ground for further analysis of the P solid forms. The sedimentary P pools were separated using the sequential extraction method as detailed in Andrieux-Loyer et al. (2008) and adapted from widespread extraction methods (Psenner et al. 1988; Ruttenberg 1992).

Fe-bound P (including the Fe-oxides bound P and the vivianite) was extracted by 0.1 mol l⁻¹ of Dithionite-Bicarbonate (8h, 20°C). In a second step, authigenic Ca-bound P (including carbonate fluorapatite Ca₅(PO₄, CO₃)₃F, biogenic hydroxyapatite Ca₁₀(PO₄)₆(OH)₂ and CaCO₃ bound P) was quantified on the remaining fraction by 1 mol l⁻¹ of Na-acetate buffer (pH = 4, 6h, 20°C) then washed by 1 mol l⁻¹ of MgCl₂ (pH = 8, 0.5h, 20°C). Then, the detrital P was extracted using 1 mol l⁻¹ of HCl (overnight). Total P was determined non-sequentially by 1 mol l⁻¹ HCl treatment overnight after sediment ignition at 550°C (4h), while the inorganic P referring to the sum of P-forms (Fe-bound P, Ca-bound P and detrital P), was extracted with 1 mol l⁻¹ HCl before sediment ignition, in order to ensure the measurements efficiency. Organic P was then quantified by the difference between total P and inorganic P (Andrieux-Loyer et al. 2008).

3.3. Model description

The modelling was performed by a steady-state version of the diagenetic model OMEXDIA (Soetaert et al. 1996; Khalil et al. 2018), implemented in the R software, using the R-package ReacTran (Soetaert and Meysman 2012) and extended to include the benthic P cycle (Slomp et al. 1996; Ait Ballagh et al. 2020). The steady-state model assumption in our study allowed to understand the relative contribution of biogeochemical processes involved in the benthic P cycle in dynamic system outside floods events. This assumption is based on a stationary state observed over several years (2007-2015) during the spring-summer season, which allows typical diagenetic situation to be investigated using a steady-state model (Rassmann et al. 2020). This condition was achieved by setting the rates of changes to 0 (i.e.

$\partial C_i / \partial t = 0$). The model outputs were fitted to the measurements of organic carbon, terminal electron acceptors (O_2 , NO_3^- , SO_4^{2-}), products of mineralization (NH_4^+ , DIP, DIC and oxygen demand units (ODU; reduced substances during anoxic mineralization)), and solid P fractions (organic P, Fe-bound P and Ca-bound P). The model solves a set of general diagenetic equations (Berner 1980; Boudreau 1997) for dissolved C_d and solid C_s concentrations (Eqs. 1 and 2), as described below:

$$-\frac{\partial}{\partial x} \left[-\phi D_s \frac{\partial C_d}{\partial x} + \omega \phi C_d \right] + \phi \sum R_d + \alpha (C_{d_0} - C_d) = 0 \quad (\text{Eq. 1})$$

$$-\frac{\partial}{\partial x} \left[-D_B \frac{\partial (1-\phi) C_s}{\partial x} + \omega (1-\phi) C_s \right] + (1-\phi) \sum R_s = 0 \quad (\text{Eq. 2})$$

Where C_d and C_s denotes the concentrations of the dissolved and solid species, respectively (in mol per volume of porewater or dry sediment⁻¹). ϕ is the sediment porosity (cm³ cm⁻³), x is the sediment depth (cm), D_s is the diffusion coefficient (cm² d⁻¹), D_B is the bioturbation coefficient (cm² d⁻¹), ω represents the sedimentation rate (cm d⁻¹), $\sum R_d$ and $\sum R_s$ denote the sum of the biogeochemical reactions occurring to dissolved and solid species, respectively. α is the depth-profile bioirrigation rate, while C_{d_0} is the bottom water concentration of dissolved phases ($\mu\text{mol l}^{-1}$).

The dissolved species are transported by molecular diffusion and bioirrigation, whereas bioturbation is the main transport process for solid phases. In addition, advection is a common process for all species (Berner 1980; Boudreau 1997; Rabouille & Gaillard 1991). The sediment compaction is represented in the model by the exponential decrease of porosity (Berner 1980; Rabouille and Gaillard 1991). Below the sediment-water interface, all the mineralization pathways produce NH_4^+ and DIP (Table 2). The labile (FDet) and semi-labile (SDet) fractions of organic carbon are mineralized in the oxic zone by oxygen (OxicMin; Table2, R1). Once the oxygen is totally consumed, the denitrification takes place in suboxic zone and is limited by

nitrate concentration (Denitrif ; Table 2, R2; Emerson et al. 2004). In the anoxic zone, metal oxides and sulfate are used as next terminal electron acceptors in the mineralization (AnoxicMin; Table 2, R3-4). The ammonium and the other reduced compounds expressed as oxygen demand units (ODU; Berner 1989; Soetaert et al. 1996) are reoxidized by oxygen in the oxic zone (Nitrif and ODUoxid, respectively; Table 2, R5-6). The stoichiometry of organic matter is represented by coefficients x and y denoting molar C/P and N/P ratios, respectively.

The diagenetic equations in Table 2 take into account all transport and reactive processes for the compounds; the labile and semi-labile fractions of organic carbon (Eq. 3 and Eq. 4), oxygen (Eq. 5), nitrate (Eq. 6), sulfate (Eq. 7), ammonium (Eq. 8), the dissolved inorganic carbon (Eq. 9) and the ODU that are produced by anoxic mineralization and may be lost from the system to form solid substances as a pyrite (Eq. 10; Berner 1970; Soetaert et al. 1996; Rassmann et al. 2020). These equations are depending on the conversion between solid and liquid units (ϑ).

$$\vartheta = \rho_s \frac{(1 - \phi)}{\phi}$$

Where ρ_s denotes the average density of the sediment particles (g cm^{-3}).

3.3.1. Benthic phosphorus cycle

The benthic P cycle is included in the model to obtain more insight into the P transformation in the Rhône prodelta sediments. The present model describes the concentration change with depth of pore water DIP ($\mu\text{mol l}^{-1}$) and three forms of particulate P (organic P, Fe-bound P and Ca-bound P ($\mu\text{mol g}^{-1}$)). All the equations are detailed in Table 3 (see also Ait Ballagh et al. (2020)). Figure 2 illustrates the biogeochemical processes of the benthic P cycle.

In the model, the P settling at the sediment-water interface is either associated to the organic matter by C/P ratios for the labile and semi labile OC fractions or bound to Fe-oxides

(Emsley 1980; Sundby et al. 1992; Ruttenberg and Berner 1993; Slomp et al. 1996; Delaney 1998; Andrieux-Loyer and Aminot 2001; Slomp et al. 2004; Ruttenberg 2014). A refractory part P_{refrac} ($\mu\text{mol g}^{-1}$) which is not degraded in the sediments is added to the labile fraction (Table 3, Eq. 11).

Below the sediment-water interface, organic matter mineralization releases DIP (cm^{-3} liquid d^{-1}) in the porewaters (Table 3, Eq. 12). In the oxic zone, the DIP may diffuse to the overlying waters or be adsorbed onto sediment Fe-oxides surfaces, which leads to Fe-bound P formation ($\mu\text{mol cm}^{-3}$ liquid d^{-1}) (Table 3, Eq. 13; (Syers et al. 1973; Froelich 1988; Fox 1990; Balls 1994; Filippelli and Delaney 1996; Anschutz et al. 1998; Slomp et al. 1998; Cade-Menun et al. 2019). The Fe-oxides are used as next terminal acceptors once oxygen, nitrate and manganese oxides are totally consumed (Canavan & Slomp 2006; Krom & Berner 1981). Therefore, the adsorbed DIP is released by the dissolution of Fe-bound P ($\mu\text{mol cm}^{-3}$ solid d^{-1}) (Table 3, Eq. 14).

In the anoxic zone, the concentrations of DIP increase through organic matter mineralization and Fe-bound P dissolution. The porewaters are supersaturated with respect to carbo-fluorapatite ($DIP_{bw} > DIP_{eq}$), and form an authigenic mineral with Ca^{2+} (Ruttenberg and Berner 1993). The authigenic Ca-bound P (Ca-bound P_{auth}) does not dissolve once it is formed, and represents a permanent sink for reactive P in the sediments ($\mu\text{mol cm}^{-3}$ liquid d^{-1}) (Table 3, Eq. 15; Slomp et al. 1996). The coupling of the biogeochemical processes takes into account the conversion from solid units to the liquid ones (ϑ) when it is necessary and stoichiometric conversion factors (Table 3, Eq. 16- Eq. 18; Soetaert et al. 1996).

3.3.2. Benthic budget calculation

The steady-state benthic budgets of all simulated dissolved and solid compounds were calculated by the model with a focus on P budget: the deposition flux, the integrated rates of

the biogeochemical processes, burial for each state variable. Total oxic mineralization, total denitrification, total anoxic mineralization, total organic P mineralization, total Fe-bound P formation, total Fe-bound P release and total Ca-bound P precipitation expressed in $\mu\text{mol m}^{-2} \text{d}^{-1}$, are the depth-integrated values of oxic mineralization (OxicMin), denitrification (Denitrif), anoxic mineralization (AnoxicMin), organic P mineralization (Porg_{min}), Fe-bound P_{formation}, Fe-bound P_{release} and Ca-bound P_{precipitation} depth profiles expressed in the model in $\text{nmol cm}^{-3} \text{d}^{-1}$, respectively.

4. Results

4.1. Model-data comparison of dissolved and solid phases

The measured and simulated concentration-depth profiles of oxygen, nitrate, sulfate, DIC, ammonium, ODU and total organic carbon (TOC) are presented in Figure 3 for all sampling stations. The O₂ concentrations depleted rapidly from 185 $\mu\text{mol l}^{-1}$ - 235 $\mu\text{mol l}^{-1}$ in bottom waters (at stations C and L, respectively; Table 5) to 0 (all stations). The O₂ penetration depth into the sediment ranged from 2 mm in the proximal domain at station Z to 12.5 mm in the distal domain at station E, a pattern which was well represented by the model (Fig. 3). Nitrate concentrations were low at all stations and showed an increase with the distance from the river mouth to the adjacent continental shelf, without exceeding 15 $\mu\text{mol l}^{-1}$ (Fig. 3). The measured nitrate concentrations in the bottom waters never exceeded 3 $\mu\text{mol l}^{-1}$ (Table 5) and showed a peak below the sediment-water interface, indicating nitrification, followed by denitrification, which was reproduced by the model (Fig. 3). The depth of the nitrate peak increased in line with the increase of oxygen penetration from the river mouth to continental shelf, an evolution that was well reproduced by the model. The measured SO₄²⁻ profiles highlighted a sharp decrease with sediment depth at stations Z and A in the proximal zone; the SO₄²⁻ concentrations ranged from about 30 mmol l^{-1} in the bottom waters to concentrations

below 5 mmol l⁻¹ in deeper layers. The SO₄²⁻ profiles showed a slighter decrease with sediment depth in the prodelta and distal domains (Fig. 3-b, 3-c; stations AK, B, K, L, C and E). The model was in good agreement with the SO₄²⁻ measurements.

The measured and simulated DIC profiles indicated an increase with depth at all stations: concentrations were extremely high in the proximal domain (around 50 mmol l⁻¹ at stations Z and A below 15 cm, Fig. 3-a). DIC concentrations in porewaters at depth decreased according to distance from the river mouth to about 5 mmol l⁻¹ in the distal domain (Fig. 3-c, station E). This pattern linked to the diagenetic intensity was also well reproduced by the model. The same biogeochemical gradient was observed for the pore water NH₄⁺ profiles, with concentrations increasing with sediment depth at all stations. A gradient of NH₄⁺ concentrations at sediment depth (15cm) from up to 3000 μmol l⁻¹ (station Z) until about 100 μmol l⁻¹ (station E) was visible, similarly to the DIC profiles. Therefore, a larger production of DIC and NH₄⁺ was clearly observed in the proximal domain compared to the prodelta and distal domains. Overall, the model was in good agreement with pore water DIC and NH₄⁺ profiles at all stations (Fig. 3). The ODU concentrations (mainly Fe²⁺) displayed an increase with sediment depth, with about 150 μmol l⁻¹ close to the river mouth, but this maximum decreased in continental shelf stations to less than 5 μmol l⁻¹ (station E; Fig. 3-c). The simulated TOC profiles never exceeded 2% at the sediment-water interface and showed a decrease with sediment depth at all stations (Fig. 3).

Figure 4 shows the measured and simulated profiles of DIP, organic P, Fe-bound P and Ca-bound P. DIP concentrations were low near the sediment-water interface, then showed an increase with sediment depth. In deeper sediment layers of the proximal domain, DIP concentrations were extremely high (about 250 μmol l⁻¹ at station Z, A and AZ). These deep DIP maximum concentrations decreased with water depths and distance from the river mouth until about 25 μmol l⁻¹ at station E (Fig. 4). Overall, the model reproduced the increasing shape

of DIP with sediment depth, except at stations Z, B and K below 10 cm depth (Fig. 3). The DIP evolution followed a similar pattern as DIC and NH_4^+ for all stations. The measured organic P profiles never exceeded $10 \mu\text{mol g}^{-1}$ at the sediment-water interface at all stations. Organic P showed a slight decrease with sediment depth and showed a generally constant concentration offshore. Stations Z and A displayed a larger spread of organic P values with sediment depth. The model showed a good agreement with average measurements for this phase but showed a large gradient near the sediment-water interface (Fig. 4). Similarly, the observed Fe-bound P concentrations at stations A, Z and AZ showed dispersed values, while the model reproduced the average Fe-bound P concentrations and showed a slight decrease with sediment depth (Fig. 4). Concerning the other stations, the average Fe-bound P concentrations were well represented by the model (Fig. 4). A similar pattern was found for Ca-bound P_{auth} concentrations and showed also dispersed values in stations A, Z and AZ. This phase showed very consistent values at the profile scale and between profiles at all stations from the river mouth to the continental shelf, which was overall reproduced by the model (Fig. 4).

4.2. Boundary conditions and model parameters

The fixed parameters of the model listed in Table 4 were ammonium adsorption coefficient (NH_3Ads), maximum oxidation of ODU (r_{ODUox}), nitrification rate constant (r_{nit}), half-saturation constants for O_2 limitation for oxic mineralization ($K_{\text{SO}_2\text{oxic}}$) and nitrification ($K_{\text{SO}_2\text{nitr}}$), nitrate limitation for nitrate reduction ($K_{\text{NO}_3\text{denit}}$), O_2 limitation for the oxidation of reduced substances ($K_{\text{SO}_2\text{duox}}$), O_2 inhibition for nitrate reduction ($K_{\text{inO}_2\text{denit}}$) and anoxic mineralization ($K_{\text{inO}_2\text{anox}}$), nitrate inhibition for anoxic mineralization ($K_{\text{inNO}_3\text{anox}}$), sulfate limitation for anoxic mineralization (K_{SSO_4}) and finally the C/N and C/P fixed for labile and semi-labile fractions of organic carbon for all the sampling stations (Table 4). The C/N_{fast} and C/N_{slow} were fixed to 7.1 (Pastor et al. 2011) and 10 (Khalil et al. 2018), respectively. The C/P

ratio was fixed by assuming that N/P was equal to 16. Therefore, the estimated C/P for labile fraction (C/P_{fast}) was averaged to 114, while the C/P for semi-labile fraction (C/P_{slow}) was 160. It is noteworthy that because the proportion of the labile fraction increases offshore (from 50% to 75%), the overall C/P ratio decreases as well from 137 in the proximal zone to 126 in the distal zone which is consistent with the aquatic nature of the organic matter degraded in these sediments.

Table 5 includes the parameters that were measured at each station. The bottom water temperature ranged between 14.7 °C (at station C) and 20 °C (at station AZ), which is typical of the area at this season. Salinity varied between 37.8 at station A and AZ and 38.5 at station K (Table 5). The bottom water concentrations of all dissolved phases did not follow a spatial pattern (Table 5). The bottom water concentrations of NO_3^- , NH_4^+ and ODU were low for all stations. The SO_4^{2-} concentrations were nearly constant in bottom waters and close to 30 mmol l^{-1} at all stations (Table 5). The porosity at surface sediment and at depth were lowest in station K (0.76 and 0.58, respectively), and highest in station C (0.86 and 0.64, respectively). The sedimentation rates adopted for this modelling study were obtained from literature (Charmasson et al. 1998; Radakovitch et al. 1999; Miralles et al. 2005; Lansard et al. 2007), ranging from 10 cm y^{-1} in proximal domain, to 0.1 cm y^{-1} in distal domain (Pastor et al. 2011).

The parameters estimated by fitting the model to the data are listed in Table 6. The reactive OC flux (OC_{flux}) followed a spatial pattern and decreased from 160 $\text{mmol m}^{-2} \text{d}^{-1}$ at station Z (the river mouth) to 10 $\text{mmol m}^{-2} \text{d}^{-1}$ at station E on the continental shelf. The first-order decay rate constants for labile fraction of OC (R_{fast}) showed a clear increase with distance to the continental shelf (0.06 d^{-1} at station Z to 0.2 d^{-1} at stations B, K, L, C, E). Contrarily, the first-order decay rate constants of the semi-labile fraction (R_{slow}) decreased with distance to the river from 0.0035 d^{-1} at station Z to 0.0002 d^{-1} at stations B and C (Table 6). The labile OC represented an important proportion of OC (P_{fast}) that increased from 50% near the river mouth

(stations Z and A) to 75% on the continental shelf. The refractory carbon varied between 0.9 and 1.4% (Table 6). The proportion of ODU deposition (P_{depo}) ranged between 90% and 99%. The bioturbation layer thickness (x_b) increased from 5 cm in stations A, Z and AZ, to 10 cm for other stations, while the bioturbation coefficient (D_{b0}) was relatively invariant (from $0.01 \text{ cm}^2 \text{ d}^{-1}$ to $0.015 \text{ cm}^2 \text{ d}^{-1}$). The bioirrigation at the sediment surface (α_{irr0}) was needed for all stations but did not indicate a clear spatial trend according to the distance from the Rhône mouth ($0.05 - 0.2 \text{ d}^{-1}$), with bioirrigation depths (x_{irr}) ranging between 6 cm and 14.5 cm of depth (Table 6).

The Fe-bound P fluxes at the sediment-water interface (Fe-bound P_{flux}) decreased offshore with values as high as $730 \mu\text{mol m}^{-2} \text{ d}^{-1}$ at station Z and decreased to $10 \mu\text{mol m}^{-2} \text{ d}^{-1}$ at stations C and E. The Fe-bound P formation rate constants ($R_{\text{Fe-bound Pform}}$) were 0.8 d^{-1} at all stations, except for stations C and E where they were only 0.1 d^{-1} . Concerning the Fe-bound P release, the rate constants ($R_{\text{Fe-bound Prel}}$) were similar at all stations between 0.0005 d^{-1} (at stations B and K) and 0.0008 d^{-1} (at the other stations; Table 6). The Fe-bound P asymptotic concentrations (Fe-bound P_{asympt}) ranged between $2 \mu\text{mol g}^{-1}$ (at station A) and $4 \mu\text{mol g}^{-1}$ (at station C). The Ca-bound P precipitation rate constants ($R_{\text{Ca-bound Pprecip}}$) ranged between 0.002 d^{-1} (stations K and C) and 0.006 d^{-1} (stations Z and L) (Table 6). Inherited concentrations of Ca-bound P (Ca-bound P_{her}) were between $3.6 \mu\text{mol g}^{-1}$ and $5.3 \mu\text{mol g}^{-1}$.

5. Discussion

The application of the model to the dataset from sediments on all stations (from Z to E) along the Rhône River prodelta allowed quantifying total organic matter mineralization rates and calculating the contribution of different diagenetic pathways to the mineralization processes. Moreover, the addition of the benthic P cycle allowed identifying the sources and sinks of internally produced DIP and calculating sedimentary P budgets in proximal, prodelta and distal domains.

5.1. Spatial variability of organic matter mineralization

5.1.1. The organic matter properties

The OC fluxes (OC_{flux}) ranged between 160 and 10 $mmol\ m^{-2}\ d^{-1}$ (Fig. 5) and were characteristic of organic-rich ecosystems (Rabouille et al. 2008). They were similar to the obtained OC fluxes for the same stations at the Rhône River prodelta by Pastor et al. (2011). Moreover, the OC fluxes decreased exponentially as a function of distance from the Rhône River mouth (Fig. 5). This decrease was already observed by Cathalot et al. (2010) for the oxygen diffusive fluxes. During flood events, occurring mostly in fall-winter, the majority of organic-rich particulate inputs from the Rhône River are accumulated in the proximal and prodelta domains. Therefore, these areas are characterized by large OC mineralization rates during spring (Lansard et al. 2009; Cathalot et al. 2010; Pastor et al. 2011; Rassmann et al. 2016). This is also evidenced in the results by complete removal of sulfate during sulfate reduction and the intense production of DIC especially in the proximal domain.

In front of the Rhône River mouth, Rassmann et al. (2016; 2020) showed that pore waters were supersaturated with Ca^{2+} , preventing carbonate dissolution in these deltaic sediments. As a consequence, the DIC was mainly produced by OC mineralization processes (i.e. bacterial respiration). The obtained fluxes ($10 - 160\ mmol\ m^{-2}\ d^{-1}$) were in the same range as those from the Louisiana shelf ($35\ mmol\ m^{-2}\ d^{-1}$; Morse and Eldridge, 2007), although higher in the proximal domain. They were higher also compared to the Guadalquivir River mouth (Ferrón et al. 2009) and the Po River (Hammond 1999).

In addition to refractory carbon, the model takes into account two fractions of organic matter with variable reactivities, the labile and semi-labile fractions. Each labile and semi-labile fraction has a first-order decay rate constant (R_{fast} and R_{slow} , respectively; Soetaert et al. 1996), while the proportion of labile OC (P_{fast}) determines the relative abundance of both fractions in the OC inputs. The reactive compound was the dominant fraction in the organic carbon input

(C_{org}) as it increased from 50% in the proximal zone to 75% in the continental shelf (Table 6), being negatively correlated to the OC fluxes as has been already mentioned in Pastor et al (2011). The first-order decay rate constant of the labile OM fraction (R_{fast}) was higher in the prodelta and distal domains compared to the proximal domain and increased with distance offshore from 0.06 d⁻¹ at the river mouth to 0.2 d⁻¹ on the continental shelf (Fig. 6a) which indicated a progressive increase in lability and proportion of labile organic matter with the distance offshore. The rate constants in the proximal zone (0.06 to 0.15 d⁻¹) were slightly higher than those calculated for the Mississippi delta and the Louisiana shelf (0.02-0.05 d⁻¹; Morse and Eldridge, 2007). However, the rate constants calculated for the continental shelf were in the same range as the eutrophic Scheldt estuary (0.15 d⁻¹; Hofmann et al. 2008) and the Thau lagoon (0.1 d⁻¹), which are characterized by labile organic matter from phytoplankton and macrophyte debris (Dedieu et al. 2007).

The first-order decay rate constant of the semi-labile OM fraction (R_{slow}) decreased with distance from the Rhône River mouth to the continental shelf by a factor of 10 (0.0035 to 0.0002 d⁻¹, respectively; Fig. 6b). This could be related to the supply of older and less reactive material as indicated by ¹⁴C dating (Cathalot et al. 2013). The R_{slow} values were in the same range as the values from the eutrophic environments, such as Scheldt estuary (0.002 d⁻¹; Hofmann et al. 2008) and the Thau lagoon (0.001 d⁻¹; Dedieu et al. 2007). This trend matched with the results of Pastor et al. (2011) and highlighted that the mix of labile and semi-labile organic matter in the proximal and prodelta domains contained essentially terrestrial fine and coarse organic matter including a small proportion of riverine organic matter (Dumoulin et al. 2018). At the same time, the sediments in the distal domains probably receive labile marine organic matter (Pozzato et al. 2018) which is entirely mineralized together with aged terrestrial organic particles (Lansard et al. 2009).

The R_{fast} , R_{slow} and P_{fast} values highlighted that the terrestrial organic matter in the proximal domain, was composed of fine and coarse continental material rich in organics (Cathalot et al. 2013). It was less reactive than those at other stations offshore probably due to prior degradation during transport or storage on land. Stations offshore receive a larger proportion of marine organic material such as phytoplankton (Pastor et al. 2011). This evolution is confirmed by the results obtained using $\delta^{13}\text{C}_{\text{OC}}$ by Lansard et al. (2009). $\delta^{13}\text{C}_{\text{OC}}$ varied from -26.8‰ in the proximal domain sediments to -23.9‰ on the shelf, showing a significant increase in the distal domain, linked to the limited dispersal of terrestrial organic matter on the continental shelf. The limited transport of terrestrial organic matter led to the highest values of R_{fast} with distance from the River mouth linked to the increase of marine material proportion (Fig. 6a).

5.1.2. Organic matter mineralization processes

The decreasing mineralization rates of organic matter from the proximal to distal domain was reflected in the penetration depths of oxygen that were less than 2 mm at stations A or Z while they reached 12.5 mm in the distal domain (Cathalot et al. 2010; Rassmann et al. 2016). At these stations, the large increase of DIC production and the high consumption of sulfate indicated the large mineralization of reactive organic matter (Fig. 3).

The pore water NH_4^+ , DIC, and DIP were indicators of organic matter mineralization, and their concentrations increased as function of sediment depth (Fig. 3; Berg et al. 2003). Overall, the prodelta stations showed lower concentrations of these compounds in deep sediments compared to the proximal domain (e.g. the NH_4^+ with 300-1000 $\mu\text{mol l}^{-1}$ and 2000-3500 $\mu\text{mol l}^{-1}$, respectively) but were higher than the distal domain (200-500 $\mu\text{mol l}^{-1}$). The same pattern was shown by the other components (Fig. 3). The observed decrease of asymptotic concentrations of DIC, NH_4^+ and DIP with distance to the shore was related to the decrease of OC mineralization rates in these areas, causing the decrease of anoxic mineralization. This is

well reflected in Table 6 where OC flux represents the metabolizable organic matter deposition which, in the model calculation, is completely mineralized in surficial sediments. The values of this “integrated mineralization” varies from over 100 mmol C m⁻² d⁻¹ in the proximal zone to 10 mmol C m⁻² d⁻¹ on the distal shelf, with a gradual decrease over the prodelta.

In the proximal domain, and after the total consumption of O₂ and NO₃⁻, the ODU mainly consisted of dissolved Fe²⁺, as H₂S is very low (Rassmann et al. 2020). The ODU showed an increase of concentrations with sediment depth, corresponding to the Fe-bound P decrease (Fig. 4). This indicated a production of dissolved Fe and a potential release of DIP through the reductive dissolution of Fe-oxides, used as a terminal electron acceptor in the anoxic zone of sediments (Krom and Berner 1981; Canavan and Slomp 2006). This Fe-bound P decrease was less important in prodelta and distal domains, as shown by rather constant Fe-bound P profiles with depth, which indicated that the released DIP in the sediments mainly originated from the organic matter release without, or with a minor contribution of net Fe-bound P release (Fig. 4).

Overall, the model calculations showed that oxic mineralization followed a spatial pattern along the biogeochemical gradient, increasing its contribution to the total mineralization from 13.7% (station A, proximal domain) to 80% (station E, the distal domain; Fig. 7). The oxic mineralization contribution showed the same pattern as Pastor et al. (2011) and was in the same range (8 – 67%). The oxygen below the sediment-water interface was used also for the reoxidation of NH₄⁺ and the reduced substances that were produced in the anoxic zone. However, the nitrification process was not very important in the Rhône prodelta sediments, and only a small fraction of ODU was reoxidized, since more than 90% of ODU was buried at depth (p_{depo}, Table 6). Therefore, the main process responsible for oxygen consumption was oxic mineralization. The denitrification represented the smallest contribution compared to the other mineralization pathways (0.7 to 3.7%) and showed an increase from proximal to distal domains

(Fig. 7). This low contribution was due to the fact that denitrified NO_3^- originated only from nitrification below the sediment-water interface, due to the low concentration of NO_3^- in bottom waters (Table 5). Anoxic mineralization was related to the Mn-oxides, Fe-oxides and sulfate reduction in the anoxic zone. It exhibited a clear spatial gradient, accounting for more than 85% of total mineralization near the river mouth, decreasing to 16% near the continental shelf (Fig. 7). Sensitivity analysis of the model showed that the variations of the sedimentation rates in the Rhône proximal and prodelta domains do not impact significantly the mineralization pathways contributions. Indeed, an increase by a factor of 10 of the sedimentation rates (from 1 to 10 cm y^{-1} in the proximal zone and 0.1 to 1 cm y^{-1} for the prodelta domain) induced slightly lower contribution of oxic mineralization (from 18 to 14% of the total mineralization) in the proximal domain and was negligible in the prodelta domain.

The model showed a decreasing loss of ODU by burial from 99% at station Z to 90% at station C (Table 6). The ODU deposition highlighted a loss of reduced products of mineralization at deeper depths as solid substances, such as pyrite (Berner 1970; Soetaert et al. 1996). Similarly, high removal rates were found previously by Pastor et al. (2011) and Rassmann et al. (2020) in the Rhône River prodelta.

5.2. Benthic P transformation along the Rhône River prodelta

At the proximal stations, DIP concentrations showed an increase with sediment depth until a maximum of 300 $\mu\text{mol l}^{-1}$. The DIP concentrations were larger compared with concentrations obtained by Rassmann et al. (2020) for the same stations (50-160 $\mu\text{mol l}^{-1}$). This was probably linked to the intensity of OC inputs from 2015 (Rassmann et al. 2020) to 2018 (this study). As pointed out in the literature, the secondary conversion of Fe-oxides to FeS certainly contributed to the net release of adsorbed DIP on Fe-oxides with sediment depth (Anschutz et al. 1998). Indeed, Rassmann et al. (2020) described a clear precipitation of FeS at

the proximal stations as shown by the spatial pattern of solid FeS, decreasing from the proximal domain to the adjacent continental shelf sediments. Both the increase of DIP concentrations and decrease of Fe-bound P with sediment depth in the proximal and prodelta zones accompanied a slight increase of Ca-bound P with depth, characterized by dispersed values of measurements (Fig. 4). The high DIP concentrations in porewaters likely contributed to the precipitation of Ca-bound P_{auth} (Fig. 4). Despite the decrease of OC fluxes to the seafloor and thus of organic P, the Fe-bound P and the Ca-bound P concentrations were almost similar at the sediment-water interface in the proximal, prodelta and distal domains (Figs. 3 and 4). This can probably be related to a compensation of flux versus sedimentation, as both decrease at a similar rate from the River mouth to offshore.

5.2.1. Sources of pore water DIP

The organic P input to the sediment was calculated using the organic C fluxes and the C/P_{fast} and C/P_{slow} assumed in this study (Table 4). The obtained integrated rates of organic P mineralization, extremely high at the River mouth ($1196 \mu\text{mol m}^{-2} \text{d}^{-1}$; Fig. 8), were characteristic of organic-rich ecosystems. Their decrease from the proximal to the distal domain (1196 to $81 \mu\text{mol m}^{-2} \text{d}^{-1}$; Fig. 8), was linked to the decrease of organic matter inputs (Pastor et al. 2011). The values obtained from the Rhône prodelta sediments were higher than the integrated rates of organic P mineralization from the eutrophic Elorn and Aulne estuaries ($262 - 757$ and $172 - 414 \mu\text{mol m}^{-2} \text{d}^{-1}$, respectively; Ait Ballagh et al. 2020). Figure 8 shows that the organic P mineralization represented the main source by up to 90% of the total DIP production, while the net Fe-bound P release had a minor contribution in benthic DIP production at all stations (Z to E).

Concerning the Fe-bound P fluxes deposited at the sediment-water interface (Fe-bound P_{flux}), Table 6 and Figure 8 highlighted their importance in the proximal domain ($730 \mu\text{mol m}^{-2} \text{d}^{-1}$ at station AZ), which were more than 10 times higher in the prodelta and distal domains

(82 and 10 $\mu\text{mol m}^{-2} \text{d}^{-1}$ at stations B and E, respectively). The Fe-bound P fluxes in the distal domain were similar to the fluxes from Vilaine Bay (4 - 25 $\mu\text{mol m}^{-2} \text{d}^{-1}$; Ratmaya, 2018), due to the similarity of sedimentation regimes at the sampling stations of the distal zone (0.1 cm yr^{-1}) and the Vilaine Bay (0.2 cm yr^{-1} ; Ratmaya, 2018).

Except at station A (Fig. 9), the model indicated that the formation of Fe-bound P in the oxic zone and its release below were very similar along the Rhône River prodelta. This indicated that, at all stations, Fe-bound P acted as a temporary sink for DIP in the oxic zone, which was thereafter released in the anoxic zone. Therefore, Fe-bound P acted as a shuttle between organic P mineralization and Ca-bound P_{auth} precipitation (sink-switching) along the Rhône River prodelta (Ruttenberg and Berner 1993; Slomp et al. 1998; Kraal et al. 2015).

5.2.2. Consumption/export of internally produced DIP

The model calculations showed that DIP produced by mineralization and net Fe-bound P release was essentially lost by recycling to the overlying waters rather than burial through Ca-bound P_{auth} formation. Figure 10 indicated that the sum of both processes, in terms of DIP consumption/export fluxes, decreased from the river mouth to the continental shelf similarly to the organic matter mineralization. A difference by a factor of 3 was observed between the proximal and the prodelta domains while a factor of 10 was calculated between the proximal and the distal domains. In the prodelta and distal domains, the recycled DIP accounted for up to 90% of total DIP losses, while the burial of reactive P as Ca-bound P_{auth} form was higher in the proximal domain but never exceeded 28% of the total DIP loss (Fig. 10).

The recycling of DIP to the overlying waters (Fig. 10) decreased from the river mouth to continental shelf (931 to 74 $\mu\text{mol m}^{-2} \text{d}^{-1}$) and occurred mainly through molecular diffusion (>70%) at all stations, while bioirrigation process accounted for a minor part. These recycled DIP fluxes to the overlying waters were in the same range as those from the Elorn and Aulne

estuaries ($85 - 738 \mu\text{mol m}^{-2} \text{d}^{-1}$; Ait Ballagh et al. 2020), the Haringvliet lake ($370 \mu\text{mol m}^{-2} \text{d}^{-1}$; Canavan and Slomp, 2006) and those from the Gulf of Finland $0 - 1032 \mu\text{mol m}^{-2} \text{d}^{-1}$ (Lehtoranta et al. 2017). The eutrophic Haringvliet lake, the Vilaine bay, the Elorn and Aulne estuaries were similar to the Rhône River prodelta in terms of the role of organic P mineralization as the main benthic DIP source, unlike the Gulf of Finland, where the DIP release mostly originated from the Fe-bound P release (Canavan and Slomp 2006; Lehtoranta et al. 2017).

5.2.3. Sedimentary biogeochemical P budgets

Figure 11 represents vertically integrated P budgets for 3 selected stations, 1 from each domain: the proximal (station A), prodelta (station K) and distal (station E) domains. For these budgets the benthic P processes were integrated over the sedimentary column of each sampling station. Overall, it is clear that DIP was mainly released by organic P mineralization, as it was the main source of DIP the Rhône prodelta sediments. However, this intensity was enhanced by the Fe-bound P phase, with minor contribution of net Fe-bound P release. As the organic P mineralization fluxes decreased from the proximal ($1004 - 1196 \mu\text{mol m}^{-2} \text{d}^{-1}$; Fig. 11, station A) to the distal domain (113 and $81 \mu\text{mol m}^{-2} \text{d}^{-1}$, station E), the Fe-bound P fluxes also decreased by a factor of 50 between the river mouth and offshore in relation with the decrease of the river material deposition offshore. Therefore, the iron related P cycle decreased in intensity towards the shelf with an overall minor contribution in the proximal zone ($8 - 81 \mu\text{mol m}^{-2} \text{d}^{-1}$) and no net Fe-bound P release contribution in the distal domain (Fig. 11).

The largest part of the intense DIP production in the sediment was recycled to the overlying waters by molecular diffusion at all domains. A significant sink was calculated as Ca-bound P_{auth} burial in the proximal zone (Fig. 11, station A) with a rapid levelling off offshore as the production of DIP in the sediment decreased. Sensitivity tests showed that uncertainty on the sedimentation rates may impact the calculated rate of DIP recycling to the water column.

Indeed, increasing sedimentation rates by a factor of 10 (from 1 to 10 cm y⁻¹ in the proximal zone) decreased DIP production in the sediment essentially by lowering the contribution of Fe-bound P dissolution to DIP, and contributed to lower overall DIP flux to the water column (-40%). This increase of sedimentation rate also induced an increase of burial as Ca-bound P (+34%) in the proximal domain. Furthermore, these sensitivity tests showed that the C/P ratios which control the input of organic P to the sediment have an influence on the mass balance of the P cycle. An increase of the C/P of semi-labile organic matter from 160 (used in this study) to 200 would decrease the recycling flux by 20% in the proximal domain, essentially because of the lower flux of organic P deposited in the sediment. Ca-bound P burial would also be lowered by 15% because of the decreased DIP availability. Lower variations were calculated for the prodelta domain with less than 10% variation of DIP recycling and Ca-bound P burial for similar variations of C/P in the semi-labile fraction. The change of C/P for the labile fraction (from 114, adopted in this study, to 100) would have a lower impact on the P mass balance with decreases of 9% for DIP recycling and 3% for Ca-bound P burial.

5.3. The P budgets at a deltaic scale

In order to assess the contribution of the Rhône River prodelta as a source/sink of DIP to or from the Mediterranean Sea, the P budgets (recycling/burial) along the Rhône River prodelta were calculated. For that purpose, we averaged the benthic DIP fluxes and authigenic Ca-bound P burial fluxes obtained using the model-data approach for each zone of interest (proximal, prodelta and continental shelf-distal zones). As these zones were rather homogeneous regarding the benthic DIP fluxes and Ca-bound P_{auth} burial ($\pm 20\%$ standard variation, see Fig. 10) compared to the overall variation between the zones (one order of magnitude between proximal and distal zones, Fig. 10), averaging did not bring significant bias in the final calculations. Surface areas of the proximal, prodelta and distal domains (Table 7)

were adopted from Lansard et al. (2009) in agreement with the Thiessen's polygon method, used in Cathalot (2008).

By using these surface areas, we obtained the total benthic DIP fluxes (a source to the water) and the sink due to precipitated Ca-bound P burial in the sediments of the Rhône River prodelta and its adjacent continental shelf. These fluxes were then compared to the riverine DIP discharge. According to Malagó et al. (2019), the Rhône River delivered 1465 tons DIP yr⁻¹ to the Mediterranean Sea from 2003 to 2007, whereas Ludwig et al. (2009) estimated a discharge of 2750 tons DIP yr⁻¹, in agreement with direct measurements by Moutin et al. (1998). These separate estimations varied by a factor of 2 which could be related to the decrease of P inputs to the rivers from the 1990s - 2000s to the end of the 2010s. It may also be due to differences in assumptions for the calculations. It is out of the scope of this paper to further discuss these issues and we used the two values for bracketing our estimates of the prodelta contribution in the DIP source and sink.

As can be seen from Table 7, the DIP fluxes recycled from the sediment accounts for 19-35% of the riverine discharge input of DIP which indicates that this source could represent an additional source corresponding to 25% of the Rhône input. We propose here that it can be considered as an additional source of DIP to the Mediterranean Sea since the largest part of this benthic flux originates from the recycling of organic P of terrestrial origin. These calculated proportions surely contain uncertainties on some assumptions such as the DIP inputs and the extrapolation of summer values of P budgets over the year. However, the calculation at the deltaic scale over the entire year was necessary in order to compare the sediment source to the river input. These calculations are important because the sediments contribution was substantial compared to riverine inputs, especially near the river mouth and no estimations were available before this study for the Rhône River prodelta. Finally, we estimate that the obtained DIP flux

from the sediment may be overestimated by less than one third due to the extrapolation over the year.

In the proximal and prodelta zones, it corresponds to the recycling of terrestrial organic matter (Cathalot et al. 2013) and is not related to internal recycling of marine (or deltaic) production based on riverine DIP input. It thus constitutes a real addition to the DIP input from the Rhône River. This is probably less true for the continental shelf sediments (33% of benthic DIP recycling) where marine organic matter produced by DIP originating from the Rhône River is partly recycled in surface sediments, therefore not totally contributing as an additional source. Considering that, the 25% addition to the riverine flux by benthic recycling of organic P can be seen as an upper boundary.

Burial of Ca-bound P_{auth} phase accounts for around 3-6% of the riverine DIP flux, which is minor sink. Furthermore, as most of this precipitation and burial of Ca-bound P_{auth} is linked to DIP production by mineralization in the sediment (see Fig. 11), it does not constitute a true sink of DIP delivered from the Rhône River. Indeed, as stated above, most of the mineralized organic P in the proximal and prodelta zones originates from the continent and is independent of the DIP source from the Rhône River. Only the share of Ca-bound P_{auth} burial in continental shelf sediments can be considered a sink of DIP originating from the Rhône River as organic matter mineralized in these sediments is of marine origin based on nutrients from the Rhône River. As it represents only 20% of the total Ca-bound P_{auth} burial, it can be considered that a very limited fraction of the DIP input by the river (<1%) is precipitated as Ca-bound P and buried in sediments.

6. Conclusion

In order to gain new insights about organic matter and phosphorus transformation in the sediments along the Mediterranean Rhône River prodelta, we used a modelling approach based on an early diagenetic model (OMEXDIA) coupled to the dataset from proximal, prodelta and distal domains. The model application highlighted the intensity of mineralization processes from the Rhône River mouth to the adjacent continental shelf. This intensity was mainly driven by high deposition of OC in the sediments, as a characteristic of organic-rich ecosystems, with a large proportion of terrestrial organic matter, hence providing a biogeochemical gradient. The combined model parameters indicated that the OC fluxes to the seafloor led to the increase of oxic mineralization rates and reduction of anoxic mineralization rates contribution, as a function of distance from the Rhône River mouth. According to the model, the anoxic mineralization contributed to the large concentrations of mineralization products at the sediment depth such as DIC, DIP and NH_4^+ , especially in the proximal domain. Our modelling approach was extended to include the benthic P cycle along the Rhône River prodelta. It showed that there was an intense production of porewater DIP, mainly by organic P mineralization. The Fe-bound P was a small net source of DIP in the proximal zone and was negligible in the prodelta and distal domains, while this phase had a key role in trapping and transferring DIP in sediment, promoting the Ca-bound P precipitation in anoxic sediments in the proximal domain. However, the sediments along the Rhône River prodelta acted mainly as secondary source of DIP to the overlying waters, corresponding to a maximum of 25% of the riverine input, as only a minor fraction was precipitated to Ca-bound P.

Data availability: The entire dataset is provided on the SeaNoe database

Ait Ballagh Fatima Ezzahra, Rabouille Christophe, Andrieux-Loyer Françoise, Soetaert Karline, Lansard Bruno, Bombled Bruno, Monvoisin Gaël, Elkalay Khalid, Khalil Karima

(2020). **Spatial variability of organic matter and phosphorus cycling in Rhône River prodelta sediments (NW Mediterranean Sea, France): a data-model approach.** SEANOE. <https://doi.org/10.17882/73204>

7. References

- Ait Ballagh, F. E., C. Rabouille, F. Andrieux-loyer, K. Soetaert, K. Elkalay, and K. Khalil. 2020. Spatio-temporal dynamics of sedimentary phosphorus along two temperate eutrophic estuaries : A data-modelling approach. *Continental Shelf Research* 193. Elsevier Ltd: 104037. <https://doi.org/10.1016/j.csr.2019.104037>.
- Andrieux-Loyer, F., and A. Aminot. 2001. Phosphorus forms related to sediment grain size and geochemical characteristics in french coastal areas. *Estuarine, Coastal and Shelf Science* 52: 617–629. <https://doi.org/10.1006/ecss.2001.0766>.
- Andrieux-Loyer, F., X. Philippon, G. Bally, R. Kérouel, A. Youenou, and J. Le Grand. 2008. Phosphorus dynamics and bioavailability in sediments of the Penzé Estuary (NW France): in relation to annual P-fluxes and occurrences of *Alexandrium Minutum*. *Biogeochemistry* 88: 213–231. <https://doi.org/10.1007/sl0533-008-9199-2>.
- Anschutz, P., S. Zhong, B. Sundby, A. Mucci, and C. Gobeil. 1998. Burial efficiency of phosphorus and the geochemistry of iron in continental margin sediments. *Limnology and Oceanography* 43: 53–64. <https://doi.org/10.4319/lo.1998.43.1.0053>.
- Antoine, D., and A. Morel. 1995. Algal pigment distribution and primary production in the eastern Mediterranean as derived from coastal zone color scanner observations. *Journal of Physical Oceanography Research* 100: 16,193-16,209.
- Antonelli, C., F. Eyrolle, B. Rolland, M. Provansal, and F. Sabatier. 2008. Suspended sediment and ¹³⁷Cs fluxes during the exceptional December 2003 flood in the Rhone River, southeast France. *Geomorphology* 95: 350–360. <https://doi.org/10.1016/j.geomorph.2007.06.007>.
- Antonelli, C., M. Provansal, and C. Vella. 2004. Recent morphological channel changes in a deltaic environment. The case of the Rhône River, France. *Geomorphology* 57: 385–402. [https://doi.org/10.1016/S0169-555X\(03\)00167-3](https://doi.org/10.1016/S0169-555X(03)00167-3).
- Arnau, P., C. Lique, and M. Canals. 2004. River mouth plume events and their dispersal in the northwestern Mediterranean Sea. *Oceanography* 17: 23–31.
- Balls, P. W. 1994. Nutrient Inputs to Estuaries from Nine Scottish East Coast Rivers; Influence of Estuarine Processes on Inputs to the North Sea. *Estuarine, Coastal and Shelf Science*. <https://doi.org/10.1006/ecss.1994.1068>.
- Batker, D., I. De la Torre, R. Costanza, J.W. Day, P. Swedeen, R. Boumans, and K.J. Bagstad. 2014. The Threats to the Value of Ecosystem Goods and Services of the Mississippi Delta.

782 In *Perspectives on the restoration of the Mississippi Delta: The once and future delta*,
783 155–173.

784 Benitez-Nelson, C. R. 2000. The biogeochemical cycling of phosphorus in marine systems.
785 *Earth Science Reviews* 51: 109–135. [https://doi.org/10.1016/S0012-8252\(00\)00018-0](https://doi.org/10.1016/S0012-8252(00)00018-0).

786 Berg, P., S. Rysgaard, and B. Thamdrup. 2003. Dynamic modeling of early diagenesis and
787 nutrient cycling. A case study in an arctic sediment. *American Journal of Science* 303:
788 905–955.

789 Berner, R. A. 1970. Sedimentary pyrite formation. *American Journal of Science* 268: 1–23.

790 Berner, R. A. 1980. Early Diagenesis: A Theoretical Approach. In *Princeton: Princeton*
791 *University Press*, 241 pp. <https://doi.org/10.1016/j.giq.2006.06.002>.

792 Berner, R. A. 1989. Biogeochemical cycles of carbon and sulfur and their effect on atmospheric
793 oxygen over phanerozoic time. *Global and Planetary Change* 1: 97–122.
794 [https://doi.org/10.1016/0921-8181\(89\)90018-0](https://doi.org/10.1016/0921-8181(89)90018-0).

795 Bianchi, T. S., and M. A. Allison. 2009. Large-river delta-front estuaries as natural “recorders”
796 of global environmental change. *Proceedings of the National Academy of Sciences* 106:
797 8085–8092. <https://doi.org/10.1073/pnas.0812878106>.

798 Boudreau, B. P. 1997. *Diagenetic models and their implementation. Modelling transport and*
799 *reactions in aquatic sediments. Springer-Verlag.* [https://doi.org/10.1016/S0264-](https://doi.org/10.1016/S0264-8172(98)80005-6)
800 8172(98)80005-6.

801 Boudreau, Bernard P. 1996. A method-of-line code for carbon and nutrient daigenesis in aquatic
802 sediments. *Computer & Geosciences* 22: 479–496.

803 Burdige, D. J. 2007. Preservation of Organic Matter in Marine Sediments: Controls,
804 Mechanisms, and an Imbalance in Sediment Organic Carbon Budgets? *Chemical Reviews*
805 107: 467–485. <https://doi.org/10.1021/cr050347q>.

806 Cade-Menun, B. J., S. Duhamel, R. J. Dodd, C. Lønborg, C. T. Parsons, and W. D. Taylor.
807 2019. Editorial: Phosphorus Along the Soil-Freshwater-Ocean Continuum. In *Frontiers in*
808 *Marine Science*, 10:1–3. <https://doi.org/10.1071/en13151>.

809 Cai, W. J., and F. L. Sayles. 1996. Oxygen penetration depths and fluxes in marine sediments.
810 *Marine Chemistry* 52: 123–131. [https://doi.org/10.1016/0304-4203\(95\)00081-X](https://doi.org/10.1016/0304-4203(95)00081-X).

811 Canavan, R. W., and C. P. Slomp. 2006. Phosphorus cycling in the sediment of a coastal
812 freshwater lake and response to salinization. In *Biogeochemical Cycling of Nutrients and*
813 *Trace Metals in the Sediment of Haringvliet Lake: Response to Salinization*, 95–113.

814 Canavan, Richard W., Caroline P. Slomp, Parisa Jourabchi, Philippe Van Cappellen, Annet M.
815 Laverman, and Gerard A. van den Berg. 2006. Organic matter mineralization in sediment
816 of a coastal freshwater lake and response to salinization. *Geochimica et Cosmochimica*
817 *Acta* 70: 2836–2855. <https://doi.org/10.1016/j.gca.2006.03.012>.

818 Canfield, D. E., E. Kristensen, and B. Thamdrup. 2005. *The Iron and Manganese Cycles*.

819 *Advances in Marine Biology*. Vol. 48. [https://doi.org/10.1016/S0065-2881\(05\)48008-6](https://doi.org/10.1016/S0065-2881(05)48008-6).

820 Cathalot, C. 2008. Devenir et impact des apports fluviaux sur les marges continentales :
821 importance biogéochimique et environnementale du recyclage dans les sédiments du
822 prodelta du Rhône. Ph.D. thesis, Université Pierre et Marie Curie, Paris, France.

823 Cathalot, C., C. Rabouille, L. Pastor, B. Deflandre, E. Viollier, R. Buscail, A. Grémare, C.
824 Treignier, and A. Pruski. 2010. Temporal variability of carbon recycling in coastal
825 sediments influenced by rivers : assessing the impact of flood inputs in the Rhone River
826 prodelta. *Biogeosciences* 7: 1187–1205.

827 Cathalot, C., C. Rabouille, N. Tisnérat-Laborde, F. Toussaint, P. Kerhervé, R. Buscail, K.
828 Loftis, et al. 2013. The fate of river organic carbon in coastal areas: A study in the Rhône
829 River delta using multiple isotopic ($\delta^{13}\text{C}$, $\delta^{14}\text{C}$) and organic tracers. *Geochimica et*
830 *Cosmochimica Acta* 118: 33–55. <https://doi.org/10.1016/j.gca.2013.05.001>.

831 Charmasson, S., O. Radakovitch, M. Arnaud, P. Bouisset, and A. S. Pruchon. 1998. Long-core
832 profiles of ^{137}Cs , ^{134}Cs , ^{60}Co and ^{210}Pb in sediment near the Rhône River (Northwestern
833 Mediterranean Sea). *Estuaries* 21: 367–378. <https://doi.org/10.1007/BF02690450>.

834 Clayton, T. D., and R. H. Byrne. 1993. Spectrophotometric seawater pH measurements : total
835 hydrogen results. *Deep-Sea Research* 40: 2115–2129. [https://doi.org/10.1016/0967-](https://doi.org/10.1016/0967-0637(93)90048-8)
836 [0637\(93\)90048-8](https://doi.org/10.1016/0967-0637(93)90048-8).

837 Cloern, J. E. 2001. Our evolving conceptual model of the coastal eutrophication problem.
838 *Marine Ecology Progress Series* 210: 223–253.

839 Dedieu, K., C. Rabouille, F. Gilbert, K. Soetaert, E. Metzger, C. Simonucci, D. Jézéquel, et al.
840 2007. Coupling of carbon, nitrogen and oxygen cycles in sediments from a Mediterranean
841 lagoon: A seasonal perspective. *Marine Ecology Progress Series* 346: 45–59.
842 <https://doi.org/10.3354/meps07031>.

843 Delaney, M L. 1998. Phosphorus accumulation in marine sediments and the oceanic phosphorus
844 cycle. *Glob. Biogeochem. Cycles*, 12: 563–572.

845 Diaz, F., P. Raimbault, B. Boudjellal, N. Garcia, and T. Moutin. 2001. Early spring phosphorus
846 limitation of primary productivity in a NW Mediterranean coastal zone (Gulf of Lions)
847 211: 51–62.

848 Diaz, R. J., and R. Rosenberg. 2008. Spreading dead zones and consequences for marine
849 ecosystems. *Science* 321: 926–929. <https://doi.org/10.1126/science.1156401>.

850 Dickson, A G, C. L. Sabine, and J. R. Christian. 2007. Guide to Best Practices for Ocean CO₂
851 Measurements. In *PICES Special Publication*, 3:191 pp.

852 Dijkstra, N., C. P. Slomp, and T. Behrends. 2016. Vivianite is a key sink for phosphorus in
853 sediments of the Landsort Deep, an intermittently anoxic deep basin in the Baltic Sea.
854 *Chemical Geology* 438. Elsevier B.V.: 58–72.
855 <https://doi.org/10.1016/j.chemgeo.2016.05.025>.

856 Downing, J. A., M. McClain, R. Twilley, J. M. Melack, J. Elser, N. N. Rabalais, W. M. Lewis,
857 et al. 1999. The impact of accelerating land-use change on the N-cycle of tropical aquatic
858 ecosystems: Current conditions and projected changes. *Biogeochemistry* 46: 109–148.
859 <https://doi.org/10.1007/BF01007576>.

860 Dugdale, R. C., and F. P. Wilkerson. 1988. Nutrient sources and primary production in the
861 Eastern Mediterranean. *Oceanologica Acta*: 179–184.

862 Dumoulin, J. P., L. Pozzato, J. Rassman, F. Toussaint, M. Fontugne, N. Tisnérat-Laborde, L.
863 Beck, et al. 2018. Isotopic signature ($\delta^{13}\text{C}$, $\delta^{14}\text{C}$) of DIC in sediment pore waters: An
864 example from the rhone river delta. *Radiocarbon* 60: 1465–1481.
865 <https://doi.org/10.1017/RDC.2018.111>.

866 Egger, M., T. Jilbert, T. Behrends, C. Rivard, and C. P. Slomp. 2015. Vivianite is a major sink
867 for phosphorus in methanogenic coastal surface sediments. *Geochimica et Cosmochimica*
868 *Acta* 169. Elsevier Ltd: 217–235. <https://doi.org/10.1016/j.gca.2015.09.012>.

869 Emerson, S., J. Hedges, H. D. Holland, and K. K. Turekian. 2004. Sediment diagenesis and
870 benthic flux. *The Oceans and Marine Geochemistry* 6: 293–319.

871 Emerson, S., and G. Widmer. 1978. Early diagenesis in anaerobic lake sediments-II.
872 Thermodynamic and kinetic factors controlling the formation of iron phosphate.
873 *Geochimica et Cosmochimica Acta* 42: 1307–1316. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-7037(78)90035-2)
874 [7037\(78\)90035-2](https://doi.org/10.1016/0016-7037(78)90035-2).

875 Emsley, J. 1980. The Phosphorus Cycle. In *The handbook of environmental chemistry*, 20:147–
876 167. [https://doi.org/10.1016/0143-1471\(82\)90111-8](https://doi.org/10.1016/0143-1471(82)90111-8).

877 Estournel, C., V. Kondrachoff, P. Marsaleix, and R. Vehil. 1997. The plume of the Rhone:
878 Numerical simulation and remote sensing. *Continental Shelf Research* 17: 899–924.
879 [https://doi.org/10.1016/S0278-4343\(96\)00064-7](https://doi.org/10.1016/S0278-4343(96)00064-7).

880 Eyrolle, F., O. Radakovitch, P. Raimbault, S. Charmasson, C. Antonelli, E. Ferrand, D. Aubert,
881 G. Raccasi, S. Jacquet, and R. Gurriaran. 2012. Consequences of hydrological events on
882 the delivery of suspended sediment and associated radionuclides from the Rhône River to
883 the Mediterranean Sea. *Journal of Soils and Sediments* 12: 1479–1495.
884 <https://doi.org/10.1007/s11368-012-0575-0>.

885 Ferrón, S, T Ortega, and J M Forja. 2009. Benthic respiration on the northeastern shelf of the
886 Gulf of Cádiz (SW Iberian Peninsula). *Marine Ecology Progress Series* 392: 69–80.
887 <https://doi.org/10.3354/meps08240>.

888 Filippelli, G. M., and M. L. Delaney. 1996. Phosphorus geochemistry of equatorial Pacific
889 sediments. *Geochimica et Cosmochimica Acta* 60: 1479–1495.
890 [https://doi.org/10.1016/0016-7037\(96\)00042-7](https://doi.org/10.1016/0016-7037(96)00042-7).

891 Fox, L. E. 1990. Geochemistry of dissolved phosphate in the Sepik River and Estuary, Papua,
892 New Guinea. *Geochimica et Cosmochimica Acta* 54: 1019–1024.
893 [https://doi.org/10.1016/0016-7037\(90\)90435-N](https://doi.org/10.1016/0016-7037(90)90435-N).

894 Froelich, P. N. 1988. Kinetic control of dissolved phosphate in natural rivers and estuaries: A
895 primer on the phosphate buffer mechanism. *Limnology and Oceanography* 33: 649–668.

896 Gatti, J., A. Petrenko, J. L. Devenon, Y. Leredde, and C. Ulses. 2006. The Rhone river dilution
897 zone present in the northeastern shelf of the Gulf of Lion in December 2003. *Continental*
898 *Shelf Research* 26: 1794–1805. <https://doi.org/10.1016/j.csr.2006.05.012>.

899 Got, H., and J. C. Aloisi. 1990. The Holocene sedimentation on the Gulf of Lions margin: a
900 quantitative approach. *Continental Shelf Research* 10: 841–855.
901 [https://doi.org/10.1016/0278-4343\(90\)90062-Q](https://doi.org/10.1016/0278-4343(90)90062-Q).

902 Grasshof, K., M. Ehrhardt, and K. Kremling. 1983. *Methods of seawater analysis, Third,*
903 *Completely Revised and Extended Edition.* Verlag Chemie GmbH.
904 [https://doi.org/10.1016/0304-4203\(78\)90045-2](https://doi.org/10.1016/0304-4203(78)90045-2).

905 Gypens, Nathalie, Christiane Lancelot, and Karline Soetaert. 2008. Simple parameterisations
906 for describing N and P diagenetic processes: Application in the North Sea. *Progress in*
907 *Oceanography* 76: 89–110. <https://doi.org/10.1016/j.pocean.2007.10.003>.

908 Hammond, D E. 1999. Diagenesis of carbon and nutrients and benthic exchange in sediments
909 of the Northern Adriatic Sea. *Marine Chemistry* 66: 53–79.

910 Hansen, H. P., and F. Koroleff. 1999. Determination of nutrients. In *Methods of Seawater*
911 *Analysis, Third Edition, Completely Revised and Extended Edition,* 159–228.
912 <https://doi.org/10.1002/9783527613984>.

913 Hofmann, A. F., K. Soetaert, and J. J. Middelburg. 2008. Present nitrogen and carbon dynamics
914 in the Scheldt estuary using a novel 1-D model. *Biogeosciences* 5: 981–1006.
915 <https://doi.org/10.5194/bg-5-981-2008>.

916 Howarth, RW, G Billen, D Swaney, A Townsend, N Jaworski, K Lajtha, JA Downing, et al.
917 1996. Regional nitrogen budgets and riverine N & P fluxes for the drainages to the North
918 Atlantic Ocean: Natural and human influences. *Biogeochemistry* 35: 75–139.
919 <https://doi.org/10.1590/2318-0331.011616030>.

920 Ibañez, C., D. Pont, and N. Prat. 1997. Characterization of the Ebre and Rhone estuaries: A
921 basis for defining and classifying salt-wedge estuaries. *Limnology and Oceanography* 42:
922 89–101. <https://doi.org/10.4319/lo.1997.42.1.0089>.

923 Jackson, J. B.C., M. X. Kirby, W. H. Berger, K. A. Bjorndal, L. W. Botsford, B. J. Bourque, R.
924 H. Bradbury, et al. 2001. Historical overfishing and the recent collapse of coastal
925 ecosystems. *Science* 293: 629–637. <https://doi.org/10.1126/science.1059199>.

926 Joshi, S. R., R. K. Kukkadapu, D. J. Burdige, M. E. Bowden, D. L. Sparks, and D. P. Jaisi.
927 2015. Organic matter remineralization predominates phosphorus cycling in the mid-bay
928 sediments in the chesapeake bay. *Environmental Science and Technology* 49: 5887–5896.
929 <https://doi.org/10.1021/es5059617>.

930 Khalil, K, A M Laverman, M Raimonet, and C Rabouille. 2018. Importance of nitrate reduction
931 in benthic carbon mineralization in two eutrophic estuaries : Modeling , observations and

laboratory experiments. *Marine Chemistry* 199. Elsevier: 24–36.
<https://doi.org/10.1016/j.marchem.2018.01.004>.

Kraal, P., E. D. Burton, A. L. Rose, B. D. Kocar, R. S. Lockhart, K. Grice, R. T. Bush, E. Tan, and S. M. Webb. 2015. Sedimentary iron-phosphorus cycling under contrasting redox conditions in a eutrophic estuary. *Chemical Geology* 392. Elsevier B.V.: 19–31.
<https://doi.org/10.1016/j.chemgeo.2014.11.006>.

Krom, M. D., and R. A. Berner. 1981. The diagenesis of phosphorus in a nearshore marine sediment. *Geochimica et Cosmochimica Acta* 45: 207–216. [https://doi.org/10.1016/0016-7037\(81\)90164-2](https://doi.org/10.1016/0016-7037(81)90164-2).

Krom, M. D., N. Kress, S. Brenner, and L. I. Gordon. 1991. Phosphorus limitation of primary productivity in the eastern Mediterranean Sea. *Limnology and Oceanography* 36: 424–432. <https://doi.org/10.4319/lo.1991.36.3.0424>.

Lansard, B., S. Charmasson, C. Gascó, M. P. Antón, C. Grenz, and M. Arnaud. 2007. Spatial and temporal variations of plutonium isotopes (²³⁸Pu and ^{239,240}Pu) in sediments off the Rhone River mouth (NW Mediterranean). *Science of the Total Environment* 376: 215–227. <https://doi.org/10.1016/j.scitotenv.2007.01.069>.

Lansard, B., C. Rabouille, L. Denis, and C. Grenz. 2008. In situ oxygen uptake rates by coastal sediments under the influence of the Rhône River (NW Mediterranean Sea). *Continental Shelf Research* 28: 1501–1510. <https://doi.org/10.1016/j.csr.2007.10.010>.

Lansard, B., C. Rabouille, L. Denis, and C. Grenz. 2009. Benthic remineralization at the land-ocean interface: A case study of the Rhône River (NW Mediterranean Sea). *Estuarine, Coastal and Shelf Science* 81. Elsevier Ltd: 544–554.
<https://doi.org/10.1016/j.ecss.2008.11.025>.

Lebo, M. E. 1991. Particle-bound phosphorus along an urbanized coastal plain estuary. *Marine Chemistry* 34: 225–246. [https://doi.org/10.1016/0304-4203\(91\)90005-H](https://doi.org/10.1016/0304-4203(91)90005-H).

Lehtoranta, J., O. P. Savchuk, J. Elken, K. Dahlbo, H. Kuosa, M. Raateoja, P. Kauppila, A. Räike, and H. Pitkänen. 2017. Atmospheric forcing controlling inter-annual nutrient dynamics in the open Gulf of Finland. *Journal of Marine Systems* 171. Elsevier B.V: 4–20. <https://doi.org/10.1016/j.jmarsys.2017.02.001>.

Lenstra, W. K., M. Egger, N. A. G. M. van Helmond, E. Kritzberg, D. J. Conley, and C. P. Slomp. 2018. Variations in river input of iron impact sedimentary phosphorus burial in an oligotrophic Baltic Sea estuary. *Biogeosciences Discussions* 15: 6979–6996.
<https://doi.org/10.1021/pr100554m>.

Lochet, F., and M. Leveau. 1990. Transfers between a eutrophic ecosystem, the river Rhône, and an oligotrophic ecosystem, the north-western Mediterranean Sea. *Hydrobiologia* 207: 95–103.

Ludwig, W., E. Dumont, M. Meybeck, and S. Heussner. 2009. River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during

970 past and future decades? *Progress in Oceanography* 80. Elsevier Ltd: 199–217.
971 <https://doi.org/10.1016/j.pocean.2009.02.001>.

972 Ludwig, W., and J. L. Probst. 1998. River sediment discharge to the oceans: Present-day
973 controls and global budgets. *American Journal of Science* 298: 265–295.
974 <https://doi.org/10.2475/ajs.298.4.265>.

975 Malagó, A., F. Bouraoui, B. Grizzetti, and A. De Roo. 2019. Modelling nutrient fluxes into the
976 Mediterranean Sea. *Journal of Hydrology: Regional Studies* 22. Elsevier: 100592.
977 <https://doi.org/10.1016/j.ejrh.2019.01.004>.

978 McKee, B. A., R. C. Aller, M. A. Allison, T. S. Bianchi, and G. C. Kineke. 2004. Transport and
979 transformation of dissolved and particulate materials on continental margins influenced by
980 major rivers: Benthic boundary layer and seabed processes. *Continental Shelf Research*
981 24: 899–926. <https://doi.org/10.1016/j.csr.2004.02.009>.

982 Meybeck, M. 1982. Carbon, nitrogen, and phosphorus transport by world rivers. *American*
983 *Journal of Science* 282: 401–450.

984 Milliman, J. D., and R. H. Meade. 1983. World-Wide Delivery of River Sediment to the Oceans.
985 *The Journal of Geology* 91: 1–21.

986 Miralles, J., O. Radakovitch, and J. C. Aloisi. 2005. 210Pb sedimentation rates from the
987 Northwestern Mediterranean margin. *Marine Geology* 216: 155–167.
988 <https://doi.org/10.1016/j.margeo.2005.02.020>.

989 Morse, J. W., and P. M. Eldridge. 2007. A non-steady state diagenetic model for changes in
990 sediment biogeochemistry in response to seasonally hypoxic/anoxic conditions in the
991 “dead zone” of the Louisiana shelf. *Marine Chemistry* 106: 239–255.
992 <https://doi.org/10.1016/j.marchem.2006.02.003>.

993 Moutin, T., P. Raimbault, H.L. Golterman, and B. Coste. 1998. The input of nutrients by the
994 Rhône river into the Mediterranean Sea: recent observations and comparison with earlier
995 data. *Hydrobiologia* 373/374: 237–246. <https://doi.org/10.1023/A>.

996 Moutin, T, and P Raimbault. 2002. Primary production , carbon export and nutrients availability
997 in western and eastern Mediterranean Sea in early summer 1996 (MINOS cruise). *Journal*
998 *of Marine Systems* 33–34: 273–288.

999 Moutin, T, F Van Wambeke, and L Prieur. 2012. Introduction to the Biogeochemistry from the
1000 Oligotrophic to the Ultraoligotrophic Mediterranean (BOUM) experiment. *Biogeosciences*
1001 9: 3817–3825. <https://doi.org/10.5194/bg-9-3817-2012>.

1002 Muller-Karger, F. E., R. Varela, R. Thunell, R. Luerssen, C. Hu, and J. J. Walsh. 2005. The
1003 importance of continental margins in the global carbon cycle. *Geophysical Research*
1004 *Letters* 32: 1–4. <https://doi.org/10.1029/2004GL021346>.

1005 Murphy, J., and J. P. Riley. 1962. A modified single solution method for determination of
1006 phosphate in natural waters. *Analytica Chimica Acta* 27: 31–36.
1007 <https://doi.org/10.1057/9781137461131>.

- 1008 Naudin, J. J., G. Cauwet, M. J. Chrétiennot-Dinet, B. Deniaux, J. L. Devenon, and H. Pauc.
1009 1997. River discharge and wind influence upon particulate transfer at the land-ocean
1010 interaction: Case study of the Rhone River plume. *Estuarine, Coastal and Shelf Science*
1011 45: 303–316. <https://doi.org/10.1006/ecss.1996.0190>.
- 1012 Nixon, S. W. 1995. Coastal marine eutrophication: A definition, social causes, and future
1013 concerns. *Ophelia* 41: 199–219. <https://doi.org/10.1080/00785236.1995.10422044>.
- 1014 Ollivier, P., B. Hamelin, and O. Radakovitch. 2010. Seasonal variations of physical and
1015 chemical erosion: A three-year survey of the Rhone River (France). *Geochimica et*
1016 *Cosmochimica Acta* 74. Elsevier Ltd: 907–927. <https://doi.org/10.1016/j.gca.2009.10.037>.
- 1017 Pastor, L., C. Cathalot, B. Deflandre, E. Viollier, K. Soetaert, F. J.R. Meysman, C. Ulses, E.
1018 Metzger, and C. Rabouille. 2011. Modeling biogeochemical processes in sediments from
1019 the Rhône River prodelta area (NW Mediterranean Sea). *Biogeosciences* 8: 1351–1366.
1020 <https://doi.org/10.5194/bg-8-1351-2011>.
- 1021 Pont, D., J. P. Simonnet, and A. V. Walter. 2002. Medium-term changes in suspended sediment
1022 delivery to the ocean: Consequences of catchment heterogeneity and river management
1023 (Rhône River, France). *Estuarine, Coastal and Shelf Science* 54: 1–18.
1024 <https://doi.org/10.1006/ecss.2001.0829>.
- 1025 Powley, H. R., M. D. Krom, and P. Van Cappellen. 2017. Understanding the unique
1026 biogeochemistry of the Mediterranean Sea: Insights from a coupled phosphorus and
1027 nitrogen model. *Global Biogeochemical Cycles* 31: 1010–1031.
1028 <https://doi.org/10.1002/2017GB005648>.
- 1029 Pozzato, L., J. Rassmann, B. Lansard, J-P. Dumoulin, P. van Breugel, C. Rabouille. 2018.
1030 Origin of remineralized organic matter in sediments from the Rhone River prodelta (NW
1031 Mediterranean) traced by $\delta^{14}\text{C}$ and $\delta^{13}\text{C}$ signatures of pore water DIC. *Progr. Oceanogr.*,
1032 163: 112-122.
- 1033 Provansal, M., O. Radakovitch, F. Sabatier, and A. Clémens. 2012. *Le Rhône aval en 21*
1034 *questions. Rapport ZABR, Graie Editors, Villeurbanne, France*
- 1035 Psenner, R., B. Bostrom, M. Dinka, K. Pettersson, R. Pucsko, and M. Sager. 1988. Fractionation
1036 of phosphorus in suspended matter and sediment. *Ergebnisse der Limnologie* 22: 219–228.
- 1037 Rabouille, C., D. J. Conley, M. H. Dai, W. J. Cai, C. T.A. Chen, B. Lansard, R. Green, et al.
1038 2008. Comparison of hypoxia among four river-dominated ocean margins: The
1039 Changjiang (Yangtze), Mississippi, Pearl, and Rhône rivers. *Continental Shelf Research*
1040 28: 1527–1537. <https://doi.org/10.1016/j.csr.2008.01.020>.
- 1041 Rabouille, C., L. Denis, K. Dedieu, G. Stora, B. Lansard, and C. Grenz. 2003. Oxygen demand
1042 in coastal marine sediments: Comparing in situ microelectrodes and laboratory core
1043 incubations. *Journal of Experimental Marine Biology and Ecology* 285–286: 49–69.
1044 [https://doi.org/10.1016/S0022-0981\(02\)00519-1](https://doi.org/10.1016/S0022-0981(02)00519-1).
- 1045 Rabouille, C., and J. F. Gaillard. 1991. Towards the EDGE: Early diagenetic global explanation.

1046 A model depicting the early diagenesis of organic matter, O₂, NO₃, Mn, and PO₄.
 1047 *Geochimica et Cosmochimica Acta* 55: 2511–2525. [https://doi.org/10.1016/0016-](https://doi.org/10.1016/0016-7037(91)90369-G)
 1048 7037(91)90369-G.

1049 Radakovitch, O, S Charmasson, M Arnaud, and P Bouisset. 1999. 210Pb and Caesium
 1050 Accumulation in the Rhône Delta. *Estuarine, Coastal and Shelf Science* 48: 77–92.

1051 Rassmann, J., E. M. Eitel, C. Cathalot, C. Brandily, B. Lansard, M. Taillefert, and C. Rabouille.
 1052 2020. Benthic alkalinity and DIC fluxes in the Rhône River prodelta generated by
 1053 decoupled aerobic and anaerobic processes. *Biogeosciences Discussions*.

1054 Rassmann, J., B. Lansard, L. Pozzato, and C. Rabouille. 2016. Carbonate chemistry in sediment
 1055 porewaters of the Rhône River delta driven by early diagenesis (northwestern
 1056 Mediterranean). *Biogeosciences* 13: 5379–5394. [https://doi.org/10.5194/bg-13-5379-](https://doi.org/10.5194/bg-13-5379-2016)
 1057 2016.

1058 Ratmaya, W. 2018. Rôle des sédiments dans le cycle des nutriments et impacts sur
 1059 l'eutrophisation des écosystèmes côtiers. Ph.D. thesis, Univ. Bretagne Occidentale, Brest,
 1060 France.

1061 Revsbech, N. P. 1989. An oxygen microsensor with a guard cathode. *Limnology and*
 1062 *Oceanography* 34: 474–478. <https://doi.org/10.4319/lo.1989.34.2.0474>.

1063 Ruttenberg, K.C. 1992. Development of a sequential extraction method for different forms of
 1064 phosphorus in marine sediments. *Limnology and Oceanography* 37: 1460–1482.
 1065 <https://doi.org/10.4319/lo.1992.37.7.1460>.

1066 Ruttenberg, K.C. 2014. 10.13 – The Global Phosphorus Cycle. *Treatise on Geochemistry*: 499–
 1067 558. <https://doi.org/10.1016/B978-0-08-095975-7.00813-5>.

1068 Ruttenberg, K.C., and R. A. Berner. 1993. Authigenic apatite formation and burial in sediments
 1069 from non-upwelling, continental margin environments. *Geochimica et Cosmochimica*
 1070 *Acta* 57: 991–1007. [https://doi.org/10.1016/0016-7037\(93\)90035-U](https://doi.org/10.1016/0016-7037(93)90035-U).

1071 Sabatier, F., G. Maillet, M. Provansal, T. J. Fleury, S. Suanez, and C. Vella. 2006. Sediment
 1072 budget of the Rhône delta shoreface since the middle of the 19th century. *Marine Geology*
 1073 234: 143–157. <https://doi.org/10.1016/j.margeo.2006.09.022>.

1074 Schnetger, B., and C. Lehnert. 2014. Determination of nitrate plus nitrite in small volume
 1075 marine water samples using vanadium(III)chloride as a reduction agent. *Marine Chemistry*
 1076 160. Elsevier B.V.: 91–98. <https://doi.org/10.1016/j.marchem.2014.01.010>.

1077 Seeberg-Elverfeldt, J., M. Schlüter, T. Feseker, and M. Kölling. 2005. Rhizon sampling of
 1078 porewaters near the sediment-water interface of aquatic systems. *Limnology and*
 1079 *Oceanography: Methods* 3: 361–371. <https://doi.org/10.4319/lom.2005.3.361>.

1080 Slomp, C. P., E. H. G. Epping, W. Helder, and W. V. Raaphorst. 1996. A key role for iron-
 1081 bound phosphorus in authigenic apatite formation in North Atlantic continental platform
 1082 sediments. *Journal of Marine Research* 54: 1179–1205.
 1083 <https://doi.org/10.1357/0022240963213745>.

- 1084 Slomp, C. P., J. F. P. Malschaert, and W. V. Raaphorst. 1998. The role of adsorption in
1085 sediment-water exchange of phosphate in North Sea continental margin sediments.
1086 *Limnology and Oceanography* 43: 832–846.
- 1087 Slomp, C. P., J. Thomson, and G. J. De Lange. 2004. Controls on phosphorus regeneration and
1088 burial during formation of eastern Mediterranean sapropels. *Marine Geology* 203: 141–
1089 159. [https://doi.org/10.1016/S0025-3227\(03\)00335-9](https://doi.org/10.1016/S0025-3227(03)00335-9).
- 1090 Smith, V H, G D Tilman, and J C Nekola. 1999. Eutrophication : impacts of excess nutrient
1091 inputs on freshwater, marine, and terrestrial ecosystems. *Environmental Pollution* 100:
1092 179–196.
- 1093 Soetaert, K., P. M. J. Herman, and J. J. Middelburg. 1996. A model of early diagenetic processes
1094 from the shelf to abyssal depths. *Geochimica et Cosmochimica Acta* 60: 1019–1040.
1095 [https://doi.org/10.1016/0016-7037\(96\)00013-0](https://doi.org/10.1016/0016-7037(96)00013-0).
- 1096 Soetaert, K., and F. Meysman. 2012. Reactive transport in aquatic ecosystems: Rapid model
1097 prototyping in the open source software R. *Environmental Modelling and Software* 32.
1098 Elsevier Ltd: 49–60. <https://doi.org/10.1016/j.envsoft.2011.08.011>.
- 1099 Stookey, L. L. 1970. Ferrozine-A New Spectrophotometric Reagent for Iron. *Analytical*
1100 *Chemistry* 42: 779–781.
- 1101 Sundby, B., C. Gobeil, N. Silverberg, and A Mucci. 1992. The phosphorus cycle in coastal
1102 marine sediments. *Limnology and Oceanography* 37: 1129–1145.
1103 <https://doi.org/10.4319/lo.1992.37.6.1129>.
- 1104 Syers, J. K., R. F. Harris, and D. E. Armstrong. 1973. Phosphate Chemistry in Lake Sediments.
1105 *Journal of Environment Quality* 2: 1–14.
1106 <https://doi.org/10.2134/jeq1973.00472425000200010001x>.
- 1107 Syvitski, J. P. M., A. J. Kettner, I. Overeem, E. W. H. Hutton, M. T. Hannon, G. R. Brakenridge,
1108 J. W. Day, et al. 2009. Sinking Deltas. *Nature Geoscience* 2: 681–686.
- 1109 Thamdrup, B., H. Fossing and B.B. Jorgensen.1994. Manganese, iron, and sulfur cycling in a
1110 coastal marine sediment, Aarhus Bay, Denmark. *Geochimica et Cosmochimica Acta*, 58,
1111 5115-5129,
- 1112 Tromp, T.K., P. Van Cappellen, and R.M. Key. 1995. A global model for the early diagenesis
1113 of organic carbon and organic phosphorus in marine sediments. *Geochimica et*
1114 *Cosmochimica Acta* 59: 1259–1284. [https://doi.org/10.1016/0016-7037\(95\)00042-X](https://doi.org/10.1016/0016-7037(95)00042-X).
- 1115 Tyrrell, T. 1999. The relative influences of nitrogen and phosphorus on oceanic primary
1116 production. *Nature* 400: 525–531.
- 1117 Van Den Broeck, N., and T. Moutin. 2002. Phosphate in the sediments of the Gulf of Lions
1118 (NW Mediterranean Sea), relationship with input by the river Rhone. *Hydrobiologia* 472:
1119 85–94. <https://doi.org/10.1023/A:1016308931115>.
- 1120 Vitousek, P. M., H. A. Mooney, J. Lubchenco, and J. M. Melillo. 1997. Human Domination of

1121 Earth's Ecosystems. *Science* 277: 494–499. <https://doi.org/10.1126/science.277.5325.494>.

1122 Wang, Y.F., and P. Van Cappellen. 1996. A multicomponent reactive transport model of early
 1123 diagenesis: Application to redox cycling in coastal marine sediments. *Geochimica et*
 1124 *Cosmochimica Acta* 60: 2993–3014. [https://doi.org/10.1016/0016-7037\(96\)00140-8](https://doi.org/10.1016/0016-7037(96)00140-8).

1125 Wijsman, J. W.M., P. M.J. Herman, J. J. Middelburg, and K. Soetaert. 2002. A model for early
 1126 diagenetic processes in sediments of the continental shelf of the Black Sea. *Estuarine,*
 1127 *Coastal and Shelf Science* 54: 403–421. <https://doi.org/10.1006/ecss.2000.0655>.

1128 Wilkinson, G. M. 2017. *Eutrophication of Freshwater and Coastal Ecosystems. Encyclopedia*
 1129 *of Sustainable Technologies*. Vol. 4. Elsevier. [https://doi.org/10.1016/B978-0-12-409548-](https://doi.org/10.1016/B978-0-12-409548-9.10160-5)
 1130 [9.10160-5](https://doi.org/10.1016/B978-0-12-409548-9.10160-5).

1131

1 **Figures**

Fig. 1 Location of sampled stations in proximal (stations A, Z, AZ), prodelta (stations AK, K, B, L) and distal (C, E) domains of the Rhône River delta

Fig. 2 Schematic representation of benthic P cycle

Fig. 3-a Simulated (grey lines) and measured (symbols (•)) O_2 , NO_3^- , SO_4^{2-} , DIC, NH_3 , ODU and TOC in the proximal domain (st Z, A and AZ)

L

17

18

Fig. 3-b. Simulated (grey lines) and measured (symbols (•)) O_2 , NO_3^- , SO_4^{2-} , DIC, NH_3 , ODU and TOC in the prodelta domain (st AK, B, K and L)

C

19

E

20

21

Fig. 3-c. Simulated (grey lines) and measured (symbols (•)) O_2 , NO_3^- , SO_4^{2-} , DIC, NH_3 , ODU and TOC in the distal domain (st C and E)

Fig. 4 Simulated (grey lines) and measured (symbols (•)) DIP, organic P, Fe-bound P and Ca-bound P in the sampled stations

Fig. 5 Metabolized OC fluxes at sediment-water interface along the Rhône River
prodelta

Fig. 6 First-order decay rate constants of labile (a) and semi-labile (b) OM fractions as a function of distance from the River mouth

Fig. 7 Mineralization pathways contribution (%) in the sampled stations

Fig. 8 Integrated rates of DIP production ($\mu\text{mol m}^{-2} \text{d}^{-1}$), for organic P mineralization and net Fe-bound P release.

Fig. 9 Integrated Fe-bound P ($\mu\text{mol m}^{-2} \text{d}^{-1}$) formation and release rates

Fig. 10 DIP losses ($\mu\text{mol m}^{-2} \text{d}^{-1}$) along the Rhône River prodelta

Fig. 11 Sedimentary P budgets ($\mu\text{mol m}^{-2} \text{d}^{-1}$) for typical stations of the different zones of the Rhône River prodelta: proximal (station A), prodelta (station K) and distal domains (station E)

Tables

Table 1 Sampling site locations and environmental parameters (bottom water temperature T_{bw} and salinity S) for each station during the benthic sampling

Site	Lat. (°N)	Long. (°E)	Dist. (km)	Depth (m)	T _{bw} (°C)	Salinity
Z	43°19.062’	4°52.034’	1.8	20.2	15.7	38
A	43°18.680’	4°51.038’	1.9	21	15.6	37.8
AZ	43°18.742’	4°51.400’	2.3	20	20	37.8
AK	43°18.426’	4°51.300’	2.7	43.7	15.6	37.9
B	43°18.262’	4°49.938’	3	48.5	15.8	37.9
K	43°18.111’	4°51.361’	3.3	57.3	15.7	38.5
L	43°18.270’	4°53.106’	4	60.8	15.1	38.1
C	43°16.237’	4°46.379’	8.6	72.5	14.7	38.2
E	43°13.333’	4°41.897’	16.6	72	14.8	38.3

Table 2 Mineralization pathways and diagenetic equations in the model

<i>Mineralization pathways</i>	<i>Reactions</i>	<i>Limitation and inhibition functions</i>		
<i>Oxic mineralization</i>	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4) + x\text{O}_2 \rightarrow x\text{CO}_2 + y\text{NH}_3 + \text{H}_3\text{PO}_4 + x \text{H}_2\text{O}$	$K_{\text{SO}_2\text{oxic}}$		R1
<i>Denitrification</i>	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4) + 0.8 * x\text{HNO}_3 \rightarrow x\text{CO}_2 + y\text{NH}_3 + \text{H}_3\text{PO}_4 + 0.4 x\text{N}_2 + 1.4 x \text{H}_2\text{O}$	$K_{\text{SNO}_3\text{denit}}$	$K_{\text{inO}_2\text{denit}}$	R2
<i>Metal reduction</i>	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4) + 1 \text{ oxidant} \rightarrow 1 \text{ ODU} + x \text{CO}_2 + y \text{NH}_3 + \text{H}_3\text{PO}_4 + 5 * x \text{H}_2\text{O}$		$K_{\text{inO}_2\text{anox}}$ $K_{\text{inNO}_3\text{anox}}$	R3
<i>Sulfate reduction</i>	$(\text{CH}_2\text{O})_x(\text{NH}_3)_y(\text{H}_3\text{PO}_4) + 0.5 \text{SO}_4^{2-} \rightarrow 0.5 \text{ ODU} + x \text{CO}_2 + y \text{NH}_3 + \text{H}_3\text{PO}_4 + 5 * x \text{H}_2\text{O}$	K_{SSO_4}		R4
<i>Nitrification</i>	$\text{NH}_3 + 2\text{O}_2 \rightarrow \text{HNO}_3 + \text{H}_2\text{O}$	$K_{\text{SO}_2\text{nitr}}, R_{\text{nitr}}$		R5
<i>Reoxidation of ODU</i>	$\text{ODU} + \text{O}_2 \rightarrow \text{oxidant}$	$K_{\text{SO}_2\text{ODUox}}, r_{\text{ODUox}}$		R6
<i>Diagenetic equations</i>				
	$\frac{d\text{FDet}}{dt} = @ - r_{\text{fast}} * \text{FDet} = 0$			Eq3
	$\frac{d\text{SDet}}{dt} = @ - r_{\text{slow}} * \text{SDet} = 0$			Eq4
	$\frac{d\text{O}_2}{dt} = @ - \text{OxicMin} - 2 * \text{Nitrif} - \text{ODUoxid} + \alpha_{\text{irr0}} * (\text{O}_{2\text{bw}} - \text{O}_2) = 0$			Eq5
	$\frac{d\text{NO}_3}{dt} = @ - 0.8 * \text{Denitrif} + \text{Nitrif} - \text{Anammox} + \alpha_{\text{irr0}} * (\text{NO}_{3\text{bw}} - \text{NO}_3) = 0$			Eq6
	$\frac{d\text{SO}_4}{dt} = @ - 0.5 * \text{AnoxicMin} + \alpha_{\text{irr0}} * (\text{SO}_{4\text{bw}} - \text{SO}_4) = 0$			Eq7
	$\frac{d\text{NH}_3}{dt} = @ + \frac{(\text{DIN}_{\text{min}} - \text{Nitrif} - \text{Anammox})}{(1 + \text{NH}_{3\text{Ads}})} + \alpha_{\text{irr0}} * (\text{NH}_{3\text{bw}} - \text{NH}_3) = 0$			Eq8
	$\frac{d\text{DIC}}{dt} = @ + \text{DIC}_{\text{min}} + \alpha_{\text{irr0}} * (\text{DIC}_{\text{bw}} - \text{DIC}) = 0$			Eq9
	$\frac{d\text{ODU}}{dt} = @ + \text{AnoxicMin} - \text{ODUoxid} - \text{ODUdepo} + \alpha_{\text{irr0}} * (\text{ODU}_{\text{bw}} - \text{ODU}) = 0$			Eq10
	<i>With</i> $\text{ODUdepo} = r_{\text{Depo}} * \text{ODU} + p_{\text{Depo}} * \text{AnoxicMin}$			
@ denotes the transport of dissolved and solid species				

Table 3 The phosphorus equations in the model

<i>P equations</i>	
$P_{org} = C_{orgfast} * \frac{P}{C_{fast}} + C_{orgslow} * \frac{P}{C_{slow}} + P_{refrac}$	Eq11
$P_{orgmin} = R_{fast} * C_{orgfast} * \frac{P}{C_{fast}} + R_{slow} * C_{orgslow} * \frac{P}{C_{slow}} * \vartheta$	Eq12
$Fe_bound P_{formation} = R_{Fe_bound P_{form}} * (DIP - DIP_s) * O_2func$	Eq13
Where $O_2func = \frac{O_2}{(O_2+0.1)}$	
$Fe_bound P_{release} = R_{P_{Fe_bound P_{rel}}} * (Fe_bound P - Fe_bound P_{asympt}) * (1 - O_2func)$	Eq14
$Ca_bound P_{precipitation} = R_{Ca_bound P_{precip}} * (DIP - DIP_{eq})$	Eq15
$\frac{dDIP}{dt} = @ + P_{orgmin} - Fe_bound P_{formation} + Fe_bound P_{release} * \vartheta$	Eq16
$- Ca_bound P_{precipitation} + \alpha_{irro} * (DIP_{bw} - DIP) = 0$	
$\frac{dFe_bound P}{dt} = @ + \frac{Fe_bound P_{formation}}{\vartheta} - Fe_bound P_{release} = 0$	Eq17
$\frac{dCa_bound P}{dt} = @ + \frac{Ca_bound P_{precipitation}}{\vartheta} = 0$	Eq18
@ denotes the transport of dissolved and solid species	

Table 4 Fixed parameters in the model are the ammonium adsorption coefficient (NH_{3Ads}), maximum oxidation of ODU (r_{ODUox}) and nitrification rate constants (r_{nit}), half-saturation constants for O_2 limitation for oxic mineralization ($K_{SO2oxic}$) and nitrification ($K_{SO2nitri}$), nitrate limitation for nitrate reduction ($K_{NO3denit}$), O_2 limitation for the oxidation of reduced substances ($K_{SO2ODUox}$), O_2 inhibition for nitrate reduction ($K_{NO2denit}$) and anoxic mineralization ($K_{NO2anox}$), nitrate inhibition for anoxic mineralization ($K_{NO3anox}$), sulfate limitation for sulfate reduction (K_{SSO4}), C/N_{fast} , C/N_{slow} , C/P_{fast} and C/P_{slow}

<i>Parameter</i>	<i>Unit</i>	<i>Value</i>	<i>Source</i>
NH_{3Ads}	-	1.3	L
r_{ODUox}	d ⁻¹	20	L
R_{nit}	d ⁻¹	10	C
$K_{SO2oxic}$	μmol O_2 l ⁻¹	1	L
$K_{SO2nitri}$	μmol O_2 l ⁻¹	10	L
$K_{NO3denit}$	μmol NO_3^- l ⁻¹	10	L
$K_{SO2ODUox}$	μmol O_2 l ⁻¹	1	L
$K_{NO2denit}$	μmol O_2 l ⁻¹	1	L
$K_{NO2anox}$	μmol O_2 l ⁻¹	1	L
$K_{NO3anox}$	μmol NO_3^- l ⁻¹	10	L
K_{SSO4}	mmol SO_4^{2-} l ⁻¹	800	L
C/N_{fast}	mol C mol N ⁻¹	7.1	L
C/N_{slow}	mol C mol N ⁻¹	10	L
C/P_{fast}	mol C mol P ⁻¹	114	C
C/P_{slow}	mol C mol P ⁻¹	160	C
DIP_s	μmol l ⁻¹	1	L

21 **Table 5 Parameters measured in every sampled station: bottom water dissolved**
22 **species concentrations C_{bw} , porosity parameters ($\emptyset_0$, $\emptyset_\infty$ and $coeff\emptyset$), bottom water**
23 **temperature, salinity and sedimentation rate (ω)**

<i>Parameter</i>	<i>Unit</i>	<i>Z</i>	<i>A</i>	<i>AZ</i>	<i>AK</i>	<i>B</i>	<i>K</i>	<i>L</i>	<i>C</i>	<i>E</i>
<i>Temperature</i>	°C	15.7	15.6	20	15.6	15.8	15.7	15.1	14.7	14.8
<i>Salinity</i>	g kg ⁻¹	38	37.8	37.8	37.9	37.9	38.2	38.1	38.2	38.3
<i>O_{2bw}</i>	μmol l ⁻¹	215	197	206	185	206	206	235	189	224
<i>NO_{3bw}</i>	μmol l ⁻¹	0.8	0	0.2	0	0	0	0.7	3	2.7
<i>NH_{3bw}</i>	μmol l ⁻¹	0	0	0.8	0.24	0.42	0.5	1.17	1.45	0
<i>DIC_{bw}</i>	μmol l ⁻¹	2372	2360	2330	2350	2359	2363	2384	2396	2377
<i>DIP_{bw}</i>	μmol l ⁻¹	0.78	4.28	2.03	1.63	1.95	2.31	0.57	0.31	1.56
<i>SO_{4bw}</i>	mmol l ⁻¹	29.9	30.2	30.5	30.6	30.7	30.3	30.6	30.5	30.6
<i>ODU_{bw}</i>	μmol l ⁻¹	0	0	0.77	0	0.31	0.18	0	0	0
<i>ω</i>	cm y ⁻¹	10	10	10	1	1	1	1	0.1	0.1
<i>∅₀</i>	-	0.81	0.83	0.8	0.84	0.83	0.76	0.84	0.86	0.85
<i>∅_∞</i>	-	0.65	0.65	0.65	0.62	0.6	0.58	0.63	0.64	0.6
<i>coeff∅</i>	cm	1	2	3	2	2	3	2	3	6

24

Table 6 Parameters estimated by the model: organic carbon flux (OC_{flux}), proportion of the labile organic carbon fraction (P_{fast}), labile and semi-labile carbon mineralization rate constants (R_{fast} and R_{slow} , respectively), bioturbation parameters (surface bio bioturbation constant D_{b0} , thickness of bioturbated layer x_b), irrigation parameters (surface irrigation concentration α_{irr0} , irrigation depth x_{irr}), Fe-bound P fluxes to sediment-water interface, rate constants for Fe-bound P formation ($R_{Fe-bound Pform}$) and release ($R_{Fe-bound Prel}$), and the Ca-bound P precipitation rate constants ($R_{Ca-bound Pprecip}$), ODU deposition rate (R_{depo}) and part of ODU deposited as solid (P_{depo}).

<i>Parameter</i>	<i>Unit</i>	<i>Z</i>	<i>A</i>	<i>AZ</i>	<i>AK</i>	<i>B</i>	<i>K</i>	<i>L</i>	<i>C</i>	<i>E</i>
<i>OC_{flux}</i>	mmol m ⁻² d ⁻¹	160	150	130	50	44	43	24	14	10
<i>R_{fast}</i>	d ⁻¹	0.06	0.1	0.15	0.15	0.2	0.2	0.2	0.2	0.2
<i>R_{slow}</i>	d ⁻¹	0.0035	0.0031	0.0016	0.0003	0.0002	0.0008	0.0003	0.0002	0.0008
<i>P_{fast}</i>	-	0.5	0.5	0.6	0.75	0.75	0.75	0.75	0.75	0.75
<i>R_{depo}</i>	d ⁻¹	0.05	0.05	0.05	0.05	0.1	0.03	0.08	0.05	0.07
<i>p_{depo}</i>	-	0.99	0.97	0.97	0.9	0.9	0.9	0.92	0.92	0.98
<i>D_{b0}</i>	cm ² d ⁻¹	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.015	0.015
<i>x_b</i>	cm	5	5	5	10	10	10	10	10	10
<i>α_{irr0}</i>	d ⁻¹	0.09	0.23	0.1	0.1	0.05	0.1	0.1	0.05	0.05
<i>x_{irr}</i>	cm	8	7	14.5	9	13	8	11	7	6
<i>C_{ref}</i>	%	1	1.35	1	1	1.25	1.4	1.2	0.9	1
<i>P_{ref}</i>	μmol g ⁻¹	4	8	5	4	2.5	5.5	5.5	5	8
<i>R_{Fe-bound Pform}</i>	d ⁻¹	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.1	0.1
<i>R_{Fe-bound Prel}</i>	d ⁻¹	0.0008	0.0008	0.0008	0.0008	0.0005	0.0005	0.0008	0.0008	0.0008
<i>R_{Ca-bound Pprecip}</i>	d ⁻¹	0.006	0.0055	0.003	0.003	0.003	0.002	0.006	0.002	0.0035
<i>Fe-bound P_{Flux}</i>	μmol m ⁻² d ⁻¹	73	57	73	8	8.2	7.5	7.5	1	1
<i>Fe-bound P_{Asymp}</i>	μmol g ⁻¹	3	2	3	3	2.5	2.5	3	4	3.5
<i>Ca-bound P_{her}</i>	μmol g ⁻¹	4	4	3.6	4.7	4.1	4.15	5.3	4.7	4.2

Table 7 Phosphorus budget for the Rhone River and prodelta. Surface area are from Lansard et al. (2009) and Cathalot et al. (2008). River DIP discharge are from Ludwig et al., (2009) and Malago et al. (2019). The benthic DIP flux and Ca-bound P burial are estimated from the model calculations of this study averaged for each homogeneous zone.

Zones	Surface area (km ²)	Benthic DIP flux (μmol m ⁻² d ⁻¹)	Ca-bound P burial (μmol m ⁻² d ⁻¹)	DIP Recycl. (kmol y ⁻¹)	Ca-bound P Burial (kmol y ⁻¹)	River DIP discharge (kmol y ⁻¹)
Proximal (A-AZ-Z)	15	864	272	4728	1488	
Prodelta (B-L-K-AK)	50	292	39	5336	703	
Contin. Shelf (C-E)	200	90	7	6606	508	
Total				16671	2699	88968^a/47258^b
Fraction of River DIP discharge (%)				19 ^a - 35 ^b %	3 ^a - 6 ^b %	

^a DIP River discharge from Ludwig et al., 2009; fractions calculated with respect to the discharge by Ludwig et al., 2009.

^b DIP River discharge from Malago et al., 2019; fractions calculated with respect to the discharge by Malago et al., 2019.