

---

## Length–weight relationships of 5 fish species from the Sine Saloum estuary, Senegal, West Africa

Sadio Oumar <sup>1,\*</sup>, Simier Monique <sup>2</sup>, Le Loch Francois <sup>3</sup>, Tito De Morais Luis <sup>3</sup>

<sup>1</sup> IRD, Univ Brest CNRS Ifremer LEMAR Dakar, Senegal

<sup>2</sup> MARBE C, Univ Montpellier, CNRS, Ifremer, IRD Sète, France

<sup>3</sup> Univ Brest, CNRS, IRD, Ifremer, UMR 6539 LEMAR, Plouzane, France

\* Corresponding author : Oumar Sadio, email address : [oumar.sadio@ird.fr](mailto:oumar.sadio@ird.fr)

---

### Abstract :

Length-weight relationships ( $W = aL^b$ ) were estimated for 5 fish species, the pearl stingray *Fontitrygon margaritella* (Compagno & Roberts, 1984), the West African spadefish *Chaetodipterus lippei* (Steindachner, 1895), the East Atlantic African spadefish *Ephippus goreensis* (Cuvier, 1831), the law croaker *Pseudolithus senegallus* (Cuvier, 1830) and the Guinean sole *Dagetichthys cadenati* (Chabanaud, 1948). Fishes were collected during several scientific fishing surveys using purse seine (length 250 m, height 20 m, 14 mm mesh size) and beach seine (180 m length, 9 m height and 25 mm mesh net) conducted from April 1990 to October 2012 in the Sine Saloum estuary, Senegal. They were weighed to the nearest g and measured to the nearest mm (disc width for *Fontitrygon margaritella* and total length for the 4 other species). The values of parameter b in the LWR equations varied from 3.048 (*Chaetodipterus lippei*) to 3.274 (*Dagetichthys cadenati*) and the coefficient of determination ( $r^2$ ) varied between 0.969 (*Fontitrygon margaritella*) and 0.998 (*Chaetodipterus lippei*). In addition, new maximum lengths are proposed for *Fontitrygon margaritella*, *Chaetodipterus lippei* and *Dagetichthys cadenati*.

**Keywords** : estuary, fish species, length–weight relationships, Senegal, sine saloum

## 46 INTRODUCTION

47 Length–weight relationships (LWR) contribute together with the monitoring of several other  
48 essential parameters (e.g. sex ratio, age of first maturity, fecundity, recruitment, natural and  
49 fishing mortality, growth patterns, estimation of biomass, population dynamics) to the  
50 required stock assessment for proper population management (Le Cren, 1951; Hampton,  
51 2000; Fromentin & Fonteneau, 2001; Froese, 2006). They help to describe a specific  
52 population, to compare fish species populations or fish life histories between seasons, years,  
53 zones and regions, provided all investigators use the same standardized sampling  
54 methodology (De La Cruz Agüero et al., 2011; Gonçalves et al., 1997; Moutopoulos &  
55 Stergiou, 2002). In this study, we assessed length-weight relationships for 5 fish species from  
56 the Sine Saloum estuary. This work complements those of Ecoutin and Albaret (2003) and  
57 Ndiaye et al. (2015) on LWR of fish species in the Saloum estuary.

## 58 MATERIALS AND METHODS

59 The Sine Saloum estuary is located 100 km south of Dakar, in Senegal, between 13°55' and  
60 14°10' North and 16°03' and 16°50' West. The Sine Saloum is an inverse estuary  
61 characterized by an increase in salinity from downstream to upstream in all seasons.  
62 Mangrove forests cover almost the entire southern portion of the system and progressively  
63 diminish in the North (Diouf, 1996; Simier et al., 2004). Data were collected during two  
64 research programs covering the whole Sine Saloum estuary in 1990-1993 and then in 2002-  
65 2003 (Diouf, 1996; Ecoutin et al., 2010) and the biological monitoring of the Bamboung  
66 MPA and its surroundings from 2003 to 2012 (Ecoutin et al., 2014; Sadio et al., 2015). All  
67 the sampling was carried out over the three main hydro-climatic seasons of the region (Simier  
68 et al., 2004). 166 fishing hauls were made using a purse seine (length 250 m, height 20 m, 14  
69 mm mesh size; see Simier et al., 2004) and 11 using a beach seine (180 m length, 9 m height  
70 and 25 mm mesh net; see Diouf, 1996). Fish were measured to the nearest mm (disc width for  
71 *Fontitrygon margaritella* and total length for the 4 other species) and weighed to the nearest  
72 g. All data are available in PPEAO database (Simier, Ecoutin, & Tito de Morais, 2019).  
73 Linear regressions of  $\log(W)$  vs.  $\log(L)$ , where  $W$  is total weight in g and  $L$  is total length  
74 or disc width in cm, were calculated to obtain the length-weight relationship of the form  $W =$   
75  $aL^b$ , where  $a$  and  $b$  are regression parameters (Froese, 2006).

## 76 RESULTS

77 Five fish species belonging to 4 families were considered in this study. Sample size, length  
78 and weight ranges, parameters of the LWR equations with 95% confidence limits and values  
79 of the determination coefficient ( $r^2$ ) were computed for the 5 fish species (Table 1). All  
80 regressions of LWRs were statistically significant ( $p < 0.001$ ).

## 81 DISCUSSION

82 All the fishes in this study were collected over a long period of time, during several scientific  
83 surveys of the fish assemblages in the Sine Saloum estuary. The sampling protocols were  
84 designed to cover the entire hydroclimatic cycle and to explore various habitats, thus  
85 providing a spatial and temporal overview of estuarine fish assemblages. The five species

86 presented here are not however among the most frequent in these assemblages. Except  
87 *Fontitrygon margaritella* and *Pseudotolithus senegallus* which are an “estuarine species from  
88 marine origin” and a “marine-estuarine species” respectively, the 3 others are “marine  
89 species, occasionally or accidentally found in estuaries” (Albaret, 1999). Thus, for *Ephippus*  
90 *goreensis* the sample size is limited and the maximum length is only 20.8 cm (for a known  
91 maximum size of 30 cm) and for *Pseudotolithus senegallus* the maximum length is 101 cm  
92 (for a known maximum length of 230 cm). Consequently, the parameters estimated here,  
93 especially for these two species, should be considered as tentative. Despite these limitations,  
94 the coefficients of determination ( $r^2$ ) were very high for the five species, indicating a good  
95 predictive power. All species had  $b$  value comprised between 2.5 and 3.5 as expected (Froese,  
96 2006). Regression results suggest that the data from this study and the results presented here  
97 are reliable and can enriched length-weight relationships in FishBase (Froese & Pauly, 2020)  
98 for the Senegal.

## 99 ACKNOWLEDGEMENTS

100 The results presented here are from the multidisciplinary program linked to the projects  
101 Narou Heuleuk (FFEM/AFD/ONG Oceanium/IRD), AMPHORE (ANR Biodiversity) and  
102 Cepia (AFD/CSRP/UICN/IRD). The author are grateful to the LEMAR team in Dakar and  
103 the fishermen and their team leader Mbaye Mbengue, also to the crew of the Diassanga  
104 research vessel and its captain François Sanseo, without whom these results could not have  
105 been obtained.

106 **Data availability statement:** Data are available at <http://ppeao.ird.fr> or upon request to  
107 [ppeao@ird.fr](mailto:ppeao@ird.fr)

## 108 CONFLICT OF INTEREST

109 The authors declare no conflict of interest.

## 110 REFERENCES

- 111 Albaret, J. J. (1999). Les peuplements des estuaires et des lagunes. In: Leveque, C., Paugy, D.  
112 (Eds.), Les poissons des eaux continentales africaines: diversité, écologie, utilisation  
113 pour l’homme. IRD, Paris, 325–350.
- 114 De La Cruz Agüero, J., Garcia Rodriguez, F. J., Cota Gomez, V. M., Chollet Villapando, J.  
115 G., & Vargara Solana, F. J. (2011). Length-weight relation of selected species of the  
116 family Gerreidae (Actinopterygii: Perciformes) from the Mexican coast. *Acta*  
117 *Ichthyologica et Piscatoria*, 41, 67-69. doi: 10.3750/aip2011.41.1.10
- 118 Diouf, P. S. (1996). Les peuplements de poissons des milieux estuariens de l’Afrique de  
119 l’Ouest : l’exemple de l’estuaire hyperhalin du Sine Saloum. Ph.D. Thesis, Université  
120 de Montpellier II, Montpellier. 177p. [http://www .bondy.ird.fr/pleins textes/pleins](http://www.bondy.ird.fr/pleins_textes/pleins_textes_7/TDM_7/010008130.pdf)  
121 [textes\\_7/TDM\\_7/010008130.pdf](http://www.bondy.ird.fr/pleins_textes/pleins_textes_7/TDM_7/010008130.pdf).
- 122 Ecoutin, J. M., & Albaret, J. J. (2003). Relation longueur-poids pour 52 espèces de poissons  
123 des estuaires et lagunes de l’Afrique de l’Ouest. *Cybium*, 27, 3-9.
- 124 Ecoutin, J. M., Simier, M., Albaret, J. J., Laë, R., Raffray, J., Sadio, O., & Tito de Morais, L.  
125 (2014). Ecological field experiment of short-term effects of fishing ban on fish  
126 assemblages in a tropical estuarine MPA. *Ocean & Coastal Management*, 100, 74-85.  
127 doi: 10.1016/j.ocecoaman.2014.08.009.
- 128 Ecoutin, J. M., Simier, M., Albaret, J. J., Laë, R., & Tito de Morais, L. (2010). Changes over

- 129 a decade in fish assemblages exposed to both environmental and fishing constraints in  
130 the Sine Saloum estuary (Senegal). *Estuarine, Coastal and Shelf Science*, 87, 284-292.  
131 doi: 10.1016/j.ecss.2010.01.009
- 132 Froese, R. (2006). Cube law, condition factor and weight–length relationships: history, meta-  
133 analysis and recommendations. *Journal of Applied Ichthyology*, 22, 241-253. doi:  
134 10.1111/j.1439-0426.2006.00805.x
- 135 Froese, R., & Pauly, D. (2020). FishBase. World Wide Web electronic publication. www.  
136 fishbase.org (accessed 09/2020).
- 137 Fromentin, J. M., & Fonteneau, A. (2001). Fishing effects and life history traits: a case study  
138 comparing tropical versus temperate tunas. *Fisheries Research*, 53, 133-150.  
139 doi: 10.1016/s0165-7836(00)00299-x
- 140 Gonçalves, J. M. S., Bentes, L., Lino, P. G. , Ribeiro, J. , Canário, A. V. M., & Erzini, K.  
141 (1997). Weight–length relationships for selected fish species of the small-scale  
142 demersal fisheries of the south and southwest coast of Portugal. *Fisheries Research*, 30,  
143 253–256. doi: 10.1016/s0165-7836(96)00569-3
- 144 Hampton, J. (2000). Natural mortality rates in tropical tunas: size really does matter.  
145 *Canadian Journal of Fisheries and Aquatic Sciences*, 57, 1002-1010.  
146 doi: 10.1139/f99-287
- 147 Le Cren, E. D. (1951). The length-weight relationship and seasonal cycle in gonad weight  
148 and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology*, 20, 201-219.  
149 doi: 10.2307/1540.
- 150 Moutopoulos, D. K., & Stergiou, K. I. (2002). Length-weight and length-length relationships  
151 of fish species from the Aegean Sea (Greece). *Journal of Applied Ichthyology*, 18, 200-  
152 203. doi: 10.1046/j.1439-0426.2002.00281.x
- 153 Ndiaye, W., Sarr, A., Diouf, M., Faye, A., & Mbodji, A. (2015). Length-weight relationships  
154 of some fish species from the Bandiala River in Saloum Delta, Senegal. *International*  
155 *Journal of Advanced Research*, 3, 132-138.
- 156 Sadio, O., Simier, M., Ecoutin, J. M., Raffray, J., Laë, R., & Tito de Morais, L. (2015). Effect  
157 of a marine protected area on tropical estuarine fish assemblages: Comparison  
158 between protected and unprotected sites in Senegal. *Ocean & Coastal Management*,  
159 116, 257-269. doi: 10.1016/j.ocecoaman.2015.08.004.
- 160 Simier, M., Blanc, L., Aliaume, C., Diouf, P. S., & Albaret, J. J. (2004). Spatial and temporal  
161 structure of fish assemblages in an “inverse estuary”, the Sine Saloum system  
162 (Senegal). *Estuarine Coastal and Shelf Science*, 59, 69-86.  
163 doi: 10.1016/j.ecss.2003.08.002
- 164 Simier, M., Ecoutin, J. M., & Tito de Morais, L. (2019). The PPEAO experimental fishing  
165 dataset: Fish from West African estuaries, lagoons and reservoirs. *Biodiversity Data*  
166 *Journal*, 7, e31374. doi: 10.3897/BDJ.7.e31374.

167  
168  
169  
170  
171  
172  
173  
174

175 Table 1. Descriptive statistics and estimated parameters of length-weight relationships ( $W = aL^b$ ) for five species from the Sine Saloum estuary, Senegal,  
 176 captured from 1990 to 2012. n, sample size; Max, maximum; Min, minimum; a and b, parameters of LWR; CL, confidence limits;  $r^2$ , coefficient of  
 177 determination. Length = Total Length except for *Fontitrygon margaritella* (Disc Width). \*, indicates that the LWR includes a new not yet reported maximum  
 178 total length.

Family	Species	n	Length (cm)	Weight (g)	Relationship parameters		
			Min – Max	Min – Max	a (95%CL)	b (95%CL)	$r^2$
Dasyatidae	<i>Fontitrygon margaritella</i> (Compagno & Roberts, 1984)*	139	10.7 – 47.0	54 – 5175	0.025 (0.018 – 0.033)	3.177 (3.081 – 3.274)	0.969
Ephippidae	<i>Chaetodipterus lippei</i> Steindachner, 1895*	73	4.4 – 32.3	3 – 1114	0.029 (0.027 – 0.033)	3.048 (3.013 – 3.082)	0.998
	<i>Ephippus goreensis</i> Cuvier, 1831	15	5.5 – 20.8	5 – 316	0.024 (0.017 – 0.032)	3.141 (3.014 – 3.269)	0.995
Sciaenidae	<i>Pseudotolithus senegallus</i> (Cuvier, 1830)	118	11.1 – 101.0	9 – 8250	0.006 (0.005 – 0.006)	3.079 (3.047 – 3.112)	0.997
Soleidae	<i>Dagetichthys cadenati</i> (Chabanaud, 1948)*	20	8.3 – 38.0	3 – 457	0.003 (0.002 – 0.005)	3.274 (3.054 – 3.494)	0.982

179