

---

## A review of the effects of contamination and temperature in *Solea solea* larvae. Modeling perspectives in the context of climate change

Sardi Adriana E. <sup>1,\*</sup>, Bégout Marie-Laure <sup>2</sup>, Cousin Xavier <sup>3,5</sup>, Labadie Pierre <sup>1</sup>, Loizeau Veronique <sup>4</sup>, Budzinski Hélène <sup>6</sup>

<sup>1</sup> University of Bordeaux, CNRS, EPOC, UMR 5805, F-33400 Talence, France

<sup>2</sup> MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, F-34250 Palavas-Les-Flots, France

<sup>3</sup> Univ. Paris-Saclay, AgroParisTech, INRAE, GABI, F-78350 Jouy-en-Josas, France

<sup>4</sup> Ifremer, LEMAR UMR 6539 - Unit of Functional Physiology of Marine Organisms, CS 10070, F-29280, Plouzané, France

<sup>5</sup> MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, F-34250 Palavas-Les-Flots, France

<sup>6</sup> Ifremer, LEMAR UMR 6539 - Unit of Functional Physiology of Marine Organisms, CS 10070, F-29280, Plouzané, France

\* Corresponding author : Adriana E. Sardi, email address : [adriana.sardi@u-bordeaux.fr](mailto:adriana.sardi@u-bordeaux.fr)

---

### Abstract :

The flatfish species *Solea solea* has been the subject of research for supporting their management in fisheries, restocking natural populations, domestication in aquaculture, and ecotoxicology research. Soles undergo a metamorphosis with drastic morphological and physiological changes and settlement, processes that make them potentially more vulnerable to pollutants than other fish. Up to date, efforts made to develop its aquaculture production in Europe have been limited. In the context of climate change (CC), this review aims i) to gather research conducted in *S. solea* larvae that summarizes the effects of increased temperature and contaminant exposures during larval development; and ii) to provide a summarized and synergistic view about its larval development. The review consists of 4 sections. Section 1 justifies the selection of this species from ecological and economic perspectives. Section 2 focuses on larvae ontogenesis, metamorphosis, rearing challenges, and further aquaculture production. Section 3 reviews studies dealing with the effects of temperature change (due to CC) and pollutants on larval development. Finally, section 4 provides a “how to go forward on ecotoxicological research” guideline, in which we highlight the methods that we found promising as tools to study the combined effects of CC and pollution. The section includes a multidisciplinary framework that proposes how existing data coming from different scientific domains can be synthesized to be useful for risk assessors and ecotoxicologists. To benefit from such a framework, it is necessary to reach consensus and nurture team effort from players that operate in different research disciplines.

---

## Highlights

► Flatfish are relevant within ecotoxicity and environmental risk assessment. ► We summarize the effects of temperature and chemical exposure on *S. solea* larvae. ► Effects on larval development are of interest for multiple disciplines. ► We present a multidisciplinary framework to improve collaboration between disciplines. ► Physiology based models are promising tools for predicting anthropogenic effects on this species.

**Keywords** : Flatfish, Climate change, Contamination, Larval stage, Multidisciplinary

37 **Introduction: Working with flatfishes in ecotoxicology, why?**

38

39 Fish are a prominent group within ecotoxicity evaluation guidelines and subsequently within environmental  
40 risk assessment for the aquatic environment. Up to date, there exist nine guidelines from the Organization for  
41 Economic Co-operation and Development (OECD) that acknowledge the importance of fish toxicology, which  
42 includes: the acute toxicity test (OECD 203, 1992), the short term toxicity test on embryo and sac-fry stages

43 (OECD 212, 1998), the fish juvenile growth test (OECD 215, 2000), the fish early life-stage toxicity test  
44 (OECD 210, 2004), the 21-day fish assay: a short-term screening for oestrogenic and androgenic activity, and  
45 vitellogenin and secondary sex characteristics (OECD 230, 2009), the fish sexual development test (OECD  
46 234, 2011), the fish short term reproduction assay (OECD 229, 2012), bioaccumulation in fish: aqueous and  
47 dietary exposure (OECD 305, 2012), and the fish embryo acute toxicity (FET) test (OECD 236, 2013).

48

49 Current integrated toxicity testing strategies consider that fish are an indispensable component, and in general  
50 terms, this is mainly because:

- 51 - fish are an essential food resource for humans;
- 52 - the aquatic environment is the final reservoir for many chemicals;
- 53 - fish are critical in aquatic food webs by top-down and bottom-up regulation of nutrient and energy flow;
- 54 - fish have been and continue to be used as sentinels of water quality used for human consumption;
- 55 - fish kills following accidents are visible to the public and encourage them on the need to protect the aquatic  
56 environment from pollutants; and
- 57 - fishing has recreational value in many cultures (Lammer *et al.*, 2009).

58

59 Among fish, flatfishes have been the subject of research supporting their management in fisheries, restocking  
60 natural populations, and domestication in aquaculture. Also, they present unique ontogeny, metamorphosing  
61 from bilateral to asymmetric, which makes them a compelling target for evolution and developmental studies  
62 (Geffen *et al.*, 2015). Furthermore, flatfishes present similar gene regulation and consequently (except for few  
63 exceptions) similar physiological responses to pollutants (Lammer *et al.*, 2009).

64

65 Specific characteristics, like metamorphosis and settlement, make flatfishes respond in a potentially different  
66 manner when exposed to pollutants. To begin with, they undergo metamorphosis, with morphological and  
67 physiological changes associated with eye migration, a 90° rotation in posture, and asymmetrical pigmentation  
68 (Geffen *et al.*, 2007). Metamorphic larvae undergo drastic remodeling of tissues and physiological changes.  
69 The transition from larval to juvenile stage involves developing most organs and tissues, the maturation of  
70 different physiological functions, and the establishment of the immune system (Ferraresso *et al.*, 2013).  
71 Flatfish larvae also undergo settlement, a behavioral and ecological change associated with a transition from a  
72 pelagic to a benthic existence. As benthic animals feed on invertebrates, flatfishes are affected by sediment-  
73 associated pollutants; so, their life history and habitat also make flatfishes valuable models for ecotoxicology  
74 research (Geffen *et al.*, 2015). The benthic lifestyle potentially exposes flatfishes to a wide range of  
75 contaminants, including -but not limited- to endocrine disruptors, heavy metals, polycyclic aromatic  
76 hydrocarbons (PAH), and dioxins (Munsch *et al.*, 2010, 2011; Wessel *et al.*, 2010; Gilliers *et al.*, 2012; Geffen  
77 *et al.*, 2015). This fact makes them useful sentinels for monitoring the biological effects of contaminants in  
78 inshore and estuarine waters.

79

80 Besides physical changes, survival and growth after recruitment are often assumed to depend on the behavioral  
81 and ecological changes associated with metamorphosis (Geffen *et al.*, 2007). Optimal cues, such as sensory  
82 perception and interpretation, guarantee the survival of early juveniles, yet it is known that swimming activity,  
83 exploration ability, and place preferences can be affected by exposure to PCBs and PBDE (Péan *et al.*, 2012).  
84 Moreover, flatfish species are widely distributed, which facilitates conducting spatio-temporal comparisons.

85  
86 The characteristics mentioned above (metamorphosis, benthic lifestyle, and behavioral adaptations) make  
87 flatfishes unique and potentially more vulnerable to anthropogenic activities and estuarine contamination. It is  
88 worth noting that the aquatic environment is the ultimate sink for many man-produced products, which are  
89 continuously being released at estuaries, either due to direct discharges or to hydrologic and atmospheric  
90 processes. The chemicals that end up in these habitats range from many thousands of organic trace pollutants  
91 (polychlorinated biphenyls (PCBs), organochlorine pesticides (OCPs), polycyclic aromatic hydrocarbons  
92 (PAHs), polychlorinated dibenzofurans (PCDFs), and dibenzo- p-dioxins (PCDDs)) to compounds of  
93 emerging interest, which include perfluoroalkyl substances (PFAS), nanoparticles, personal care products,  
94 detergents, estrogen-like products, and flame retardants. All these substances are potentially interacting and  
95 impacting the health of flatfish and represent a risk for the health of benthonic fish populations (van der Oost  
96 *et al.*, 2003).

97  
98 Finally, Geffen *et al.* (2015) argued that one of the most rapidly developing areas that would benefit from the  
99 synergy of flatfish research in aquacultured and wild populations is the ecotoxicology domain. Unfortunately,  
100 research on flatfishes is distributed across journals in several (sub)disciplines, and because the reference lists  
101 of individual papers rarely extend across these disciplines, in order to gain a full benefit of knowledge  
102 exchange, it is necessary to read widely.

103

## 104 **Objectives**

105

106 This review aims to gather works primarily focused on ecotoxicological research conducted in the flatfish  
107 species *Solea solea* along with their larval development and early juvenile phase. We draw attention to this  
108 life period because it is critical for fisheries and aquaculture development and for understanding anthropogenic  
109 impacts on *S. solea* populations and ecosystem degradation. Additionally, a secondary objective is to provide  
110 a summarized and synergistic view on research conducted in *S. solea* larvae that serves different disciplines  
111 such as aquaculture, fisheries, marine ecology, and ecotoxicology.

112

113 We divided the review into four sections. The first section justifies the selection of this species and reviews *S.*  
114 *solea* ecologic and economic relevance.

115

116 Section two focuses on the main particularities of this species. Soles, like all flatfishes, undergo an extreme  
117 metamorphosis which translates into a susceptible larval stage. The section summarizes *S. solea* regular larvae  
118 development by citing the most recent literature describing the optimal conditions for *S. solea* laboratory

119 rearing and growth. The main questions addressed in section 2 include: What are the main bottlenecks on the  
120 aquaculture production of *S. solea*? Furthermore, what have been the scientific advances that have been made  
121 to overcome these problems?

122  
123 In the third section, we will focus on reviewing works studying the effects of climate change (CC, we only  
124 focus on temperature change) and contamination on larval development. This section comprises the main focus  
125 of the review and will include works from the molecular level until the population level, exploring and  
126 comparing these different “tools” employed to understand, estimate, and predict the effects of human activities  
127 on this species. The questions we intend to answer in section 3 include: What are the individual and combined  
128 effects of CC and environmental pollution on larval development of *S. solea*? Furthermore, can we predict or  
129 anticipate the effects of combined stresses?

130  
131 Finally, in the last section, we will review the methods and efforts made to integrate different types of data to  
132 1) have a holistic view on the subject and 2) to be able to extrapolate (if possible) to different levels of  
133 biological organization or other species. This section is meant as a “how to go forward on ecotoxicological  
134 research” guideline, in which we primarily highlight the methods that we found promising as tools to study  
135 the combined effects of CC and pollution. The main questions addressed in section 4 include: What methods  
136 allow us to understand the combined effects of CC and contamination? How can we reduce animal testing for  
137 new chemicals and improve risk assessment? And finally, how much can we rely on numerical models?

138

## 139 **Section 1 - Why *Solea solea*?**

140

### 141 **1.1 Common sole life-cycle main events**

142 The flatfish species, *Solea solea*, also known as the common or Dover sole, and from now on simply denoted  
143 as sole, is an important commercial species found in shallow waters (between 10 to 60 m depth but also down  
144 to 200 m) of the eastern Atlantic Ocean. Its distribution range extends from the north-western coast of Africa  
145 in the Mediterranean Sea to the southern coast of Norway (Zambonino-Infante *et al.*, 2013).

146

147 The common sole is a batch-spawner, and increases in temperature trigger spawning. Depending on the  
148 geographical location, spawning events might occur from late winter to spring. In the Mediterranean Sea,  
149 spawning occurs between January and April, with two peaks in February, while in the Bay of Biscay, spawning  
150 occurs between December and May (FAO, 2019), depending on the latitude. In the North Sea, spawning  
151 happens between April and June, and different sole stocks present differences in the average date of peak  
152 spawning, with spawns occurring from late March in the Bristol Channel to mid-May in the Irish Sea and in  
153 late May in the eastern-central North Sea (Fincham *et al.*, 2013). The optimal temperature for spawning ranges  
154 from 7 to 12 °C (Lacroix *et al.*, 2013).

155

156 Spawning behavior in *S. solea* involves three stages, courtship, spawning rise, and a post-spawning return to  
157 the seafloor. During courtship, the male follows a female on the bottom and then swims below the female. If

158 courtship is successful, the pair swims vertically towards the water surface with closely synchronized body  
159 movements, and gametes are released at the apex of the rise, followed by a return to the seafloor (Baynes *et*  
160 *al.*, 1994).

161  
162 Fertilized eggs and larvae are both pelagic (Fig. 1), which allows for wide dispersion and exposes them to  
163 environmental variability, notably temperature and food abundance. Fertilized eggs hatch after few days (5  
164 days at 16 °C, Imsland *et al.*, 2003), and the larval stage -from hatching until complete metamorphosis- can  
165 last between 26 and 15 days, at temperatures of 16 °C and 19 °C, respectively (De Swaef *et al.*, 2017; Richard *et*  
166 *al.*, 1991).

167  
168 The larval stage comprises a metamorphic period, where significant ontogenic, morphological, physiological,  
169 and behavioral changes occur (Geffen *et al.*, 2007). The length of metamorphosis depends on temperature and  
170 food availability, and by the end of it, juvenile soles are ready to change from pelagic to benthic lifestyle to  
171 colonize shallow coastal waters in estuaries and bays (Zambonino-Infante *et al.*, 2013).

172  
173 Newly metamorphosed juveniles settle in coastal areas for around two years until they reach maturity (Post *et*  
174 *al.*, 2017). These areas serve as nursery grounds, where abundant food (epi and endobenthic prey) favors  
175 growth. Further, the pelagic larval and the juvenile stage on nursery grounds are critical for common sole  
176 recruitment, *i.e.*, the number of fish surviving to enter a fishery and contribute to the next generation  
177 (Zambonino-Infante *et al.*, 2013). After settlement on estuarine nursery areas, *S. solea* individuals undergo  
178 further ontogenic habitat shift (in some populations, juvenile soles progressively migrate tens of kilometers  
179 from shallow estuarine waters to deeper coastal water).

180  
181 At the adult stage, they migrate offshore (few or up to hundreds of kilometers depending on location) in  
182 particular for reaching spawning grounds, and they eventually come back closer to the shore out of  
183 reproduction season to reach feeding grounds (Vasconcelos *et al.*, 2014). Regardless of this migration, Diopere  
184 and collaborators (2017) argue that the wide geographical area at which this species is found and the fact that  
185 they do not conduct long migrations makes it reasonable to assume that sole in European waters belongs to  
186 more than one population. However, the precise habitat range of this species remains unclear (Post *et al.*,  
187 2017). Moreover, and because spawning grounds are distant from the nursery areas, the planktonic larvae have  
188 a drifting period which ideally should end at an area that is suitable as a nursery for the juvenile stage. The  
189 question is whether they do anything to maximize their probability of arriving there or whether it is a random  
190 process (Duffy-Anderson *et al.*, 2015). Factors such as release time, location, stage duration, climate, and  
191 hydrodynamic circulation play a crucial role in larval mortality rates and determine the transport of eggs, larvae  
192 and the population renewal (Vinagre, 2007; Duffy-Anderson *et al.*, 2015).

193  
194 Such a complex life cycle (Fig. 1) with progressive habitat shifts along *S. solea* ontogenesis makes it  
195 challenging to determine how population dynamics are affected by habitat quality. In this regard, Vasconcelos

196 and collaborators (2014) showed in their review how little is known about habitat-specific demographic rates  
197 on species of commercial interest, including the common sole. Knowledge about the relationship between  
198 habitat characteristics and population dynamics is particularly valuable from a conservation and management  
199 perspective since the vulnerability of coastal habitats to anthropogenic stressors is increasing (Rochette *et al.*,  
200 2013). Specific recommendations include emphasizing the spatiotemporal scales in studies addressing the  
201 reproduction, mortality, immigration, and emigration dynamics of *S. solea*. By this means, it will be possible  
202 to define settlement habitats, the degree of connectivity between primary and secondary settlement habitats,  
203 and the level of dependence of juvenile and adult populations on these essential coastal habitats (Vasconcelos  
204 *et al.*, 2014).

205

## 206 **1.2 Ecological importance and habitat selection**

207 *Solea solea* is an r strategist species, characterized by a strong fecundity -with hundreds of thousands of eggs  
208 spawned (per kg of female)- and low survival rates. The environmental fluctuations strongly influence survival  
209 rates, egg and larval life history along the larval dispersion process, and the transition from a pelagic to a  
210 benthic lifestyle (Savina *et al.*, 2016). At first, eggs derive towards nursery areas thanks to the action of winds,  
211 tides, and currents (Savina *et al.*, 2010). After hatching, larvae start showing active vertical migration in the  
212 water column triggered when feeding starts (Lagardère *et al.*, 1999). Larvae use vertical migration to enter,  
213 leave, or remain in estuaries by either selecting the seaward outflow at the surface or the inflow near the bottom  
214 (Duffy-Anderson *et al.*, 2015). Therefore, the successful arrival in nursery areas is dependent on hydroclimatic  
215 conditions (Lagardère *et al.*, 1999).

216

217 Recently metamorphosed individuals settle in nursery areas and switch to a benthic lifestyle starting their  
218 critical juvenile stage when their movements are limited, which will last for up to 2 years (Eichinger *et al.*,  
219 2010). According to the “nursery size hypothesis” (Van der Veer *et al.*, 2000), juvenile survival depends on  
220 the surface area of nurseries; as each nursery can only sustain a population in proportion to the size available  
221 (Duffy-Anderson *et al.*, 2015), and it is during this period that soles become top predators (Amara *et al.*, 2001).  
222 Indeed, early juvenile diet is mainly based on copepods, progressively changing to include short-lived, highly  
223 productive benthic organisms, such as polychaete worms, mollusks, and small crustaceans (Amara *et al.*, 2001;  
224 Rijnsdorp *et al.*, 2004).

225

226 Juvenile fish distribution is influenced, among other factors, by sediment type, food availability, and movement  
227 ability, and the trade-off outcome of these factors determines their habitat selection. For *S. solea*, estuarine and  
228 shallow coastal systems act as nurseries where the trade-off between growth and survival is optimized due to  
229 abundant food resources, few predators, and favorable environmental conditions (Freitas *et al.*, 2012). This  
230 fish-habitat interaction is of utmost importance at early-life stages when successful recruitment and  
231 development of fish from larval to juvenile stages depends on the quality and the characteristics of nursery  
232 areas (Post *et al.*, 2017).

233



234 For flatfishes, the type of sediment is also a decisive factor, as it can inhibit burial behavior or affect prey  
235 availability (Post *et al.*, 2017). That is why habitat degradation, such as beach sand nourishment, directly  
236 impacts recruitment success and species fitness. Coarsening the sediment in the coastal zone would increase  
237 the energy requirement for juvenile sole through prolonged digging behavior, consequently impacting  
238 individual growth and increasing their vulnerability to predation due to their smaller size (Post *et al.*, 2017).  
239 Post and collaborators (2017) measured the sediment preference of soles by determining their relative  
240 distribution over the sediment classes. Preference -defined as selecting a specific sediment type through their  
241 presence at the end of the experimental period (4 days)- was tested in a circular preference chamber. Their  
242 results demonstrated that juvenile soles prefer fine grains, and they suggest that such marked preference could  
243 be a consequence of higher food availability and lower energy to bury (Howell and Canario, 1987; Post *et al.*,  
244 2017).

245

246 The juvenile stage, where soles have a benthic lifestyle and inhabit coastal nursery areas, is the period at which  
247 individuals are exposed to a broader range of contaminants, which includes not only hydrophilic compounds  
248 dissolved in the water column but also hydrophobic contaminants that can be either directly absorbed from the  
249 sediment or ingested from contaminated food (Rochette, 2011).

250

251 Given its characteristics and ecological importance, *S. solea* has been included within the EU Marine Strategy  
252 Framework Directive (MSFD) as a model and indicator species for evaluating the good environmental status  
253 in the marine environment in 2020.

254

255 Finally, readers seeking for more detailed information on niche ecology and habitat selection of *S. solea* are  
256 encouraged to revise the works from Vinagre *et al.* (2006) that evaluates the species distribution and determines  
257 the habitat quality using habitat suitability index models or the study by Vinagre and collaborators (2005),  
258 which evaluates the niche overlap between European flounder and *S. solea*. Furthermore, there is an important  
259 body of research that focuses on comparing *Solea senegalensis* and *S. solea* and their differential use of  
260 estuaries (Cabral and Costa 1999), species distribution (Vinagre *et al.*, 2009), prey consumption (Cabral 2000;  
261 Vinagre and Cabral 2008) and habitat-specific growth rates (Vinagre *et al.*, 2008) from populations inhabiting  
262 the Tagus estuary, Portugal. The list of publications on related topics goes on, including Martinho *et al.* (2008),  
263 where the authors compared the timing of estuarine colonization, the population structure, growth, and diet  
264 composition of flatfish juveniles from the species *Dicentrarchus labrax*, *Platichthys flesus*, *S. solea*.

265

### 266 **1.3 Economic importance of soles**

267 The sole species, *Solea solea*, and *Solea senegalensis*, are characterized by high flesh quality and high market  
268 value making them very promising candidates for European aquaculture (Howel *et al.*, 1997; Bonaldo *et al.*,  
269 2011). Also, fisheries landings for *Solea spp.* have shown a reduction of over 43%, and the average size of  
270 wild sole has also declined (Morais *et al.*, 2014). This reduction in the offer has promoted average prices of  
271 different sole products (fresh, frozen, and farmed) to increase and consistently attain high prices in European

272 markets (Morais *et al.*, 2014). For example, in 2013, farmed products attained 12.25 € kg<sup>-1</sup> in MercaMadrid  
273 (Morais *et al.*, 2014), and in May 2019, *S. solea* reached an average price of 23.8 € kg<sup>-1</sup> in France  
274 (FranceAgriMer, <https://rnm.franceagrimer.fr/prix?SOLE&12MOIS>). Their high commercial value, the  
275 growing market demand for adults and juveniles, and the increasing fishing pressure -not restricted to the coast  
276 but also present in the estuary- make sole fisheries an attractive business from both the socio and economic  
277 perspectives. These reasons have reinforced the interest and promise of developing commercial aquaculture  
278 with these species. In fact, *S. solea* and *S. senegalensis* have been repeatedly identified as the most promising  
279 species for aquaculture diversification in Europe for at least four decades now (Morais *et al.*, 2014).

280

281 As a result, during the last 30 years, sole species have attracted considerable research attention, and significant  
282 investments have been made to develop rearing methodologies (Howell 1997; Dinis *et al.*, 1999). Nevertheless,  
283 the production of *Solea spp.* has failed to reach successful commercial development until recently, and this  
284 success is limited to *S. senegalensis*. The three major bottlenecks for *Solea spp.* production are 1) high larval  
285 mortality rates related to nutrition and growth dispersion, 2) sub-optimal larval weaning strategies that result  
286 in weaning onto formulated diets and lead to low growth and high juvenile mortalities; and 3) inadequate  
287 control measures for common diseases. Additionally, the techniques for intensive rearing of the sole are yet to  
288 be optimized, which limits the final volume of fish production (FAO, 2019; Morais *et al.*, 2014). Other  
289 problems include a decreased growth rate at high stocking densities, pigmentation abnormalities,  
290 malformations related to eye migration (Dinis *et al.*, 1999). Finally, the reproduction of cultured specimens is  
291 complicated in F1 and later generations (especially in *S. senegalensis*, Ibarra-Zatarain *et al.*, 2020). In the next  
292 section, we will summarize some of the research advances made to solve the above-mentioned problems.

293

## 294 **Section 2 - Importance of studying *Solea solea* larvae: development, the cost, and challenge of** 295 **metamorphosis**

296

297 External biotic and abiotic factors affecting larval development have consequences at later life stages (Howel  
298 and Baynes, 2004). Fish larval stages represent a transitional period at which both differentiation and growth  
299 lead to substantial changes in structure, physiology, size, and morphology. Compared to any other life stages  
300 of a marine fish species, individuals at their larval stage will have the highest potential for growth, weight-  
301 specific metabolic rates, natural mortality rates, and the highest sensitivity to environmental stressors  
302 (Hutchinson *et al.*, 1998; Belanger *et al.*, 2010).

303

304 These factors highlight the importance of studying fish larvae. Early stages are usually characterized by strict  
305 environmental requirements, where factors such as nutrition, microbial environment, and physical/chemical  
306 conditions determine the healthy development and the survival rate. Moreover, all organs develop during the  
307 embryonic and larval stages, and often, causes of permanent abnormal development occur during this early  
308 stage, ultimately influencing how fish will perform later in life (The Research Council of Norway, 2009).

309

310 Up to date, only a few works have focused on reviewing research conducted on this species. Such works have  
311 mainly focused on the potential of *Solea solea* and the closely related species *S. senegalensis* for commercial

312 aquaculture. These include the work by Baynes *et al.* (1993), which reviews the egg production by sole in  
313 captivity, and Howel (1997) summarizes the challenges and benefits of cultivation of sole. However, the  
314 information provided is limited to assessing the technical rather than the economic feasibility of farming sole.  
315 More recently, the works by Imsland *et al.* (2003), Morais *et al.* (2014), and Munoz-Cueto *et al.* (2019) focused  
316 on presenting the biology and discussing the potential use of *Solea spp.* -*S. solea* and *S. senegalensis* for  
317 aquaculture. In this last reference, a chapter by Solé (2019) walks the reader through the use of these species  
318 within ecotoxicological studies, particularly by measuring biomarkers in sole tissues.

319

320 To our knowledge, this is the first review that exclusively focuses on *S. solea* larval biology and ecotoxicology.  
321 In our opinion, this could be partly explained given that only until recently, advances in culturing marine fish  
322 species and in developmental biology -notably on model species such as zebrafish- have allowed for the entire  
323 understanding of many biological processes and have made the field of larval ontogenesis more accessible for  
324 experimental exploration (The Research Council of Norway, 2009). Added to this, the threats imposed by  
325 climate change and pollution justify the need for summarizing those research works devoted to understanding  
326 the individual or combined effects of CC and anthropogenic activities on larvae.

327

328 For this review, we searched various academic search services (Thomson Reuters, Web of Knowledge, and  
329 Web of Science) for works published in the period between 2003 and 2020 using combinations of the following  
330 topics: *Solea solea*, flatfish, larvae, larviculture, ontogenesis, chemical contamination, and climate change. We  
331 choose to start our searching from 2003 onwards to avoid duplicating information already available in Imsland  
332 *et al.* (2003).

333

## 334 **2.1 Why does fundamental research on larval biology matter?**

335 Fundamental research in larval biology is essential for assuring the ecosystem functioning of nursery areas.  
336 Larvae biology is also crucial for marine fish farming. Because all organs and biological systems develop  
337 during the embryonic and larval stages, early development strongly influences how the fish performs later in  
338 life (Howel and Baynes, 2004). Therefore, knowledge about optimal larval development can substantially  
339 impact further stages and make a big difference in economic and sustainable terms. An appropriate example  
340 was reducing halibut discards due to incomplete eye migration, which accounted for up to 40-60% of halibut  
341 fry. Such a problem was solved by only applying photoperiods during larval rearing (Solbakken and Pittman,  
342 2004).

343

344 There is potential for improvements in aquaculture cost efficiency by increasing growth and survival rates and  
345 reducing the prevalence of deformities, and the key for such improvements might be related to larval  
346 development. Moreover, for wild stocks, knowledge about early life stages provides insight into how the fish  
347 survive under various conditions by adapting to their environment. Several reasons pinpoint fish early-life-  
348 stages (ELS) as more sensitive to pollutants than fully developed individuals from the same species  
349 (Hutchinson *et al.*, 1998). These reasons are related to a combination of critical developments occurring during

350 ELS, the limited possibilities to compensate for the impact of physiological defects, and significant changes  
351 in the composition of the developing tissue (Foekema *et al.*, 2012).

352

## 353 **2.2 Advances made on larvae survival during rearing**

354 Before standardizing a massive farming production of the sole, several problems are to be overcome, including  
355 larval mortality, feeding behavior, susceptibility to disease, stocking density, and reproduction. Besides, there  
356 are high mortality rates during juvenile production around metamorphosis and weaning (Bonaldo *et al.*, 2011).

357

358 For successful larvae rearing, the first factor to consider is related to the quality of the eggs, which can be  
359 defined as the egg's ability to be fertilized, survive, and develop into a normal embryo (Lund *et al.*, 2008;  
360 Parma *et al.*, 2015). As the egg must satisfy the nutritional needs of embryonic and larvae stages, and because  
361 healthy larvae depend largely but not exclusively (hormones and other molecules transferred in eggs are also  
362 crucial for embryonic and larval development), on the amount of fatty acids (FAs), the quality of eggs is usually  
363 assessed by studying its lipid and FAs composition (Lund *et al.*, 2008; Parma *et al.*, 2015). Besides,  
364 broodstocks' nutrition and life history has been shown to affect egg nutrients qualitatively and quantitatively  
365 (Foekema *et al.*, 2012; Parma *et al.*, 2015).

366

367 In order to study the relationship between hatching rate, survival, and growth of larvae and egg quality, Parma  
368 *et al.* (2015) analyzed and characterized the FAs composition of *S. solea* eggs obtained from domesticated  
369 broodstock during an entire spawning season. According to their results, the authors conclude that the feed  
370 supplied to broodstock before and during vitellogenesis has a significant role in determining the eggs' FA  
371 profile in sole. Another objective of their work included defining the relationship between FAs and egg  
372 viability parameters and larval survival as a means to provide information for improving -if necessary-  
373 broodstock feed in common sole. Regarding this last objective, the authors observed that, on the one hand,  
374 saturated and monounsaturated (MUFA) fatty acids were positively correlated with the hatching rate. On the  
375 other hand, TL, 22:6n-3 (DHA), 20:4n-6 (ARA), polyunsaturated FAs of the (n-3) series (n-3 PUFA), and  
376 polyunsaturated FAs of the (n-6) series were negatively correlated with the hatching rate. MUFA, 20:5n-3  
377 (EPA), n-6/n-3 were positively correlated with survival rate, while DHA, n-3 PUFA, DHA/EPA were  
378 negatively correlated. The relationships found between total lipids and FAs with egg and larval viability  
379 parameters showed a positive correlation with egg quality, while n-3 PUFA, DHA, DHA/EPA, and ARA  
380 content were inversely correlated with egg larval survival. This is typically different from many other farmed  
381 marine fish species, suggesting the need for a specific broodstock feed for this species.

382

383 In order to elucidate if offspring quality was affected by culture-related conditions, Lund *et al.* (2008)  
384 compared the eggs' quality -via FAs composition- and its relationship to egg size, fertilization, and hatching  
385 rate in eggs coming from cultured and wild broodstock. The FAs profile allowed discriminating between  
386 culture and wild origin. Eggs from wild broodstock were bigger, had higher levels of C16:1(n-7), C20:4(n-6),  
387 and C20:5(n-3), and larval survival was significantly higher than that from cultured broodstock. Cultured

388 broodstock eggs had high levels of C20:1(n-9), C18:2(n- 6), and C18:3(n-3), and their fertilization and  
389 hatching rates were generally lower than eggs from wild broodstock but not related to FAs eggs composition  
390 (Lund *et al.*, 2008).

391  
392 In a more recent series of experiments, De Swaef *et al.* (2017, 2019) aimed to achieve a more sustainable sole  
393 larviculture. They developed several research tools designed for tackling some key issues hampering  
394 commercial sole aquaculture. These tools include the design of a 24-well microplate rearing system (De Swaef  
395 *et al.*, 2017), the evaluation of several probiotic and prebiotic candidates to increase sole larval health (De  
396 Swaef, 2019), and the evaluation of disinfection protocols to control the bacterial disease vibriosis (De Swaef  
397 *et al.*, 2018).

398  
399 For the development of a 24 well microplate rearing system, the authors argue that by rearing larvae  
400 individually, problems related to group housing, such as variation between replicates and changes in the water  
401 quality of the system after the death of one larva, can be avoided (De Swaef *et al.*, 2017; De Swaef, 2019) and  
402 consequently larvae mortality during rearing decreased. Consequently, developing a system for rearing fish  
403 larvae individually guarantees that a dead individual will not affect the health of an adjacent one and facilitates  
404 the monitoring of their health status, growth, and behavior. It also improves the experiment's reproducibility  
405 and finally (because each well is considered as a replicate), it reduces the number of experimental animals  
406 needed (De Swaef *et al.*, 2017).

407  
408 Many of the problems faced during larval sole rearing relate to the fact that the molecular mechanisms  
409 underlying physiological and morphological changes during the larval-to-juvenile transition are still unknown.  
410 Because of this, efforts to sequence and characterize the transcriptome of *S. solea* have been done (Ferraresso  
411 *et al.*, 2013; Benzekri *et al.*, 2014). Ferraresso *et al.* (2013) revealed that pre-metamorphic larvae display a  
412 distinctive transcriptomic landscape compared to previous and later stages. The highest number of  
413 differentially expressed genes was found between 1 day-post-hatching (dph) and 4 dph, with a total of 1,539  
414 significant genes (974 over and 565 under-expressed in 4 dph larvae), while only 120 genes (81 up- and 39  
415 down-regulated) displayed a change in expression between 11 and 13 dph. In detail, genes related to the  
416 development of the visual system and neuronal processes are up-regulated during the early stages of larval  
417 development, while muscle development and anaerobic energy pathways increase in expression over time.  
418 Moreover, the expression profile of genes within the thyroid hormones cascade, the growth hormone, and the  
419 insulin-like growth factor genes revealed their crucial role in fish growth and initiation of metamorphosis  
420 (Ferraresso *et al.*, 2013).

421  
422 The development of visual perception is essential for larval feeding. At hatching, the eyes are slightly  
423 pigmented or non-pigmented and most likely non-functional. Therefore, successful first-feeding depends on a  
424 well-developed eye and visual system since fish larvae are essentially visual feeders. Light intensity is among  
425 the variables that could affect the survival of larvae during rearing. Bonvini *et al.* (2016) studied the effect of

426 different light regimes on *S. solea* larvae reared in the laboratory. They prepared 4 different rearing treatments  
427 with the following light intensities 1000 Lux, 500 Lux, 50 Lux, and 3 Lux. Larvae were exposed to the different  
428 light intensities from fertilization until metamorphosis, and effects on development, metamorphosis success,  
429 and growth were evaluated. The authors conclude that sole larvae require different light intensities along  
430 ontogenesis. High light intensity (between 50-500 Lux) is recommended during the pelagic stage (4–12 dph),  
431 in which first feeding larvae would benefit from light to initiate feeding. Also, histological examination  
432 revealed the importance of vision and light in the first feeding of this species (Bonvini *et al.*, 2016). After  
433 metamorphosis, the full development of other sensory organs indicates that feeding activity is also mediated  
434 by chemosensory perception, and thus low light intensity (3-50 Lux) is recommended to ensure better growth  
435 from the onset of metamorphosis to the benthic phase (Bonvini *et al.*, 2016).

436

### 437 **2.3 Advances made on larvae weaning, nutrition, and susceptibility to disease**

438 Nutrition plays a key role in sustainable aquaculture development, where fish health and wellness are of prime  
439 importance. Problems such as reduced growth, low survival, malpigmentation, and malformations are often  
440 encountered when fish are subjected to stress or deficient diet (Piccinetti *et al.*, 2012). Moreover, standard  
441 feeding regimes during larval and early juvenile stages represent a challenge for fish farmers due to the required  
442 administration of live feed, which is usually characterized by variable availability, high price fluctuations, poor  
443 hygienic conditions, and high levels of pathogenic bacteria (Bonaldo *et al.*, 2011). Thus, the feasibility of early  
444 weaning in common sole is a key factor in the industrialization of farming processes of this promising  
445 aquaculture species. In this context, Bonaldo and collaborators (2011) experimented with different feeding  
446 regimes and evaluated the effects that weaning *Solea solea* larvae as early as 13 days post-hatching (dph) has  
447 over metamorphosis success, survival, and fish size. The experiment lasted until 29 dph and consisted of two  
448 groups, a control one fed with live feed from 4 dph until 27 dph; and a second one weaned after 13 dph using  
449 three different microdiets (MDs), AgloNorse, Ewos (FR-A), Gemma, Skretting (FR-G) and O.range Large,  
450 INVE (FR-O) (Bonaldo *et al.*, 2011). The authors did not find significant differences in survival nor the  
451 metamorphosis development between control and the early-weaned group. However, the larvae that were  
452 weaned earlier showed lower weight and length than the control ones (Bonaldo *et al.*, 2011). Besides, they  
453 obtained size variations regarding the type of MD administrated, with FR-G showing the lowest standard  
454 growth rate. The authors conclude that it is feasible to conduct very early weaning on common sole and that  
455 this practice is advantageous as it reduces the number of *Vibrio spp.* in the water, preventing disease outbreaks  
456 (Bonaldo *et al.*, 2011). Finally, because larvae length and fatty acid body composition were significantly  
457 influenced (reduced) by MDs, growth was negatively affected by the early introduction of MDs. Thus, the  
458 authors hinted at the need for further research linked to nutrient utilization and the digestive capability of sole  
459 larvae as a means to further improve larvae performances.

460

461 On this last note, Parma *et al.* (2013) studied the molecular ontogeny and expression of digestive enzyme  
462 precursors on early-weaned larvae. They aimed to gain further knowledge of the effects of early weaning on  
463 the performances and physiological mechanisms during larval development of common sole, optimize

464 weaning protocols further and reduce mortality around this period (Parma *et al.*, 2013). For that, the authors  
465 cloned partial sequences of the enzymes trypsin (trypsinogen 1 (tryp1), trypsinogen 3 (tryp3), trypsinogen Y  
466 (trypY), chymotrypsin (chymotrypsinogen (chymt)), amylase (amy), and hepatic lipase (HL), which are among  
467 the most important pancreatic enzymes in teleost fish (Parma *et al.*, 2013). The expression of the former genes  
468 was assessed from the onset of exogenous feeding until the end of metamorphosis. Four weaning protocols  
469 (WPs) were evaluated, a control one with weaning occurring only at 27 dph, two intermediate treatments at  
470 which live food was provided until 13 and 18 dph, and a final treatment with larvae weaned to micro diets at  
471 4 dph. Based on their results, Parma and collaborators conclude that the different WPs did not greatly influence  
472 the expression of selected genes. On the contrary, survival, growth rate, and metamorphosis success were  
473 influenced by the WP, notably for larvae weaned at 4 dph, which showed the lowest survival rate (18.6% vs.  
474 a mean survival rate of  $52.8\% \pm 3.1$  for the 3 remaining protocols) and the lowest growth rate for the first 13  
475 days (SGR of 4.4 vs. a mean specific growth rate of  $17.3 \pm 0.4$  for the 3 remaining protocols). Finally,  
476 regardless that most larvae started metamorphosis on day 19 (71% of larvae), none had reached complete  
477 metamorphosis by the end of the experiment in the early-weaned group (Parma *et al.*, 2013).

478  
479 Regarding nutrient utilization, Piccinetti *et al.* (2012) tested the effects of a total or partial replacement of live  
480 feed (mainly consisting of rotifers and *Artemia nauplii*, group A/control group) with a copepod-based diet  
481 (mixed diet of live feed and dry copepods, named group B) on common sole growth, survival, lipid  
482 composition, lipid assimilation, defense to stress and malpigmentation (Piccinetti *et al.*, 2012). Their motivation  
483 was based on the observation that traditionally used live preys like rotifers and *Artemia* have nutritional  
484 deficiencies that result in a general decrease in fish health, development, growth, and pigmentation (Piccinetti  
485 *et al.*, 2012 and references therein). To develop rearing schedules suitable for market production, the authors  
486 conducted an experiment in which live preys were replaced with preserved copepods - the natural food of  
487 larvae- (Piccinetti *et al.*, 2012). For that, starting from day 9 post-hatching, group B larvae were fed a mixed  
488 diet (co-feeding), while group A and C were fed on live feed and copepod exclusive diets, respectively. Their  
489 results showed that including a complement of dry copepods (group B) within the diet increased survival by  
490 over 8% compared with group A (fed with rotifers and *Artemia nauplii*). In addition, larvae fed with a mixed  
491 diet started and ended metamorphosis faster than the group fed with live prey (group A: 16 dph- 25 dph vs.  
492 group B: 13 dph - 20 dph). Malpigmentation was significantly lower in group B than group A. Finally, larvae  
493 from group B had a higher standard growth rate both in terms of body weight (BW) and total length (TL) than  
494 individuals from group A. The authors explain that live prey motion is an essential attribute in stimulating  
495 larval predatory activity, which explains why the best results, in terms of survival, growth, and pigmentation,  
496 were achieved when the preserved copepods were administered together with live preys (rotifers and *Artemia*).  
497 Diet with only preserved copepods (group C) resulted in 100% mortality after 7 dph (Piccinetti *et al.*, 2012).

498  
499 The outbreak of infectious diseases can impact survival in marine fish larviculture (De Swaef *et al.*, 2018;  
500 Dinis *et al.*, 1999; Morais *et al.*, 2014; FAO, 2019). Microbiota can help to prevent and control diseases.  
501 However, knowledge about the establishment and action of microflora in fish's gut during larval stages is still  
502 scarce (Vino *et al.*, 2006). The most common members of the microflora of healthy marine fish are *Vibrio*

503 *spp.*, *Pseudomonas spp.*, and *Acinetobacter spp.* (Vino *et al.*, 2006). Vibriosis, a bacterial infection caused by  
504 abnormal/unbalanced colonization of *Vibrio spp.*, is among the diseases of primary concern for *S. solea*, being  
505 particularly challenging to tackle during early life stages (Austin and Austin, 2016; Novriadi, 2016).

506  
507 Thus, another research subject in fish nutrition that has been primarily studied as an alternative for improving  
508 disease resistance and nutrient absorption is the effects that dietary manipulation can have over fish gut  
509 microbiota, mainly by the administration of prebiotics and probiotics (Ringø *et al.*, 2014; De Swaef, 2019).

510  
511 During the last decade, many works have studied the effects of prebiotics and probiotics, not only for their role  
512 in improving disease resistance (as an immunostimulant) but also as a functional dietary supplement. Details  
513 on this subject escape from the scope of this work, and we refer the interested reader to consult the many other  
514 reviews available on the subject (Bricknell and Dalmo, 2005; Avella *et al.*, 2011; Ringø *et al.*, 2014; Torrecillas  
515 *et al.*, 2014; Vine *et al.*, 2006). However, in general lines, manipulation of gut microbiota has proven beneficial  
516 in adult fish, enhancing growth and feed efficiency and immunity and disease resistance (Ringø *et al.*, 2014).  
517 Regardless of many studies describing such beneficial effects, literature on the effects of prebiotics and  
518 probiotics administration to fish larvae is somewhat limited (De Swaef, 2019). In this regard, and to reduce  
519 losses in early stages of larval development and maximize production of captive common sole, Avella *et al.*  
520 (2011) studied the effects of administrating *Enterococcus faecium* IMC 511, isolated from sole broodstocks,  
521 as a probiotic candidate during *S. solea* larval development. During the first 50 days post-hatching, the  
522 probiotic was administrated daily through live feeds. The authors evaluated the intestinal presence of *E.*  
523 *faecium* IMC 511 during the experiment and the effects on intestinal microbial load, fish survival, welfare, and  
524 growth. For that, morphometric and molecular approaches were conducted. As markers of animal welfare, the  
525 authors monitored gene expression of cortisol and the 70 kDa Heat Shock Protein (HSP70). Their results  
526 showed reduced *Vibrio* populations in *S. solea* larvae intestines treated with the probiotic *E. faecium* IMC 511,  
527 but survival was not affected. The expression of stress markers (cortisol and HSP70) was significantly lower  
528 in probiotic-fed larvae than control larvae. By the end of the experiment, sole juveniles treated with probiotics  
529 had significantly higher body weight (BW) and total length (TL). Based on their results, the authors conclude  
530 that 50 days of probiotic treatment can improve common sole larval growth (Avella *et al.*, 2011).

531  
532 Finally, De Swaef (2019) assessed the impact of dietary supplementation in *S. solea* larvae of two prebiotic  
533 candidates derived from the aquatic environment, the mannanoligosaccharide and alginic acid. Contrary to the  
534 results previously cited, the tested prebiotics did not increase larval growth or survival and did not shift the  
535 proliferation and apoptosis rate of the intestinal epithelium (De Swaef, 2019). On the contrary, the  
536 administration of alginic acid had a negative effect on standard body length and the larvae supplied with 0.2%  
537 alginic acid followed by a challenge test (consisting of infection by *V. anguillarum* WT at a final concentration  
538 of  $1 \times 10^7$  colony-forming units/mL at 10 DAH) displayed a lower survival in comparison with the challenged  
539 larvae without any supplementation. These results reveal a possible negative impact of alginic acid  
540 supplementation on common sole larvae (De Swaef, 2019) and highlight the need for further research on the



541 benefits of prebiotics, probiotics, or synbiotics (nutritional supplements combining a mixture of probiotics and  
542 prebiotics in the form of synergism, Ringø *et al.*, 2014) in *S. solea* larvae.

543

544 Because fish are free-living organisms since their initial stages, they are exposed to numerous pathogens even  
545 before their lymphoid organs, and consequently, their adaptive immunity has fully developed (Ferraresso *et al.*  
546 *al.*, 2016). Therefore, to defend the host from infection, newly hatched fish rely on their innate immune  
547 repertoire that acts in a non-specific manner. Ferraresso *et al.* (2016) argued that for developing effective  
548 approaches for disease management at early life stages of *S. solea* is necessary to understand the mechanism  
549 related to innate immune responses. For that, in their study, the authors characterized and assessed the  
550 transcriptional onset of unexplored relevant genes of both innate and adaptive immune systems during the *S.*  
551 *solea* ontogenesis, from hatching to the accomplishment of the juvenile form (Ferrareso *et al.*, 2016). Their  
552 results showed that several genes involved in the innate immune repertoire are up-regulated during the first  
553 stages (such as lysozyme, AMPs (hepcidin, b-defensin), PPRs, and complement components), supporting the  
554 hypothesis of protection from environmental pathogens during early development. Moreover, the onset of the  
555 expression of adaptive immune genes (i.e., Class I and class II MHC, TCRs) coincides with metamorphosis  
556 and larvae-to-juvenile transition. Finally, data collected suggest that full maturation of the cell-mediated  
557 immune system of *S. solea* is not completely mature by the end of metamorphosis (Ferraresso *et al.*, 2016).

558

#### 559 **2.4 Pigmentation and eye migration**

560 Other developmental problems related to the successful commercialization of soles are problems related to the  
561 appearance of the fish product, which can be insufficient for the market standards, such as malpigmentation  
562 and incomplete eye migration.

563

564 Pigmentation problems are known to be related to vitamin A, fatty acid composition, and thyroid hormones.  
565 However, in most cases, it is an unbalanced fatty acid composition of the feed the main reason for  
566 malpigmentation of farmed flatfish, and the thresholds for concentrations of these essential fatty acids are  
567 species-specific (The Research Council of Norway, 2009). In this regard, Lund and collaborators (2008)  
568 studied the effects that diets supplemented with arachidonic acid (ARA) and eicosapentaenoic acid (EPA) have  
569 on sole survival, metamorphic success, and notably pigmentation. They hypothesized that the induction of  
570 malpigmentation (*i.e.*, hypomelanosis) was linked to a specific larval period or “a pigmentation window” at  
571 which larvae sensitivity to ARA increases. Their results confirmed their hypothesis of the presence of an early  
572 pigmentation window at pre-metamorphosis. Also, they revealed that in common sole larvae, ARA induced  
573 malpigmentation, but no effect was associated with diets enriched with EPA (Lund *et al.*, 2008). Regarding  
574 metamorphosis success, the initiation or advance of eye migration was not related to dietary ARA, EPA, but  
575 the size of the individuals (Lund *et al.*, 2008).

576

### 577 **Section 3 - Effects of temperature and contamination on *Solea solea* larvae**

578

579 Temperature is the main factor regulating flatfishes' species range. Human-induced climate change has played  
580 a central role in larval dispersal variability, leading to species losses and invasions, biogeographical and regime  
581 shifts, and changes in the adaptation capabilities of organisms (Lacroix *et al.*, 2016). Also, temperature  
582 modulates the duration of the egg and larval stages and, therefore, determines the distance between spawning  
583 and nursery grounds (Gibson, 2015). Further, the species range and the average level and the annual variability  
584 in recruitment regulate sole populations. The variability in recruitment is governed by two distinct processes,  
585 the effect of food availability on the adult condition at spawning time and the density-dependent mortality of  
586 juveniles on the nursery grounds (Gibson, 2015). Finally, temperature also affects fish growth rate and ultimate  
587 size.

588

589 Chemical exposure from anthropogenic activities negatively impacts marine habitats, notably on coastal  
590 habitats, where most of the human population is established (60% at 100 km from the water; Brown, 2006).  
591 Moreover, several species of economic interest expend part of their life cycle on coastal habitats and estuaries,  
592 where specific characteristics in temperature and physicochemical conditions guarantee their healthy  
593 development (Seitz *et al.*, 2014). However, estuaries can accumulate and stock several pollutants in the  
594 sediment compartment (Ridgway and Shimmield, 2002). Previous works that conducted inter-estuarine  
595 comparisons showed that the chemical contamination status notably affects individuals' growth and survival  
596 and thus the quality of the nursery area (Gilliers *et al.*, 2006; Amara *et al.*, 2007). In the case of *Solea solea*,  
597 unwanted changes in nursery ecosystems could affect (i) juvenile settlement, (ii) survival of juvenile  
598 specimens, and (iii) healthy development and growth, which in the long term could impact the renewal capacity  
599 of the population –recruitment-, and impact the species fitness.

600

### 601 **3.1 Changes in spawning, larval development and survival, and recruitment related to** 602 **temperature**

603

604 In recent years, fish spawning patterns and growth rates have been changing. Increased temperatures are  
605 believed to have various effects on fish populations, including effects on their phenology by altering growth,  
606 the survival of juveniles and adults, and reproductive traits, such as the timing of spawning (Alix *et al.*, 2020).  
607 Shifts in the timing of spawning will have both direct and indirect consequences for fishery management and  
608 could have significant implications on species recruitment and population size. These effects include –among  
609 other things– mismatches between the timing of larval development and the availability of planktonic food  
610 sources, and a reduction in the effectiveness of fishery closures during spawning seasons (Fincham *et al.*, 2013;  
611 Alix *et al.*, 2020).

612

613 In their work, Fincham *et al.* (2013) evaluated temperature data from coastal monitoring sites situated close to  
614 spawning grounds, together with data on *Solea solea* timing of spawning. The last data came from market  
615 sampling programs in ports in England and the Netherlands, where random samples of fish landed by  
616 commercial fishing vessels are taken, and biological data, such as the proportion of spawning females amongst  
617 the total number of sampled adult females, was calculated for each week of the year for a time series of 40

618 years (Fincham *et al.*, 2013). Their results showed that four out of the seven stocks they studied had a  
619 significant shift in the time of spawning in response to increased sea surface temperature (SST). In detail, the  
620 stocks in the Irish Sea, east-central North Sea, southern North Sea, and the eastern English Channel spawned  
621 in earlier weeks of the year at a rate of 1.5 weeks per decade (Fincham *et al.*, 2013). They observed both spatial  
622 and temporal differences, which were strongly correlated to temperature. They also found that winter SST is  
623 inversely correlated with the timing of spawning, which is caused by the faster maturing of gonads due to  
624 warmer temperatures during the winter. The authors conclude that in a warmer world, *S. solea* populations  
625 from the English Channel will spawn earlier (Fincham *et al.*, 2013). However, as recruitment depends on food  
626 availability, changes in the phenology can have important implications for the population dynamics (Gibson,  
627 2015). Evidence suggests that the timing of peaks in larval food abundance may not necessarily advance with  
628 warming temperatures (Wiltshire *et al.*, 2008). A mismatch between the timing of spawning and hatching  
629 periods with the peak of the phytoplankton or copepods will reduce the recruitment success and abundance of  
630 the species.

631  
632 Although temperature seems to be the main driver to alter sole spawning phenology from 1970 until 2010, the  
633 authors do not dismiss the effect of other factors, including population density, growth, and maturity at age.  
634 However, and because the temperature has a positive influence on the growth of age-0 sole (Teal *et al.*, 2008),  
635 if phenology changes were due to changes in growth rate, then the rising temperature should delay the timing  
636 of spawning, resulting in bigger but younger fish at first spawning (Fincham *et al.*, 2013). Nevertheless, there  
637 is evidence that sole size and age at first spawning have decreased (Mollet *et al.*, 2007), while the timing of  
638 maturation may be delayed in smaller, relatively more abundant fish in their first year of maturation (Ramsay  
639 and Witthames, 1996). Thus, if growth affects phenology, the rising temperature would lead to the reverse  
640 effect of that observed by the authors, *i.e.*, delayed timing of spawning. This reasoning strongly suggests that  
641 the observed phenological shifts are less likely to be the indirect result of changes in the size-age composition  
642 of spawning fish but a more direct effect of winter temperatures on gonad maturation (Fincham *et al.*, 2013).

643  
644 After the spawning, the transport of eggs and larvae from spawning grounds to nursery areas is primarily driven  
645 by hydrodynamic processes. However, behavioral and environmental factors might also influence the final  
646 dispersal pattern as well as the larval survival and abundance (Lacroix *et al.*, 2013). The survival of the early  
647 life stages is dependent on environmental constraints -notably temperature- the availability of food, and the  
648 presence of predators (Lacroix *et al.*, 2013). After that, fish recruitment is dependent on larval abundance and  
649 success of metamorphosis at nursery areas. Therefore, understanding larval dispersal is of utmost importance  
650 as it might explain recruitment patterns and has direct consequences for the conservation of fish, stock  
651 management, the design of marine reserves, the management of non-indigenous invasive species, and the  
652 prediction of climate change effects (Lacroix *et al.*, 2016).

653  
654 On the question about larval dispersal, Lacroix and collaborators have notably contributed with a series of  
655 works (Savina *et al.*, 2010; Lacroix *et al.*, 2013, 2016; Barbut *et al.*, 2019) for which the main focus has been  
656 to understand the impact of interannual variability of hydrodynamics and temperature on larval dispersal and

657 supply to the nurseries. The works also aim to predict the potential changes in *S. solea* dispersal patterns under  
658 warmer scenarios due to climate change in the English Channel and the North Sea. Other relevant works related  
659 to sole larval dispersal in other regions include the works by Vinagre *et al.* (2007) in the Tagus Estuary in  
660 Portugal and the study by Tanner *et al.* (2017) in the Iberian coast.

661

662 For evaluating larval dispersal, a larval transport model for sole coupled with a 3D hydrodynamic model of  
663 the North Sea was implemented, and interannual variability of the transport of fish larvae during different  
664 periods was investigated (Savina *et al.*, 2010; Lacroix *et al.*, 2013, 2016). Shortly, the study of 2013 showed  
665 important interannual variability (coefficient of variation of 29.2%) of the total larval abundance at the  
666 nurseries. The larvae dispersal patterns were broader than those reported in the first generation model by Savina  
667 *et al.* (2010). In their work from 2016, the inclusion of wind speed and directions as factors in the model  
668 together with simulations based on predicted scenarios from the “Warm+” IPCC showed an increase in the  
669 dispersal distance (+70%) and an increase in the pelagic larval duration (+22%) which was in response to  
670 reduced temperature (-9%) endured by larvae that hatched earlier (Lacroix *et al.*, 2016).

671

672 In recent work, Barbut *et al.* (2019) evaluated the relationship between dispersal capacity and larval traits, such  
673 as behavior, pelagic larval duration, seasonality of reproduction, and structure of the water column and  
674 currents, from populations of six commercially exploited flatfish species in the North Sea. Their results  
675 demonstrate considerable variation in both connectivity and spatial distribution of all studied species, with two  
676 groups emerging with similar life-history traits and dispersal patterns features (Barbut *et al.*, 2019).

677

678 The set of papers from this group tell a coherent story in which knowledge about larval dispersal increases  
679 progressively. However, this group of researchers' methods and the model are common, and since they are  
680 based on a single model, the level of uncertainty was not included. On this last point, we direct the reader to  
681 Hufnagl *et al.* (2017 and references therein) work in which the variability that can be expected concerning  
682 connectivity, uncertainty, and the order of magnitude of variability resulting from choosing different physical  
683 models, was addressed by comparing a suite of 11 North Sea models (Hufnagl *et al.*, 2017).

684

685 On a final note, warmer scenarios may improve the survival of juveniles. Earlier spawning will prolong the  
686 growing season of 0-group fish and may result in an increase in the body size during the 1st winter (Teal *et al.*,  
687 2008), and since winter survival is positively related to body size, 0-group cohorts with enhanced energy  
688 stores are more likely to survive (Post and Parkinson, 2001). However, evidence suggests that fish size may  
689 shrink in response to climate-induced changes in temperature and oxygen (Baudron *et al.*, 2014). The  
690 physiology behind the pattern is well established for aquatic ectotherms, higher temperatures decrease the  
691 aerobic capacity, and individuals with smaller body sizes have a reduced risk of oxygen deprivation (Baudron  
692 *et al.*, 2014). For *S. solea*, there is evidence showing that exposure to elevated temperatures during early  
693 developmental stages alters the environmental adaptation performance of fish at later stages, making juveniles  
694 more resistant to later in life hypoxia events (Zambonino *et al.*, 2013). However, and in contradiction to

695 previous results, Zambonino *et al.*'s (2013) results showed that higher temperatures during early stages resulted  
696 in larger-bodied individuals.

697

### 698 **3.2 Changes in spawning, larval development, and survival, related to chemical exposure**

699

700 The presence of significant levels of pollutants in sole tissues can affect their physiological performance and  
701 harm the health of consumers. Given this, several studies assessing the bioaccumulation capacity and toxicity  
702 of different chemical contaminants (organic pollutants and metals) have been conducted using *Solea solea* as  
703 target species (Dierking *et al.*, 2009; Wessel *et al.*, 2010; Trisciani *et al.*, 2011; Cannas *et al.*, 2013; Sanchez-  
704 Nogué *et al.*, 2013; Sister *et al.*, 2013; Solé *et al.*, 2013; Cuevas and Zorita, 2018; Mounier *et al.*, 2020a,  
705 2020b), some other study focused on differential mortality in age-0 juveniles (Guinand *et al.* 2011). However,  
706 all these studies have focused on the juvenile stage and the ecotoxicity of contaminants at the early life stages  
707 of *S. solea* it is still, up to date, understudied.

708

709 Although working with juvenile fish in laboratory conditions requires specific facilities and is logistically  
710 challenging, most ecotoxicity tests conducted with soles are designed at this developmental stage because it  
711 guarantees enough tissue for running tests in target organs (*i.e.*, biomarker and bioaccumulation  
712 analyses). Moreover, working with juveniles reduces sex-related endpoints variability and does not consider  
713 the influence of reproduction on bioaccumulation. Larval stages (*i.e.*, egg, yolk-sac larvae, and free-feeding  
714 pre-metamorphic larvae) are considered to be the most sensitive stages of soles life cycle (Ferraresso *et al.*,  
715 2013). As discussed in the previous sections, mortality under control conditions still reaches up to 60%  
716 (Foekema *et al.*, 2008, 2014), and protocols for rearing larvae in the lab still need to be refined (De Swaef,  
717 2019). At the same time, the fields of biology and toxicology have seen some important developments.  
718 Advances in measurement technologies, computational capabilities, fundamental toxicological understanding  
719 at the molecular level, and *in vitro* technologies have increased the amount of information produced following  
720 ecotoxicity tests (Ankley *et al.*, 2010). This fact, together with the refinement of analytical protocols (Olsson  
721 *et al.*, 1978; Thomson and Budzinski, 2000) and the proposal of easier to manipulate rearing methods (such as  
722 the microplate early life stages adapted protocol presented in De Swaef *et al.*, 2017) opens the door for  
723 conducting ecotoxicity tests with sole larvae.

724

725 There are three sources for contaminant exposure during the larval stages. The first is the result of parental  
726 exposure. During fish vitellogenesis, most of the lipid resources are stored in the yolk, and during this process,  
727 lipophilic pollutants are also transferred to the eggs (Foekema *et al.*, 2012). The second exposure route occurs  
728 after hatching when pelagic larvae start the uptake and elimination (through detoxification mechanisms) of  
729 pollutants from the water. Once the larvae deplete the yolk reserves, it will start feeding, and therefore, the  
730 third exposure route is activated through ingestion of large amounts of lipid-rich -and potentially contaminated-  
731 zooplankton.

732

733 As the first route of exposure involves maternally transferred contamination, there is the question of the effects  
734 of lipophilic persistent organic pollutants (POPs) on early life stages (ELS).. For predicting sole's  
735 bioaccumulation of POPs, such as polychlorinated biphenyls (PCBs), Foekema *et al.* (2012) adapted the  
736 bioaccumulation model OMEGA (Hendriks *et al.*, 2001) to the ELS of sole turning it to ELS-OMEGA. The  
737 model was validated with experimental data and proved effective in predicting the development of PCBs'  
738 concentrations in the tissue of developing soles. The authors measured and modeled the concentrations of  
739 PCBs (52, 110, 101, 118, 138, and 153) in water and tissue during the development of an embryo into a juvenile  
740 fish. The model showed good fits to the measured data, and according to the simulations, tissue concentrations  
741 of pollutants with log octanol-water partition ratio ( $K_{ow}$ ) > 5 peaks when the lipid reserves are depleted at the  
742 moment that larvae become free-feeding, revealing that maternally transferred highly lipophilic compounds  
743 are barely excreted (Foekema *et al.*, 2012). Moreover, these peaks may exceed two to four times the  
744 concentration in the spawning parent, concentrations that could result in extreme body burdens during critical  
745 stages of larval development, with potentially severe consequences for larval survival. Once the fish starts  
746 feeding externally, fish increase rapidly in weight, and dilution by growth plays an essential role in tissue  
747 concentration decrease (Foekema *et al.*, 2012).

748  
749 The authors highlight the risk of underestimating the effects of maternally transferred PCBs (especially those  
750 with  $K_{ow}$  > 5) at the early-life-stages, principally because of the exposure to waterborne substances of a fish  
751 embryo inside the egg is limited compared with the post-hatch situation (Foekema *et al.*, 2012). Unfortunately,  
752 the ELS toxicity test focus on exposing eggs via the water, and very often, the experiment will end just before  
753 the end of the yolk-sac stage, the moment at which the internal concentration will peak to its maximum  
754 (Foekema *et al.*, 2012). The authors argue that terminating ELS tests before the yolk-sac stage ends is valid  
755 for testing substances with log  $K_{ow}$  < 5. They, however underline that when testing more hydrophobic  
756 substances, mixtures of substances, or unknown mixtures (such as those present in effluents or contaminated  
757 sediments), the toxic effects of the lipophilic compounds will be seriously underestimated (Foekema *et al.*,  
758 2012).

759  
760 The previous statement is supported by data from two more works by Foekema *et al.* (2008, 2014), in which  
761 delayed effects after exposure during the egg stage were observed on *S. solea* larvae. In their earlier work, the  
762 authors exposed *S. solea* eggs to a concentration series of PCB 126 in seawater until 4, 8, 10, and 15 days post  
763 fertilization (dpf), and the development of the larvae was followed up under unexposed conditions until larvae  
764 were fully metamorphosed (up to 50 dpf) (Foekema *et al.*, 2008). Their second work aimed to recreate the  
765 effects of maternally transferred POPs in early larvae development (Foekema *et al.*, 2014). The results from  
766 both studies indicate that ELS fish tests underestimate the effect of highly lipophilic compounds, which  
767 continue to affect survival and metamorphosis success long after the exposure, showing that, at least for sole,  
768 the moment of observation has a significant impact on the determination of the effect concentration  
769 (approximated by the  $LC_{50}$  in Foekema's works), even without further exposure (Foekema *et al.*, 2008; 2014).

770

771 **Section 4 - Methods that integrate the data -bridging the gap-**

772

773 Studying the effects of temperature and contamination along *Solea solea* larvae ontogenesis is relevant for  
774 several disciplines (Fig. 2). Researchers in fisheries and ecology want to know if the stocks will be secured in  
775 future years and, what will be the species biogeographical distribution, the impact of invasive species (e.g.,  
776 Marras *et al.*, 2015), and how spawning and larval dispersal might be affected by increased temperatures  
777 (Pankhurst & Munday, 2011). Aquaculture research is interested in understanding larvae's physiology and  
778 developing protocols for rearing them while minimizing the cost and maximizing survival. Environmental  
779 chemists develop analytical protocols to detect smaller and smaller concentrations of environmental pollutants  
780 that might be present in several matrixes (water, pore water, sediments, and biota). Their efforts orientate  
781 towards understanding and predicting bioaccumulation and transfer of pollutants in sole into the trophic web.  
782 Moreover, their results are input for fate and transport models. Finally, ecotoxicologists aim to predict and  
783 understand the effects of combined stress from exposure to a mixture of chemicals under future climate change  
784 scenarios to provide relevant data for chemical regulation.

785

786 Generating relevant data for risk assessment and chemical regulation is the major challenge that  
787 ecotoxicologists have and keep facing up to today. In short, the ecotoxicology domain aims to answer two  
788 impossible questions i) which of the millions of species are at risk from pollution? And ii) which of the tens  
789 of thousands of chemicals contribute most to the risk? As Ashauer and Jager (2018) stated, the great pitfall in  
790 ecotoxicology is that there is no theory to link the effects of toxicants at the molecular or cellular level to  
791 changes in life-history traits. Moreover, at the cellular level, there are plenty of examples that demonstrate that  
792 responses are cell-type and species-specific and conditioned by age, size, gender, season, and temperature,  
793 among many other biotic and abiotic variables (van der Oost *et al.*, 2003; Au, 2004; Martínez-Álvarez *et al.*,  
794 2005; Hardman, 2006). Also, multiple types of effects depend on the pollutant or the mixture of pollutants,  
795 and that can vary depending on the exposure route (Hardman, 2006). Even with a fast pace of research  
796 production, there is no way that we can catch up with the speed at which the industry produces new chemicals.  
797 So, how can we catch up with xenobiotics production? Furthermore, which are the tools that we need to bring  
798 together to bear on climate change and pollution effects?

799

800 Our poor knowledge of physiological modes of action (pMoA) and how they vary across species and toxicants  
801 has been recently identified as a major knowledge gap (Ashauer and Jager, 2018). Physiological MoA refers  
802 to how chemical stress (single compound or mixture) affects an organism's energy allocation and,  
803 consequently, life-history traits. It can be determined using Dynamic Energy Budgets (DEB) models, which  
804 have strong theoretical foundations (Kooijman, 2010). In their paper, Ashauer and Jager (2018) proposed  
805 pMoA as a key element to extrapolate from small to larger biological levels. We believe that the authors'  
806 proposition is correct and that researching pMoA is a promising venture.

807

808 According to the authors, knowing pMoAs is an excellent way to boost and complete the information that the  
809 two main approaches in ecotoxicology -data from chemical stress ecology and high-throughput bioassays of  
810 cellular or molecular markers ELS- offer to ecotoxicologists (Ashauer and Jager, 2018). Shortly, chemical  
811 stress ecology treats pollution as one of the many factors that influence individuals, populations, communities,  
812 and ecosystems. On the one hand, this approach aims to provide ecological relevance and realism to  
813 environmental risk assessments, but it fails to discriminate if the observed effects are due to a single pollutant  
814 or a mixture of them, giving little relevant information from the regulatory point of view. On the other hand,  
815 high-throughput bioassays orbits on the promises to upscale results from bioassay to organisms and beyond,  
816 rallying around the idea of quantitative adverse outcome pathways (AOPs) first proposed by Ankley *et al.* in  
817 2010. The AOP framework aims to provide a useful structure within which existing knowledge can be  
818 organized, from which key uncertainties and research priorities can be identified, and through which we can  
819 improve predictive approaches needed to advance regulatory ecotoxicology. It integrates evidence from *in*  
820 *vitro* assays linking a molecular initiating event (e.g., a molecular interaction between a xenobiotic and a  
821 specific biomolecule) and an adverse outcome in the context of a pathway that leads to an adverse outcome at  
822 a biological level of organization relevant to risk assessment (Ankley *et al.*, 2010). However, to answer which  
823 of the millions of species are at risk from pollution, it is necessary to extrapolate quantitative AOPs across  
824 species. For that, we need to assume that molecular pathways are qualitatively and quantitatively conserved,  
825 and data suggest that a significant fraction of receptors and target sites are not conserved between species  
826 (Gunnarsson *et al.*, 2008; Rand-Weaver *et al.*, 2013).

827  
828 For covering the present needs in ecotoxicology, focusing on determining the pMoA would be a strong tool as  
829 it would represent a middle point between AOPs and stress ecology. As organisms require resources to grow,  
830 develop, and reproduce, it is these traits that we ultimately require to link AOPs to ecological theory and  
831 upscale to the population level and higher (Ashauer and Jager, 2018). Changes due to toxicant exposure or  
832 environmental stress in energy-demanding traits -such as growth and reproduction- logically imply changes in  
833 the energy budget and model parameters. By using energy-budget models and linking the related parameters  
834 to AOPs, we will be able to 1) identify where energy allocation has changed due to a stressor -the physiological  
835 mode of action- and by how much, and 2) justify the observed changes -if any- with data from AOPs, which  
836 will therefore allow us to interpolate from the individual to the subcellular level. Further, pMoA acts at the  
837 individual level, and there are theories available in ecology that allow extrapolating from individual to  
838 population and community level (e.g., Physiologically Structured Population model (PSP), Caswell *et al.*,  
839 1997; Ghwila and Willms 2019). Thus, pMoAs could be the missing link between subcellular levels and higher  
840 levels of biological organization. Nevertheless, it is important to highlight that, on the possible types of pMoAs  
841 within a CC and contaminated scenario, one important dichotomy involves toxicant-induced changes that alter  
842 the ability of an organism to respond to CC stressors (toxicant-induced climate susceptibility [TICS]), in  
843 contrast to climate-induced toxicant sensitivities (CITS), which is the scenario where CC affects the toxicity  
844 of chemicals (Hooper *et al.*, 2013). We invite the reader to seek further details regarding pMoAs and AOPs in  
845 Ashauer and Jager (2018) and Perkins *et al.* (2019), respectively.



846 Finally, the reason why studying combined effects of temperature and chemical exposure is challenging is that,  
847 as previously mentioned, the main approaches followed by ecotoxicologists are incapable to either discriminate  
848 the separate effect of each stressor (as in chemical stress ecology) or measuring the mixed effect and translate  
849 it to a relevant level of biological organization (when using high-throughput bioassays). The implementation  
850 of DEB models provides the means to overcome this issue while taking advantage (or making use) of the  
851 information provided by these two ecotoxicological approaches. The ecological theory is indispensable to  
852 extrapolate from the individual to the population level, and subcellular data complements the information from  
853 model parameters, allowing us to understand the molecular mechanisms behind the pMoA. Further,  
854 concentrating efforts from different disciplines into a common modeling goal directly translate in reducing  
855 animal testing (by reusing data from tests already conducted), and since DEB models offer a real opportunity  
856 for species extrapolation (see Kooijman, 2010 for examples of model parameters variation related to body size;  
857 Baas *et al.*, 2015; Sardi *et al.*, 2019), it also translates in reducing costs for research associated to risk  
858 assessment.

859

860 We hope that within this section, we have managed to convey the following message: existing data can be  
861 synthesized in a useful way to risk assessors and ecotoxicologists, but it demands consensus and team effort  
862 from players that operate in different ways disciplines of research.

863

#### 864 **Further research recommendations and conclusions**

865

866 Working with DEB and derived models (GUTS see Jager and Ashauer, 2018a and b, and DEBtox, Kooijman  
867 and Bedeaux, 1996 or Jager and Zimmer, 2012) offers an excellent opportunity to answer (individually or by  
868 coupling them to other models) to several -if not all- the research questions asked in many different disciplines,  
869 and shortly stated at the beginning of the previous section (Fig. 2).

870

871 In DEB models, temperature and food availability are forcing variables (Kooijman, 2010). The last means that  
872 we could run simulations for higher temperatures and provide insights about the effects of sea warming on  
873 energy budgets allocation, growth, maturation, and reproduction success. Besides, DEB models can be coupled  
874 with larval dispersal and hydrodynamic models (Huret *et al.*, 2012), which will provide information about the  
875 time and size of fish at spawning for a given temperature and the dispersal of hatched larvae. Also, by  
876 combining DEB models with ecosystem models, it is possible to build habitat quality maps (*e.g.*, Teal *et al.*,  
877 2012) at which the effect of climate change on the spatio-temporal fish dynamics can be explored. Moreover,  
878 this type of approach can be conducted in similar species with different temperature optima (such as it is the  
879 case between *S. solea* and *S. senegalensis*) and provide insights about changes in species biogeographical range  
880 or native species displacement by invasive ones (see Solé, 2019).

881

882 In ecotoxicology, coupling DEB with TKTD models (DEBtox model, Kooijman and Bedeaux, 1996; Billoir  
883 *et al.*, 2007; Jager *et al.*, 2004; Augustine *et al.*, 2012; Sadoul *et al.*, 2019) allows for depicting the pMoAs  
884 which are potentially the missing link between subcellular processes and effects on higher levels of biological

885 organization. DEB and DEBtox models are so far the most powerful available tool that could allow for species  
886 and chemical extrapolation, and their standard implementation could be the necessary information for  
887 improving environmental risk assessments and chemical regulation.

888

889 In aquaculture research, larvae rearing methods could benefit from the information coming from simulations  
890 done with DEB models, in which growth is fitted for different temperature and food conditions. Also, DEBtox  
891 or GUTS models can be applied to evaluate the effect of antibiotics, probiotics, and disinfection methods often  
892 applied for reducing larvae mortality. This practice will potentially reduce the experimental cost and render  
893 the business profitable. This last point is not only crucial from a commercial perspective. Improving rearing  
894 systems and developing the aquaculture on this species also impact the food security and human health (by  
895 preventing humans from consuming fish that have bioaccumulated pollutants), and decrease the fishing  
896 pressure over natural populations, which might face difficulties in their renewal owing to the changes in  
897 temperature and pollution projected for future years (Fig. 2).

898

899 Finally, DEB models coupled to bioaccumulation models would provide the means to predict contaminant  
900 dilution by growth effect and to estimate the concentration of chemicals ingested by individuals along  
901 ontogenesis, both common questions in environmental chemistry research (Arnot and Gobas, 2006).

902

903 In a nutshell, even though the development of DEB models requires significant effort and investment -as it  
904 requires the user to understand DEB theory, its mathematical foundation, notation, and interpretation of model  
905 parameters- its application is potentially beneficial for many disciplines, opening the door for interdisciplinary  
906 research. In this line of thought, we would like to propose an interdisciplinary framework that aims to bring  
907 together and take full advantage of already existing data (often coming from different research fields) and tools  
908 (often operating at different levels of biological organization) to bear on climate change and pollution effects  
909 on *S. solea* larvae ontogenesis. Figure 3 illustrates such a framework. At least five different disciplines have  
910 an interest in studying the combined effects of temperature and chemical contamination in sole larvae, though  
911 their research questions (outer blue circle) differ. Answering those questions requires collecting/generating  
912 different types of data (purple circle), which are often used as input for calibrating models (dark pink circle)  
913 that operate at different levels of biological organization. Finally, all these models can be used either  
914 individually or in combination, but we can only achieve full understanding regarding the effects of  
915 contamination or temperature stress at the individual level after determining the pMoAs (light pink circle) and  
916 linking these to processes at the sub-cellular level with AoPs (yellow circle, Fig. 3).

917

918 To conclude, models are simply tools to deduce quantitative conclusions from a set of assumptions and data.  
919 In this case, the DEB family of models offers an incredible opportunity to ecotoxicologists searching for the  
920 missing link between effects at subcellular level and effects at the population and community level and,  
921 consequently, producing and presenting data in a way that is useful to environmental risk assessors.


922

923 **Acknowledgments**

924 This work was funded by the IdEx Bordeaux International Post-doctorates program attributed to A. E. Sardi.


925  
926 Figure 1. *Solea solea* life cycle. Soles have a complex life cycle that depends on biotic and abiotic variables.  
927 In the continental shelf, temperature triggers spawning from benthonic adults. After hatching, the pelagic  
928 larvae undergo extreme metamorphosis and migrate towards the nursery and coastal areas, all while  
929 undergoing metamorphosis. Migration and metamorphosis success depends on hydrodynamics and food  
930 availability. The length of metamorphosis varies depending on temperature and food availability, but it can  
931 last up to 15 days at 16 °C. The abbreviation dph stands for days-post-hatching.

932  
933




934  
935  
936  
937  
938

939 Figure 2. Scheme illustrating some of the research questions related to sole and larval ontogenesis for  
 940 different research disciplines. See the text for proposed tools and solutions.



941  
 942  
 943  
 944  
 945

946 Figure 3. Interdisciplinary framework to enhance ecotoxicology and climate change-related research in  
 947 fish larvae. Abbreviations stand for Climate Change (CC), GUTS (General Unified Threshold model  
 948 of Survival), DEB (Dynamic Energy Budgets), TKTD (Toxico-kinetic toxico-dynamics), IBM  
 949 (Individual-Based Model), AoPs (Adverse outcome pathways), and pMoAs (physiological modes of  
 950 actions).  
 951  
 952



953 **References**

- 954
- 955 Amara, R., Laffargue, P., Dewarumez, J. M., Maryniak, C., Lagardère, F., & Luczac, C. (2001). Feeding  
956 ecology and growth of O-group flatfish (sole, dab and plaice) on a nursery ground (Southern Bight of  
957 the North Sea). *J Fish Biol*, 58(3), 788–803. <https://doi.org/10.1006/jfbi.2000.1498>
- 958 Amara R., Meziane, T., Gilliers, C., Hermel, G., & Laffargue, P. (2007). Growth and condition indices in  
959 juvenile sole *Solea solea* measured to assess the quality of essential fish habitat. *Mar Ecol Prog Ser*,  
960 351: 201-208.
- 961 Ankley, G. T., Bennett, R. S., Erickson, R. J., Hoff, D. J., Hornung, M. W., Johnson, R. D., *et al.* (2010).  
962 Adverse outcome pathways: A conceptual framework to support ecotoxicology research and risk  
963 assessment. *Environ Toxicol Chem*, 29(3), 730–741. <https://doi.org/10.1002/etc.34>
- 964 Alix, M., Kjesbu, O. S., & Anderson, K. C. (2020). From gametogenesis to spawning: How climate-driven  
965 warming affects teleost reproductive biology. *Journal of Fish Biology*, (June).  
966 <https://doi.org/10.1111/jfb.14439>
- 967 Arnot, J. A., & Gobas, F. A. P. C. (2006). A review of bioconcentration factor (BCF) and bioaccumulation  
968 factor (BAF) assessments for organic chemicals in aquatic organisms. *Environ Rev*, 14(4), 257–297.  
969 <https://doi.org/10.1139/A06-005>
- 970 Au, D. W. T. (2004). The application of histo-cytopathological biomarkers in marine pollution monitoring:  
971 A review. *Mar Pollut Bull*, 48(9–10), 817–834. <https://doi.org/10.1016/j.marpolbul.2004.02.032>
- 972 Austin, B., & Austin, D. A. (2016). Bacterial fish pathogens: Disease of farmed and wild fish, sixth edition.  
973 Bacterial Fish Pathogens: Disease of Farmed and Wild Fish, Sixth Edition.  
974 <https://doi.org/10.1007/978-3-319-32674-0>
- 975 Augustine, S., Gagnaire, B., Adam-Guillermin, C., & Kooijman, S. A. L. M. (2012). Effects of uranium on  
976 the metabolism of zebrafish, *Danio rerio*. *Aquat Toxicol*, 118-119 :9–26.  
977 <https://doi.org/10.1016/j.aquatox.2012.02.029>
- 978 Avella, M. A., Olivotto, I., Silvi, S., Ribecco, C., Cresci, A., Palermo, F., *et al.* (2011). Use of *Enterococcus*  
979 *faecium* to improve common sole (*Solea solea*) larviculture. *Aquaculture*, 315(3–4), 384–393.  
980 <https://doi.org/10.1016/j.aquaculture.2011.02.046>
- 981 Baas, J., Spurgeon, D., & Broerse, M. (2015). A simple mechanistic model to interpret the effects of  
982 narcotics. SAR and QSAR. *Environ Res*, 26(3), 165–180.  
983 <https://doi.org/10.1080/1062936X.2015.1018940>
- 984 Baynes, S. M., Howell, B. R., & Beard, T. W. (1993). A review of egg production by captive sole, *Solea*  
985 *solea* (L.). *Aquac Res*, 24(2), 171–180. <https://doi.org/10.1111/j.1365-2109.1993.tb00538.x>
- 986 Baynes, S.M., Howell, B.R., Beard, T.W., & Hallam, J.D. (1994). A description of spawning behaviour of  
987 captive dover sole, *Solea solea* (L.). *J Sea Res*, 32 (3–4), 271-275. [https://doi.org/10.1016/0077-](https://doi.org/10.1016/0077-7579(94)90004-3)  
988 [7579\(94\)90004-3.](https://doi.org/10.1016/0077-7579(94)90004-3)
- 989 Benzekri, H., Armesto, P., Cousin, X., Rovira, M., Crespo, D., Merlo, M. A., *et al.* (2014). De novo  
990 assembly, characterization and functional annotation of Senegalese sole (*Solea senegalensis*) and  
991 common sole (*Solea solea*) transcriptomes: Integration in a database and design of a microarray.  
992 *BMC Genom*, 15(1). <https://doi.org/10.1186/1471-2164-15-952>
- 993 Belanger, S. E., Balon, E. K., & Rawlings, J. M. (2010). Saltatory ontogeny of fishes and sensitive early  
994 life stages for ecotoxicology tests. *Aquat Toxicol*, 97(2), 88-95
- 995 Billoir, E., Péry, A. R. R., & Charles, S. (2007). Integrating the lethal and sublethal effects of toxic  
996 compounds into the population dynamics of *Daphnia magna* : A combination of the DEBtox and  
997 matrix population models. *Ecol Model*, 203(3-4) :204–214. [10.1016/j.ecolmodel.2006.11.021](https://doi.org/10.1016/j.ecolmodel.2006.11.021)
- 998 Bonaldo, A., Parma, L., Badiani, A., Serratore, P., & Gatta, P. P. (2011). Very early weaning of common  
999 sole (*Solea solea* L.) larvae by means of different feeding regimes and three commercial microdiets:  
1000 Influence on performances, metamorphosis development and tank hygiene. *Aquaculture*, 321(3–4),  
1001 237–244. <https://doi.org/10.1016/j.aquaculture.2011.09.007>

- 1002 Bonvini, E., Parma, L., Gatta, P. P., Mandrioli, L., Sirri, R., Martelli, G., et al. (2016). Effects of light  
1003 intensity on growth, feeding activity and development in common sole (*Solea solea* L.) larvae in  
1004 relation to sensory organ ontogeny. *Aquac Res*, 47(6), 1809–1819. <https://doi.org/10.1111/are.12639>
- 1005 Brown, C., 2006. *Marine and Coastal Ecosystems and Human Well-Being: Synthesis*. United Nations  
1006 Publications.
- 1007 Cabral, H., & Costa, M.J. (1999). Differential use of nursery areas within the Tagus estuary by sympatric  
1008 soles, *Solea solea* and *Solea senegalensis*. *Environ Biol Fishes* 56, 389–397.  
1009 <https://doi.org/10.1023/A:1007571523120>
- 1010 Cabral, H.N. (2000). Comparative feeding ecology of sympatric *Solea solea* and *S. senegalensis*, within the  
1011 nursery areas of the Tagus estuary, Portugal. *J Fish Biol*, 57, 1550–1562.  
1012 <https://doi.org/10.1006/jfbi.2000.1408>
- 1013 Cannas, M., Atzori, F., Rupsard, F., Bustamante, P., Loizeau, V., & Lefrançois, C. (2013). PCBs  
1014 contamination does not alter aerobic metabolism and tolerance to hypoxia of juvenile sole (*Solea*  
1015 *solea* L. 1758). *Aquat Toxicol*, 127, 54–60. <https://doi.org/10.1016/j.aquatox.2012.04.017>
- 1016 Caswell H., Nisbet R.M., de Roos A.M., Tuljapurkar S. (1997) Structured-Population Models: Many  
1017 Methods, a Few Basic Concepts. In: Tuljapurkar S., Caswell H. (eds) *Structured-Population*  
1018 *Models in Marine, Terrestrial, and Freshwater Systems*. Population and Community Biology  
1019 Series, vol 18. Springer, Boston, MA. [https://doi.org/10.1007/978-1-4615-5973-3\\_1](https://doi.org/10.1007/978-1-4615-5973-3_1)
- 1020 De Swaef, E., Demeestere, K., Boon, N., Van den Broeck, W., Haesebrouck, F., & Decostere, A. (2017).  
1021 Development of a reliable experimental set-up for Dover sole larvae *Solea solea* L. and exploring the  
1022 possibility of implementing this housing system in a gnotobiotic model. *Res Vet Sci*, 115(July), 418–  
1023 424. <https://doi.org/10.1016/j.rvsc.2017.07.025>
- 1024 De Swaef, E., Vercauteren, M., Duchateau, L., Haesebrouck, F., & Decostere, A. (2018). Experimental  
1025 infection model for vibriosis in Dover sole (*Solea solea*) larvae as an aid in studying its pathogenesis  
1026 and alternative treatments. *Vet Res*, 49(1), 1–9. <https://doi.org/10.1186/s13567-018-0520-3>
- 1027 De Swaef, E. (2019). *Towards sustainable dover sole (Solea solea) larviculture. Contributing to the*  
1028 *exploration of factors influencing its success*. Dissertation, Ghent University.
- 1029 Dinis, M. T., Ribeiro, L., Soares, F., & Sarasquete, C. (1999). A review on the cultivation potential of  
1030 *Solea senegalensis* in Spain and in Portugal. *Aquaculture*, 176(1–2), 27–38.  
1031 [https://doi.org/10.1016/S0044-8486\(99\)00047-2](https://doi.org/10.1016/S0044-8486(99)00047-2)
- 1032 Diopere, E., Vandamme, S. G., Hablützel, P. I., Cariani, A., Van Houdt, J., Rijnsdorp, A., et al. (2018).  
1033 Seascape genetics of a flatfish reveals local selection under high levels of gene flow. *ICES J Mar Sci*,  
1034 75(2), 675–689. <https://doi.org/10.1093/icesjms/fsx160>
- 1035 Duffy-Anderson J. T., Bailey, K. M., Cabral, H. N., Nakata, H. and van der Veer H. W. (2015). The  
1036 planktonic stages of flatfishes: physical and biological interactions in transport processes in Gibson,  
1037 R. N., Nash, R. D. M., Geffen, A. J., & Veer, H. W. van der (Eds.) *Flatfishes Biology and*  
1038 *Exploitation (Second edi)*. Wiley. Retrieved from <http://ir.obihiro.ac.jp/dspace/handle/10322/3933>
- 1039 FAO 2019 Cultured Aquatic Species Information Programme. *Solea solea*. Cultured Aquatic Species  
1040 Information Programme. Text by Colen, R., Ramalho, A., Rocha, F., and Dinis, M.T. In: FAO  
1041 Fisheries and Aquaculture Department [online]. Rome. Updated 18 February 2014. [Cited 21 June  
1042 2019].
- 1043 Ferrarresso, S., Bonaldo, A., Parma, L., Cinotti, S., Massi, P., Bargelloni, L., & Gatta, P. P. (2013).  
1044 Exploring the larval transcriptome of the common sole (*Solea solea* L.). *BMC Genom*, 14(1), 1–22.  
1045 <https://doi.org/10.1186/1471-2164-14-315>
- 1046 Ferrarresso, S., Bonaldo, A., Parma, L., Buonocore, F., Scapigliati, G., Gatta, P. P., & Bargelloni, L. (2016).  
1047 Ontogenetic onset of immune-relevant genes in the common sole (*Solea solea*). *Fish and Shellfish*  
1048 *Immun*, 57, 278–292. <https://doi.org/10.1016/j.fsi.2016.08.044>
- 1049 Fincham, J. I., Rijnsdorp, A. D., & Engelhard, G. H. (2013). Shifts in the timing of spawning in sole linked  
1050 to warming sea temperatures. *J Sea Res*, 75, 69–76. <https://doi.org/10.1016/j.seares.2012.07.004>

- 1051 Foekema, E. M., Deerenberg, C. M., & Murk, A. J. (2008). Prolonged ELS test with the marine flatfish  
 1052 sole (*Solea solea*) shows delayed toxic effects of previous exposure to PCB 126. *Aquat Toxicol*,  
 1053 90(3), 197–203. <https://doi.org/10.1016/j.aquatox.2008.08.015>
- 1054 Foekema, E. M., Fischer, A., Parron, M. L., Kwadijk, C., de Vries, P., & Murk, A. J. (2012). Toxic  
 1055 concentrations in fish early life stages peak at a critical moment. *Environ Toxicol Chem*, 31(6), 1381–  
 1056 1390. <https://doi.org/10.1002/etc.1836>
- 1057 Foekema, E. M., Lopez Parron, M., Mergia, M. T., Carolus, E. R. M., vd Berg, J. H. J., Kwadijk, C., et al.  
 1058 (2014). Internal effect concentrations of organic substances for early life development of egg-exposed  
 1059 fish. *Ecotox Environ Safe*, 101(1), 14–22. <https://doi.org/10.1016/j.ecoenv.2013.12.006>
- 1060 FranceAgriMer, <https://rnm.franceagrimer.fr/prix?SOLE&12MOIS>
- 1061 Freitas, V., Kooijman, S. A. L. M., & Van Der Veer, H. W. (2012). Latitudinal trends in habitat quality of  
 1062 shallow-water flatfish nurseries. *Mar Ecol Prog Ser*, 471, 203–214.  
 1063 <https://doi.org/10.3354/meps10025>
- 1064 Geffen, A. J., van der Veer, H. W., & Nash, R. D. M. (2007). The cost of metamorphosis in flatfishes. *J Sea*  
 1065 *Res*, 58(1), 35–45. <https://doi.org/10.1016/j.seares.2007.02.004>
- 1066 Geffen, A. J., Pittman K., & Imsland A. K. (2015) Synergies between aquaculture and fisheries in Gibson,  
 1067 R. N., Nash, R. D. M., Geffen, A. J., & Veer, H. W. van der (Eds.) *Flatfishes Biology and*  
 1068 *Exploitation* (Second edi). Wiley. Retrieved from <http://ir.obihiro.ac.jp/dspace/handle/10322/3933>
- 1069 Ghwila, M., & Willms, A. R. (2019). A physiologically-structured fish population model with size-  
 1070 dependent foraging. *Math Biosci*, 315, 108233
- 1071 Gibson R. N. (2015) Introduction in Gibson, R. N., Nash, R. D. M., Geffen, A. J., & Veer, H. W. van der  
 1072 (Eds.) *Flatfishes Biology and Exploitation* (Second edi). Wiley. Retrieved from  
 1073 <http://ir.obihiro.ac.jp/dspace/handle/10322/3933>
- 1074 Gilliers, C., Le Pape O., Morin J., Desaunay Y., & R. Amara, 2006. Are growth and density quantitative  
 1075 estimators of essential fish habitat quality? An application to the common sole *Solea solea* nursery  
 1076 grounds. *Estuar Coas Shelf S*, 69: 96. <https://doi.org/10.1016/j.ecss.2006.02.006>
- 1077 Gilliers, C., Claireaux, G., Galois, R., Loizeau, V., & Pape, O. Le. (2012). Influence of Hydrocarbons  
 1078 Exposure on Survival, Growth and Condition of Juvenile Flatfish: A Mesocosm Experiment. *J Lif*  
 1079 *Sci*, 4(2), 113–122. <https://doi.org/10.1080/09751270.2012.11885203>
- 1080 Guinand, B., Durieux, E. D., Dupuy, C., Cerqueira, F., & Bégout, M. L. (2011). Phenotypic and genetic  
 1081 differentiation in young-of-the-year common sole (*Solea solea*) at differentially contaminated nursery  
 1082 grounds. *Mar Environ Res*, 71(3), 195-206. <https://doi.org/10.1016/j.marenvres.2011.01.003>
- 1083 Gunnarsson, L., Jauhiainen, A., Kristiansson, E., Nerman, O., & Larsson, D. G. J. (2008). Evolutionary  
 1084 conservation of human drug targets in organisms used for environmental risk assessments. *Environ*  
 1085 *Sci Technol*, 42(15), 5807–5813. <https://doi.org/10.1021/es8005173>
- 1086 Hardman, R. (2006). A toxicologic review of quantum dots: Toxicity depends on physicochemical and  
 1087 environmental factors. *Environ Health Persp*, 114(2), 165–172. <https://doi.org/10.1289/ehp.8284>
- 1088 Hooper, M. J., Ankley, G. T., Cristol, D. A., Maryoung, L. A., Noyes, P. D., & Pinkerton, K. E. (2013).  
 1089 Interactions between chemical and climate stressors: A role for mechanistic toxicology in assessing  
 1090 climate change risks. *Environ Toxicol Chem*, 32(1), 32–48. <https://doi.org/10.1002/etc.2043>
- 1091 Howell, B. R., & Canario, A. V. M. (1987). The influence of sand on the estimation of resting metabolic  
 1092 rate of juvenile sole, *Solea solea* (L.). *J Fish Biol*, 31(2), 277-280. <https://doi.org/10.1111/j.1095-8649.1987.tb05231.x>
- 1094 Howell, B. R. (1997). A re-appraisal of the potential of the sole, *Solea solea* (L.), for commercial  
 1095 cultivation. *Aquaculture*, 155(1–4), 355–365. [https://doi.org/10.1016/S0044-8486\(97\)00103-8](https://doi.org/10.1016/S0044-8486(97)00103-8)
- 1096 Hufnagl, M., Payne, M., Lacroix, G., Bolle, L. J., Daewel, U., Dickey-Collas, M., et al. (2017). Variation  
 1097 that can be expected when using particle tracking models in connectivity studies. *J Sea Res*, 127, 133–  
 1098 149. <https://doi.org/10.1016/j.seares.2017.04.009>



- 1099 Huret M., Vandromme P., Petitgas P., Pecquerie L. (2012). Connectivity patterns of anchovy larvae in the  
 1100 Bay of Biscay from a coupled transport-bioenergetic model forced by size-structured zooplankton.  
 1101 ICES CM 2012/E:14.
- 1102 Hutchinson, T. H., Solbé, J., & Kloepper-Sams, P. J. (1998). Analysis of the ECETOC Aquatic Toxicity  
 1103 (EAT) database. III - Comparative toxicity of chemical substances to different life stages of aquatic  
 1104 organisms. *Chemosphere*, 36(1), 129–142. [https://doi.org/10.1016/S0045-6535\(97\)10025-X](https://doi.org/10.1016/S0045-6535(97)10025-X)
- 1105 Ibarra-Zatarain, Z., Martín, I., Rasines, I., Fatsini, E., Rey, S., Chereguini, O., & Duncan, N. (2020).  
 1106 Exploring the relationship between stress coping styles and sex, origin and reproductive success, in  
 1107 Senegalese sole (*Solea senegalensis*) breeders in captivity. *Physiol Behav*, 220, 112868.  
 1108 <https://doi.org/https://doi.org/10.1016/j.physbeh.2020.112868>
- 1109 Imsland, A. K., Foss, A., Conceição, L. E. C., Dinis, M. T., Delbare, D., Schram, E., et al. (2003). A review  
 1110 of the culture potential of *Solea solea* and *S. senegalensis*. *Rev Fish Biol Fisher*, 13(4), 379–408.  
 1111 <https://doi.org/10.1007/s11160-004-1632-6>
- 1112 Jager, T., Crommentuijn, T., van Gestel, C. A. M., & Kooijman, S. A. L. M. (2004). Simultaneous  
 1113 Modeling of Multiple End Points in Life-Cycle Toxicity Tests. *Environ Sci Technol*, 38(10) :2894–  
 1114 2900. <https://doi.org/10.1021/es0352348>
- 1115 Jager, T., & Ashauer, R. (2018a). Modelling survival under chemical stress to the GUTS framework.  
 1116 Leanpub. Retrieved from [https://leanpub.com/guts\\_book](https://leanpub.com/guts_book)
- 1117 Jager, T., & Ashauer, R. (2018b). How to Evaluate the Quality of Toxicokinetic—Toxicodynamic Models  
 1118 in the Context of Environmental Risk Assessment. *Integr Environ Assess*, 14(5), 604–614.  
 1119 <https://doi.org/10.1002/ieam.2026>
- 1120 Jager, T., & Zimmer, E. I. (2012). Simplified Dynamic Energy Budget model for analysing ecotoxicity  
 1121 data. *Ecol Model*, 225, 74–81. <https://doi.org/10.1016/j.ecolmodel.2011.11.012>
- 1122 Kooijman, S. A. L. M. (2010). Dynamic Energy Budget theory for metabolic organisation. Water (Third  
 1123 edit, Vol. 365). New York: Cambridge University Press. <https://doi.org/10.1098/rstb.2010.0167>
- 1124 Kooijman, S. A. L. M., & Bedaux, J. J. M. (1996). Analysis of toxicity tests on Daphnia survival and  
 1125 reproduction. *Water Res*, 30(7), 1711–1723. [https://doi.org/10.1016/0043-1354\(96\)00054-1](https://doi.org/10.1016/0043-1354(96)00054-1)
- 1126 Lacroix, G., Maes, G. E., Bolle, L. J., & Volckaert, F. A. M. (2013). Modelling dispersal dynamics of the  
 1127 early life stages of a marine flatfish (*Solea solea* L.). *J Sea Res*, 84, 13–25.  
 1128 <https://doi.org/10.1016/j.seares.2012.07.010>
- 1129 Lacroix, G., Barbut, L., & Volckaert, F. A. M. (2016). Complex effect of projected sea temperature and  
 1130 wind change on flatfish dispersal. *Global Change Biol*, 24(1), 85–100.  
 1131 <https://doi.org/10.1111/gcb.13915>
- 1132 Lagardère, F., Amara, R., & Joassard, L. (1999). Vertical distribution and feeding activity of  
 1133 metamorphosing sole, *Solea solea*, before immigration to the Bay of Vilaine nursery (northern Bay of  
 1134 Biscay, France). *Environ Biol Fish*, 56, 213–228. <https://doi.org/10.1023/A:1007581818941>
- 1135 Lund, I., Steinfeldt, S. J., Banta, G., & Hansen, B. W. (2008). The influence of dietary concentrations of  
 1136 arachidonic acid and eicosapentaenoic acid at various stages of larval ontogeny on eye migration,  
 1137 pigmentation and prostaglandin content of common sole larvae (*Solea solea* L.). *Aquaculture*, 276(1–  
 1138 4), 143–153. <https://doi.org/10.1016/j.aquaculture.2008.01.004>
- 1139 Marras, S., Cucco, A., Antognarelli, F., Azzurro, E., Milazzo, M., Bariche, M., et al. (2015). Predicting  
 1140 future thermal habitat suitability of competing native and invasive fish species: From metabolic scope  
 1141 to oceanographic modelling. *Conserv Physiol*, 3(1), 1–14. <https://doi.org/10.1093/conphys/cou059>
- 1142 Martínez-Álvarez, R. M., Morales, A. E., & Sanz, A. (2005). Antioxidant defenses in fish: Biotic and  
 1143 abiotic factors. *Rev Fish Biol Fisher*, 15(1–2), 75–88. <https://doi.org/10.1007/s11160-005-7846-4>
- 1144 Martinho, F., Leitão, R., Neto, J.M., Cabral, H., Lagardère, F., & Pardal, M.A. (2008). Estuarine  
 1145 colonization, population structure and nursery functioning for 0-group sea bass (*Dicentrarchus*  
 1146 *labrax*), flounder (*Platichthys flesus*) and sole (*Solea solea*) in a mesotidal temperate estuary. *J Appl*  
 1147 *Ichthyol*, 24, 229–237. <https://doi.org/10.1111/j.1439-0426.2007.01049.x>

- 1148 Mollet, F. M., Kraak, S. B. M., & Rijnsdorp, A. D. (2007). Fisheries-induced evolutionary changes in  
 1149 maturation reaction norms in North Sea sole *Solea solea*. *Mar Ecol Prog Ser*, 351, 189–199.  
 1150 <https://doi.org/10.3354/meps07138>
- 1151 Morais, S., Aragão, C., Cabrita, E., Conceição, L. E. C., Constenla, M., Costas, B., et al. (2014). New  
 1152 developments and biological insights into the farming of *Solea senegalensis* reinforcing its  
 1153 aquaculture potential. *Reviews in Aquaculture*, 8(3), 227–263. <https://doi.org/10.1111/raq.12091>
- 1154 Mounier, F., Pecquerie, L., Lobry, J., Sardi, A. E., Labadie, P., Budzinski, H., & Loizeau, V. (2020a).  
 1155 Dietary bioaccumulation of persistent organic pollutants in the common sole *Solea solea* in the context  
 1156 of global change. Part 1: Revisiting parameterisation and calibration of a DEB model to consider inter-  
 1157 individual variability in experimental and natural conditions. *Ecol Model*, 433.  
 1158 <https://doi.org/10.1016/j.ecolmodel.2020.109224>
- 1159 Mounier, F., Loizeau, V., Pecquerie, L., Drouineau, H., Labadie, P., Budzinski, H., & Lobry, J. (2020b).  
 1160 Dietary bioaccumulation of persistent organic pollutants in the common sole *Solea solea* in the context  
 1161 of global change. Part 2: Sensitivity of juvenile growth and contamination to toxicokinetic parameters  
 1162 uncertainty and environmental conditions variability in estuaries. *Ecol Model*, 431.  
 1163 <https://doi.org/10.1016/j.ecolmodel.2020.109196>
- 1164 Munschy, C., Héas-Moisan, K., Tixier, C., Pacepavicius, G., & Alaei, M. (2010). Dietary exposure of  
 1165 juvenile common sole (*Solea solea* L.) to polybrominated diphenyl ethers (PBDEs): Part 2. Formation,  
 1166 bioaccumulation and elimination of hydroxylated metabolites. *Environ Pollut*, 158(12), 3527–3533.  
 1167 <https://doi.org/10.1016/j.envpol.2010.08.021>
- 1168 Munschy, C., Héas-moisan, K., Tixier, C., Olivier, N., Gastineau, O., Bayon, N. Le, & Buchet, V. (2011).  
 1169 Dietary exposure of juvenile common sole (*Solea solea* L.) to polybrominated diphenyl ethers  
 1170 (PBDEs): Part 1. Bioaccumulation and elimination kinetics of individual congeners and their  
 1171 debrominated metabolites. *Environ Pollut*, 159(1), 229–237.  
 1172 <https://doi.org/10.1016/j.envpol.2010.09.001>
- 1173 Munoz-Cueto, J. A., Mañanós-Sánchez, E., & Sánchez-Vázquez, F. J. (Eds.). (2019). *The Biology of Sole*.  
 1174 CRC Press.
- 1175 Novriadi, R. (2016). Vibriosis in aquaculture. *Omni-Akuatika*, 12(1), 1–12.  
 1176 <https://doi.org/10.20884/1.oa.2016.12.1.24>
- 1177 Pankhurst, N. W., & Munday, P. L. (2011). Effects of climate change on fish reproduction and early life  
 1178 history stages, *Mar Freshwater Res*, 1015–1026. <https://doi.org/10.1071/MF10269>
- 1179 Parma, L., Bonaldo, A., Massi, P., Yúfera, M., Martínez-Rodríguez, G., & Gatta, P. P. (2013). Different  
 1180 early weaning protocols in common sole (*Solea solea* L.) larvae: Implications on the performances  
 1181 and molecular ontogeny of digestive enzyme precursors. *Aquaculture*, 414–415, 26–35.  
 1182 <https://doi.org/10.1016/j.aquaculture.2013.07.043>
- 1183 Parma, L., Bonaldo, A., Pirini, M., Viroli, C., Parmeggiani, A., Bonvini, E., & Gatta, P. P. (2015). Fatty  
 1184 Acid Composition of Eggs and its Relationships to Egg and Larval Viability from Domesticated  
 1185 Common Sole (*Solea solea*) Breeders. *Reprod Domest Anim*, 50(2), 186–194.  
 1186 <https://doi.org/10.1111/rda.12466>
- 1187 Péan, S. (2012). Effets des polluants organiques persistants sur le comportement des poissons. Dissertation,  
 1188 Université de La Rochelle
- 1189 Perkins, E. J., Ashauer, R., Burgoon, L., Conolly, R., Landesmann, B., Mackay, C., et al. (2019). Building  
 1190 and Applying Quantitative Adverse Outcome Pathway Models for Chemical Hazard and Risk  
 1191 Assessment. *Environ Toxicol Chem*, 38(9), 1850–1865. <https://doi.org/10.1002/etc.4505>
- 1192 Piccinetti, C. C., Ricci, L. A., Tokle, N., Radaelli, G., Pascoli, F., Cossignani, L., et al. (2012). Malnutrition  
 1193 may affect common sole (*Solea solea* L.) growth, pigmentation and stress response: Molecular,  
 1194 biochemical and histological implications. *Comp Biochem Physiol A*, 161(4), 361–371.  
 1195 <https://doi.org/10.1016/j.cbpa.2011.12.009>
- 1196 Post, M. H. M., Blom, E., Chen, C., Bolle, L. J., & Baptist, M. J. (2017). Habitat selection of juvenile sole  
 1197 (*Solea solea* L.): Consequences for shoreface nourishment. *J Sea Res*, 122, 19–24.  
 1198 <https://doi.org/10.1016/j.seares.2017.02.011>

- 1199 Ramsay, K., & Witthames, P. (1996). Using oocyte size to assess seasonal ovarian development in *Solea*  
1200 *solea* (L.). *J Sea Res*, 36(3–4), 275–283. [https://doi.org/10.1016/S1385-1101\(96\)90796-0](https://doi.org/10.1016/S1385-1101(96)90796-0)
- 1201 Rand-Weaver, M., Margiotta-Casaluci, L., Patel, A., Panter, G. H., Owen, S. F., & Sumpter, J. P. (2013).  
1202 The read-across hypothesis and environmental risk assessment of pharmaceuticals. *Environ Sci*  
1203 *Technol*, 47(20), 11384–11395. <https://doi.org/10.1021/es402065a>
- 1204 Richard, P., Bergeron, J.-P., Boulhic, M., Galois, R., & Person-Le Ruyet, J. (1991). Effect of starvation on  
1205 RNA, DNA and protein content of laboratory-reared larvae and juveniles of *Solea solea*. *Mar Ecol*  
1206 *Prog Ser*, 72, 69–77. <https://doi.org/10.3354/meps072069>
- 1207 Ridgway, J. & G. Shimmield. (2002). Estuaries as Repositories of Historical Contamination and their Impact  
1208 on Shelf Seas. *Estuar Coast Shelf S.* 55, 903. <https://doi.org/10.1006/ecss.2002.1035>
- 1209 Ringø, E., Olsen, R. E., Jensen, I., Romero, J., & Lauzon, H. L. (2014). Application of vaccines and dietary  
1210 supplements in aquaculture: possibilities and challenges. *Rev Fish Biol Fisher*, 24(4), 1005–1032.  
1211 <https://doi.org/10.1007/s11160-014-9361-y>
- 1212 Rochette, S., O. Le Pape, J. Vigneau & E. Rivot (2013). A hierarchical Bayesian model for embedding  
1213 larval drift and habitat models in integrated life cycles for exploited fish. *Ecol Appl*, 23(7), 1659-1676.  
1214 <https://doi.org/10.1890/12-0336.1>
- 1215 Sadoul, B., Augustine, S., Zimmer, E., Bégout, M. L., & Vijayan, M. M. (2019). Prediction of long-term  
1216 variation in offspring metabolism due to BPA in eggs in rainbow trout using the DEB model. *J Sea*  
1217 *Res*, 143, 222-230.
- 1218 Sardi, A. E., Augustine, S., Olsen, G. H., & Camus, L. (2019). Exploring inter-species sensitivity to a  
1219 model hydrocarbon, 2-Methylnaphtalene, using a process-based model. *Environ Sci Pollut Res*,  
1220 26(11), 11355–11370. <https://doi.org/10.1007/s11356-019-04423-8>
- 1221 Savina, M., Lunghi, M., Archambault, B., Baulier, L., Huret, M., & Le Pape, O. (2016). Sole larval supply  
1222 to coastal nurseries: Interannual variability and connectivity at interregional and interpopulation  
1223 scales. *J Sea Res*, 111, 1–10. <https://doi.org/10.1016/j.seares.2015.11.010>
- 1224 Savina, M., Lacroix, G. & Ruddick, K. (2010) Modelling the transport of common sole larvae in the southern  
1225 North Sea: influence of hydrodynamics and larval vertical movements. *J Mar Syst*, 81, 86–98.  
1226 <https://doi.org/10.1016/j.jmarsys.2009.12.008>
- 1227 Seitz, R. D., Wennhage, H., Bergstro, U., Lipcius, R. N., & Ysebaert, T. (2014). Ecological value of coastal  
1228 habitats for commercially and ecologically important species. *ICES J Ma Sci*, 71(3), 648–665.  
1229 <https://doi.org/10.1093/icesjms/fst152>
- 1230 Solé (2019). *Ecotoxicology in Munoz-Cueto, J. (Ed.), Mañanós-Sánchez, E. (Ed.) & Sánchez-Vázquez, F.*  
1231 *(Ed.). The Biology of Sole, first edition. CRC Press, Boca Raton, 302-318.*  
1232 <https://doi.org/10.1201/9781315120393>
- 1233 Solé, M., Manzanera, M., Bartolomé, A., Tort, L., & Caixach, J. (2013). Persistent organic pollutants (POPs)  
1234 in sediments from fishing grounds in the NW Mediterranean: Ecotoxicological implications for the  
1235 benthic fish *Solea sp.* *Mar Pollut Bull*, 67(1–2), 158–165.  
1236 <https://doi.org/10.1016/j.marpolbul.2012.11.018>
- 1237 Teal, L. R., De Leeuw, J. J., Van Der Veer, H. W., & Rijnsdorp, A. D. (2008). Effects of climate change on  
1238 growth of 0-group sole and plaice. *Marine Ecology Progress Series*, 358, 219–230.  
1239 <https://doi.org/10.3354/meps07367>
- 1240 Teal, L. R., van Hal, R., van Kooten, T., Ruardij, P., & Rijnsdorp, A. D. (2012). Bio-energetics underpins  
1241 the spatial response of North Sea plaice (*Pleuronectes platessa* L.) and sole (*Solea solea* L.) to climate  
1242 change. *Global Change Biology*, 18(11), 3291–3305. <https://doi.org/10.1111/j.1365-2486.2012.02795.x>
- 1244 The Research Council of Norway (2009). The fish larva: a transitional life form, the foundation for  
1245 aquaculture and fisheries. The Research Council of Norway. Oslo.
- 1246 Torrecillas S, Montero D, Izquierdo M (2014) Improved health and growth of fish fed  
1247 mannanoligosaccharides: potential mode of action. *Fish Shellfish Immunol* 36:525–544.  
1248 <https://doi.org/10.1016/j.fsi.2013.12.029>

- 1249 van der Oost, R., Beyer, J., & Vermeulen, N. P. E. (2003). Fish bioaccumulation and biomarkers in  
1250 environmental risk assessment: a review. *Environ Toxicol Phar*, 13(2), 57–149.  
1251 [https://doi.org/10.1016/S1382-6689\(02\)00126-6](https://doi.org/10.1016/S1382-6689(02)00126-6)
- 1252 Van Der Veer, H. W., Berghahn, R., Miller, J. M., & Rijnsdorp, A. D. (2000). Recruitment in flatfish, with  
1253 special emphasis on North Atlantic species: Progress made by the Flatfish Symposia. *ICES J Mar Sci*,  
1254 57(2), 202–215. <https://doi.org/10.1006/jmsc.1999.0523>
- 1255 Vasconcelos, R. P., Eggleston, D. B., Pape, O. Le, & Tulp, I. (2014). Patterns and processes of habitat-  
1256 specific demographic variability in Exploited Marine Species, *ICES J Mar Sci* 71(3), 638–647.  
1257 <https://doi.org/10.1093/icesjms/fst136>
- 1258 Vinagre, C., França, S., Costa, M.J., & Cabral, H.N. (2005). Niche overlap between juvenile flatfishes,  
1259 *Platichthys flesus* and *Solea solea*, in a southern European estuary and adjacent coastal waters. *J.*  
1260 *Appl. Ichthyol.* 21, 114–120. <https://doi.org/10.1111/j.1439-0426.2004.00639.x>
- 1261 Vinagre, C., Fonseca, V., Cabral, H., & Costa, M.J. (2006). Habitat suitability index models for the  
1262 juvenile soles, *Solea solea* and *Solea senegalensis*, in the Tagus estuary: Defining variables for  
1263 species management. *Fish Res* 82, 140–149. <https://doi.org/10.1016/j.fishres.2006.07.011>
- 1264 Vinagre, C., Costa, M. J., & Cabral, H. N. (2007). Impact of climate and hydrodynamics on sole larval  
1265 immigration towards the Tagus estuary, Portugal. *Estuarine, Coastal and Shelf Science*, 75(4), 516–  
1266 524. <https://doi.org/10.1016/j.ecss.2007.05.035>
- 1267 Vinagre, C., & Cabral, H.N. (2008). Prey consumption by the juvenile soles, *Solea solea* and *Solea*  
1268 *senegalensis*, in the Tagus estuary, Portugal. *Estuar Coast Shelf Sci*, 78, 45–50.  
1269 <https://doi.org/10.1016/j.ecss.2007.11.009>
- 1270 Vinagre, C., Fonseca, V., Maia, A., Amara, R. & Cabral H. (2008). Habitat specific growth rates and  
1271 condition indices for the sympatric soles *Solea solea* (Linnaeus, 1758) and *Solea senegalensis* Kaup  
1272 1858, in the Tagus estuary, Portugal, based on otolith daily increments and RNA-DNA ratio. *J Appl*  
1273 *Ichthyol*, 24(2), 163-169. <https://doi.org/10.1111/j.1439-0426.2007.01048.x>
- 1274 Vinagre, C., Maia, A., Reis-Santos, P., Costa, M. J. and H. N. Cabral (2009). Small-scale distribution of  
1275 *Solea solea* and *Solea senegalensis* juveniles in the Tagus estuary (Portugal). *Estuar Coast Shelf Sci*,  
1276 81, 296-300. <https://doi.org/10.1016/j.ecss.2008.11.008>
- 1277 Vine, N. G., Leukes, W. D., & Kaiser, H. (2006). Probiotics in marine larviculture. *FEMS Microbiol Rev*,  
1278 30(3), 404–427. <https://doi.org/10.1111/j.1574-6976.2006.00017.x>
- 1279 Wessel, N., Santos, R., Menard, D., Le Menach, K., Buchet, V., Lebayon, N., *et al.* (2010). Relationship  
1280 between PAH biotransformation as measured by biliary metabolites and EROD activity, and  
1281 genotoxicity in juveniles of sole (*Solea solea*). *Mar Environ Res*, 69(SUPPL. 1), S71–S73.  
1282 <https://doi.org/10.1016/j.marenvres.2010.03.004>
- 1283 Wiltshire, K.H., Malzahn, A.M., Wirtz, K., Greve, W., Janisch, S., Mangelsdorf, P., *et al.* (2008).  
1284 Resilience of North Sea phytoplankton spring bloom dynamics: an analysis of long-term data at  
1285 Helgoland Roads. *Limnol Oceanogr* 53, 1294–1302. <https://doi.org/10.4319/lo.2008.53.4.1294>
- 1286 Zambonino-Infante, J. L., Claireaux, G., Ernande, B., Jolivet, A., Quazuguel, P., Severe, A., *et al.* (2013).  
1287 Hypoxia tolerance of common sole juveniles depends on dietary regime and temperature at the larval  
1288 stage: evidence for environmental conditioning. *P Roy Soc Lond B Bio*, 280(1758), 1–9.  
1289 <https://doi.org/10.1098/rspb.2012.3022>