

Supplementary Table S1

Table I. Growth parameters used for the calculation of the residual body condition index (Gulf of Lions area, France).

Species	Current study		Other studies		References
	<i>a</i>	<i>b</i>	<i>a</i>	<i>b</i>	
Spotted flounder	0,001	3,43	0,003	3,3	Merella <i>et al.</i> (1997)
Horned octopus	0,012	2,19		2,61	Moriyasu (1983)
Grey gurnard	0,007	3,04	0,0029	3,49	Merella <i>et al.</i> (1997)
Blackbelly rosefish	0,053	2,80	0,013	3,04	Merella <i>et al.</i> (1997)
Shortfin squid	0,000	2,50	0,019	3,2	Merella <i>et al.</i> (1997)
Four-spot megrim	0,004	3,13	0,003	3,05	Crec'hriou <i>et al.</i> (2013)
Blackbellied angler	0,026	2,90	0,022	2,86	Garcia-Rodriguez <i>et al.</i> (2005)
Angler	0,044	2,83	0,0036	3,43	Merella <i>et al.</i> (1997)
European hake	0,004	3,14	0,00596	3,04	Mellon-Duval <i>et al.</i> (2010)
Blue whiting	0,004	3,13	0,0007	3,69	Merella <i>et al.</i> (1997)
Red mullet	0,012	3,00	0,00713	3,08	Bensebaini <i>et al.</i> (2019)
Surmullet	0,014	2,99	0,0157	2,95	Bensebaini <i>et al.</i> (2019)
Norway lobster	0,470	3,12	0,613	3,04	Sardá <i>et al.</i> (1998)
Common octopus	0,088	2,01		2,98	Guerra (1979)
Axillary seabream	0,020	2,97	0,012	3,04	Crec'hriou <i>et al.</i> (2013)
Blackspot seabream	0,009	3,09	0,008	3,17	Campillo (1992)
Common pandora	0,033	2,85	0,016	2,905	Busalacchi <i>et al.</i> (2014)
Greater forkbeard	0,012	2,92	0,0026	3,27	Merella <i>et al.</i> (1997)
Mediterranean horse mackerel	0,015	2,90	0,008	2,98	Crec'hriou <i>et al.</i> (2013)
Atlantic horse mackerel	0,011	2,96	0,0069	3,065	Aksissou <i>et al.</i> (2013)
Capelan	0,011	3,00	0,0056	3,2029	Ragonese <i>et al.</i> (1998)
John dory	0,024	2,92	0,013	3,02	Crec'hriou <i>et al.</i> (2013)

References

- Aksissou, M., Hernando Casal, J.A., and Kerkich, M. 2013. Age and growth of the horse mackerel *Trachurus trachurus* (Linnaeus,1758) catches in the bay of M'diq (Mediterranean coast of Morocco). IRACST – Engineering Science and Technology: An International Journal (ESTIJ), Vol.3.
- Bensebaini, C. M., Billet, N., Certain, G., Jadaud, A., and Le Roy, E. 2019. Poissons, céphalopodes et crustacés du Golfe du Lion. Courbes de croissance. Taille et poids. Maturité sexuelle. <https://archimer.ifremer.fr/doc/00505/61688/> (Accessed 1 July 2021).
- Busalacchi, B., Bottari, T., Giordano, D., Profeta, A., and Rinelli, P. 2014. Distribution and biological features of the common pandora, *Pagellus erythrinus* (Linnaeus, 1758), in the southern Tyrrhenian Sea (Central Mediterranean). Helgoland Marine Research, 68: 491–501.
- Campillo, A. 1992. Les pêcheries françaises de Méditerranée. Synthèse des connaissances. IFREMER. <https://archimer.ifremer.fr/doc/00393/50467/> (Accessed 26 May 2020).
- Crec'hriou, R., Neveu, R., and Lenfant, P. 2013. Length-weight relationship of main commercial fishes from the French Catalan coast. Journal of Applied Ichthyology, 29: 1191–1192.

- García-Rodríguez, M., Pereda, P., Landa, J., and Esteban, A. 2005. On the biology and growth of the anglerfish *Lophius budegassa* Spinola, 1807 in the Spanish Mediterranean: A preliminary approach. *Fisheries Research*, 71: 197–208.
- Guerra, A. 1979. Fitting a von Bertalanffy expression to *Octopus vulgaris* growth. *Investigación Pesquera*, 43: 319–326.
- Mellon-Duval, C., De Pontual, H., Metral, L., and Quemener, L. 2010. Growth of European hake (*Merluccius merluccius*) in the Gulf of Lions based on conventional tagging. *Ices Journal Of Marine Science*, 67: 62–70.
- Merella, P., Quetglas, A., Alemany, F., and Carbonell, A. 1997. Length-weight relationship of fishes and cephalopods from the Balearic Islands (Western Mediterranean). *Naga, the ICLARM Quarterly*, 20: 66–68.
- Moriyasu, M. 1983. Étude biométrique de la croissance d'*Eledone cirrhosa* [LAM. 1978 (Cephalopoda, Octopoda)] du golfe du Lion. *Oceanologica Acta*, 6: 35–41.
- Ragonese, S., and Bianchini, M. L. 1998. Growth, mortality and yield-per-recruit of the poor cod, *Trisopterus minutus capelanus*, from the Strait of Sicily. *Naga, the ICLARM Quarterly*, 21: 61–70.
- Sardá, F., Lleonart, J., and Cartes, J. E. 1998. An analysis of the population dynamics of *Nephrops norvegicus* (L.) in the Mediterranean Sea. *Scientia Marina*, 62: 135–143.