



Continental-scale geographic change across Zealandia during Paleogene subduction initiation

R. Sutherland¹, G.R. Dickens², P. Blum³, C. Agnini⁴, L. Alegret⁵, G. Asatryan⁶, J. Bhattacharya², A. Bordenave⁷, L. Chang⁸, J. Collot⁷, M.J. Cramwinckel⁹, E. Dallanave¹⁰, M.K. Drake¹¹, S.J.G. Etienne⁷, M. Giorgioni¹², M. Gurnis¹³, D.T. Harper¹¹, H.-H.M. Huang¹⁴, A.L. Keller¹⁵, A.R. Lam¹⁶, H. Li¹⁷, H. Matsui¹⁸, H.E.G. Morgans¹⁹, C. Newsam²⁰, Y.-H. Park²¹, K.M. Pascher¹⁹, S.F. Pekar²², D.E. Penman²³, S. Saito²⁴, W.R. Stratford¹⁹, T. Westerhold²⁵ and X. Zhou²⁶

¹SGEES, Victoria University of Wellington, P.O. Box 600, Wellington 6140, New Zealand

²Earth, Environmental and Planetary Sciences, Rice University, Houston, Texas 77005, USA

³International Ocean Discovery Program, Texas A&M University, College Station, Texas 77845-9547, USA

⁴Dipartimento di Geoscienze, Università di Padova, 35131 Padova, Italy

⁵Departamento de Ciencias de la Tierra & IUCA, Universidad de Zaragoza, 50009 Zaragoza, Spain

⁶Museum für Naturkunde, Leibniz-Institut für Evolutions und Biodiversitätsforschung, 10115 Berlin, Germany

⁷Geological Survey of New Caledonia, Noumea BP 465, New Caledonia

⁸School of Earth and Space Sciences, Peking University, Beijing, China

⁹Department of Earth Sciences, Utrecht University, 3584 CB Utrecht, The Netherlands

¹⁰Faculty of Geosciences, Universität Bremen, 28359 Bremen, Germany

¹¹Ocean Sciences Department, University of California, Santa Cruz, California 95064, USA

¹²Instituto de Geociência, Universidade de Brasília, Brasília, Brazil

¹³Seismological Laboratory, California Institute of Technology, Pasadena, California 91125, USA

¹⁴Atmosphere and Ocean Research Institute, University of Tokyo, Tokyo 113-8654, Japan

¹⁵Department of Earth Sciences, University of California, Riverside, California 92521, USA

¹⁶Department of Geosciences, University of Massachusetts, Amherst, Massachusetts 01003-9297, USA

¹⁷Institute of Oceanology, Chinese Academy of Sciences, Qingdao 266071, China

¹⁸Department of Earth Science, Tohoku University, Sendai 980-8572, Japan

¹⁹GNS Science, P.O. Box 30368, Lower Hutt 5040, New Zealand

²⁰Department of Earth Sciences, University College London, London WC1E 6BT, UK

²¹Department of Oceanography, Pusan National University, Busan 46421, Republic of Korea

²²School of Earth and Environmental Sciences, Queens College (CUNY), Flushing, New York 11451, USA

²³Department of Geology and Geophysics, Yale University, New Haven, Connecticut 06511, USA

²⁴Research and Development Center for Ocean Drilling Science, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, 236-0001, Japan

²⁵Center for Marine Environmental Sciences (MARUM), University of Bremen, 28359 Bremen, Germany

²⁶Institute of Marine and Coastal Sciences, Rutgers University, Rutgers, New Jersey 08854, USA

ABSTRACT

Data from International Ocean Discovery Program (IODP) Expedition 371 reveal vertical movements of 1–3 km in northern Zealandia during early Cenozoic subduction initiation in the western Pacific Ocean. Lord Howe Rise rose from deep (~1 km) water to sea level and subsided back, with peak uplift at 50 Ma in the north and between 41 and 32 Ma in the south. The New Caledonia Trough subsided 2–3 km between 55 and 45 Ma. We suggest these elevation changes resulted from crust delamination and mantle flow that led to slab formation. We propose a “subduction resurrection” model in which (1) a subduction rupture event activated lithospheric-scale faults across a broad region during less than ~5 m.y., and (2) tectonic forces evolved over a further 4–8 m.y. as subducted slabs grew in size and drove plate-motion change. Such a subduction rupture event may have involved nucleation and lateral propagation of slip-weakening rupture along an interconnected set of preexisting weaknesses adjacent to density anomalies.

INTRODUCTION

Major global plate-motion change occurred between 52 and 43 Ma, as manifested by the Emperor-Hawaii bend (Steinberger et al., 2004; O'Connor et al., 2013), reorientation of mid-ocean ridges (Muller et al., 2000; Steinberger et al., 2004; Cande et al., 2010), and rifting of Antarctica (Cande et al., 2000). This coincided with subduction initiation (Fig. 1) in the Izu-Bonin-Mariana (IBM) system (Arculus et al., 2015; Reagan et al., 2017), and nascent collision of the Indian and Asian plates (Aitchison et al., 2007). Development of western Pacific




Figure 1. (A) Global continents on a shaded elevation model (ETOPO1, <https://www.ngdc.noaa.gov/mgg/global/>), Izu-Bonin-Mariana margin (IBM), Papua New Guinea (PNG), and “Pacific Ring of Fire” (red dots). Arrows show plate movement during formation of Emperor-Hawaii seamounts. **(B)** Hypsometric profiles of continents (area in legend, $\times 10^6 \text{ km}^2$). Ice (dashed) and rock (solid) surfaces are shown for Antarctica and North America (includes Greenland). **(C)** Crustal thickness (CRUST1.0, <https://igppweb.ucsd.edu/~gabi/crust1.html>) versus surface elevation. Ellipses show one standard deviation for each continent (same legend as in B).

slab pull explains the sense of plate-motion changes (Gurnis et al., 2004).

Zealandia, a distinct but mostly submerged continent (Mortimer et al., 2017), has a low median elevation (Fig. 1), which is primarily an isostatic response to its relatively thin crust ($\sim 18 \text{ km}$ on average). Between 83 Ma and 79 Ma, Zealandia separated from Gondwana (Gaina et al., 1998; Sutherland, 1999), and much of the continent has been below sea level since, as documented by now-uplifted marine stratigraphic records in New Zealand and New Caledonia (Paris, 1981; Laird and Bradshaw, 2004)

and submarine sections recovered by the Deep Sea Drilling Program (DSDP; Fig. 2; Burns and Andrews, 1973).

However, the topographic history of Zealandia is not straightforward. Seismic reflection surveys and geological mapping reveal widespread Eocene deformation in northern Zealandia (Bache et al., 2012; Browne et al., 2016). This has been coined the “Tectonic Event of the Cenozoic in the Tasman Area” (TECTA; Sutherland et al., 2017), and it appears to have begun about the same time as cessation of spreading in the Tasman Sea (Gaina et al., 1998) and sub-

duction initiation near or east of Norfolk Ridge (Fig. 2; Gurnis et al., 2004; Cluzel et al., 2006; Sutherland et al., 2010; Bache et al., 2012; Matthews et al., 2015).

International Ocean Discovery Program (IODP) Expedition 371 (Sutherland et al., 2019) was designed to determine the Cenozoic paleogeography of northern Zealandia, and how and why this large region ($\sim 3 \times 10^6 \text{ km}^2$) evolved over time. We discuss the evidence collected and reasons for topographic change, and we propose a new framework for understanding subduction initiation.

DRILLING RESULTS

Before IODP Expedition 371, only three boreholes, DSDP Sites 206–208, each with limited core recovery, penetrated strata in northern Zealandia beneath the TECTA unconformity. We drilled six sites in the context of seismic reflection surveys (Fig. 2; Figs. DR1 and DR2 in the GSA Data Repository¹). We classified paleodepth (meters below modern sea level [mbsl]) into the following categories (Van Morkhoven et al., 1986): neritic (<200 mbsl), upper bathyal (200–600 mbsl), middle bathyal (600–1000 mbsl), lower bathyal (1000–2000 mbsl), and abyssal (>2000 mbsl). We discovered Paleogene fossils indicative of nearby neritic conditions at sites now far below sea level (Fig. DR3).

Parts of northern Zealandia were transiently uplifted and then subsided. IODP Site U1506 on northern Lord Howe Rise rose close to sea level with a shallow carbonate platform at ca. 50 Ma, and subsided to a bathyal environment (~600 mbsl) by 45 Ma. Neritic fossils of Eocene (ca. 50 Ma) age at Site U1506 are now ~1770 mbsl. At DSDP Site 208 (Fig. 2), middle Eocene (45–43 Ma) cores contain benthic foraminifers indicative of middle bathyal conditions, but planktic-benthic ratios in Paleocene (65–56 Ma) cores indicate shallower conditions, and unconformities separate Paleocene from older and younger strata (Burns et al., 1973).

Southern Lord Howe Rise experienced later transient uplift. Beneath an unconformity at IODP Site U1510 (1238 mbsl), upper Eocene (41–37 Ma) siliceous chalk was deposited at middle bathyal depths, and neritic fossils indicate downslope transport. Site U1510 is ~80 km from DSDP Site 592 (1088 mbsl), where an unconformity separates lower Miocene (23–19 Ma) chalk from lower Oligocene (33–32 Ma) ooze (Kennett et al., 1986). Fossils indicate a lower or middle bathyal environment since the late Eocene at Site 592, but lower Oligocene strata contain layers of coarse (1–4 cm) mollusk (*Ostrea*) fragments (Kennett et al., 1986), consistent with nearby shallow water. DSDP Site 207 (Fig. 2) subsided from upper bathyal to middle bathyal depths during the Paleocene to middle Eocene, but an unconformity separates Eocene (43–38 Ma) from middle Miocene (15–13 Ma) strata, and inclusion of slumped upper Eocene (38–36 Ma) material along the unconformity is consistent with peak regional uplift in the latest Eocene and early Oligocene (36–32 Ma). The crest of southern Lord Howe Rise has a current depth of 900–1000 mbsl.

Bioclastic limestone dredged from ~1750 mbsl in southwest Reinga Basin (Fig. 2) con-




Figure 2. (A) Zealandia bathymetry (m), new (stars) and existing (circles) drill sites, New Caledonia Trough (NCT), Norfolk Ridge (NR), D'Entrecasteaux Ridge (DER), Reinga Basin (RB), and outline of Zealandia (dotted). B.—Basin. (B) Timing of events inferred from integrated analysis (see the Data Repository [see footnote 1]). Plio.—Pliocene; Pleist.—Pleistocene; DSDP—Deep Sea Drilling Project.

tains neritic benthic foraminifers with ages of 36–30 Ma, and reworked Eocene (43–38 Ma) planktic species (Browne et al., 2016; Sutherland et al., 2017). At IODP Site U1508 (1609 mbsl), in the eastern Reinga Basin, onlap indicates deformation started at ca. 39 Ma (Figs. DR1, DR2, and DR4), and Oligocene (26–23 Ma) chalk contains a middle to lower bathyal fauna mixed with shallow-water ostracods and benthic foraminifers, along with palynoflora indicating downslope transport from land. Reinga Basin and Lord Howe Rise sample locations have erosional unconformities identified on seismic reflection profiles (Fig. DR4).

At IODP Site U1509 (2911 mbsl), in the southern New Caledonia Trough, we drilled into Cretaceous Fairway–Aotea Basin strata (Fig. 2; Collot et al., 2009). Pleistocene to Oligocene ooze and chalk contain lower bathyal to abyssal benthic foraminifers. Eocene assemblages indicate lower bathyal paleodepths. Paleocene and Cretaceous assemblages indicate a paleo-water depth of ~1000 m. Cretaceous claystones contain plant fragments and fern spores that indicate coastal proximity. Combined data suggest ~2000 m of Cenozoic subsidence, with most accomplished after 59 Ma and before 45 Ma.

¹GSA Data Repository item 2020110, geological and geophysical data; New Caledonia geology; paleontological evidence used to infer paleogeography; and Figures DR1–DR4, is available online at <http://www.geosociety.org/datarepository/2020/>, or on request from editing@geosociety.org.

At IODP Site U1507 (3568 mbsl), we drilled sediments of the northern New Caledonia Trough for the first time. Fossils from the oldest drilled sediments (864 m below seafloor [mbsf]) indicate lower bathyal depths at 41 Ma (Sutherland et al., 2019). Sedimentation rates increase downhole from 10 m/m.y. to 40 m/m.y. (Fig. DR2), and extrapolation to the base of the unit, determined from seismic reflection data to be ~1300 mbsf, indicates a Paleogene age for the basin.

GEOGRAPHIC CHANGE

Cretaceous rifting from Gondwana likely thinned the crust of Zealandia, but large elevation changes (1–3 km) across a wide region (Fig. 3)

occurred during the Paleogene. Lord Howe Rise uplifted by at least 1 km, with a southeast migration in this motion from 50 to 35 Ma. New Caledonia Trough subsided 1–3 km, starting at ca. 55–45 Ma, with no resolvable difference in timing between north and south. The East Reinga Basin records deformation at ca. 39 Ma with peak uplift at ca. 26–23 Ma (Fig. 2; details of fossil evidence are given in the “Paleogeography” section of the Data Repository).

Flexure would not produce the magnitude, wavelength, nor timing of observed elevation changes, so we suggest crustal delamination and slab formation by reactivation of a west-dipping Cretaceous subduction zone (Fig. 4; Sutherland et al., 2010) to explain the observed features.

Thermal isostatic and dynamic forces (upwelling) are inferred to have driven uplift of Lord Howe Rise, while delamination of basaltic lower crust, minor local extension, and dynamic forces (downwelling) caused New Caledonia Trough to subside.

SUBDUCTION INITIATION

Subduction initiation can be spontaneous if gravitational instability and a weakness are juxtaposed (Stern, 2004), or it can be induced if gravitational instability grows during convergence across a fault (Gurnis et al., 2004). We propose an additional case: A stable gravitational anomaly may exist but will founder to produce a slab if failure occurs. Time scales,




Figure 3. Paleogeographic reconstructions at 55, 50, 45, 35, and 25 Ma. Dark blue is >2000 m water depth, cyan is 2000–200 m, pink is 200–0 m, and white is land. Stars are new drill sites. Present coastline (gray) is shown for reference. Filled arrows show uplift (up) or subsidence (down). Open arrows show active convergent or divergent crustal deformation. Triangles show active (red) or extinct (gray) arc volcanic activity. Black line with teeth is a suggestion for trench location, but alternate hypotheses exist. Plat.—Plateau; R.—Ridge; IODP—International Ocean Discovery Program; DSDP—Deep Sea Drilling Project.




Figure 4. Cartoon cross sections illustrating the transition from Paleocene rifted margin to Eocene subduction. Line is between northern Lord Howe Rise and southern New Caledonia (Fig. 3A). Pink—continental crust; red—arc plutons; green—Cretaceous subducted slab (dark blue where it is eclogite); purple—ocean crust, gray—mantle lithosphere, and yellow—asthenosphere.

length scales, and processes may differ, but the idea has similarities to velocity-weakening behavior on a fault during an earthquake. We propose the term subduction rupture event (SRE) to describe the nucleation and lateral propagation of the onset of fault slip and slip-weakening on lithospheric faults during subduction initiation. Induced subduction initiation requires regional forcing, whereas an SRE requires only local forcing (nucleation of initial failure) and slip-weakening processes that facilitate lateral propagation.

Extension and volcanic activity associated with IBM subduction started at 53–52 Ma (Arculus et al., 2015), and the onset of metamorphism in New Caledonia was at 55–50 Ma (Pirard and Spandler, 2017; Vitale-Brovarone et al., 2018; Fig. 2; see the Data Repository). In the Tonga forearc, the oldest plagiogranites have ages ca. 51–50 Ma, and arc activity is evident after ca. 48 Ma (Meffre et al., 2012). Seafloor spreading in the Tasman Sea also ended at ca. 52 Ma (Gaina et al., 1998). The Emperor-Hawaii bend records onset of Pacific plate-motion change at ca. 50 Ma, with a time of maximum curvature at ca. 48–47 Ma (O'Connor et al., 2013), and magnetic anomalies record rapid seafloor spreading south of Australia and New Zealand after 45–43 Ma (Sutherland, 1995; Cande and Stock, 2004). These major tectonic changes in the western Pacific were broadly synchronous with subsidence of the New Caledonia Trough

and transient uplift of the northern Lord Howe Rise, and we interpret them to have been caused by initial slab formation.

Geological evidence from New Zealand, New Caledonia, and magnetic data show that a fossil Mesozoic arc lies beneath the New Caledonia Trough, and the Norfolk Ridge contains forearc accretionary complexes (Paris, 1981; Sutherland, 1999; Mortimer, 2004). Collision of a young large igneous province caused Cretaceous flat-slab subduction death and hence underplating of a thick basaltic lower crust (Davy et al., 2008). We propose that metamorphism of delaminated lower crust to eclogite provided a density anomaly that led to slab formation during the Eocene (Fig. 4).

Suitable conditions for reactivation might exist in an extinct subduction zone, including weakness of the subduction fault zone, and gravitational instability of buoyant arc rocks and serpentinized mantle set against dense eclogite of the slab and/or root of the arc, in addition to thermal contrasts (Leng and Gurnis, 2015). Subducted sediment or continent slivers may also play some role. Subduction zones have high continuity, so resurrection could propagate over a large distance.

Subduction initiated along ~10,000 km of the western Pacific between 55 and 50 Ma and seemingly preceded major plate-motion change. In our SRE hypothesis, subduction was induced as slip laterally propagated to resurrect extinct

subduction zones. The rate of lateral propagation (>1 m/yr) was about two orders of magnitude faster than typical plate-motion rates.

After the SRE, forces, topography, and volcanism evolved in response to local conditions. The ca. 48 Ma change in Pacific plate direction and speed toward the western Pacific occurred earlier than the ca. 44 Ma acceleration in Australia toward Indonesia. The ~4–8 m.y. delay before plate motions changed reflects progressive growth of slabs and reductions in fault resistance. The North Pacific evolved fastest, but the SRE may have nucleated elsewhere. The oldest evidence for SRE activity that we are aware of is in Papua New Guinea, where ophiolites were emplaced at ca. 58 Ma (Lus et al., 2004).

There has been one major kinematic change during Earth history for which we know plate motions through magnetic anomalies, hotspots, and now regional topographic changes: the Eocene event of the western Pacific. We suggest that subduction initiation involves: (1) an SRE, and (2) development of forces as slabs grow and faults weaken. Fossil subduction margins provide the right ingredients for this to happen: lateral continuity, weakness, and density contrasts. The Pacific “subduction resurrection” context contrasts with the Mediterranean, where subduction initiation was induced by Oligocene continental collision (Handy et al., 2010), but prolonged (>30 m.y.) slab foundering had limited impact on global plate motions, perhaps due to limitations of suitability and continuity of inherited geology.

Subduction initiation beneath Zealandia altered geography, crustal thickness, and likely also crustal composition. It may be that other continents were shaped at a similar time (e.g., thin continental parts of Indonesia and South China Sea). As there has been only one major plate-motion change event since 83 Ma, the frequency-magnitude relationship for SRE events over geological time is hard to determine. It is plausible, though, that between 40 and 100 such events have occurred since the onset of plate tectonics. The unstable dynamical behavior we infer challenges the principle of uniformitarianism for plate tectonics because there is likely no modern analogue for the SRE processes that occurred at ca. 55–50 Ma. However, the geographical, geological, and geochemical evolution of continents and mantle probably was affected by these infrequent events. The records in Zealandia provide unique insight for recognizing and understanding them, and may even be a basis for prediction of favorable geological conditions: subduction resurrection.

ACKNOWLEDGMENTS

We thank the International Ocean Discovery Program (IODP); the personnel of R/V *JOIDES Resolution* on Expedition 371; proponents unable to sail on Expedition 371; and everyone on surveys TAN1312, TAN1409, and TECTA. This work was funded by the

U.S. National Science Foundation; IODP participating countries; New Zealand, France, and New Caledonia (site surveys); the Spanish Ministry of Economy and Competitiveness and Fondo Europeo de Desarrollo Regional (FEDER) funds project CGL2017-84693-R and a Leonardo Grant, BBVA Foundation (Alegría); Korean IODP (K-IODP) (Park); China grant NSFC 41473029, 91958110 (He Li); and Brazil grant 183/2017-CII/CGPE/DPB/CAPES (Giorgioni).

REFERENCES CITED

- Aitchison, J.C., Ali, J.R., and Davis, A.M., 2007, When and where did India and Asia collide?: *Journal of Geophysical Research—Solid Earth*, v. 112, B05423, <https://doi.org/10.1029/2006JB004706>.
- Arculus, R.J., et al., 2015, A record of spontaneous subduction initiation in the Izu-Bonin-Mariana arc: *Nature Geoscience*, v. 8, p. 728–733, <https://doi.org/10.1038/ngeo2515>.
- Bache, F., Sutherland, R., Stagpoole, V., Herzer, R., Collot, J., and Rouillard, P., 2012, Stratigraphy of the southern Norfolk Ridge and the Reinga Basin: A record of initiation of Tonga-Kermadec-Northland subduction in the southwest Pacific: *Earth and Planetary Science Letters*, v. 321–322, p. 41–53, <https://doi.org/10.1016/j.epsl.2011.12.041>.
- Browne, G.H., Lawrence, M.J.F., Mortimer, N., Clowes, C.D., Morgans, H.E.G., Hollis, C.J., Beu, A.G., Black, J.A., Sutherland, R., and Bache, F., 2016, Stratigraphy of Reinga and Aotea Basins, NW New Zealand: Constraints from dredge samples on regional correlations and reservoir character: *New Zealand Journal of Geology and Geophysics*, v. 59, p. 396–415, <https://doi.org/10.1080/00288306.2016.1160940>.
- Burns, R.E., and Andrews, J.E., 1973, Regional aspects of deep sea drilling in the southwest Pacific, in Burns, R.E., et al., eds., *Initial Reports of the Deep Sea Drilling Project, Volume 21: Washington, D.C., U.S. Government Printing Office*, p. 897–906, <https://doi.org/10.2973/dsdp.proc.21.128.1973>.
- Burns, R.E., Andrews, J.E., van der Lingen, G.J., Churkin, M., Galehouse, J.S., Packham, G.H., Davies, T.A., Kennett, J.P., Dumitrica, P., Edwards, A.R., Von Herzen, R.P., Burns, D., and Webb, P.N., 1973, Site 208, in Burns, R.E., et al., eds., *Initial Reports of the Deep Sea Drilling Project, Volume 21: Washington, D.C., U.S. Government Printing Office*, p. 271–281.
- Cande, S.C., and Stock, J.M., 2004, Pacific-Antarctic-Australia motion and the formation of the Macquarie plate: *Geophysical Journal International*, v. 157, p. 399–414, <https://doi.org/10.1111/j.1365-246X.2004.02224.x>.
- Cande, S.C., Stock, J.M., Mueller, R.D., and Ishihara, T., 2000, Cenozoic motion between East and West Antarctica: *Nature*, v. 404, p. 145–150, <https://doi.org/10.1038/35004501>.
- Cande, S.C., Patriat, P., and Dyment, J., 2010, Motion between the Indian, Antarctic and African plates in the early Cenozoic: *Geophysical Journal International*, v. 183, p. 127–149, <https://doi.org/10.1111/j.1365-246X.2010.04737.x>.
- Cluzel, D., Meffre, S., Maurizot, P., and Crawford, A.J., 2006, Earliest Eocene (53 Ma) convergence in the Southwest Pacific: Evidence from pre-obduction dikes in the ophiolite of New Caledonia: *Terra Nova*, v. 18, p. 395–402, <https://doi.org/10.1111/j.1365-3121.2006.00704.x>.
- Collot, J., Herzer, R.H., Lafoy, Y., and Géli, L., 2009, Mesozoic history of the Fairway-Aotea Basin: Implications regarding the early stages of Gondwana fragmentation: *Geochemistry Geophysics Geosystems*, v. 10, Q12019, <https://doi.org/10.1029/2009GC002612>.
- Davy, B.W., Hoernle, K., and Werner, R., 2008, Hikurangi Plateau: Crustal structure, rifted formation, and Gondwana subduction history: *Geochemistry Geophysics Geosystems*, v. 9, Q07004, <https://doi.org/10.1029/2007GC001855>.
- Gaina, C., Mueller, D.R., Royer, J.-Y., Stock, J., Hardbeck, J.L., and Symonds, P., 1998, The tectonic history of the Tasman Sea: A puzzle with 13 pieces: *Journal of Geophysical Research*, v. 103, p. 12413–12433.
- Gurnis, M., Hall, C.E., and Lavier, L.L., 2004, Evolving force balance during incipient subduction: *Geochemistry, Geophysics, Geosystems*, v. 5, Q07001, <https://doi.org/10.1029/2003GC000681>.
- Handy, M.R., Schmid, S.M., Bousquet, R., Kissling, E., and Bernoulli, D., 2010, Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological-geophysical record of spreading and subduction in the Alps: *Earth-Science Reviews*, v. 102, p. 121–158, <https://doi.org/10.1016/j.earscirev.2010.06.002>.
- Kennett, J.P., et al., 1986, *Initial Reports of the Deep Sea Drilling Project Leg 90: Washington, D.C., U.S. Government Printing Office*, 1517 p., <https://doi.org/10.2973/dsdp.proc.90.1986>.
- Laird, M.G., and Bradshaw, J.D., 2004, The breakup of a long-term relationship: The Cretaceous separation of New Zealand from Gondwana: *Gondwana Research*, v. 7, p. 273–286, [https://doi.org/10.1016/S1342-937X\(05\)70325-7](https://doi.org/10.1016/S1342-937X(05)70325-7).
- Leng, W., and Gurnis, M., 2015, Subduction initiation at relic arcs: *Geophysical Research Letters*, v. 42, p. 7014–7021, <https://doi.org/10.1002/2015GL064985>.
- Lus, W.Y., McDougall, I., and Davies, H.L., 2004, Age of the metamorphic sole of the Papuan ultramafic belt ophiolite, Papua New Guinea: *Tectonophysics*, v. 392, p. 85–101, <https://doi.org/10.1016/j.tecto.2004.04.009>.
- Mathews, K.J., Williams, S.E., Whittaker, J.M., Müller, R.D., Seton, M., and Clarke, G.L., 2015, Geologic and kinematic constraints on Late Cretaceous to mid Eocene plate boundaries in the southwest Pacific: *Earth-Science Reviews*, v. 140, p. 72–107, <https://doi.org/10.1016/j.earscirev.2014.10.008>.
- Meffre, S., Falloon, T.J., Crawford, T.J., Hoernle, K., Hauff, F., Duncan, R.A., Bloomer, S.H., and Wright, D.J., 2012, Basalts erupted along the Tongan fore arc during subduction initiation: Evidence from geochronology of dredged rocks from the Tonga fore arc and trench: *Geochemistry Geophysics Geosystems*, v. 13, Q12003, <https://doi.org/10.1029/2012GC004335>.
- Mortimer, N., 2004, New Zealand's geological foundations: *Gondwana Research*, v. 7, p. 261–272, [https://doi.org/10.1016/S1342-937X\(05\)70324-5](https://doi.org/10.1016/S1342-937X(05)70324-5).
- Mortimer, N., Campbell, H.J., Tulloch, A.J., King, P.R., Stagpoole, V.M., Wood, R.A., Rattenbury, M.S., Sutherland, R., Adams, C.J., and Collot, J., 2017, Zealandia: Earth's hidden continent: *GSA Today*, v. 27, no. 3, p. 27–35.
- Muller, R.D., Gaina, C., Tikku, A., Mihut, D., Cande, S.C., and Stock, J.M., 2000, Mesozoic/Cenozoic tectonic events around Australia, in Richards, M.A., Gordon, R.G., and Van Der Hilst, R.D., eds., *The History and Dynamics of Global Plate Motions: American Geophysical Union Geophysical Monograph 121*, p. 161–188.
- O'Connor, J.M., Steinberger, B., Regelous, M., Koppers, A.A., Wijbrans, J.R., Haase, K.M., Stoffers, P., Jokat, W., and Garbe-Schönberg, D., 2013, Constraints on past plate and mantle motion from new ages for the Hawaiian-Emperor Seamount Chain: *Geochemistry Geophysics Geosystems*, v. 14, p. 4564–4584, <https://doi.org/10.1002/ggge.20267>.
- Paris, J.-P., 1981, *Geologie de la Nouvelle-Calédonie; un Essai de Synthèse (Geology of New-Caledonia; A Synthetic Text): Mémoires du Bureau de Recherches Géologiques et Minières 113*, 278 p.
- Pirard, C., and Spandler, C., 2017, The zircon record of high-pressure metasedimentary rocks of New Caledonia: Implications for regional tectonics of the south-west Pacific: *Gondwana Research*, v. 46, p. 79–94, <https://doi.org/10.1016/j.gr.2017.03.001>.
- Reagan, M. K., et al., 2017, Subduction initiation and ophiolite crust: New insights from IODP drilling: *International Geology Review*, v. 59, p. 1439–1450, <https://doi.org/10.1080/00206814.2016.1276482>.
- Steinberger, B., Sutherland, R., and O'Connell, R.J., 2004, Prediction of Emperor-Hawaii seamount locations from a revised model of global plate motion and mantle flow: *Nature*, v. 430, p. 167–173, <https://doi.org/10.1038/nature02660>.
- Stern, R.J., 2004, Subduction initiation: Spontaneous and induced: *Earth and Planetary Science Letters*, v. 226, p. 275–292, [https://doi.org/10.1016/S0012-821X\(04\)00498-4](https://doi.org/10.1016/S0012-821X(04)00498-4).
- Sutherland, R., 1995, The Australia-Pacific boundary and Cenozoic plate motions in the SW Pacific: Some constraints from Geosat data: *Tectonics*, v. 14, p. 819–831, <https://doi.org/10.1029/95TC00930>.
- Sutherland, R., 1999, Basement geology and tectonic development of the greater New Zealand region: An interpretation from regional magnetic data: *Tectonophysics*, v. 308, p. 341–362, [https://doi.org/10.1016/S0040-1951\(99\)00108-0](https://doi.org/10.1016/S0040-1951(99)00108-0).
- Sutherland, R., Collot, J., Lafoy, Y., Logan, G.A., Hackney, R., Stagpoole, V., Uruski, C., Hashimoto, T., Higgins, K., Herzer, R.H., Wood, R., Mortimer, N., and Rollet, N., 2010, Lithosphere delamination with foundering of lower crust and mantle caused permanent subsidence of New Caledonia Trough and transient uplift of Lord Howe Rise during Eocene and Oligocene initiation of Tonga-Kermadec subduction, western Pacific: *Tectonics*, v. 29, TC2004, <https://doi.org/10.1029/2009TC002476>.
- Sutherland, R., Collot, J., Bache, F., Henrys, S., Barker, D., Browne, G., Lawrence, M., Morgans, H., Hollis, C., and Clowes, C., 2017, Widespread compression associated with Eocene Tonga-Kermadec subduction initiation: *Geology*, v. 45, p. 355–358, <https://doi.org/10.1130/G38617.1>.
- Sutherland, R., Dickens, G.R., Blum, P., and the Expedition 371 Scientists, 2019, *Tasman frontier subduction initiation and Paleogene climate: Proceedings of the International Ocean Discovery Program*, v. 371: College Station, Texas, International Ocean Discovery Program, <http://publications.iodp.org/proceedings/371/371title.html>.
- Van Morkhoven, F.P., Berggren, W.A., Edwards, A.S., and Oertli, H., 1986, Cenozoic cosmopolitan deep-water benthic foraminifera: *Elf Aquitaine Memoir*, v. 11, p. 1–421.
- Vitale-Brovarone, A., Agard, P., Monié, P., Chauvet, A., and Rabaute, A., 2018, Tectonometamorphic architecture of the HP belt of New Caledonia: *Earth-Science Reviews*, v. 178, p. 48–67, <https://doi.org/10.1016/j.earscirev.2018.01.006>.

Printed in USA