


Reproductive behavior of Atlantic halibut (*Hippoglossus hippoglossus*) interpreted from electronic tags

Rachel C. Marshall^{1,2}  | Jonathan A. D. Fisher¹ | Anthony L. Einfeldt³ | Paul Gatti^{1,4} | Dominique Robert⁵ | Arnault Le Bris^{1,6}

¹Memorial University of Newfoundland, Fisheries and Marine Institute, Center for Fisheries Ecosystem Research, St. John's, Newfoundland, Canada

²University of Rhode Island, Graduate School of Oceanography, Narragansett, Rhode Island, USA

³Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada

⁴UMR DECOD (Ecosystem Dynamics and Sustainability), IFREMER, INRAE, Institut Agro, Plouzané, France

⁵Université du Québec à Rimouski, Institut des Sciences de la Mer de Rimouski, Rimouski, Quebec, Canada

⁶Institute of Marine Research, Bergen, Norway

Correspondence

Rachel C. Marshall and Arnault Le Bris, Memorial University of Newfoundland, Fisheries and Marine Institute, Center for Fisheries Ecosystem Research, 155 Ridge Road, St. John's, NL A1C 5R3, Canada. Email: rachel.marshall@mi.mun.ca and Arnault.LeBris@mi.mun.ca

Dominique Robert, Université du Québec à Rimouski, Institut des Sciences de la Mer de Rimouski 310, allée des Ursulines CP 3300, station, A Rimouski, QC G5L 3A1 Canada. Email: Dominique_Robert@uqar.ca

Funding information

Canada First Research Excellence Fund; Canada Research Chair program; Department of Fisheries and Aquaculture, Government of Newfoundland and Labrador; Department of Fisheries and Land Resources of Newfoundland and Labrador; Development Corporation of Newfoundland and Labrador Ignite Grants; John R. Evans Leaders Fund of the Canada Foundation for Innovation; Ministère de l'agriculture, des pêcheries et de l'alimentation du Québec; Natural Sciences and Engineering Research Council of Canada Strategic Partnership Grant, Grant/Award Number: STPGP 506993-17; Ressources Aquatiques Québec

Abstract

Reproductive timing, location, and behavior are important characteristics that determine marine population dynamics, structure, and resilience to threats, including fishing and climate change. It is challenging to evaluate factors driving variability in these reproductive traits in wild fishes because of the difficulty observing individuals in their natural environments. In the present study, we used high-resolution depth, temperature, and acceleration time series recorded by pop-up satellite archival tags to (1) identify and characterize patterns in depth and acceleration that may be indicative of spawning events in large Atlantic halibut (*Hippoglossus hippoglossus*), and (2) estimate the effects of individual traits (body size and sex) and environmental factors (location and temperature) on spawning time and frequency. Unique rapid rises observed in the winter depth profiles were interpreted as spawning events. The initiation of the first presumed spawning rise was negatively correlated to water temperature experienced during the prespawning season, suggesting that currently increasing water temperature in the Gulf of St. Lawrence may induce phenological change in halibut spawning time. The number of rises of batch-spawning females was unrelated to female body size. The present study demonstrates how electronic tagging can be used for in-depth characterization of timing, location, and behaviors associated with spawning in a large flatfish species. Such information can inform spatiotemporal management and conservation measures aiming to protect species from directed fishing and by-catch during spawning.

KEYWORDS

Atlantic halibut, electronic tag, fish spawning, Gulf of St. Lawrence, pop-up satellite archival tag

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Journal of Fish Biology* published by John Wiley & Sons Ltd on behalf of Fisheries Society of the British Isles.

1 | INTRODUCTION

Reproductive timing, location, and behavior are important factors determining fish population structure, dynamics, and resilience to anthropogenic threats such as fishing and climate change. Understanding variability in these traits is necessary for developing effective management and conservation measures (Le Bris et al., 2013; Zemeckis et al., 2014). However, characterizing the spawning behaviors of marine fish in the wild is challenging, especially for species that are difficult to observe in their natural environments, such as species that are highly migratory, or that inhabit deep and dark environments far offshore. Spawning time and location of marine fish have traditionally been determined by visual and/or histological characterization of the maturity stage of fish gonads (Kohler, 1967; McBride et al., 2022), or by the presence of fish eggs and larvae in planktonic surveys (Ghinter et al., 2023; Pepin & Helbig, 1997). Laboratory tank studies have revealed physiological characteristics of reproduction, including quality and quantity of eggs, spawning frequency, and environmental effects on reproductive timing (Brown et al., 2006; Norberg et al., 2001). Courting and spawning behaviors have also been described using laboratory studies (Brawn, 1961; Smith et al., 1999), as well as active acoustic surveys (Grabowski et al., 2012; Rose, 1993).

Electronic data storage tags have become more prevalent as a method for studying spawning and other behaviors in elusive marine species because of their capacity to record high-resolution time series of individual environmental habitat use (e.g., Grabowski et al., 2015; Siwicke et al., 2022; Tsuda et al., 2006; Yasuda et al., 2013). Time series from electronic tags can reveal unique patterns of depth use that are indicative of potential spawning behaviors, such as the oscillatory diving patterns with prolonged surface intervals exhibited by bluefin tuna (Block et al., 2001), the rapid rises off the seafloor in Pacific halibut (Seitz et al., 2005), or the periodic short-term shallow water activity in small spotted catsharks (Wearmouth et al., 2013). When combined with reconstructed migratory tracks, these patterns can help reveal unknown spawning areas (Gatti et al., 2020) and ultimately inform on the risks of interaction with fisheries or other ocean uses (Marshall et al., 2023).

Pop-up satellite archival tags (PSATs) are a type of electronic archival tag widely used to study marine species, especially large migratory fish. They are attached externally, record ambient oceanographic conditions (temperature, depth, light intensity, etc.), and can be programmed to release from their host fish, float to the surface, and transmit data via satellite, allowing access to data summaries without the need to recapture the tagged fish (Block et al., 2005; Hussey et al., 2015). The physical recovery of PSATs also provides access to the full high-resolution data sets archived in the tags, including three-dimensional acceleration data, which can reveal behavioral patterns not necessarily observable from lower-resolution satellite transmitted data (Fisher et al., 2017; Nielsen et al., 2018).

In recent years, several studies have used PSATs to study the migratory behavior of Atlantic halibut (*Hippoglossus hippoglossus*) in the Northwest Atlantic (Armsworthy et al., 2014; Gatti et al., 2020; Gatti et al., 2021; James et al., 2020; Le Bris et al., 2018; Liu et al., 2019; Murphy et al., 2017), encouraged by the steady increase in fisheries landings observed in the past two decades (DFO, 2021a, 2021b). Several of these PSAT studies observed rapid ascents and descents during the presumed spawning seasons and suggested that these behaviors were spawning rises (Armsworthy et al., 2014; Le Bris et al., 2018; Murphy et al., 2017) because of their consistency with the spawning rises described from PSAT time series for the closely related Pacific halibut (*Hippoglossus stenolepis*, Seitz et al., 2005) and directly observed for several other flatfish species (Carvalho et al., 2003; Konstantinou & Shen, 1995; Manabe et al., 2000; Manabe & Shinomiya, 2001; Moyer et al., 1985). Although the reproductive behavior of halibut has not been directly observed in the wild, some inferences have been possible from indirect methods. Egg sampling, gonad histology, and laboratory tank studies have demonstrated that halibut are deep-water batch-spawners, releasing 0.5–7 million mesopelagic eggs (Cargnelli et al., 1999; Haug, 1990; Haug & Gulliksen, 1988a) during varying periods between November and June depending on the region (Armsworthy et al., 2014; Björnsson et al., 1998; Haug, 1990; Kjørsvik et al., 1987; Kohler, 1967; Neilson et al., 1993). Females follow regular batch-spawning intervals based on egg development timing (Finn et al., 2002; Norberg et al., 1991), and males are capable of spawning much more frequently over longer periods (Methven et al., 1992; Norberg et al., 1991). However, despite a rapidly expanding body of research on Atlantic halibut ecology (Shackell et al., 2022), the variability in spawning timing and sex-specific behaviors of the species in the wild remains largely unstudied, in part because of the difficulty observing halibut in the winter spawning season, especially in the Gulf of St. Lawrence that is seasonally covered by sea ice.

In the present study, we used high-resolution depth, temperature, and acceleration data from recovered PSATs to characterize the spawning behavior of Atlantic halibut in the Gulf of St. Lawrence. Building on previous studies observing presumed spawning rises in Atlantic halibut and other flatfish, we identified presumed spawning rises in female and male halibut and determined their dates of occurrence, amplitude, and frequency. We then evaluated the role of temperature, fish body size, and location on both the timing of the first presumed spawning rise and the number of spawning rises per individual female. We were particularly interested in the timing and number of spawning rises because they have direct applications to the management of the fishery. Spawning time can directly inform the timing of fishery closures during the spawning period and may also provide a better understanding of population structure (Zemeckis et al., 2014). Information on the numbers of spawning rises, and whether they are dependent on fish size, is relevant to estimating the total egg production in a population and understanding the capacity of a population to sustain various levels of fishing pressure or patterns of fishing selectivity (Hixon et al., 2014; Le Bris et al., 2015).

2 | MATERIALS AND METHODS

2.1 | Ethics statement

The care and use of experimental animals complied with the Canadian Council on Animal Care's animal welfare laws, guidelines, and policies as approved by Memorial University's Animal Care Committee, protocol 19-02-AL.

2.2 | Electronic tagging

As part of a research effort to understand the ecology of Atlantic halibut in the Northwest Atlantic, PSATs were deployed on 119 individual halibut of legal size (>85 cm) landed in good condition across the Gulf of St. Lawrence in September and October 2013, 2014, 2015, and 2017 (Gatti et al., 2020), several months before the spawning season. Following the methodology developed and used by the International Pacific Halibut Commission (Seitz et al., 2003), PSATs were attached to halibut using sterilized titanium darts and 180-kg-test monofilament tethers. The dart was inserted under the pterygiophores on the dorsal, eyed side of the fish. This methodology was reported to have no long-term effect on the behavior of tagged individuals (Seitz et al., 2003). Tagging operations were carried out exclusively aboard chartered commercial fishing vessels, and fish were caught using commercial demersal longlines (for more information on tagging operations, see Gatti et al., 2020; James et al., 2020; Le Bris et al., 2018; Murphy et al., 2017). Two models of tags were used: X-Tag PSATs from Microwave Telemetry, Inc., which recorded depth (± 0.34 to 5.38 m resolution from 0 to 1296 m), temperature (± 0.16 to 0.23°C resolution), and light intensity level ($\pm 4 \times 10^{-5}$ lx resolution at 555 nm) every 2 min (Table S1), and MiniPATs from Wildlife Computers Inc., which recorded depth (± 0.5 m), temperature ($\pm 0.1^\circ\text{C}$), and light intensity (5×10^{-12} to 5×10^{-2} W/cm²) at 5–15 s resolution throughout the deployment period (Table S1). MiniPATs used in the 2017 deployment also recorded acceleration ($n = 36$). Tags were preprogrammed to release from their host fish at a specific date ~ 1 year following tag deployment. High-resolution archived data sets were available from tags that were physically recovered. Tag recoveries were conducted aboard chartered commercial fishing vessels using a CLS ARGOS RXG-134 goniometer (CLS America Inc., Lanham, MD, USA) (Fisher et al., 2017; Gatti et al., 2020). Of the 119 tags deployed, 99 popped off and transmitted to satellite, and 64 of those were physically recovered. Two recovered tags showed stationary depth time series after a few days, and the tagged fish was presumed dead. Two recovered tags also popped off early, so the time series were removed from analyses.

The depth time series from 60 physically recovered tags were visually inspected for evidence of spawning behavior: consecutive rapid ascents and descents similar to patterns previously documented as presumed spawning behavior for halibut species (Murphy et al., 2017; Seitz et al., 2005). Forty of the 60 physically recovered tags showed potential spawning behavior and were selected for sex determination analyses.

TABLE 1 Sex conclusively assigned to 36 halibut (Table S1) using one of three methods with enough confidence to include in analyses.

	Female	Male	Total
Ultrasound	4	5	9
Genetics	10	1	11
Visual consensus	12	4	16
Total	26	10	36

2.3 | Sex determination

Sex was determined in one of three ways (Table 1). (1) A veterinary ultrasound, available only during the 2014 and 2015 tagging studies, was used during tagging operations to non-invasively inspect the shape of the gonads (Loher & Stephens, 2011) for nine halibut tagged in 2014 and 2015. (2) Taking fin clip samples for genetic analysis was introduced into the tagging protocol in 2017. Fin clip samples were not taken before 2017 because the technology to identify sex genetically from fin clip samples was not yet developed (Einfeldt et al., 2021). Genetic analysis using RADseq-derived single nucleotide polymorphisms associated with a region of chromosome 12 containing the presumed master sex-determining gene *GSDF* was used to assign sex for 11 halibut displaying presumed spawning rises on PSAT depth profiles (Einfeldt et al., 2021). (3) Sex of 16 additional halibut was identified based on visual inspection of the PSAT depth profiles following the description by Murphy et al. (2017) of distinctive female and male patterns: females show a very regular pattern of one clear rise every few days, whereas males are characterized by a greater number of rises at irregular intervals over a longer period (Figure 1). Depth profiles, including those with confirmed sex from ultrasound or genetics, were distributed to nine research collaborators without communicating the known sex of any profiles. By using this approach, the 20 profiles with known sex could be used to validate the visual inspection method. Responses were reviewed, and sex was assigned to 16 profiles for which there was greater than 75% agreement among the participants. Eighteen of the 20 profiles with known sex were also correctly assigned with greater than 75% agreement among visual inspection responses, reinforcing confidence in the visual assignment method. The final tally of sexed halibut used for further analyses was 36 (Table 1; Table S1).

2.4 | Spawning rise detection

Halibut presumed spawning rises have been identified as rises off of and returning to the seafloor that are extreme both in distance traveled and in rate of ascent and descent, compared to other vertical movement events during the year (Fisher et al., 2017; Seitz et al., 2005). Building on this idea that spawning rises are outliers in the movement of halibut, several methods were attempted to detect spawning rises as objectively as possible (Marshall, 2020: Appendix A). Because of the marked difference in spawning behavior between females and males, two different approaches were chosen for females




FIGURE 1 Examples of depth profiles from recovered PSATs (pop-up satellite archival tags) showing the presumed spawning behavior of (a) a female halibut (determined by ultrasound) showing rises at very regular intervals, (b) a female halibut determined by visual consensus of the depth profile and comparison to the depth profiles of known females, (c) a male halibut (genetically determined) showing a greater number of rises than the females and at much more irregular intervals, (d) a male halibut determined by visual consensus of the depth profile and comparison to the depth profiles of known males, and (e and f) two examples of halibut showing presumed spawning behavior for which sex could not be conclusively determined.

and males. For the 26 female halibut, we favored a four-step automated procedure using the rate of change in depth. (1) Rates of depth change were estimated by calculating first-order difference in the depth time series. (2) Absolute values of the rates of depth change were filtered using a 5-min moving average to remove noise caused by tag sensor errors or random vertical movements. (3) The mean and standard deviation of the filtered time series of depth changes were calculated. (4) Spawning events were identified as any depth change (absolute value) per time step greater than the mean rate of depth changes plus 12 standard deviations (an arbitrary limit selected after extensive preliminary analysis) that occurred between November 1 and April 30, the period of halibut spawning in the Northwest Atlantic suggested by previous studies (Kohler, 1967; McCracken, 1958; Neilson et al., 1993).

Automated spawning rise detection, including the rate of depth change method described earlier for females, was also attempted for males, but because spawning behavior of males was more irregular (Figure 1), likely due to the ability of males to spawn more readily

without the physiological constraints associated with developing eggs (Coleman & Jones, 2011), as well as possible activity associated with courting, the automated detection did not work. Instead, presumed spawning rises of males could be detected only by visual inspection of the depth profiles. Any ascents, of duration less than 4 h from leaving the seafloor to returning to the seafloor, reaching distances ≥ 25 m from the seafloor, occurring between November 1 and April 30 were considered spawning rises. The 25-m threshold was somewhat arbitrary, but it was consistent with characterization of halibut spawning rise minimum distances by Armsworthy et al. (2014) and Fisher et al. (2017), and with the minimum spawning rise distances detected for females using the aforementioned automated process.

2.5 | Spawning rise characterization

To provide a complete characterization of each presumed spawning rise, the depths and times were recorded at the start, end, and peak of

every rise for both female and male halibut. Time between successive rises was calculated by subtracting the time at the peak of one spawning rise from the time at the peak of the next spawning rise. Furthermore, the effects of several environmental and individual factors on two characteristics of special interest, the date of the first presumed spawning rise and the total number of spawning rises, were evaluated using generalized linear models (GLM). GLMs were run only on the female halibut spawning rise data because of the higher confidence in the female spawning rise detection, and the understanding that females constrain reproductive potential more than males due to greater physiological requirements for producing eggs (Bateman, 1948; Coleman & Jones, 2011; McBride et al., 2015).

GLMs were run using the R statistical analysis software package “stats.” The first GLM was used to analyse the variability in the date of the first presumed spawning rise of females, representative of the start of the spawning season. The model tested the effects of water temperature recorded by PSATs during the season prior to the occurrence of spawning rises, water temperature at the beginning of the first spawning rise, fish size (cm fork length), presumed spawning location, year, and tagging location (indicative of halibut origins before the winter spawning migration) on the start of presumed spawning. The pre-season period was fixed from December 1 to January 15 based on previous indications that the latest stage of halibut gonad maturity (Haug & Gulliksen, 1988b) and the highest vitellogenin levels (Björnsson et al., 1998) occur 1–2 months prior to spawning events. To summarize the change in temperature during the fixed pre-season, temperature was expressed in terms of growing degree days (GDD):

$$\text{GDD} = \sum_{i=1}^n \frac{T_{i\max} + T_{i\min}}{2} - T_0$$

Degree days were calculated using the temperature maximum (T_{\max}) and minimum (T_{\min}) recorded by the PSATs for each day (i) of the specified period. The reference temperature (T_0) was set at 0°C. Presumed spawning locations were estimated using a hidden-Markov geolocation model that inferred daily fish position by comparing the PSAT-recorded depth data with known Gulf of St. Lawrence


bathymetry (a full description and validation of the geolocation model is available in Gatti et al., 2021). The geolocation model estimated the probability distribution of an individual fish across the study grid for each day. Probability distributions were averaged for the numbers of days during the presumed spawning period, and the cell with the highest probability was considered to be the spawning location. Then these locations were categorized based on the Gulf of St. Lawrence oceanographic regions described by Galbraith et al. (2020). The GLM error was assumed to follow a normal distribution, and this assumption was confirmed during model validation by plotting residuals versus fitted values, and versus each predictor in and not in the model (Zuur & Ieno, 2016). Model selection was implemented using the AIC (Akaike, 1974) corrected for a small sample size (AIC_c; Burnham & Anderson, 2002).

A second GLM was used to analyse the variability in the number of presumed spawning rises per female halibut. The model tested the effect of mean water temperature experienced by the individual during its spawning period (defined as the period between and including the dates of the first and last presumed spawning rises), the date of the first spawning rise, fish size, spawning location, year, and tagging location on the number of spawning rises. The GLM error was assumed to follow a Poisson error distribution with a log link function, and this assumption and risk of overdispersion were confirmed during model validation (Zuur & Ieno, 2016). Model selection was implemented using the AIC_c (Akaike, 1974; Burnham & Anderson, 2002).

2.6 | Acceleration

MiniPATs deployed in 2017 ($n = 36$) were equipped with three-axis accelerometer sensors. Accelerometer data at a frequency of 0.2 Hz were archived in the tag and were accessible from physically recovered tags ($n = 25$). A previous study has shown that acceleration data from externally attached PSATs can be used to infer fish activity (Nielsen et al., 2018). In the present study, we aimed to determine whether presumed spawning rises were associated with noticeable acceleration patterns. To do so, the total acceleration relative to

FIGURE 2 Proportion of halibut across all tagging years, grouped by sex, considered to be within their spawning period on a given day. Only halibut exhibiting presumed spawning behavior were included in this analysis. Presumed spawning periods were defined as the period between and including the first spawning rise and last spawning rise for each individual.



gravity (units of g), or magnitude, was estimated using values from the three axes:

$$\text{Magnitude} = \sqrt{x^2 + y^2 + z^2}$$

The time series of the acceleration magnitudes for female halibut showing presumed spawning rises ($n = 17$) were visually inspected to quantify the frequency of spawning rises accompanied by a sudden increase in magnitude of acceleration. To further explore whether there were consistent patterns of acceleration relative to spawning rises, the maximum magnitude of acceleration between the start time and end time of each rise was recorded. Acceleration magnitude in relation to male halibut presumed spawning rises was not explored because of lower confidence in the spawning rise detection for males than for females.

3 | RESULTS

3.1 | Spawning behavior characteristics

The depth profiles from 36 PSATs, 26 females and 10 males, were examined for spawning rises (Table S1). A total of 830 presumed spawning rises were identified. Females exhibited between 5 and 13 spawning rises (median: 7) per individual, and males exhibited between 22 and 182 spawning rises (median: 57) per individual. Presumed spawning periods for females and males overlapped (Figure 2). The date of the first spawning rise ranged from January 7 to February 26 among females and from January 6 to 29 among males. Spawning period per individual lasted between 9 and 37 days (median: 16 days) for females, and between 31 and 82 days (median: 61 days) for males.

The spatial distribution of females and males during the period of presumed spawning, estimated from the geolocation model, also overlapped, with greater than 50% of total females and males co-occurring in the same oceanographic region (Figure 3). Grouping

the spawning locations by oceanographic region (Galbraith et al., 2020) put 4 females in the Esquiman Channel, 1 female in the Anticosti Channel, 6 females in the Northwest Gulf, 13 females in the Central Gulf, and 2 females in the Cabot Strait. One male was estimated to be in the Esquiman Channel region just on the border of the Central Gulf region, two males were estimated to be in the Cabot Strait region, and the other seven males were in the Central Gulf.




FIGURE 4 Number of days between successive presumed spawning rises for female and male halibut. Violin plots show the frequency of between-rise times for each sex. Box plots show data medians and first and third quartiles, and whiskers extend to the last values ≤ 1.5 times the interquartile range. Mean values are indicated by single points.




FIGURE 3 Estimated location of individual female (circles) and male (triangles) halibut during the period of presumed spawning rises, estimated by Gatti et al. (2020). The black lines delineate the oceanographic regions of the Gulf of St. Lawrence as defined by Galbraith et al. (2020). Halibut remained in their general presumed spawning areas for several months (Gatti et al., 2020).

TABLE 2 Presumed spawning rise depth use by female and male halibut at each spawning location.

Sex	Presumed spawning location	Fish count	Maximum		Minimum		Mean	
			Start depth	Peak depth	Start depth	Peak depth	Start depth	Peak depth
Female	Anticosti Channel	1	326.0	213.0	250.0	153.5	295.1	185.1
	Cabot Strait	2	503.5	472.0	476.5	279.5	490.7	386.2
	Central Gulf	13	482.5	403.5	303.5	158.5	407.0	274.5
	Esquiman Channel	4	376.5	289.5	253.5	151.0	299.9	223.6
	Northwest Gulf	6	421.5	333.0	251.5	122.5	361.3	256.7
Male	Anticosti Channel	0	-	-	-	-	-	-
	Cabot Strait	2	449.0	421.5	328.0	259.0	422.8	371.4
	Central Gulf	7	459.5	433.5	283.0	144.0	412.3	353.4
	Esquiman Channel	1	329.8	293.2	278.4	168.1	308.1	265.4
	Northwest Gulf	0	-	-	-	-	-	-

Note: Starting depth is assumed to indicate the seafloor depth at an individual's location.




FIGURE 5 An example of detailed presumed spawning rise observation for one female halibut. (a) The full depth profile recorded by the electronic tag. (b) A close look at the depth profile during an interval of 1 month within the winter spawning period showing seven clear spawning rises. (c) The depth profile for a single spawning rise showing a vertical ascent and descent of 227.00 m over the course of an hour. (d) Magnitude of the acceleration revealing a burst 4.42 minutes before the peak of the spawning rise.




FIGURE 6 Time between the maximum acceleration values observed during a female presumed spawning rise and the peak of the spawning rise ($n = 118$) as a function of acceleration magnitude. Negative times indicate when the maximum acceleration occurred before the peak of the spawning rise.

The duration of individual presumed spawning rises from start to end took between 3.25 and 201.58 min (median: 21.00 min) for females, and between 3.00 and 237.25 min (median: 32.12 min) for males. Data exploration did not show any obvious associations with specific periods of the day (day, night, dawn, and dusk). Spawning rises occurred at any time of day, between 00:11:40 and 23:57:45 (median: 12:51:22) for females and between 00:03:15 and 23:55:30 (median: 11:46:47) for males. The amount of time between rises ranged from 0.02 to 14.80 days (median: 3.15 days) for females and from 0.01 to 20.61 days (median: 0.23 days) for males (Figure 4).

The starting depths of presumed spawning rises were between 250.00 and 503.50 m (median: 396.25 m) for females, and between 278.39 and 459.50 m (median: 417.00 m) for males. The depths at the peaks of spawning rises were between 122.50 and 472.00 m (median: 267.00 m) for females, and between 144.00 and 433.50 m (median: 356.75 m) for males. Summaries of depth use at each spawning location are presented in Table 2. Rise distances ranged from 24.54 to 234.50 m (median: 111.25 m) for females and from 25.00 to 284.00 m (median: 40.00 m) for males. The median rise distance for males is expectedly low compared to the female median because the 25-m minimum rise distance criteria strongly influenced male spawning rise detection.

3.2 | Effects on timing and number of spawning rises

The date of the first presumed spawning rise was best explained by a GLM that included only the effect of pre-season GDD based on model selection with AIC_c (Table S2). The model predicted a significant negative relationship between the date of the first female presumed spawning rise and pre-season GDD ($\beta = -5.318 \pm 2.345$, $t_{1,24} = -2.268$, $p < 0.034$), suggesting that female halibut spawn

earlier when they experience warmer temperature during the pre-spawning season. No covariates were found to be significant predictors of the number of female presumed spawning rises. Four models with one covariate each, including mean temperature during the spawning period, fork length, date of the first spawning rise, and tagging location, were selected as the lowest AIC_c models, comparable within two units of each other (Burnham & Anderson, 2002; Table S2), but in no case was there any significance (Table S2).

3.3 | Acceleration

Visual inspection of the magnitude of the acceleration of 17 females showed that 97.5% (115 out of 118) of the presumed spawning rises were accompanied by spikes in acceleration magnitude (Figure 5). Various spikes in acceleration magnitude were observed for each single spawning rise, but the most pronounced acceleration was often observed close to the peak of the spawning rises (Figure 6).

4 | DISCUSSION

In the present study, we used the high-resolution time series recorded by PSATs, geolocation of individual tagged fish, and information on halibut sex to characterize the presumed spawning behavior of Atlantic halibut in the Gulf of St. Lawrence. An inherent limitation of electronic tagging data to study fish behavior is the necessity to infer behaviors from patterns observed in the time series data. Analyses of depth time series recorded by PSATs deployed on Atlantic halibut revealed unique rises (up to 284.00 m) off the seafloor by females and males with peak occurrence in February. We interpreted these unique ascents as spawning rises following previous interpretations of similar rises observed in depth time series recorded by electronic tags deployed on several groundfish species, including Pacific halibut (Seitz et al., 2005), Atlantic halibut (Armsworthy et al., 2014; Murphy et al., 2017), Greenland halibut (Siwicke et al., 2022), and Atlantic cod (Grabowski et al., 2014). Furthermore, spawning rises have been directly observed for several flatfish (Carvalho et al., 2003; Konstantinou & Shen, 1995; Manabe et al., 2000; Manabe & Shinomiya, 2001; Moyer et al., 1985).

Although direct observations of halibut spawning would be necessary to validate our interpretation that the unique rises in the PSAT depth time series are spawning behavior, several characteristics of the rises provide strong support. First, the timing and location of the presumed spawning rises identified in the present study match with the recent first capture of larval Atlantic halibut in the Gulf of St. Lawrence (Ghinter et al., 2023). Second, the occurrences of female and male spawning rises overlap spatially and temporally, a necessity for successful mating and egg fertilization. The third piece of evidence is the consistency between the frequency and period of rises observed in the present study and the reproductive physiology of female and male halibut documented in laboratory and aquaculture studies. Female presumed spawning rises generally occurred at 2–4 day intervals, which is consistent with a documented mean

ovulatory rhythm of 72–80 h (Norberg et al., 1991) and a mean hydration cycle of sequential egg batches of 50 h at 7°C (Finn et al., 2002). The periodic rises seen on the depth time series of female halibut may therefore correspond to the releases of egg batches at intervals constrained by ovulation and oocyte hydration. Male halibut do not show comparable fluctuations in levels of androgenic steroids and have a more continuous spawning capacity (Methven et al., 1992; Norberg et al., 1991), which is consistent with the greater number and higher frequency of presumed spawning rises by males than by females observed in the present study. Additionally, milt release in male halibut can continue for several months (Methven et al., 1992), which is consistent with our observation of a longer period of presumed spawning in males (median: 61 days) than females (median: 16 days).

The accelerometer data available from physically recovered tags may provide additional support to the spawning rise interpretation. We observed sudden bursts in the magnitude of acceleration recorded by PSATs close to the peak of female presumed spawning rises in 97.5% of cases (115 out of 118). Although little is known about halibut mating behavior, detailed description of Atlantic cod mating behavior showed that cod release eggs while at the top of a rapid rise (Brawn, 1961). Furthermore, previous studies of flatfish spawning have observed eggs being released at the peaks of spawning rises (Carvalho et al., 2003; Konstantinou & Shen, 1995). Acceleration bursts may therefore correspond to release of eggs by female Atlantic halibut. These bursts occurred while the halibut were in the water column, at water depths between 404.5 and 141 m, consistent with observations from Haug et al. (1984, 1986), who found a mesopelagic distribution of Atlantic halibut eggs in Norwegian Fjords. In the present study, we simply explored a general correspondence between sudden changes in the magnitude of acceleration and timing of presumed spawning rises, although accelerometer data archived in PSATs have previously been used to characterize swimming activity of Pacific (Nielsen et al., 2018) and Atlantic halibut (Gatti et al., 2021). Future studies could further investigate the potential of using accelerometer data to identify and characterize the spawning events of wild marine fish, but this may require the use of higher-frequency accelerometers than the one used in the present study (i.e., 0.2 Hz).

Directly observed flatfish have been seen completing spawning rises during which one female and one male swim together up into the water column, simultaneously or sequentially release eggs and a milt cloud, and then immediately return to the seafloor (Carvalho et al., 2003; Konstantinou & Shen, 1995; Manabe et al., 2000; Manabe & Shinomiya, 2001; Moyer et al., 1985). If halibut are performing spawning rises in pairs, the much higher number of male rises than female rises observed in the present study and the observation that male rises are not restricted to female egg batch intervals would support the interpretation that males are engaging in spawning rises with multiple females over the course of the presumed spawning period. Males may also be employing alternative reproductive tactics such as attempting to “sneak in” and fertilize eggs being spawned as a result of a paired rise between a female and another male, as has been seen in other species (Bekkevold et al., 2002; Habrun & Sancho, 2012; Taborsky, 2008). It is also possible that

some of the male rises seen on the time series may be part of courting behavior to attract females, as observed for the flounder *Bothus podas* (Carvalho et al., 2003).

An important question in the reproductive biology of batch-spawning species, relevant to fisheries assessment and management, is whether the number of eggs released by a female during the spawning season depends on fish size. Batch-spawning in marine species may be an adaptation to increase individual fecundity otherwise limited by body cavity size (Fordham & Trippel, 1999), or a bet-hedging strategy to increase long-term fitness by exposing offspring to a broader range of environmental conditions and thus reducing the risk of recruitment failure caused by variable environmental conditions (Lambert & Ware, 1984). Population dynamics predict that if the number of egg batches increases with female size, then a population with preserved size structure (i.e., presence of larger older females) should have higher relative total egg production and/or be more resilient to environmental fluctuations (Hixon et al., 2014; Le Bris et al., 2015). A review of 41 articles evaluating relationships between the number of egg batches and species age or size found a positive relationship for 21 out of 33 species investigated (Fitzhugh et al., 2012). In the present study, we did not find a statistically significant relationship between the number of presumed spawning rises, assumed to correspond to the release of egg batches, and female body size, although our results do not reject the idea that spawning potential increases with fish size. Exponentially higher fecundity at larger body sizes (Barneche et al., 2018) may be expressed in other ways not interpretable on PSAT time series, such as a higher number of eggs per batch, larger egg size, or improved egg viability (Hixon et al., 2014). The size of presumed spawning females in the present study ranged from 130 to 202 cm fork length, and it is possible that the lack of females potentially maturing at a smaller size (halibut females mature at ~119 cm; DFO, 2021b) may have biased the analysis. However, previous studies of flatfish reproduction have demonstrated that fish size is linked to fecundity (Cooper et al., 2007; Rideout & Morgan, 2007, 2010; Schmitt & Skud, 1978), including for Atlantic halibut (Haug & Gulliksen, 1988a).

Phenological changes in fish reproduction caused by warming waters are increasingly documented, with warmer waters accelerating gonadal development, inducing earlier spawning (Haug & Gulliksen, 1988b; Kjesbu et al., 2010), and possibly shortening the spawning season (Rogers & Dougherty, 2019; Slesinger et al., 2021). In the present study, the date of the first presumed spawning rise observed in female halibut was negatively correlated with the temperature experienced during the prespawning season. This result contradicts the findings of the aquaculture study by Brown et al. (2006), which found halibut spawning period to be delayed after warmer ambient conditions. However, the halibut studied by Brown et al. (2006) were kept in overall warmer water than that which was recorded by PSATs for halibut in the wild, suggesting a possible threshold water temperature above which additional warming has no effect on spawning phenology (Rogers & Dougherty, 2019). The present study therefore suggests that the current rapid and steady increase in water temperature (~2°C increase in sea surface

temperature since the 1990s) observed in the Gulf of St. Lawrence (Galbraith et al., 2012; Galbraith et al., 2022) might lead to an earlier spawning season for Atlantic halibut, which could have consequences for larvae survival and subsequent recruitment to the fishery (Durant et al., 2007). However, our model of presumed spawning start date as a function of pre-season temperature, while significant, explained little of the total variance, suggesting that other factors are likely contributing to the variation in spawning time, including natural interindividual variability. This limited explanatory power might also reflect the fact that bottom temperatures in the Gulf channels are relatively constant with little seasonal variation throughout the year (Galbraith et al., 2022). Additionally, we also tested for the effects of temperature during the spawning period on the number of presumed spawning rises and found no relationship, suggesting that temperature does not affect the duration of the spawning period in halibut. This result is consistent with the results from Brown et al. (2006), who found a non-significant difference in annual spawning period length between two groups of Atlantic halibut kept at ambient and chilled water temperatures.

Importantly, our analysis of presumed spawning rises indicates a later peak spawning period (mid-February) within the Gulf of St. Lawrence than has been reported for the adjacent and separately managed stock on the Scotian Shelf and Grand Banks (October–January) (Armsworthy et al., 2014; Neilson et al., 1993). This temporal difference in peak spawning may be a mechanism contributing to the maintenance of subtle but statistically significant population genetic structuring quantified between these adjacent stocks in comparison across the Northwest Atlantic (Kess et al., 2021).

The findings of the present study demonstrate that although the spawning behavior of marine fish species can remain elusive due to the difficulty observing fish in their often dark and deep natural habitats, high-resolution time series recorded by electronic tags can enable a detailed characterization of otherwise unobservable presumed spawning behaviors, such as the unique rapid rises off the seafloor undertaken by female and male Atlantic halibut during the winter in the Gulf of St. Lawrence. By interpreting these rises as spawning behavior, we were able to identify with unprecedented precision the spawning period for Atlantic halibut in the Gulf of St. Lawrence, and also to evaluate some of the factors driving interindividual variability in halibut spawning time and duration. However, although ample evidence exists to support the assumption that these rises are spawning behaviors, without direct observation, there is always uncertainty. Using promising novel telemetry technologies, future work may validate interpretations of spawning behavior. By this, fisheries and ocean managers can gain the understanding of the reproductive behavior of valuable marine species that is necessary to the development of effective management and conservation measures.

AUTHOR CONTRIBUTION

Project conceptualization was undertaken by Arnault Le Bris, Jonathan A. D. Fisher, and Rachel C. Marshall. Data collection was overseen and performed by Arnault Le Bris and Jonathan A. D. Fisher;

formal data analysis was done by Rachel C. Marshall, with tag geolocation done by Paul Gatti and Arnault Le Bris. The original manuscript was written by Rachel C. Marshall, with Arnault Le Bris, Jonathan A. D. Fisher, Paul Gatti, and Dominique Robert contributing to the review and editing process. Dominique Robert was responsible for funding acquisition.

ACKNOWLEDGMENTS

We thank all the Gulf of St. Lawrence fish harvesters involved in project planning, tag deployment, and tag recovery. Major funding for this research was provided by the Natural Sciences and Engineering Research Council of Canada via the Strategic Partnership Grants for Projects Program's contributions to "Quantifying spatial dynamics and stock structure of Atlantic halibut within the Gulf of St. Lawrence to improve sustainable exploitation and management," and Dominique Robert was supported by the Canada Research Chairs programme.

FUNDING INFORMATION

Tag deployment and tag recovery operations between 2013 and 2016 were funded by the Newfoundland and Labrador Department of Fisheries and Aquaculture and by the Research and Development Corporation of Newfoundland and Labrador Ignite Grants to Jonathan A. D. Fisher and Dominique Robert. The Gulf-wide tagging deployment and recovery programmes of 2017–2018, and the postdoctoral position held by postgraduate and master's degree programme completed by Rachel C. Marshall, were largely supported by a Strategic Partnership Grant from the Natural Sciences and Engineering Research Council (NSERC) of Canada (grant number: STPGP 506993 002D 17), with further support from the John R. Evans Leaders Fund of the Canada Foundation for Innovation, the Department of Fisheries and Land Resources of Newfoundland and Labrador, Ressources Aquatiques Québec, and the Ministère de l'agriculture, des pêcheries et de l'alimentation du Québec. This research was also supported by the Ocean Frontier Institute, a network funded by the Canada First Research Excellence Fund. Dominique Robert was supported by the Canada Research Chair programme.

ORCID

Rachel C. Marshall  <https://orcid.org/0000-0001-8239-8953>

REFERENCES

- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, 19, 716–723. <https://doi.org/10.1109/TAC.1974.1100705>
- Armsworthy, S., Trzcinski, M. K., & Campana, S. (2014). Movements, environmental associations, and presumed spawning locations of Atlantic halibut (*Hippoglossus hippoglossus*) in the Northwest Atlantic determined using archival satellite pop-up tags. *Marine Biology*, 161, 645–656. <https://doi.org/10.1007/s00227-013-2367-5>
- Barneche, D. R., Robertson, D. R., White, C. R., & Marshall, D. J. (2018). Fish reproductive-energy output increases disproportionately with body size. *Science*, 360, 642–645. <https://doi.org/10.1126/science.aao6868>
- Bateman, A. J. (1948). Intra-sexual selection in *drosophila*. *Heredity*, 2, 349–368.

- Bekkevold, D., Hansen, M. M., & Loeschcke, V. (2002). Male reproductive competition in spawning aggregations of cod (*Gadus morhua*, L.). *Molecular Ecology*, 11, 91–102. <https://doi.org/10.1046/j.0962-1083.2001.01424.x>
- Björnsson, B. T., Halldórsson, Ó., Haux, C., Norberg, B., & Brown, C. L. (1998). Photoperiod control of sexual maturation of the Atlantic halibut (*Hippoglossus hippoglossus*): Plasma thyroid hormone and calcium levels. *Aquaculture*, 166, 117–140. [https://doi.org/10.1016/S0044-8486\(98\)00276-2](https://doi.org/10.1016/S0044-8486(98)00276-2)
- Block, B. A., Dewar, H., Blackwell, S. D., Williams, T., Prince, E., Farwell, C., Boustany, A., Teo, S., Seitz, A., Walli, A., & Fudge, D. (2001). Migratory movements, depth preferences, and thermal biology of Atlantic bluefin tuna. *Science*, 293, 1310–1314. <https://doi.org/10.1126/science.1061197>
- Block, B. A., Teo, S. L. H., Walli, A., Boustany, A., Stokesbury, M. J. W., Farwell, C. J., Weng, K. C., Dewar, H., & Williams, T. D. (2005). Electronic tagging and population structure of Atlantic bluefin tuna. *Nature*, 434, 1121–1127. <https://doi.org/10.1038/nature03463>
- Brawn, V. M. (1961). Reproductive behaviour of the cod (*Gadus callarias* L.). *Behaviour*, 18, 177–197. <https://doi.org/10.1163/156853961X00114>
- Brown, N. P., Shields, R. J., & Bromage, N. R. (2006). The influence of water temperature on spawning patterns and egg quality in the Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture*, 261, 993–1002. <https://doi.org/10.1016/j.aquaculture.2006.08.025>
- Burnham, K., & Anderson, D. (2002). *Model selection and multimodel inference: A practical information-theoretic approach* (2nd ed.). Springer.
- Cargnelli, L. M., Griesbach, S. J., & Morse, W. W. (1999). Atlantic halibut, *Hippoglossus hippoglossus*, life history and habitat characteristics. *Essential Fish Habitat Source Document NMFS-NE*, 125, 1–17.
- Carvalho, N., Afonso, P., & Santos, R. S. (2003). The harem mating system and mate choice in the wide-eyed flounder, *Bothus podas*. *Environmental Biology of Fishes*, 66, 249–258.
- Coleman, S. W., & Jones, A. G. (2011). Patterns of multiple paternity and maternity in fishes. *Biological Journal of the Linnean Society*, 103, 735–760. <https://doi.org/10.1111/j.1095-8312.2011.01673.x>
- Cooper, D. W., Maslennikov, K. P., & Gunderson, D. R. (2007). Natural mortality rate, annual fecundity, and maturity at length for Greenland halibut (*Reinhardtius hippoglossoides*) from the northeastern Pacific Ocean. *Fishery Bulletin*, 105, 296–304.
- DFO. (2021a). Stock Assessment of Gulf of St. Lawrence (4RST) Atlantic Halibut in 2020. DFO Canadian Science Advisory Secretariat Science Advisory Report 2021/034.
- DFO. (2021b). Stock Status Update of Atlantic Halibut (*Hippoglossus hippoglossus*) on the Scotian Shelf and Southern Grand Banks in NAFO Divisions 3NOPs4VWX5Zc for 2020. DFO Canadian Science Advisory Secretariat Science Response 2021/024.
- Durant, J. M., Hjermmann, D. Ø., Ottersen, G., & Stenseth, N. C. (2007). Climate and the match or mismatch between predator requirements and resource availability. *Climate Research*, 33, 271–283. <https://doi.org/10.3354/cr033271>
- Einfeldt, A. L., Kess, T., Messmer, A., Duffy, S., Wringe, B. F., Fisher, J., den Heyer, C., Bradbury, I. R., Ruzzante, D., & Bentzen, P. (2021). Chromosome level reference of Atlantic halibut *Hippoglossus hippoglossus* provides insight into the evolution of sexual determination systems. *Molecular Ecology Resources*, 21, 1686–1696. <https://doi.org/10.1111/1755-0998.13369>
- Finn, R. N., Ostby, G. C., Norberg, B., & Fyhn, H. J. (2002). In vivo oocyte hydration in Atlantic halibut (*Hippoglossus hippoglossus*): proteolytic liberation of free amino acids, and ion transport, are driving forces for osmotic water influx. *Journal of Experimental Biology*, 205, 211–224.
- Fisher, J. A. D., Robert, D., Le Bris, A., & Loher, T. (2017). Pop-up satellite archival tag (PSAT) temporal data resolution affects interpretations of spawning behaviour of a commercially important teleost. *Animal Biotelemetry*, 5, 1–10. <https://doi.org/10.1186/s40317-017-0137-8>
- Fitzhugh, G. R., Shertzer, K. W., Kellison, G. T., & Wyanski, D. M. (2012). Review of size- and age-dependence in batch spawning: Implications for stock assessment of fish species exhibiting indeterminate fecundity. *Fishery Bulletin*, 110, 413–425.
- Fordham, B. S. E., & Trippel, E. A. (1999). Feeding behaviour of cod (*Gadus morhua*) in relation to spawning. *Journal of Applied Ichthyology*, 15, 1–9. <https://doi.org/10.1046/j.1439-0426.1999.00098.x>
- Galbraith, P. S., Chassé, J., Dumas, J., Shaw, J.-L., Caverhill, C., Lefavre, D., & Lafleur, C. (2022). Physical Oceanographic Conditions in the Gulf of St. Lawrence during 2021. DFO Canadian Science Advisory Secretariat Science Research Document 2022/034, iv + 83.
- Galbraith, P. S., Chassé, J., Shaw, J.-L., Dumas, J., Caverhill, C., Lefavre, D., & Lafleur, C. (2020). Physical Oceanographic Conditions in the Gulf of St. Lawrence during 2019. DFO Canadian Science Advisory Secretariat Science Research Document 2020/030, iv + 84.
- Galbraith, P. S., Larouche, P., Chassé, J., & Petrie, B. (2012). Sea-surface temperature in relation to air temperature in the Gulf of St. Lawrence: Interdecadal variability and long term trends. *Deep Sea Research Part II: Topical Studies in Oceanography, Satellite Oceanography and Climate Change*, 77–80, 10–20. <https://doi.org/10.1016/j.jsr.2012.04.001>
- Gatti, P., Fisher, J. A. D., Cyr, F., Galbraith, P. S., Robert, D., & Le Bris, A. (2021). A review and tests of validation and sensitivity of geolocation models for marine fish tracking. *Fish and Fisheries*, 22, 1041–1066. <https://doi.org/10.1111/faf.12568>
- Gatti, P., Le Bris, A., Fisher, J. A. D., Marshall, R. C., & Robert, D. (2020). Stock scale electronic tracking of Atlantic halibut reveals summer site fidelity and winter mixing on common spawning grounds. *ICES Journal of Marine Science*, 77, 2890–2904.
- Ghinter, L., Anderson, C., Robert, D., Winkler, G., Bernatchez, L., & Audet, C. (2023). A first glimpse of larval ecology of halibut species in the Gulf of St. Lawrence, Canada. *Journal of Fish Biology*, 1–6, 712–717. <https://doi.org/10.1111/jfb.15298>
- Grabowski, T. B., Boswell, K. M., McAdam, B. J., Wells, R. J. D., & Marteinsdóttir, G. (2012). Characterization of Atlantic cod spawning habitat and behavior in Icelandic coastal waters. *PLoS One*, 7, e51321. <https://doi.org/10.1371/journal.pone.0051321>
- Grabowski, T. B., McAdam, B. J., Thorsteinsson, V., & Marteinsdóttir, G. (2015). Evidence from data storage tags for the presence of lunar and semi-lunar behavioral cycles in spawning Atlantic cod. *Environmental Biology of Fishes*, 98, 1767–1776. <https://doi.org/10.1007/s10641-015-0396-2>
- Grabowski, T. B., Thorsteinsson, V., & Marteinsdóttir, G. (2014). Spawning behavior in Atlantic cod: Analysis by use of data storage tags. *Marine Ecology Progress Series*, 506, 279–290. <https://doi.org/10.3354/meps10787>
- Habrún, C. A., & Sancho, G. (2012). Spawning ascent durations of pelagic spawning reef fishes. *Current Zoology*, 58, 95–102. <https://doi.org/10.1093/czoolo/58.1.95>
- Haug, T. (1990). Biology of the Atlantic halibut, *Hippoglossus hippoglossus* (L., 1758). In J. H. S. Blaxter & A. J. Southward (Eds.), *Advances in marine biology* (pp. 1–70). Academic Press. [https://doi.org/10.1016/S0065-2881\(08\)60198-4](https://doi.org/10.1016/S0065-2881(08)60198-4)
- Haug, T., & Gulliksen, B. (1988a). Fecundity and oocyte sizes in ovaries of female Atlantic halibut, *Hippoglossus hippoglossus* (L.). *Sarsia*, 73, 259–261. <https://doi.org/10.1080/00364827.1988.10413411>
- Haug, T., & Gulliksen, B. (1988b). Variation in liver and body condition during gonad development of Atlantic halibut, *Hippoglossus hippoglossus* (L.). *Fiskeridirektoratets Skrifter Serie Havundersøkelser*, 18, 351–363.
- Haug, T., Kjörsvik, E., & Solemdal, P. (1984). Vertical distribution of Atlantic halibut (*Hippoglossus hippoglossus*) eggs. *Canadian Journal of Fisheries and Aquatic Sciences*, 41, 798–804. <https://doi.org/10.1139/f84-092>
- Haug, T., Kjörsvik, E., & Solemdal, P. (1986). Influence of some physical and biological factors on the density and vertical distribution of

- Atlantic halibut *Hippoglossus hippoglossus* eggs. *Marine Ecology Progress Series*, 33, 207–216. <https://doi.org/10.3354/meps033207>
- Hixon, M. A., Johnson, D. W., & Sogard, S. M. (2014). BOFFFFs: On the importance of conserving old-growth age structure in fishery populations. *ICES Journal of Marine Science*, 71, 2171–2185. <https://doi.org/10.1093/icesjms/fst200>
- Hussey, N. E., Kessel, S. T., Aarestrup, K., Cooke, S. J., Cowley, P. D., Fisk, A. T., Harcourt, R. G., Holland, K. N., Iverson, S. J., Kocik, J. F., Flemming, J. E. M., & Whoriskey, F. G. (2015). Aquatic animal telemetry: A panoramic window into the underwater world. *Science*, 348. <https://doi.org/10.1126/science.1255642>
- James, T. L., Landsman, S. J., Ramsay, L. L., Giffin, M. D., Le Bris, A., & van den Heuvel, M. R. (2020). Migration patterns of Atlantic halibut captured in the southern gulf of St. Lawrence as assessed with pop-up satellite archival and floy tags. *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 1233–1242. <https://doi.org/10.1139/cjfas-2019-0262>
- Kess, T., Einfeldt, A. L., Wringe, B., Lehnert, S. J., Layton, K. K. S., McBride, M. C., Robert, D., Fisher, J., Le Bris, A., den Heyer, C., Shackell, N., Ruzzante, D. E., Bentzen, P., & Bradbury, I. R. (2021). A putative structural variant and environmental variation associated with genomic divergence across the Northwest Atlantic in Atlantic halibut. *ICES Journal of Marine Science*, 78, 2371–2384. <https://doi.org/10.1093/icesjms/fsab061>
- Kjesbu, O. S., Righton, D., Krüger-Johnsen, M., Thorsen, A., Michalsen, K., Fonn, M., & Witthames, P. R. (2010). Thermal dynamics of ovarian maturation in Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences*, 67, 605–625. <https://doi.org/10.1139/F10-011>
- Kjørsvik, E., Haug, T., & Tjemsland, J. (1987). Spawning season of the Atlantic halibut (*Hippoglossus hippoglossus*) in northern Norway. *ICES Journal of Marine Science*, 43, 285–293. <https://doi.org/10.1093/icesjms/43.3.285>
- Kohler, A. C. (1967). Size at maturity, spawning season, and food of Atlantic halibut. *Journal of the Fisheries Research Board of Canada*, 24, 53–66. <https://doi.org/10.1139/f67-006>
- Konstantinou, H., & Shen, D. C. (1995). The social and reproductive behavior of the eyed flounder, *Bothus ocellatus*, with notes on the spawning of *Bothus lunatus* and *Bothus ellipticus*. *Environmental Biology of Fishes*, 44, 311–324. <https://doi.org/10.1007/BF00008245>
- Lambert, T. C., & Ware, D. M. (1984). Reproductive strategies of demersal and pelagic spawning fish. *Canadian Journal of Fisheries and Aquatic Sciences*, 41, 1565–1569. <https://doi.org/10.1139/f84-194>
- Le Bris, A., Fisher, J. A. D., Murphy, H. M., Galbraith, P. S., Castonguay, M., Loher, T., & Robert, D. (2018). Migration patterns and putative spawning habitats of Atlantic halibut (*Hippoglossus hippoglossus*) in the Gulf of St. Lawrence revealed by geolocation of pop-up satellite archival tags. *ICES Journal of Marine Science*, 75, 135–147. <https://doi.org/10.1093/icesjms/fsx098>
- Le Bris, A., Fréchet, A., & Wroblewski, J. S. (2013). Supplementing electronic tagging with conventional tagging to redesign fishery closed areas. *Fisheries Research*, 148, 106–116. <https://doi.org/10.1016/j.fishres.2013.08.013>
- Le Bris, A., Pershing, A. J., Hernandez, C. M., Mills, K. E., & Sherwood, G. D. (2015). Modelling the effects of variation in reproductive traits on fish population resilience. *ICES Journal of Marine Science*, 72, 2590–2599. <https://doi.org/10.1093/icesjms/fsv154>
- Liu, C., Bank, C., Kersula, M., Cowles, G. W., Zemeckis, D. R., Cadrin, S. X., & McGuire, C. (2019). Movements of Atlantic halibut in the Gulf of Maine based on geolocation. *ICES Journal of Marine Science*, 76, 2020–2032. <https://doi.org/10.1093/icesjms/fsz169>
- Loher, T., & Stephens, S. M. (2011). Use of veterinary ultrasound to identify sex and assess female maturity of Pacific halibut in nonspawning condition. *North American Journal of Fisheries Management*, 31, 1034–1042. <https://doi.org/10.1080/02755947.2011.635240>
- Manabe, H., Ide, M., & Shinomiya, A. (2000). Mating system of the lefteye flounder, *Engyprosopon grandisquama*. *Ichthyological Research*, 47, 69–74. <https://doi.org/10.1007/BF02674315>
- Manabe, H., & Shinomiya, A. (2001). Two spawning seasons and mating system of the bastard halibut, *Tarphops oligolepis*. *Ichthyological Research*, 48, 421–424. <https://doi.org/10.1007/s10228-001-8167-9>
- Marshall, R. C. (2020). *Multidisciplinary perspectives on Atlantic halibut spawning behavior and vulnerability in the Gulf of St Lawrence* (MSc in Fisheries Science). Memorial University of Newfoundland, Newfoundland and Labrador.
- Marshall, R. C., Fisher, J. A. D., Gatti, P., Carruthers, E. H., Robert, D., & Le Bris, A. (2023). Combining electronic tag data and fish harvesters' knowledge to identify ocean management concerns. *Marine Policy*, 147, 105393. <https://doi.org/10.1016/j.marpol.2022.105393>
- McBride, R. S., Maynard, G. A., Elzey, S. P., Hennen, D. R., Tholke, E. K., Runnebaum, J. M., & McGuire, C. H. (2022). Evaluating growth dimorphism, maturation, and skip spawning of Atlantic halibut in the Gulf of Maine using a collaborative research approach. *Journal of Northwest Atlantic Fishery Science*, 53, 57–77. <https://doi.org/10.2960/J.v53.m736>
- McBride, R. S., Somarakis, S., Fitzhugh, G. R., Albert, A., Yaragina, N. A., Wuenschel, M. J., Alonso-Fernández, A., & Basilone, G. (2015). Energy acquisition and allocation to egg production in relation to fish reproductive strategies. *Fish and Fisheries*, 16, 23–57. <https://doi.org/10.1111/faf.12043>
- McCracken, F. D. (1958). On the biology and fishery of the Canadian Atlantic halibut, *Hippoglossus hippoglossus* L. *Journal of the Fisheries Research Board of Canada*, 15, 1269–1311. <https://doi.org/10.1139/f58-070>
- Methven, D. A., Crim, L. W., Norberg, B., Brown, J. A., Goff, G. P., & Huse, I. (1992). Seasonal reproduction and plasma levels of sex steroids and vitellogenin in Atlantic halibut (*Hippoglossus hippoglossus*). *Canadian Journal of Fisheries and Aquatic Sciences*, 49, 754–759. <https://doi.org/10.1139/f92-084>
- Moyer, J. T., Yogo, Y., Zaiser, M. J., & Tsukahara, H. (1985). Spawning behavior and social organization of the flounder *Crossorhombus kobensis* (Bothidae) at Miyake-jima, Japan. *Japanese Journal of Ichthyology*, 32, 363–367. <https://doi.org/10.1007/BF02905443>
- Murphy, H. M., Fisher, J. A. D., Bris, A. L., Desgagnés, M., Castonguay, M., Loher, T., & Robert, D. (2017). Characterization of depth distributions, temperature associations, and seasonal migrations of Atlantic halibut in the Gulf of St. Lawrence using pop-up satellite archival tags. *Marine and Coastal Fisheries*, 9, 1–356. <https://doi.org/10.1080/19425120.2017.1327905>
- Neilson, J. D., Kearney, J. F., Perley, P., & Sampson, H. (1993). Reproductive biology of Atlantic halibut (*Hippoglossus hippoglossus*) in Canadian waters. *Canadian Journal of Fisheries and Aquatic Sciences*, 50, 551–563. <https://doi.org/10.1139/f93-064>
- Nielsen, J. K., Rose, C. S., Loher, T., Drobny, P., Seitz, A. C., Courtney, M. B., & Gauvin, J. (2018). Characterizing activity and assessing bycatch survival of Pacific halibut with accelerometer pop-up satellite archival tags. *Animal Biotelemetry*, 6, 10. <https://doi.org/10.1186/s40317-018-0154-2>
- Norberg, B., Valkner, V., Huse, J., Karlsen, I., & Grung, G. L. (1991). Ovulatory rhythms and egg viability in the Atlantic halibut (*Hippoglossus hippoglossus*). *Aquaculture*, 97, 365–371. [https://doi.org/10.1016/0044-8486\(91\)90328-5](https://doi.org/10.1016/0044-8486(91)90328-5)
- Norberg, B., Weltzien, F.-A., Karlsen, Ø., & Holm, J. C. (2001). Effects of photoperiod on sexual maturation and somatic growth in male Atlantic halibut (*Hippoglossus hippoglossus* L.). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 4TH International Symposium on Fish Endocrinology, 129, 357–365. [https://doi.org/10.1016/S1096-4959\(01\)00320-7](https://doi.org/10.1016/S1096-4959(01)00320-7)
- Pepin, P., & Helbig, J. A. (1997). Distribution and drift of Atlantic cod (*Gadus morhua*) eggs and larvae on the Northeast Newfoundland Shelf.

- Canadian Journal of Fisheries and Aquatic Sciences*, 54, 670–685. <https://doi.org/10.1139/cjfas-54-3-670>
- Rideout, R. M., & Morgan, M. J. (2007). Major changes in fecundity and the effect on population egg production for three species of north-West Atlantic flatfishes. *Journal of Fish Biology*, 70, 1759–1779. <https://doi.org/10.1111/j.1095-8649.2007.01448.x>
- Rideout, R. M., & Morgan, M. J. (2010). Relationships between maternal body size, condition and potential fecundity of four north-west Atlantic demersal fishes. *Journal of Fish Biology*, 76, 1379–1395. <https://doi.org/10.1111/j.1095-8649.2010.02570.x>
- Rogers, L. A., & Dougherty, A. B. (2019). Effects of climate and demography on reproductive phenology of a harvested marine fish population. *Global Change Biology*, 25, 708–720. <https://doi.org/10.1111/gcb.14483>
- Rose, G. A. (1993). Cod spawning on a migration highway in the north-West Atlantic. *Nature*, 366, 458–461. <https://doi.org/10.1038/366458a0>
- Schmitt, C. C., & Skud, B. E. (1978). *Relation of fecundity to long-term changes in growth, abundance and recruitment* (Vol. 66, pp. 1–31). International Pacific Halibut Commission Scientific Report.
- Seitz, A. C., Norcross, B. L., Wilson, D., & Nielsen, J. L. (2005). Identifying spawning behavior in Pacific halibut, *Hippoglossus stenolepis*, using electronic tags. *Environmental Biology of Fishes*, 73, 445–451. <https://doi.org/10.1007/s10641-005-3216-2>
- Seitz, A. C., Wilson, D., Norcross, B. L., & Nielsen, J. L. (2003). Pop-up archival transmitting (PAT) tags: A method to investigate the migration and behavior of Pacific halibut (*Hippoglossus stenolepis*) in the Gulf of Alaska. *Alaska Fishery Research Bulletin*, 10, 124–136.
- Shackell, N. L., Fisher, J. A. D., den Heyer, C. E., Hennen, D. R., Seitz, A. C., Le Bris, A., Robert, D., Kersula, M. E., Cadrin, S. X., McBride, R. S., McGuire, C. H., Kess, T., Ransier, K. T., Liu, C., Czich, A., & Frank, K. T. (2022). Spatial ecology of Atlantic halibut across the Northwest Atlantic: A recovering species in an era of climate change. *Reviews in Fisheries Science & Aquaculture*, 30, 281–305. <https://doi.org/10.1080/23308249.2021.1948502>
- Siwicke, K. A., Seitz, A. C., Rodgveller, C. J., & Echave, K. B. (2022). Characterizing spawning behavior of Greenland halibut (*Reinhardtius hippoglossoides*) in the eastern Bering Sea and Aleutian Islands. *Fishery Bulletin*, 120, 55–67. <https://doi.org/10.7755/FB.120.1.5>
- Slesinger, E., Jensen, O. P., & Saba, G. (2021). Spawning phenology of a rapidly shifting marine fish species throughout its range. *ICES Journal of Marine Science*, 78, 1010–1022. <https://doi.org/10.1093/icesjms/fsaa252>
- Smith, T. I. J., McVey, D. C., Jenkins, W. E., Denson, M. R., Heyward, L. D., Sullivan, C. V., & Berlinsky, D. L. (1999). Broodstock management and spawning of southern flounder, *Paralichthys lethostigma*. *Aquaculture*, 176, 87–99. [https://doi.org/10.1016/S0044-8486\(99\)00053-8](https://doi.org/10.1016/S0044-8486(99)00053-8)
- Taborsky, M. (2008). Alternative reproductive tactics in fish. In R. F. Oliveira, M. Taborsky, & H. J. Brockmann (Eds.), *Alternative reproductive tactics: An integrative approach* (pp. 251–299). Cambridge University Press.
- Tsuda, Y., Kawabe, R., Tanaka, H., Mitsunaga, Y., Hiraishi, T., Yamamoto, K., & Nashimoto, K. (2006). Monitoring the spawning behaviour of chum salmon with an acceleration data logger. *Ecology of Freshwater Fish*, 15, 264–274. <https://doi.org/10.1111/j.1600-0633.2006.00147.x>
- Wearmouth, V. J., Southall, E. J., Morritt, D., & Sims, D. W. (2013). Identifying reproductive events using archival tags: Egg-laying behaviour of the small spotted catshark *Scyliorhinus canicula*. *Journal of Fish Biology*, 82, 96–110. <https://doi.org/10.1111/j.1095-8649.2012.03473.x>
- Yasuda, T., Katsumata, H., Kawabe, R., Nakatsuka, N., & Kurita, Y. (2013). Identifying spawning events in the Japanese flounder *Paralichthys olivaceus* from depth time-series data. *Journal of Sea Research*, Proceedings of the 8th International Symposium on Flatfish Ecology, Part I, 75, 33–40. <https://doi.org/10.1016/j.seares.2012.04.001>
- Zemeckis, D. R., Dean, M. J., & Cadrin, S. X. (2014). Spawning dynamics and associated management implications for Atlantic cod. *North American Journal of Fisheries Management*, 34, 424–442. <https://doi.org/10.1080/02755947.2014.882456>
- Zuur, A. F., & Ieno, E. N. (2016). A protocol for conducting and presenting results of regression-type analyses. *Methods in Ecology and Evolution*, 7, 636–645. <https://doi.org/10.1111/2041-210X.12577>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Marshall, R. C., Fisher, J. A. D., Einfeldt, A. L., Gatti, P., Robert, D., & Le Bris, A. (2023). Reproductive behavior of Atlantic halibut (*Hippoglossus hippoglossus*) interpreted from electronic tags. *Journal of Fish Biology*, 103(5), 1031–1043. <https://doi.org/10.1111/jfb.15501>