

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/369033362>

Physically Constrained Covariance Inflation from Location Uncertainty

Preprint · March 2023

DOI: 10.13140/RG.2.2.18433.53603

CITATIONS

0

READS

178

3 authors, including:



Yicun Zhen

College of Oceanography, Hohai University

20 PUBLICATIONS 161 CITATIONS

[SEE PROFILE](#)



Valentin Resseguier

Scalian

21 PUBLICATIONS 430 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Red LUM (Reduced Location Uncertainty Model) [View project](#)



Full LUM (Full Location Uncertainty Model) [View project](#)

Physically Constrained Covariance Inflation from Location Uncertainty

Yicun Zhen¹, Valentin Resseguier^{2,3}, and Bertrand Chapron⁴

¹College of Oceanography, Hohai University, Nanjing, China

²LAB SCALIAN DS, Rennes, France

³INRAE, OPAALE, Rennes, France

⁴Laboratoire d’Océanographie Physique et Spatiale, Ifremer, Plouzaé, France

Correspondence: Yicun Zhen (zhenyicun@proton.me)

Abstract. Motivated by the concept of “location uncertainty”, initially introduced in Mémin (2014), a scheme is sought to perturb the “location” of a state variable at every forecast time step. Further considering Brenier’s theorem Brenier (1991), asserting that the difference of two positive density fields on the same domain can be represented by a transportation map, perturbations are demonstrated to consistently define a SPDE from the original PDE. It ensues that certain quantities, up to the user, are conserved at every time step. Remarkably, derivations following both the SALT Holm (2015) and LU Mémin (2014); Resseguier et al. (2016) settings, can be recovered from this perturbation scheme. Still, it opens broader applicability since it does not explicitly rely on Lagrangian mechanics or Newton’s laws of force. For illustration, a stochastic version of the thermal shallow water equation is presented.

Copyright statement. TEXT

10 1 Introduction

Data assimilation is meant to extract information from measurements to improve the state estimate. Kalman-filter-based and particle-filter-based methods are now commonly used for academical studies and operational forecasts. For both methods, the estimate of state variable and the uncertainty quantification of the state estimate are repeated at each data assimilation cycle. In the classical Kalman filter, this uncertainty is represented by a covariance matrix. In Monte-Carlo-based methods (i.e. the ensemble Kalman filters and particle filters, etc.), it is represented by the spread of the ensemble members or particles. The uncertainty of the state estimate is further part of the input for the next data assimilation cycle. Frequently observed, the uncertainty can be underestimated in nonlinear numerical experiments when there is no model noise Schlee et al. (1966); Harlim and Majda (2010); Franzke et al. (2015). As a consequence, the state estimate in the subsequent time steps may not be efficiently adjusted by the physical measurements: the system is over-confident about its current state estimate. This phenomenon is usually referred to as filter divergence, possibly associated to the “curse of dimensionality”.

To address the latter issue, "covariance localization" has been developed for both Kalman-filter-based methods and particle filters Houtekamer and Mitchell (2001); Poterjoy (2016). To further mitigate filter divergence, a practical strategy is to inflate the uncertainty estimate at each forecast time step or each data assimilation cycle Anderson (2007); Tibshirani and Knight (1999); Li et al. (2009); Kotsuki et al. (2017); Ying and Zhang (2015); Miyoshi (2011); Raanes et al. (2019); Zhen and Harlim
25 (2015). For geophysical applications, the uncertainty is then often inflated by rescaling the ensemble covariance in order to match bias and variance. A natural alternative is the addition of noises in the dynamical equations.

In the context of ensemble/particle-based methods, the uncertainty is usually inflated by artificially perturbing each ensemble member/particle. We refer the reader to Resseguier et al. (2021) for a review on the subject. It is then a natural question to ask: is there a mathematical principle to guide this uncertainty inflation? In the fluid dynamics community, random forcings are not
30 introduced for inflation, but to mimic the intermittent back-scattering of energy from small scales toward large scales. Among those approaches, we may mention the stochastic Lagrangian models Pope (1994) and the Eulerian Gaussian backscatterings of EDQNM Orszag (1970); Leith (1971). Additive noise models, like the linear inverse models Penland and Sardeshmukh (1995), have then also been proposed for filtering purposes, and thoroughly reviewed by Tandeo et al. (2020). Most methods
35 mainly focus on comparing the estimated uncertainty and the statistics of the innovation process, but ignore other mathematical/physical aspects (for instance, the conservation laws, etc.). Other empirical approaches, referred to as SPPT Buizza et al. (1999) and SKEBS Berner et al. (2009), introduce multiplicative noises, with success in operational weather and climate forecast centers Franzke et al. (2015). Still many drawbacks have been reported, above all violations of conservation laws Reynolds et al. (2016); Leutbechner et al. (2016). Recently, the operational ocean circulation model NEMO has also been randomized (e.g. Leroux et al., 2022), but again, without conservative considerations.

Several authors proposed schemes specifically to enforce energy conservation or at least a given energy budget (e.g. Sapsis and Majda, 2013; Gugole and Franzke, 2019; Resseguier et al., 2021). To better constrain non-Gaussian schemes, many authors
40 rely on physics and possibly on time-scale separation. Introduced by Hasselmann (1976), it is generally associated with the rigorous theories of averaging and homogenization. Majda et al. (1999) decomposed the state variable into slowly-varying modes x_j and fast-varying modes y_j . The authors demonstrated that the interaction term between x_j and y_j , in the equation
45 for x_j , can be modeled as a stochastic process solely in terms of x_j 's, as the ratio of the time scales of x_j and y_j tends to 0. Nevertheless, homogenization methods, like Majda et al. (1999), may also lead to violation of energy conservation, even though some workarounds exist Gottwald and Melbourne (2013); Jain et al. (2014).

In Brzeźniak et al. (1991), later modified in Mikulevicius and Rozovskii (2004); Flandoli (2011) and Mémin (2014); Resseguier et al. (2016, 2021), preservation of kinetic energy is specifically emphasized. The true velocity of an incompressible
50 flow is decomposed into a regular component and a turbulent one, and the latter modeled by a stochastic noise. Mikulevicius and Rozovskii (2004) and Mémin (2014) further derived stochastic Navier-Stokes equations. For these two approaches, the large-scale advecting velocity differs, induced by different regularisation of the Newton' second law. Following an other path, considering the Hamilton's principle with a stochastic advection constraint on Lagrangian fluid trajectories, Holm (2015) also proposed a consistent stochastic setting, i.e stochastic advection by Lie transport (SALT). In particular, this derivation preserves

55 Kelvin’s circulation. Similarities and differences between these different stochastic frameworks are discussed in Resseguier et al. (2020).

From another perspective, the classical optimal transport theory suggests that the difference of two smooth positive density fields (ρ_1 and ρ_2) on a bounded domain Ω can be described by a transportation map: $T : \Omega \rightarrow \Omega$. More specifically, there exists a diffeomorphism T of Ω to transform ρ_1 to ρ_2 under the diffeomorphism T with a minimal cost. Broadly speaking, T can
60 be interpreted as how much ρ_2 differs from ρ_1 , and T operates as a location correction. Indeed, starting from the same initial condition $\rho(t)$, suppose that $\rho_1 = \rho^{\text{model}}(t + \Delta t)$ is the model forecast and $\rho_2 = \rho(t + \Delta t)$ is the true forecast. The additional uncertainty of ρ_1 due to model error can then be represented by a random T . It further suggests that the inflation of uncertainty can be achieved by casting a random T on each ensemble member/particle.

Motivated by such an optimal transport perspective and the concept of “location uncertainty”, proposed in Mémin (2014), a
65 new strategy can thus seek to design a well constrained “location perturbation” of the state variable. Specifically, the idea of covariance inflation can be informally generalized to physical fields that are not always positive, i.e. physical fields other than the density field. Mathematically, a density field ρ is naturally associated to a differential n -form θ_ρ , where $n = \dim \Omega$. The statement “ ρ_1 transforms to ρ_2 under the diffeomorphism T ” is equivalent to the mathematical relation $\theta_{\rho_1} = T^* \theta_{\rho_2}$, where T^* , acting on all differential forms, is the pull-back operator induced by T , or equivalently, $\theta_{\rho_2} = (T^{-1})^* \theta_{\rho_1}$. Therefore, a
70 random T (or equivalently, T^{-1}) could induce a perturbation of any differential k -form.

To implement a physically-constrained perturbation scheme, the state variable S under consideration must then be associated to some differential form θ , i.e. construct a 1-1 correspondence between snapshots of S and snapshots of θ . Note, this can be generalized to other types of tensor fields. It will be demonstrated (section 5) that it is indeed sometimes helpful to choose θ to be a contravariant tensor field other than differential forms. Yet, it must be stressed that associating the state variable S to a
75 differential form θ is a key important step.

Correspondingly, at each forecast time step, the covariance inflation should follow 4 steps:

- Step 1, find $\theta(t)$ based on $S(t)$.
- Step 2, construct a random diffeomorphism $T : \Omega \rightarrow \Omega$.
- Step 3, replace $\theta(t)$ with $T^* \theta(t)$ and calculate $S(t)$ based on the new value of $\theta(t)$.
- 80 – Step 4, calculate the forecast $S(t + \Delta t)$ based on the new value of $S(t)$.

Associating S to different θ shall then be constrained by different conservation laws for the perturbation scheme. More precisely, certain physical quantities are conserved in step 3, no matter how T is constructed or realized in step 2. We emphasize that the conservation law of the perturbation scheme merely depends on the choice of θ , but is independent of the dynamics of the original deterministic system. A resulting SPDE will conserve a given quantity only if both the perturbation scheme
85 and the original deterministic system conserve that quantity. We also remark that this scheme can not conserve all the physical quantities at the same time unless additional constraints upon the parameters are imposed. Hence the users must choose by themselves which physical quantity to conserve.

In sum, this manuscript provides with the perspective that the displacement vector field of physical state fields should be determined by the tensor fields associated to the physical fields. The advantage of this perspective is that certain physical quantities can be conserved while applying a displacement vector field to transfer the original physical field. A direct application of this perspective is the physically constrained covariance inflation scheme proposed in this manuscript. When the tensor fields are positive n -forms on a bounded domain that have the same total mass, Brenier's theorem shows that the 'optimal' displacement vector field exists and is unique, for a given cost function. In this case, the optimality of displacement vector field is well-defined. In other cases, the issue of 'optimality' together with the existence and uniqueness of 'optimal' displacement vector field need to be carefully explored. We reserve this to the future study.

This paper is organized as follows. Section 2 is a brief introduction of optimal transport theory. In section 3 we present the perturbation scheme in detail, including the motivation, the specific techniques in derivation, and several examples. In section 4, the resulting perturbation scheme is then compared with the stochastic advection by Lie transport (SALT) equations Holm (2015) and the location uncertainty (LU) equations Mémín (2014). For properly chosen θ and T_t , it is demonstrated that both SALT and LU settings are recovered within the proposed framework. To illustrate our purpose, a stochastic version of the thermal shallow water equation is then derived in section 5. Final conclusion and discussion are given in section 6.

Convention of notation:

- The letter i only refer to the i -th independent Brownian motion. The letters p, q, j, k refer to the components if p, q, j, k are upper indices.
- Einstein's convention on summation (applies to all indices except i, j): if indice p show in both upper and lower indices, then the summation over p automatically applies.
- Summation over i, j, p automatically applies in all equations. For instance, e_i refers to $\sum_i e_i$, and y_j refers to $\sum_j y_j$

2 Monge's formulation of optimal transport problem and Brenier's answer

Hereafter we briefly summarize some necessary concepts and results in optimal transport theory. Let Ω be a bounded domain in a n -dimensional Euclidean space.

Definition 2.0.1 (Monge's optimal transport problem). *Given cost function $c(x, y) \geq 0$ and probability measures $\mu, \nu \in \mathcal{P}(\Omega)$,*

$$\text{minimize } \mathbb{M}(T) = \int_{\Omega} c(x, T(x)) d\mu(x) \tag{1}$$

over μ measurable maps $T : \Omega \rightarrow \Omega$ subject to $\nu = T_{\#}\mu$.

Here the probability measures μ and ν are interpreted as mass distributions with total mass equal to 1. The map T is called a transport plan which moves the mass $d\mu(x)$ at location x to location $T(x)$, with the cost $c(x, T(x))$ per unit of mass. Therefore the quantity $\mathbb{M}(T)$ is the total cost of the transport plan T . The constraint $\nu = T_{\#}\mu$ is interpreted as that T transports the mass

distribution μ to the mass distribution ν . In the case that T is a diffeomorphism and that both ν and μ have smooth densities, i.e. assume that $d\nu(x) = f(x)d^n x$ and $d\mu(x) = g(x)d^n x$ for some smooth functions f, g on Ω ,

$$\nu = T_{\#}\mu \iff g(x) = f(T(x))|J_T(x)|, \quad (2)$$

120 where $J_T(x)$ refers to the Jacobian matrix of T at x . If we associate ν and μ to differential n -forms $\theta_\nu = f dx^1 \wedge \dots \wedge dx^n$ and $\theta_\mu = g dx^1 \wedge \dots \wedge dx^n$, then

$$\nu = T_{\#}\mu \iff \theta_\mu = T^*\theta_\nu. \quad (3)$$

Brenier Brenier (1991) proved the existence and uniqueness of the solution to the Monge's optimal transport problem for $c(x, y) = |x - y|^2$. To better illustrate how optimal transport theory motivates us, we consider the following simplified version of Brenier's theorem.

Theorem 2.1 (Brenier, simplified version). *Let μ and ν be measures with bounded smooth density on a bounded domain $\Omega \subset \mathbb{R}^n$. Let $c(x, y) = |x - y|^2$. Then there is a convex function $\phi : \Omega \rightarrow \mathbb{R}$, such that $(\nabla\phi)_{\#}\mu = \nu$. And $\nabla\phi : x \rightarrow x + \nabla\phi|_x$, defined μ -almost everywhere, is the unique solution to the Monge's optimal transport problem.*

The convexity of ϕ implies that the map $\nabla\phi$ is one-to-one. Broadly speaking, Brenier's theorem implies that the difference of two density fields can be represented by a transportation map T .

3 The Perturbation Scheme

Consider a compressible flow on a bounded domain Ω . Let ρ denote the density field. Let $\rho^{\text{model}}(t + \Delta t)$ and $\rho^{\text{true}}(t + \Delta t)$ be the model forecast and the true forecast starting from the same density field at time t . If we assume that the model forecast and the truth have the same total mass, Brenier's theorem says that there exists a diffeomorphism $T : \Omega \rightarrow \Omega$ so that

$$135 \quad \rho^{\text{true}}(x, t + \Delta t) = \rho^{\text{model}}(T(x), t + \Delta t)J_T(x). \quad (4)$$

Note that the transportation T hereinafter is equivalent to the mapping T^{-1} used in the introduction. Eq.(4) can further be written in terms of differential form. Let $\theta_\rho = \rho dx^1 \wedge \dots \wedge dx^n$, then Eq.(4) is equivalent to

$$T^*\theta_\rho^{\text{model}}(t + \Delta t) = \theta_\rho^{\text{true}}(t + \Delta t). \quad (5)$$

For general differential forms θ , it is unclear whether a diffeomorphism T always exists that satisfies Eq.(5). However, Eq.(5) provides us with a tool for covariance inflation by constructing a random T at every infinitesimal time step. At each time step we construct a small perturbation T :

$$T_t(x) = x + a(t, x)\Delta t + e_i(t, x)\Delta\eta_i(t), \quad (6)$$

where $a(t, x), e_i(t, x) \in \mathbb{R}^n$, $\Delta\eta_i(t) \sim \mathcal{N}(0, \Delta t)$ is a random number. Essentially, $T_t(x) - x$ can be interpreted as a "location error" caused by the model error. In Eq.(6), $a(t, x)\Delta t$ refers to a systematic location error, and $e_i\Delta\eta_i$ refers to a random location

145 error. Stated in the introduction, the state variable S must first be associated to a differential form θ . Then at every time step, T_t induces a perturbation of $\theta(t)$ by $\theta(t) \rightarrow T_t^* \theta(t)$. It hence induces a perturbation of the state variable $S(t)$. A forecast is then performed based on the perturbed state. Consequently, this perturbation scheme derives a SPDE from the original PDE.

This procedure can also be generalized to other types of tensor fields. We refer to Chern et al. (1999) for a rigorous definition of the tensor fields and the wedge algebra. For instance, we may choose $\theta = \rho \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^n}$, where $\{\frac{\partial}{\partial x^i}\}_{i \leq n}$ forms a global
 150 basis of the tangent field. Then T_t induces a perturbation of θ by $\theta(t) \rightarrow T_{t*} \theta$, where T_{t*} is the push-forward operator induced by T_t . In section 5, such a generalization is found useful in the example of thermal shallow water equation.

Remark 1. When θ is a mixture of covariant and contravariant tensor fields, the perturbation scheme is slightly more complicated. Assume that $T_t : \Omega_1 \rightarrow \Omega_2$ is a diffeomorphism, and $\theta = v \otimes \omega$ where v and ω are contravariant or covariant tensor fields respectively on Ω_2 . Then $T_t^* \omega$ is a covariant tensor field on Ω_1 . However, T_t can not directly induce a contravariant
 155 tensor field on Ω_1 . In order to get a tensor field on Ω_1 , we consider $T_t^{-1} : \Omega_2 \rightarrow \Omega_1$, and apply the push-forward operator on v . In sum, we may define the perturbation to be

$$\theta(t) \rightarrow \left((T_t^{-1})_* v \right) \otimes \left(T_t^* \omega \right). \quad (7)$$

Appendix A derives the expression of T_t^{-1} directly from the expression of T_t .

3.1 Calculation of $T_t^* \theta$ (or $T_{t*} \theta$)

160 A rigorous mathematical definition and calculation of T_t and T_t^* should be given in terms of stochastic flows of diffeomorphisms and its Lie derivatives. A brief discussion of the relationship between T_t^* and the Lie derivative is given in section 4.1. We further refer to Leon (2021) for detailed definition of the Lie derivative. Yet, to rapidly assess $T_t^* \theta$ (or $T_{t*} \theta$), a Taylor expansion and usage of Ito's lemma can be used.

Given coordinates (x^1, \dots, x^n) , when θ is a differential k -form, it can be written as

$$165 \quad \theta = \sum_{i_1 < \dots < i_k} f^{i_1, \dots, i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}. \quad (8)$$

Then

$$T_t^* \theta = \sum_{i_1 < \dots < i_k} f^{i_1, \dots, i_k}(T_t(x)) T_t^* (dx^{i_1} \wedge \dots \wedge dx^{i_k}). \quad (9)$$

Given in appendix B, Taylor expansion and Ito lemma are applied to expand $T_t^* \theta$, leading to compactly write

$$T_t^* \theta = \theta + \mathcal{M}(\theta) \Delta t + \mathcal{N}_i(\theta) \Delta \eta_i, \quad (10)$$

170 for some differential k -forms $\mathcal{M}(\theta)$ and $\mathcal{N}_i(\theta)$. Hereafter, several examples of $T_t^* \theta$ are presented.

The full derivation of these examples are skipped. We further express all the terms in coordinates. For instance, we replace $\langle \nabla f, a \rangle$ with $a^j \partial_{x^j} f$, where, by convention of notation, $a^j \partial_{x^j} f = \sum_j a^j \frac{\partial f}{\partial x^j}$. Similarly, $e_i^\top H_f e_i$ is replaced with $e_i^p e_i^q \partial_{x^p} \partial_{x^q} f$.

Remark 2. When $\theta = f \frac{\partial}{\partial x^{i_1}} \wedge \cdots \wedge \frac{\partial}{\partial x^{i_k}}$ is a contravariant tensor field,

$$T_{t*}\theta = f(T_t^{-1}(x))T_{t*}\left(\frac{\partial}{\partial x^{i_1}} \wedge \cdots \wedge \frac{\partial}{\partial x^{i_k}}\right). \quad (11)$$

175 The formula for T_t^{-1} is derived in appendix A. Then the expression of $f(T_t^{-1}(x))$, $T_{t*}\frac{\partial}{\partial x^{i_1}} \wedge \cdots \wedge \frac{\partial}{\partial x^{i_k}}$ and $T_{t*}\theta$ can be derived step by step in a similar way as in appendix B.

Example 3.1.1. When $\theta = f$ is a function (differential 0–form),

$$(T_t^*\theta) = f + \left(a^j \partial_{x^j} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f\right) \Delta t + e_i^p \partial_{x^p} f \Delta \eta_i \quad (12)$$

Example 3.1.2. When $\theta = dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n$,

$$180 \quad T_t^*\theta = \left\{1 + \left(\partial_{x^p} a^p + \frac{1}{2} J_i\right) \Delta t + \partial_{x^p} e_i^p \Delta \eta_i\right\} \theta, \quad (13)$$

where $J_i = \partial_{x^p} e_i^p \partial_{x^q} e_i^q - \partial_{x^p} e_i^q \partial_{x^q} e_i^p$.

Example 3.1.3. When $\theta = f dx^1 \wedge \cdots \wedge dx^n$,

$$\begin{aligned} T_t^*\theta = & \left\{ f + \left((\partial_{x^p} a^p + \frac{1}{2} J_i) f + (a^p + e_i^p \partial_{x^q} e_i^q) \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) \Delta t \right. \\ & \left. + (\partial_{x^p} e_i^p f + e_i^p \partial_{x^p} f) \Delta \eta_i \right\} dx^1 \wedge \cdots \wedge dx^n \end{aligned} \quad (14)$$

185 **Example 3.1.4.** When $\theta = f^j dx^j$ (note that by the convention of notation, $f^j dx^j = \sum_{j=1}^n f^j dx^j$),

$$\begin{aligned} T_t^*\theta = & \left\{ f^j + (a^p \partial_{x^p} f^j + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f^j + \partial_{x^j} a^p f^p + \partial_{x^j} e_i^p e_i^q \partial_{x^q} f^p) \Delta t \right. \\ & \left. + (e_i^p \partial_{x^p} f^j + \partial_{x^j} e_i^p f^p) \Delta \eta_i \right\} dx^j \end{aligned} \quad (15)$$

Example 3.1.5. When $\theta = f \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^n}$,

$$\begin{aligned} 190 \quad T_{t*}\theta = & \left\{ f + \left((\partial_{x^p} a^p + \frac{1}{2} J_i) f + (-(a^p + e_i^p \partial_{x^q} e_i^q) + \partial_{x^q} e_i^p e_i^q) \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) \Delta t \right. \\ & \left. + (\partial_{x^p} e_i^p f - e_i^p \partial_{x^p} f) \Delta \eta_i \right\} \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^n} \end{aligned} \quad (16)$$

3.2 Derivation of the Stochastic PDE

Suppose S is the full state variable of the dynamical system:

$$\frac{\partial S}{\partial t} = g(S). \quad (17)$$

Let f be a component or a collection of components of S . We then associate f to a differential form θ in the perturbation
195 scheme, i.e. there is an invertible map \mathcal{F} that maps the space of f to the space of θ , such that $\mathcal{F}(f) = \theta$. Suppose the propagation equation for f is

$$df = g^f(S) dt. \quad (18)$$

This implies a propagation equation for θ :

$$d\theta = g^\theta(S)dt. \quad (19)$$

200 The discrete-time perturbed forecast at each time step consists of the following two steps:

$$\begin{cases} \tilde{\theta}(t + \Delta t) = \theta(t) + g^\theta(S(t))\Delta t & (20) \\ \theta(t + \Delta t) = T_t^* \tilde{\theta}(t + \Delta t) & (21) \end{cases}$$

with $T_t^* \tilde{\theta}(t + \Delta t) = \tilde{\theta}(t + \Delta t) + \mathcal{M}(\tilde{\theta}(t + \Delta t))\Delta t + \mathcal{N}_i(\tilde{\theta}(t + \Delta t))\Delta\eta_i + o(\Delta t)$ for some differential forms $\mathcal{M}(\tilde{\theta})$ and $\mathcal{N}_i(\tilde{\theta})$.

The physical PDE (20) being deterministic, $\|\tilde{\theta}(t + \Delta t) - \theta(t)\|$ scales in $O(\Delta t)$. Indeed, there is no noise term to induce a scaling in $O(\sqrt{\Delta t})$. Therefore, it can be assumed that there exists $C > 0$ so that $\|\mathcal{M}(\tilde{\theta}(t + \Delta t)) - \mathcal{M}(\theta(t))\| < C\Delta t$ and

205 $\|\mathcal{N}_i(\tilde{\theta}(t + \Delta t)) - \mathcal{N}_i(\theta(t))\| < C\Delta t$, for Δt small enough. Then

$$\begin{aligned} T_t^* \tilde{\theta}(t + \Delta t) &= \tilde{\theta}(t + \Delta t) + \left(\mathcal{M}(\theta(t)) + \mathcal{O}(\Delta t)\right)\Delta t + \left(\mathcal{N}_i(\theta(t)) + \mathcal{O}(\Delta t)\right)\Delta\eta_i + o(\Delta t) \\ &= \tilde{\theta}(t + \Delta t) + \mathcal{M}(\theta(t))\Delta t + \mathcal{N}_i(\theta(t))\Delta\eta_i + o(\Delta t) \end{aligned} \quad (22)$$

Therefore,

$$\theta(t + \Delta t) = \theta(t) + g^\theta(S(t))\Delta t + \mathcal{M}(\theta(t))\Delta t + \mathcal{N}_i(\theta(t))\Delta\eta_i + o(\Delta t). \quad (23)$$

210 This suggests the following stochastic propagation equation for θ :

$$d\theta = g^\theta(S)dt + \mathcal{M}(\theta)dt + \mathcal{N}_i(\theta)d\eta_i. \quad (24)$$

Since there is a 1-1 correspondence between θ and f , Eq.(19) also suggests a stochastic propagation equation for f , which can be written as

$$df = g^f(S)dt + \mathcal{M}^f(f)dt + \mathcal{N}_i^f(f)d\eta_i. \quad (25)$$

215 We denote the additional terms in Eq.(25) by

$$d_s f := \mathcal{M}^f(f)dt + \mathcal{N}_i^f(f)d\eta_i. \quad (26)$$

Then Eq.(25) can be written as:

$$df = g^f(S)dt + d_s f. \quad (27)$$

Remark 3 ($d_s f$ is not directly related to the original dynamics). $d_s f$ is completely determined by $T_t^* \theta$, but is not directly related to the original dynamics Eq.(18). Therefore, once the expression of T in Eq.(6) and the choice of θ is determined, the perturbation term $d_s f$ is prescribed. However, the choice of θ is up to the user, and may then be related to the original dynamics.

Remark 4. In particular, there is no noise in the the original dynamics Eq.(18) which could be correlated with the noise of the resulting stochastic scheme (21). That is why the Itô lemma directly applies in the Taylor development (B4) of f , and then in
 225 the equation (22), leading to (23) and the final SPDE. Indeed, unlike the Itô-Wentzell formula Kunita (1997) – a cornerstone of the LU scheme – there is no additional cross-correlation term between T_t^* and $\tilde{\theta}(t + \Delta t)$. The final SPDE (24) makes clear the link between the solution θ and the Brownian motions η_i . But, at a given time step t , since (18) has no noise term, $\tilde{\theta}(t + \Delta t)$ is correlated with the $t' \mapsto \eta_i(t')$ for $t' < t$ only, and is independent of the new Brownian increment $\Delta\eta_i(t)$ generating T_t . Therefore, there is no cross-correlation term between T_t^* and $\tilde{\theta}(t + \Delta t)$.

230 **Example 3.2.1.** When $\theta = f$, example 3.1.1,

$$T_t^* \theta - \theta = \left(a^p \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) \Delta t + e_i^p \partial_{x^p} f \Delta \eta_i \quad (28)$$

This implies that

$$d_s f = \left(a^p \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) dt + e_i^p \partial_{x^p} f d\eta_i \quad (29)$$

To physically interpret this equation, we rewrite:

$$235 \quad \frac{d_s f}{dt} + V^p \partial_{x^p} f = \partial_{x^p} \left(\left(\frac{1}{2} e_i^p e_i^q \right) \partial_{x^q} f \right) \quad (30)$$

where

$$V^p = -a^p + \frac{1}{2} \partial_{x^q} (e_i^p e_i^q) - e_i^p \frac{d\eta_i}{dt} \quad (31)$$

Terms of advection and diffusion are recognized. The matrix $\frac{1}{2} e_i^p e_i^q$ is symmetric non-negative and represents a diffusion matrix. The p -th component of the advecting velocity V^p is composed of the drift $-a^p$, a correction $\frac{1}{2} \partial_{x^q} (e_i^p e_i^q)$, and a stochastic
 240 advecting velocity $-e_i^p \frac{d\eta_i}{dt}$.

If the original deterministic PDE (18) is an advection diffusion equation, with advecting velocity u and diffusion coefficient D , the final SPDE to simulate (Eq. (25)) is now a stochastic advection-diffusion equation, with advecting velocity $u + V$ and diffusion matrix $DI_d + \frac{1}{2} e_i^p e_i^q$:

$$\frac{df}{dt} + (u^p + V^p) \partial_{x^p} f = \partial_{x^p} \left((D \delta_{pq} + \frac{1}{2} e_i^p e_i^q) \partial_{x^q} f \right) \quad (32)$$

245 This type of SPDE appears in the LU framework, detailed in section 4.2.1.

Example 3.2.2. When $\theta = f dx^1 \wedge \dots \wedge dx^n$, example 3.1.3,

$$T_t^* \theta - \theta = \left\{ \left((\partial_{x^p} a^p + \frac{1}{2} J_i) f + (a^p + e_i^p \partial_{x^q} e_i^q) \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) \Delta t \right. \\ \left. + (\partial_{x^p} e_i^p f + e_i^p \partial_{x^p} f) \Delta \eta_i \right\} dx^1 \wedge \dots \wedge dx^n \quad (33)$$

This implies that

$$250 \quad \mathbf{d}_s f = \left((\partial_{x^p} a^p + \frac{1}{2} J_i) f + (a^p + e_i^p \partial_{x^q} e_i^q) \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) dt \\ + (\partial_{x^p} e_i^p f + e_i^p \partial_{x^p} f) d\eta_i \quad (34)$$

Rewritten, it leads to:

$$\frac{\mathbf{d}_s f}{dt} + \partial_{x^p} (\tilde{V}^p f) = \partial_{x^p} \left((\frac{1}{2} e_i^p e_i^q) \partial_{x^q} f \right) \quad (35)$$

where

$$255 \quad \tilde{V}^p = V^p - (e_i^p \partial_{x^q} e_i^q) = -a^p + \frac{1}{2} (\partial_{x^q} e_i^p e_i^q - e_i^p \partial_{x^q} e_i^q) - e_i^p \frac{d\eta_i}{dt} \quad (36)$$

Again a advection-diffusion equation is recognized, but of different nature. Indeed, as expected for a n-form, the PDE is similar to a density conservation equation. Moreover, the advecting drift is slightly different to take into account the cross-correlations between $f(T_t(x))$ and $T_t^*(dx^1 \wedge \dots \wedge dx^n)$.

Recall, in fluid dynamics, the Reynolds transport theorem provide an integral conservation equation for the transport of
260 any conserved quantity within a fluid, connected to its corresponding differential equation. The Reynolds transport theorem is central to the LU setting. The present example thus already outlines a closed link between the proposed perturbation approach and the LU formulation. Accordingly, the SPDE (35) naturally appears in the LU framework, as detailed in section 4.2.2.

Example 3.2.3. When $\theta = f^j dx^j$, example 3.1.4,

$$265 \quad T_t^* \theta - \theta = \left\{ (a^p \partial_{x^p} f^j + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f^j + \partial_{x^j} a^p f^p + \partial_{x^j} e_i^p e_i^q \partial_{x^q} f^p) \Delta t \\ + (e_i^p \partial_{x^p} f^j + \partial_{x^j} e_i^p f^p) \Delta \eta_i \right\} dx^j \quad (37)$$

For each j , the coefficients of dx^j in $T_t^* \theta - \theta$ and those in θ can be compared, to lead to

$$\mathbf{d}_s f^j = (a^p \partial_{x^p} f^j + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f^j + \partial_{x^j} a^p f^p + \partial_{x^j} e_i^p e_i^q \partial_{x^q} f^p) dt \\ + (e_i^p \partial_{x^p} f^j + \partial_{x^j} e_i^p f^p) d\eta_i \quad (38)$$

Regrouping the terms for physical interpretation, it writes:

$$270 \quad \frac{\mathbf{d}_s f^j}{dt} + V^p \partial_{x^p} f^j + \partial_{x^j} \left(-a^p - e_i^p \frac{d\eta_i}{dt} \right) f^p - \partial_{x^j} e_i^p e_i^q \partial_{x^q} f^p = \partial_{x^p} \left((\frac{1}{2} e_i^p e_i^q) \partial_{x^q} f^j \right) \quad (39)$$

Two additional terms complete the advection-diffusion term. The first one, $\partial_{x^j} \left(-a^p - e_i^p \frac{d\eta_i}{dt} \right) f^p$, is reminiscent to the additional terms appearing in SALT momentum equations Holm (2015); Resseguier et al. (2020). The second term, $-\partial_{x^j} e_i^p e_i^q \partial_{x^q} f^p$, comes from cross-correlation in Itô notation.

Example 3.2.4. When $\theta = f \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^n}$, example 3.1.5,

$$275 \quad T_{t*}\theta - \theta = \left\{ \left((\partial_{x^p} a^p + \frac{1}{2} J_i) f + (- (a^p + e_i^p \partial_{x^q} e_i^q) + \partial_{x^q} e_i^p e_i^q) \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) \Delta t \right. \\ \left. + (\partial_{x^p} e_i^p f - e_i^p \partial_{x^p} f) \Delta \eta_i \right\} \frac{\partial}{\partial x^1} \wedge \cdots \wedge \frac{\partial}{\partial x^n} \quad (40)$$

This implies

$$\mathbf{d}_s f = \left((\partial_{x^p} a^p + \frac{1}{2} J_i) f + (- (a^p + e_i^p \partial_{x^q} e_i^q) + \partial_{x^q} e_i^p e_i^q) \partial_{x^p} f + \frac{1}{2} e_i^p e_i^q \partial_{x^p} \partial_{x^q} f \right) dt \\ + (\partial_{x^p} e_i^p f - e_i^p \partial_{x^p} f) d\eta_i \quad (41)$$

280 It can then be verified that:

$$\frac{\mathbf{d}_s f}{dt} + \partial_{x^p} \tilde{V}^p f - \tilde{V}^p \partial_{x^p} f = \partial_{x^p} \left(\left(\frac{1}{2} e_i^p e_i^q \right) \partial_{x^q} f \right) \quad (42)$$

where

$$\tilde{V}^p = \tilde{V}^p - (e_i^p \partial_{x^q} e_i^q) = V^p - 2(e_i^p \partial_{x^q} e_i^q) \quad (43)$$

It is recognized the diffusion term, $\partial_{x^p} \left(\left(\frac{1}{2} e_i^p e_i^q \right) \partial_{x^q} f \right)$, the divergence term, $\partial_{x^p} \tilde{V}^p f$, comparable to the density equation, and
 285 the advection term, $-\tilde{V}^p \partial_{x^p} f$. However, the velocity fields appearing in the divergent and advecting terms do not coincide. Indeed, they are even opposite for divergence-free noise ($\partial_{x^q} e_i^q = 0$). This type of equation may appear uncommon but will be shown useful when applied to randomized thermal shallow water equations.

3.3 Conservation laws related to $\mathbf{d}_s f$

A major advantage of the proposed perturbation scheme is to possibly prescribe θ to ensure that certain quantities are conserved.

290 Define the discrete time version of $\mathbf{d}_s f$ as:

$$\Delta_s f = \mathcal{M}^f(f) \Delta t + \mathcal{N}_i^f(f) \Delta \eta_i. \quad (44)$$

In general, conservation laws can be derived from the following two identities about the pull-back operator:

$$(T_t^* \theta_1) \wedge (T_t^* \theta_2) = T_t^* (\theta_1 \wedge \theta_2) \quad (45)$$

$$dT_t^* \theta = T_t^* d\theta, \quad (46)$$

295 where d refers to the differential operator acting on differential forms. Hereafter, we present how to derive the conservation laws for two particular examples.

Example 3.3.1. Suppose $\theta_1 = f dx^1 \wedge \cdots \wedge dx^n$ and define

$$\hat{\theta}_1 = T_t^* \theta_1 \quad (47)$$

$$\hat{f} = f + \Delta_s f. \quad (48)$$

300 Then $\hat{\theta}_1 = \hat{f}dx^1 \wedge \dots \wedge dx^n$. Therefore

$$\begin{aligned} \int_{\Omega} \hat{f}dx^1 \dots dx^n &= \int_{\Omega} \hat{\theta}_1 = \int_{\Omega} T_t^* \theta_1 = \int_{T_t(\Omega)} \theta_1 = \int_{\Omega} \theta_1 \\ &= \int_{\Omega} f dx^1 \dots dx^n. \end{aligned} \quad (49)$$

Eq.(49) implies that the total integral of f is not changed by the perturbation scheme. Next suppose that $\theta_2 = g$ is a function. Similarly we define

$$305 \quad \hat{\theta}_2 = T_t^* \theta_2 \quad (50)$$

$$\hat{g} = g + \Delta_s g. \quad (51)$$

Applying Eq.(45),

$$\begin{aligned} \int_{\Omega} \hat{f}\hat{g}dx^1 \dots dx^n &= \int_{\Omega} \hat{\theta}_1 \wedge \hat{\theta}_2 = \int_{\Omega} T_t^*(\theta_1 \wedge \theta_2) = \int_{T_t(\Omega)} \theta_1 \wedge \theta_2 \\ &= \int_{\Omega} \theta_1 \wedge \theta_2 = \int_{\Omega} fg dx^1 \dots dx^n \end{aligned} \quad (52)$$

310 The total integral of fg is thus also conserved by the perturbation scheme. Similarly for any integer $m \geq 0$, fg^m is conserved by the perturbation scheme.

Example 3.3.2. Suppose $n = 2$ and $\theta = udx + vdy$, where $\mathbf{u} = (u, v)$ is the velocity field. The vorticity $\omega = \partial_x v - \partial_y u$ corresponds to the differential 2-form $d\theta$:

$$d\theta = \omega dx^1 \wedge dx^2. \quad (53)$$

315 Define $\hat{\theta} := T_t^* \theta = \hat{u}dx^1 + \hat{v}dx^2$ and $\hat{\omega} = \partial_x \hat{v} - \partial_y \hat{u}$. Then $d\hat{\theta} = \hat{\omega}dx^1 \wedge dx^2$, and

$$\begin{aligned} \int_{\Omega} \hat{\omega}dx^1 dx^2 &= \int_{\Omega} d\hat{\theta} = \int_{\omega} dT_t^* \theta = \int_{\Omega} T_t^* d\theta = \int_{T_t(\Omega)} d\theta \\ &= \int_{\Omega} \omega dx^1 dx^2. \end{aligned} \quad (54)$$

Therefore the vorticity is conserved by the perturbation scheme.

Example 3.3.3. Suppose $n = 3$ and $\theta = udx + vdy + wdz$, where $\mathbf{u} = (u, v, w)$ is the velocity field. The vorticity $\omega = (\partial_y w - \partial_z v, \partial_z u - \partial_x w, \partial_x v - \partial_y u)$ corresponds to the differential 2-form $d\theta$:

$$d\theta = (\partial_y w - \partial_z v)dy \wedge dz + (\partial_x w - \partial_z v)dz \wedge dx + (\partial_x v - \partial_y u)dx \wedge dy. \quad (55)$$

The helicity $\Theta = u(\partial_y w - \partial_z v) + v(\partial_x v - \partial_y u) + w(\partial_x v - \partial_y u)$ corresponds to the differential 3-form:

$$d\theta \wedge \theta = \left(u(\partial_y w - \partial_z v) + v(\partial_x v - \partial_y u) + w(\partial_x v - \partial_y u) \right) dx \wedge dy \wedge dz. \quad (56)$$

Similarly, we define $\hat{\Theta}$ by $d\hat{\theta} \wedge \hat{\theta} = \hat{\Theta} dx \wedge dy \wedge dz$. Then

$$\begin{aligned} 325 \quad \int_{\Omega} \hat{\Theta} dx dy dz &= \int_{\Omega} d\hat{\theta} \wedge \hat{\theta} = \int_{\Omega} (dT_t^* \theta) \wedge (T_t^* \theta) \\ &= \int_{\Omega} (T_t^* d\theta) \wedge (T_t^* \theta) = \int_{\Omega} T_t^* (d\theta \wedge \theta) = \int_{T_t(\Omega)} d\theta \wedge \theta = \int_{\Omega} \Theta dx dy dz. \end{aligned} \quad (57)$$

Hence, in this case, the total amount of helicity is conserved.

Example 3.3.4. Suppose that $\theta_1 = f dx^1 \wedge \dots \wedge dx^n$ and that $\theta_2 = g \frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n}$. There exists a pairing $\langle \cdot, \cdot \rangle$ for the differential n -forms and the contravariant n -vectors, i.e. $\langle \theta_1, \theta_2 \rangle = fg$ is a function on Ω . Define

$$330 \quad \hat{\theta}_1 = T_t^* \theta_1 = \hat{f} dx^1 \wedge \dots \wedge dx^n \quad (58)$$

$$\hat{\theta}_2 = (T_t^{-1})_* \theta_2 = \hat{g} \frac{\partial}{\partial x^1} \wedge \dots \wedge \frac{\partial}{\partial x^n} \quad (59)$$

Then we have

$$\hat{f} \hat{g} (T_t^{-1}(x)) = \langle \hat{\theta}_1, \hat{\theta}_2 \rangle \Big|_{T_t^{-1}(x)} = \langle \theta_1, \theta_2 \rangle \Big|_x = fg(x), \quad (60)$$

and that

$$335 \quad \int_{\Omega} \hat{f}^2 \hat{g} dx^1 \dots dx^n = \int_{\Omega} \langle \hat{\theta}_1, \hat{\theta}_2 \rangle \theta_1 = \int_{\Omega} \langle \theta_1, \theta_2 \rangle \theta_1 = \int_{\Omega} f^2 g dx^1 \dots dx^n \quad (61)$$

Remark 5 (The conservation law of the perturbation scheme is independent of the conservation law of the original dynamical system). *The derivation of Eqs.(49) (52), (54), (57), and (61) is based on the generic properties of the pull-back and push-forward operator of tensor fields. Since the choice of θ is not directly determined by the dynamical system, the conservation law of the perturbation scheme is independent of the original dynamical system. Recall that the perturbed forecast consists of*

340 *two steps: Eq.(20) and (21). The conservation law of the perturbation scheme implies that certain quantities are conserved in the second step. On the other hand, the original dynamical system Eq.(20) might enjoy some other conservation law. If a quantity is conserved by both the original dynamical system and the perturbation scheme, then this quantity must be conserved by the final stochastic PDE. If a quantity is conserved by only one of Eqs.(20) and (21), then it can not be concluded that this quantity is conserved by the final SPDE.*

345 4 Comparison with other perturbation schemes

In this section, we demonstrate that both the stochastic advection by Lie transport (SALT) equation Holm (2015) and the location uncertainty (LU) equation Mémin (2014); Resseguier et al. (2016, 2020) can be recovered using the proposed perturbation scheme and properly choosing θ and the parameters a, e_i .

4.1 Comparison with SALT equation

350 The original SALT equation Holm (2015) is derived based on a stochastically constrained variational principle $\delta S = 0$, for which

$$\begin{cases} S(u, q) = \int \ell(u, q) dt \\ dq + \mathcal{L}_{dx_t} q = 0. \end{cases} \quad (62)$$

where $\ell(u, q)$ is the Lagrangian of the system, \mathcal{L} is the Lie derivative, and $x_t(x)$ is defined by (using our notation)

$$x_t(x) = x_0(x) + \int_0^t u(x, s) ds - \int_0^t e_i(x) \circ d\eta_i(s), \quad (63)$$

355 in which u is the velocity vector field, and the \circ means that the integral is defined in the Stratonovich sense, instead of in the Ito sense. Hence, $dx_t = u(x, t)dt - e_i \circ d\eta_i$ refers to an infinitesimal stochastic tangent field on the domain. Broadly speaking, we can express $dx_t = T_t(x) - x + udt$. Note the difference between Ito's notation and Stratonovich's notation, i.e. $e_i \circ d\eta_i \neq e_i d\eta_i$. Our expression of T_t essentially follows Ito's notation, and $T_t(x) \neq x - e_i \Delta\eta_i$ in this subsection. Instead, it becomes $T_t(x) = x + \frac{1}{2} e_i^p \partial_{x_p} e_i \Delta t - e_i \Delta\eta_i$.

360 In the second equation of Eq.(62), q is assumed to be a quantity advected by the flow. q can correspond to any differential form that is not uniquely determined by the velocity (since the SALT equation for the velocity is usually determined by the first equation of Eq.(62)). In Holm (2015), the Lie derivative $\mathcal{L}_{dx_t} q$ is calculated using Cartan's formula:

$$\mathcal{L}_{dx_t} q = d(i_{dx_t} q) + i_{dx_t} dq. \quad (64)$$

Essentially, the Lie derivative $\mathcal{L}_{dx_t} q$ corresponds to $T_t^* q - q + f^q(S)dt$, if we assume that the deterministic forecast of q is simply the advection of q by u . More generally, $\mathcal{L}_{dx_t - udt} q = T_t^* q - q$. Therefore, the SALT equation for q is the same as our
365 equation for q . We remark that the Cartan's formula can not be directly applied to calculate the Lie derivative if the expression of dx_t is in Ito's notation.

The SALT equation regarding the velocity u comes from the first equation of Eq.(62). For most cases, the velocity u is associated with the momentum, a differential 1-form $m = u^j dx^j = u^1 dx^1 + \dots + u^n dx^n$. In the examples discussed in Holm
370 (2015), it is observed that, when the Lagrangian includes the kinetic energy, the stochastic noises contribute a term $\mathcal{L}_{dx_t} \theta$, where θ is a differential 1-form related to the momentum 1-form. For instance, $\theta = m$ in the example of "Stratonovich stochastic Euler-Poincaré flow" in Holm (2015), and $\theta = m + R^j dx^j$ in the example of "Stochastic Euler-Boussinesq equations of a rotating stratified incompressible fluid" in Holm (2015). Already pointed out, the operator \mathcal{L}_{dx_t} is closely related to T_t^* , and the momentum equation in SALT can be derived using our proposed scheme by properly choosing θ .

375 Holm (2015) requires that q to be a differential form since Cartan's formula is only useful for differential forms q . This restriction can be relaxed by employing the original definition of Lie derivative with respect to a deterministic/stochastic flow of diffeomorphism discussed in Leon (2021), so that $\mathcal{L}_{dx_t} q$ can be generalized to the case where q is a mixed tensor field. This corresponds to our Eq.(7).

Compared with Holm (2015); Leon (2021), the proposed perturbation approach seems more flexible and does not have to
 380 rely on the Lagrangian mechanics. In particular, the velocity field can be associated to other tensor fields than the momentum
 1-form. The perturbation, not directly related to the physics, can then be applied to any PDE. Moreover, our approach provides
 a new interpretation of $\mathcal{L}_{dx_t - u dt}$ in terms of the optimal transportation associated with the infinitesimal forecast error at each
 time step. This interpretation certainly suggests practical numerical methods to infer a, e_i . Given a long sequence of reanalysis
 data or simulated high-resolution data, the one-step forecast can be evaluated using the low resolution model, with the high
 385 resolution state at each time step being the initial condition. T_t is then estimated at each time step by comparing the low
 resolution forecast and the high resolution forecast. Finally, a and e_i could be learnt from these samples of T_t .

4.2 Comparison with the LU equation

Mentioned above, the Reynolds transport theorem is central to the LU setting, and we already outlines a closed link between
 the proposed perturbation approach and the LU formulation. This link – related to differential n -forms – will be precised later
 390 in this subsection. But, before this, we focus on another key ingredient of LU: the stochastic material derivative of functions
 (differential 0-forms).

4.2.1 0-forms in the LU framework

Dropping the forcing terms, LU equation for compressible and incompressible flow writes Resseguier et al. (2016).

$$\partial_t f + \mathbf{w}^* \cdot \nabla f = \nabla \cdot \left(\frac{1}{2} \mathbf{a} \nabla f \right) - \boldsymbol{\sigma} \dot{\mathbf{B}} \cdot \nabla f \quad (65)$$

$$395 \quad \mathbf{w}^* = \mathbf{w} - \frac{1}{2} (\nabla \cdot \mathbf{a})^\top + \boldsymbol{\sigma} (\nabla \cdot \boldsymbol{\sigma})^\top, \quad (66)$$

where f can be any quantity that is assumed to be transported by the flow, i.e. $Df/Dt = 0$ where D/Dt is the Itô material
 derivative. For instance, f could be the velocity (dropping forces in the SPDE), the temperature, or the buoyancy. Compared to
 SALT notations, $-e_i d\eta_i$ is denoted $\boldsymbol{\sigma} d\mathbf{B} = \boldsymbol{\sigma}_{\bullet i} dB_i$. We refer to (Resseguier et al., 2020, Appendix A) for the complete table
 of SALT-LU notations correspondences. Derived in (Resseguier, 2017, Appendix 10.1) and (Resseguier et al., 2021, 6.1.3), we
 400 can rewrite it as

$$\partial_t f + \mathbf{w}_S \cdot \nabla f = \frac{1}{2} (\boldsymbol{\sigma}_{\bullet i} \cdot \nabla) (\boldsymbol{\sigma}_{\bullet i} \cdot \nabla f) - (\boldsymbol{\sigma} \dot{\mathbf{B}}) \cdot \nabla f, \quad (67)$$

$$= - (\boldsymbol{\sigma} \circ \dot{\mathbf{B}}) \cdot \nabla f, \quad (68)$$

$$\mathbf{w}_S = \mathbf{w} + \mathbf{w}_S^c \quad (69)$$

$$\mathbf{w}_S^c = -\frac{1}{2} (\nabla \cdot \mathbf{a})^\top + \frac{1}{2} \boldsymbol{\sigma} (\nabla \cdot \boldsymbol{\sigma})^\top, \quad (70)$$

$$405 \quad = -\frac{1}{2} (\boldsymbol{\sigma}_{\bullet i} \cdot \nabla) \boldsymbol{\sigma}_{\bullet i}, \quad (71)$$

where $\boldsymbol{\sigma} \circ \dot{\mathbf{B}}$ is the Stratonovich noise of the SPDE, \mathbf{w} and \mathbf{w}_S (denoted u in the SALT framework) are respectively the Itô
 drift and the Stratonovich drift of the fluid flow. Separating the terms of the SPDE related to the deterministic dynamics from

the term associated to the stochastic scheme, it comes

$$d^{\text{LU}} f = g^f(S)dt + d_s^{\text{LU}} f, \quad (72)$$

410 where

$$g^f(S) = -\mathbf{w} \cdot \nabla f \quad (73)$$

$$d_s^{\text{LU}} f = -\mathbf{w}_S^c \cdot \nabla f dt + \frac{1}{2}(\boldsymbol{\sigma}_{\bullet i} \cdot \nabla)(\boldsymbol{\sigma}_{\bullet i} \cdot \nabla f)dt - (\boldsymbol{\sigma} d\mathbf{B}) \cdot \nabla f \quad (74)$$

Terms in Eqs.(65) and (66) translate to our notation in the following way:

$$\begin{aligned} -\mathbf{w}_S^c \cdot \nabla f dt &= \frac{1}{2} e_i^q \partial_{x_q} e_i^p \partial_{x_p} f \\ 415 \quad \frac{1}{2}(\boldsymbol{\sigma}_{\bullet i} \cdot \nabla)(\boldsymbol{\sigma}_{\bullet i} \cdot \nabla f) &= \frac{1}{2} e_i^p \partial_{x_p} (e_i^q \partial_{x_q} f) \\ &= \frac{1}{2} (e_i^p \partial_{x_p} e_i^q \partial_{x_q} f + e_i^p e_i^q \partial_{x_p} \partial_{x_q} f) \\ -\boldsymbol{\sigma} d\mathbf{B} \cdot \nabla f &= e_i^p \partial_{x_p} f d\eta_i \end{aligned}$$

Hence

$$d_s^{\text{LU}} f = (e_i^q \partial_{x_q} e_i^p \partial_{x_p} f + \frac{1}{2} e_i^p e_i^q \partial_{x_p} \partial_{x_q} f)dt + e_i^p \partial_{x_p} f d\eta_i \quad (75)$$

420 Direct calculation yields that Eq.(75) coincides with Eq.(29) when

$$T_t(x) = x + e_i^q \partial_{x_q} e_i \Delta t + e_i \Delta \eta_i = x - \mathbf{w}_S^c \Delta t + (-\mathbf{w}_S^c \Delta t - \boldsymbol{\sigma} \Delta \mathbf{B}). \quad (76)$$

The LU equation can thus be derived by choosing $\theta = f$ and T_t by Eq.(76). At the first glance, it seems not straightforward to make such a choice. Nevertheless, it can be recognized that the term $(-\mathbf{w}_S^c \Delta t - \boldsymbol{\sigma} \Delta \mathbf{B}) = (\frac{1}{2} e_i^q \partial_{x_q} e_i \Delta t + e_i \Delta \eta_i)$ is the Itô noise plus its Itô-to-Stratonovich correction. Hence, it corresponds to the Stratonovich noise $e_i \circ d\eta_i$ of the flow associated to T_t . The additional drift $-\mathbf{w}_S^c \Delta t$ is different in nature. It is related to the advection correction $\mathbf{w}_S^c \cdot \nabla f$ in the LU setting. Indeed, in the LU framework, the Itô drift, \mathbf{w} , is seen as the resolved large-scale velocity. That is why, in this framework, the deterministic dynamics (74) involves the Itô drift, \mathbf{w} . This is also the reason why, under the LU derivation, the advected velocity is assumed to be given by the Itô drift, \mathbf{w} . It differs from the Stratonovich drift $\mathbf{w}_S = \mathbf{w} + \mathbf{w}_S^c$, used as advected velocity in SALT approach or in Mikulevicius and Rozovskii (2004) (where the Stratonovich drift is denoted u). Interested readers are referred to (Resseguier et al., 2020, Appendix A) for a discussion on these assumptions. Note however that in all these approaches, the advecting velocity is always the Stratonovich drift. This can be seen e.g., in the Stratonovich form of LU equations (68).

To also understand (76), the inverse flow can be considered. According to appendix A,

$$T_t^{-1}(x) = x - e_i \Delta \eta_i = x + \boldsymbol{\sigma} \Delta \mathbf{B}. \quad (77)$$

435 Considering T_t to represent how much the model forecast differs from the true forecast at every time step, T_t^{-1} can be understood to represent how much the true forecast differs from the model forecast at each time step. Therefore, the LU equation can be derived using the proposed perturbation scheme, choosing $\theta = f$ and assuming that the true forecast differs from the model forecast by a displacement prescribed by Eq.(77).

4.2.2 n-forms in the LU framework

440 The LU physical justification relies on a stochastic interpretation of fundamental conservation laws, typically conservation of extensive properties (i.e. integrals of functions over a spatial volume) like momentum, mass, matter and energy Resseguier et al. (2016). These extensive properties can be expressed by integrals of differential n -forms. For instance, the mass and the momentum are integrals of the differential n -forms $\rho dx^1 \wedge \dots \wedge dx^n$ and $\rho \mathbf{w} dx^1 \wedge \dots \wedge dx^n$, respectively. In the LU framework, a stochastic version of the Reynolds transport theorem (Resseguier et al., 2016, Eq. (28)) is used to deal with these differential

445 n -forms $\theta = f dx^1 \wedge \dots \wedge dx^n$. Assuming an integral conservation $\frac{d}{dt} \int_{V(t)} f = 0$ on a spatial domain $V(t)$ transported by the flow, that theorem leads to the following SPDE:

$$\frac{Df}{Dt} + \nabla \cdot (\mathbf{w}^* + \boldsymbol{\sigma} \dot{\mathbf{B}})f = \frac{d}{dt} \left\langle \int_0^t D_t f, \int_0^t \nabla \cdot \boldsymbol{\sigma} \dot{\mathbf{B}} \right\rangle = (\nabla \cdot \boldsymbol{\sigma}_{\bullet i})(\nabla \cdot \boldsymbol{\sigma}_{\bullet i})^T f \quad (78)$$

where D/Dt denotes the Itô material derivative. Here again, forcing terms are dropped for the sake of readability. This SPDE can be rewritten using the expression of that material derivative (Eq. (9) and (10) of Resseguier et al. (2016)):

$$450 \quad \partial_t f + \nabla \cdot (\mathbf{w}_S f) = \frac{1}{2} \nabla \cdot (\mathbf{a} \nabla f) + \frac{1}{2} \nabla \cdot (\boldsymbol{\sigma}_{\bullet i} (\nabla \cdot \boldsymbol{\sigma}_{\bullet i})^T f) - \nabla \cdot (\boldsymbol{\sigma} \dot{\mathbf{B}} f) \quad (79)$$

$$= \frac{1}{2} \nabla \cdot (\boldsymbol{\sigma}_{\bullet i} (\nabla \cdot (\boldsymbol{\sigma}_{\bullet i} f))^T) - \nabla \cdot (\boldsymbol{\sigma} \dot{\mathbf{B}} f) \quad (80)$$

$$= -\nabla \cdot (\boldsymbol{\sigma} \circ \dot{\mathbf{B}} f) \quad (81)$$

The original deterministic equation and stochastic perturbation correspond to

$$g^f(S) = -\nabla \cdot (\mathbf{w} f) \quad (82)$$

$$455 \quad d_s^{\text{LU}} f = (-\nabla \cdot (\mathbf{w}_S^c f) + \frac{1}{2} \nabla \cdot (\mathbf{a} \nabla f) + \frac{1}{2} \nabla \cdot (\boldsymbol{\sigma}_{\bullet i} (\nabla \cdot \boldsymbol{\sigma}_{\bullet i})^T f)) dt - \nabla \cdot (\boldsymbol{\sigma} d\mathbf{B} f) \quad (83)$$

$$= \nabla \cdot (((\frac{1}{2} \nabla \cdot \mathbf{a})^T dt - \boldsymbol{\sigma} d\mathbf{B}) f) + \nabla \cdot (\frac{1}{2} \mathbf{a} \nabla f) dt \quad (84)$$

Identifying $\mathbf{a} = \boldsymbol{\sigma}_{\bullet i} \boldsymbol{\sigma}_{\bullet i}^T = e_i e_i^T$ and $\boldsymbol{\sigma} \dot{\mathbf{B}} = -e_i d\eta_i$, Eq. (35) corresponds to example 3.2.2 about n -forms, with

$$\tilde{V} = -a^p + \frac{1}{2} (\partial_{x^q} e_i^p e_i^q - e_i^p \partial_{x^q} e_i^q) - e_i^p \frac{d\eta_i}{dt} = -(\frac{1}{2} \nabla \cdot \mathbf{a})^T + \boldsymbol{\sigma} \dot{\mathbf{B}} \quad (85)$$

i.e.

$$460 \quad a^p = \partial_{x^q} (e_i^p e_i^q) - (e_i^p \partial_{x^q} e_i^q) = e_i^q \partial_{x^q} e_i^p. \quad (86)$$

Again the remapping is obtained

$$T_i(x) = x + e_i^q \partial_{x^q} e_i \Delta t + e_i \Delta \eta_i = x - \mathbf{w}_S^c \Delta t + (-\mathbf{w}_S^c \Delta t - \boldsymbol{\sigma} \Delta \mathbf{B}), \quad (87)$$

previously derived for differential 0-form in LU framework (Eq. (76)). Therefore, the proposed approach also generalizes the LU framework for n -forms, and its capacity – given by the Reynolds transport theorem – to deal with extensive properties.

465 **Remark 6.** For incompressible flows, LU equation further imposes that

$$\begin{cases} \nabla \cdot \boldsymbol{\sigma} = 0 \\ \nabla \cdot \nabla \cdot \mathbf{a} = 0 \end{cases} \quad (88)$$

Translating it into our notation, it reads as

$$\begin{cases} \partial_{x_p} e_i^p = 0 \text{ for each } i \\ \partial_{x_p} \partial_{x_q} (e_i^p e_i^q) = 0 \end{cases}$$

470 Applying the result in example 3.1.2, straightforward calculation gives Eq.(88) to be equivalent to that $T_t^* \theta = \theta$ for $\theta = dx^1 \wedge \cdots \wedge dx^n$. Such a result was expected since constraints Eq. (88) are obtained from the LU density conservation.

5 A stochastic version of thermal shallow water equation

In this section, the proposed approach is applied to derive a stochastic version of thermal shallow water equation. Another stochastic version of thermal shallow water equation can be found in Holm and Luesink (2021). The thermal shallow water equation is derived in Warnford and Dellar (2013):

$$475 \quad \frac{\partial h}{\partial t} + \nabla \cdot (h\bar{u}) = 0, \quad (89)$$

$$\frac{\partial \Theta}{\partial t} + (\bar{u} \cdot \nabla) \Theta = -\kappa(h\Theta - h_0\Theta_0), \quad (90)$$

$$\frac{\partial \bar{u}}{\partial t} + (\bar{u} \cdot \nabla) \bar{u} + f\hat{z} \times \bar{u} = -\nabla(h\Theta) + \frac{1}{2}h\nabla\Theta \quad (91)$$

480 This model can be used to describe a two-layer system under equivalent barotropic approximation. The upper layer is active but with a spatio-temporal varying density $\rho(x, t)$, while the lower layer is quiescent with a fixed constant density ρ_0 . The state variable h represents the height of the active layer, and $\Theta = g(\rho_0 - \rho)/\rho_0$ is the density contrast. \bar{u} is the averaged horizontal velocity of the active layer at each column. Note that $\rho < \rho_0$ (hence $\Theta > 0$) in the scenario of equivalent barotropic approximation Warnford and Dellar (2013).

Stated in Warnford and Dellar (2013), the following physical quantities are conserved up to the forcing:

$$\text{Total energy: } E = \int_{\Omega} \frac{1}{2} (h|\bar{u}|^2 + h^2\Theta) d^2x \quad (92)$$

$$485 \quad \text{Total mass: } \mathcal{M} = \int_{\Omega} h d^2x \quad (93)$$

$$\text{Total momentum: } \mathcal{M} = \int_{\Omega} h\bar{u} d^2x \quad (94)$$

The objective is thus to choose proper tensor fields $\theta_{\bar{u}}, \theta_h$, and θ_{Θ} for the state variables \bar{u}, h , and Θ , respectively, so that E, \mathcal{M} , and \mathcal{M} are conserved by the perturbation scheme. Again, it must be emphasized that the conservation law of the perturbation scheme does not directly imply that the same quantities are conserved by the final SPDE.

490 The domain is 2-dimensional. To conserve mass, the only choice for θ_h is $\theta_h = h dx^1 \wedge dx^2$, which is a differential 2-form. It plays the role of density. In order to conserve the momentum, we need the momentum to be a differential 2-form as well. Hence we must choose $\theta_{\bar{u}}$ to be a function (differential 0-form). Therefore, the only choice for $\theta_{\bar{u}}$ is $\theta_{\bar{u}} = \bar{u}$. This choice of $\theta_{\bar{u}}$ and θ_h implies that $h|\bar{u}|^2$ also corresponds to a 2-form $|\bar{u}|^2\theta_h$. Hence the kinetic energy is automatically conserved by the perturbation scheme. This means that if we want E to be conserved, we must select θ_Θ so that $h^2\Theta$ corresponds to a differential

495 2-form. Note that θ_h is already a 2-form. We must thus select θ_Θ so that $h\Theta$ corresponds to a function. The only choice for θ_Θ is the contravariant tensor $\theta_\Theta = \Theta \frac{\partial}{\partial x^1} \wedge \frac{\partial}{\partial x^2}$. In this case, $h\Theta$ corresponds to the differential 0-form $\langle \theta_h, \theta_\Theta \rangle = h\Theta$, where $\langle \cdot, \cdot \rangle$ in this section is the natural pairing of covariant n -tensor fields and contravariant n -tensor fields.

In sum, we have chosen the following tensor fields:

$$\theta_h = h dx^1 \wedge dx^2 \quad (95)$$

$$500 \quad \theta_{\bar{u}^j} = \bar{u}^j \quad (\text{for } j = 1, 2) \quad (96)$$

$$\theta_\Theta = \Theta \frac{\partial}{\partial x^1} \wedge \frac{\partial}{\partial x^2}. \quad (97)$$

For

$$T_t(x) = x + a\Delta t + e_i \Delta \eta_i, \quad (98)$$

we have

$$505 \quad T_t^{-1}(x) = x + (-a + e_i^p \partial_{x_p} e_i) \Delta t - e_i \Delta \eta_i. \quad (99)$$

Then $T_t^* \theta_h$, $T_t^* \theta_{\bar{u}}$, and $(T_t^{-1})_* \theta_\Theta$ can be calculated following examples 3.1.3, 3.1.1, and 3.1.5. This further implies $d_s h$, $d_s \bar{u}$, and $d_s \Theta$, as shown in examples 3.2.2, 3.2.1, and 3.2.4. Note that T_t^{-1} instead of T_t is applied to θ_Θ as shown in Eq.(7). Finally, we end up with the following SPDE:

$$510 \quad dh = -\nabla(h\bar{u})dt + \left(h(\partial_{x_p} a^p + \frac{1}{2} J_i) + a^p \partial_{x_p} h + \frac{1}{2} e_i^p e_i^q \partial_{x_p} \partial_{x_q} h + \partial_{x_p} h e_i^p \partial_{x_q} e_i^q \right) dt \\ + (h \partial_{x_p} e_i^p + \partial_{x_p} h e_i^p) d\eta_i \quad (100)$$

$$d\Theta = \{ -(\bar{u} \cdot \nabla) \Theta - \kappa(h\Theta - h_0 \Theta_0) \} dt \\ + \left(\Theta(-\partial_{x_p} a^p + \partial_{x_p} (\partial_{x_q} e_i^q e_i^p) + \frac{1}{2} J_i) + \partial_{x_p} \Theta a^p + \frac{1}{2} e_i^p e_i^q \partial_{x_p} \partial_{x_q} \Theta - \partial_{x_p} \Theta e_i^p \partial_{x_q} e_i^q \right) dt \\ - (\Theta \partial_{x_p} e_i^p - \partial_{x_p} \Theta e_i^p) d\eta_i \quad (101)$$

$$515 \quad d\bar{u}^j = -\{ (\bar{u} \cdot \nabla) \bar{u} - f \hat{z} \times \bar{u} - \nabla(h\Theta) + \frac{1}{2} h \nabla \Theta \}^j dt \\ + \left(\partial_{x_p} \bar{u}^j a^p + \frac{1}{2} e_i^p e_i^q \partial_{x_p} \partial_{x_q} \bar{u}^j \right) dt + \partial_{x_p} \bar{u}^j e_i^p d\eta_i, \quad (102)$$

where $J_i = \partial_{x_p} e_i^p \partial_{x_q} e_i^q - \partial_{x_q} e_i^p \partial_{x_p} e_i^q$. And the total mass, total momentum and the total energy shall all be conserved by the perturbation scheme.

6 Conclusions

The starting point of this work is to question “how to consistently perturb the location of the state variable?”, motivated by Brenier’s theorem Brenier (1991) which suggests that the difference of two density fields can be represented by a transport map T . Noting that optimal transportation has a clean representation in terms of differential n -forms, we proposed to perturb the “location” of the state variable S , at every forecast time step, by perturbing the corresponding differential k -forms θ by $\theta \leftarrow T_t^* \theta$, where T_t is a random diffeomorphism which deviates from the identity map infinitesimally.

Under this framework, we end up with a stochastic PDE of the state variable S in the form

$$dS = f(S)dt + d_s S, \quad (103)$$

where $f(S)dt$ is the incremental of S given by the original deterministic system. The term $d_s S$ is the additional stochastic incremental of S caused by the perturbation scheme.

In this paper, we generalize this scheme to mixed type of tensor fields θ . A key point is indeed to link the state variable S with some tensor field θ . The choice of θ can then correspond to the conservation laws of certain quantities. We describe in detail how to calculate T_t^* and T_{t*} , and present results for several examples corresponding to different choices of θ . We also discussed about the conservation laws for these examples. We emphasize that Brenier’s theorem merely serves as the motivation but not the theoretical foundation of the proposed scheme, since the ‘optimality’ of the displacement vector field need to be rigorously defined for general tensor fields θ that are not positive differential n -forms.

Interestingly, similarities and differences can be studied between the proposed perturbation scheme and the existing stochastic physical SALT and LU settings Holm (2015); Mémin (2014); Resseguier et al. (2016). In particular, both SALT and LU equations can be recovered using a prescribed definition of the random diffeomorphism T_t used by the perturbation scheme. For illustration, a stochastic version of the thermal shallow water equation is presented. Compared with SALT and LU settings Holm (2015); Mémin (2014); Resseguier et al. (2016), the proposed perturbation scheme does not directly rely on the physics. Hence it is more flexible and can be applied to any PDE. Yet, the proposed derivation also provides interesting means to interpret the operator $\mathcal{L}_{d_{x_t} - u dt}$, appearing in the SALT equation. In terms of the optimal transportation, this term represents the infinitesimal forecast error at every forecast time step.

In order to apply the proposed perturbation scheme to any specific model, the parameters a and e_i must be determined specifically. Hence it is necessary to learn these parameters from existing data, experimental runs, or additional physical considerations Resseguier et al. (2020, 2021). We anticipate this framework naturally provides a new perspective on how to learn these parameters. Likely, this task will invoke the need of numerical algorithms to estimate the optimal transportation map for general differential k -forms or even mixed type of tensor fields. This will be subjects of future investigations.

Appendix A: Calculation of T_t^{-1}

Suppose that

$$T_t(x) = x + a\Delta t + e_i\Delta\eta_i. \quad (\text{A1})$$

550 We assume that T_t^{-1} has the following form of expression:

$$T_t^{-1}(x) = x + z\Delta t + b_i\Delta\eta_i. \quad (\text{A2})$$

Our goal is to find z and b_i . Then we have

$$\begin{aligned} x &= T_t(T_t^{-1}(x)) = T_t(x + z\Delta t + b_i\Delta\eta_i) \\ &= x + z\Delta t + b_i\Delta\eta_i + a \Big|_{x+z\Delta t+b_i\Delta\eta_i} \Delta t + e_i \Big|_{x+z\Delta t+b_i\Delta\eta_i} \Delta\eta_i \end{aligned} \quad (\text{A3})$$

555 Similar to the derivation in section (3.1), we apply Taylor expansion and Ito's lemma, and drop the terms of higher-order infinitesimal:

$$\begin{aligned} a \Big|_{x+a\Delta t+b_i\Delta\eta_i} \Delta t &= a \Big|_x \Delta t + o(\Delta t) \\ e_i \Big|_{x+z\Delta t+b_i\Delta\eta_i} \Delta\eta_i &= e_i \Big|_x \Delta\eta_i + e_{ip} b_i^p \Big|_x \Delta t + o(\Delta t). \end{aligned} \quad (\text{A4})$$

Therefore

$$560 \quad x = T_t(T_t^{-1}(x)) = x + (z + a + e_{ip}b_i^p)\Delta t + (b_i + e_i)\Delta\eta_i + o(\Delta t). \quad (\text{A5})$$

This implies that

$$b_i + e_i = 0 \quad (\text{A6})$$

$$z + a + e_{ip}b_i^p = 0 \quad (\text{A7})$$

Therefore

$$565 \quad b_i = -e_i \quad (\text{A8})$$

$$z = -a + e_{ip}e_i^p, \quad (\text{A9})$$

or equivalently,

$$T_t^{-1}(x) = x + (-a + e_{ip}e_i^p)\Delta t - e_i\Delta\eta_i \quad (\text{A10})$$

Appendix B: Derivation of $T_t^*\theta$

570 Given coordinates (x^1, \dots, x^n) , when θ is a differential k -form, it can be written as

$$\theta = \sum_{i_1 < \dots < i_k} f^{i_1, \dots, i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k}. \quad (\text{B1})$$

Since T_t^* is linear, we may assume that

$$\theta = f dx^{i_1} \wedge \cdots \wedge dx^{i_k} \tag{B2}$$

for some $1 \leq i_1 < \cdots < i_k \leq n$. Let $T_t(x) = (T_t^1(x), \dots, T_t^n(x))$, then

$$575 \quad (T_t^* \theta)(x) = f(T_t(x)) dT_t^{i_1} \wedge \cdots \wedge dT_t^{i_k}. \tag{B3}$$

We calculate $f(T_t(x))$ and $dT_t^{i_1} \wedge \cdots \wedge dT_t^{i_k}$ separately. We denote $\Delta x = T_t(x) - x = a\Delta t + e_i \Delta \eta_i$, and H_f the Hessian matrix of f . At a given time t , f is assumed independent from the noises $\Delta \eta_i(t)$. Then

$$f(T_t(x)) = f(x + \Delta x) = f(x) + \langle \nabla f, \Delta x \rangle + \frac{1}{2} (\Delta x)^\top H_f \Delta x + o((\Delta x)^2) \tag{B4}$$

$$= f(x) + \langle \nabla f, a\Delta t + e_i \Delta \eta_i \rangle + \frac{1}{2} e_i^\top H_f e_i (\Delta \eta_i)^2 \tag{B5}$$

$$580 \quad + \mathcal{O}((\Delta t)^2) + \mathcal{O}(\Delta t \Delta \eta_i) + o((\Delta t)^2) + o((\Delta \eta_i)^2) + o(\Delta t \Delta \eta_i) \tag{B6}$$

According to Ito's lemma $d\eta_i d\eta_j = \delta_{ij} dt$, and we can replace $(\Delta \eta_i)^2$ with Δt . Hence

$$f(T_t(x)) = f(x) + \langle \nabla f, a \rangle \Delta t + \langle \nabla f, e_i \rangle \Delta \eta_i + \frac{1}{2} e_i^\top H_f e_i \Delta t + o(\Delta t) \tag{B7}$$

$$= f(x) + \left(\langle \nabla f, a \rangle + \frac{1}{2} e_i^\top H_f e_i \right) \Delta t + \langle \nabla f, e_i \rangle \Delta \eta_i + o(\Delta t). \tag{B8}$$

Next,

$$585 \quad T_t^* (dx^{i_1} \wedge \cdots \wedge dx^{i_k}) = dT_t^{i_1} \wedge \cdots \wedge dT_t^{i_k} \\ = (dx^{i_1} + da^{i_1} \Delta t + de_i^{i_1} \Delta \eta_i) \wedge \cdots \wedge (dx^{i_k} + da^{i_k} \Delta t + de_i^{i_k} \Delta \eta_i). \tag{B9}$$

Note that da^{ij} and de_i^{ij} refer to the spatial differentiation. Again, we apply the "discrete version" of Ito's rule $(\Delta \eta_i)^2 = \Delta t$, and collect all the terms of order $\mathcal{O}(\Delta t)$ and $\mathcal{O}(\Delta \eta_i)$:

$$590 \quad T_t^* (dx^{i_1} \wedge \cdots \wedge dx^{i_k}) = dx^{i_1} \wedge \cdots \wedge dx^{i_k} + \left(\sum_{s=1}^k dx^{i_1} \wedge \cdots \wedge da^{i_s} \wedge \cdots \wedge dx^{i_k} \right) \Delta t \\ + \left(\sum_{s=1}^k dx^{i_1} \wedge \cdots \wedge de_i^{i_s} \wedge \cdots \wedge dx^{i_k} \right) \Delta \eta_i \\ + \left(\sum_{s < r} dx^{i_1} \wedge \cdots \wedge de_i^{i_s} \wedge \cdots \wedge de_i^{i_r} \wedge \cdots \wedge dx^{i_k} \right) \Delta t \\ + o(\Delta t) \tag{B10}$$

According to the chain rule, $da^{i_s} = \partial_{x^j} a^{i_s} dx^j$, $de_i^{i_s} = \partial_{x^j} e_i^{i_s} dx^j$. Note that $\partial_{x^j} e_i^{i_s}$ refers to the i_s -th component of $\partial_{x^j} e_i$, where $\partial_{x^j} e_i = \frac{\partial e_i}{\partial x^j}$ and $e_i(x) \in \mathbb{R}^n$ is the i -th basis vector field of T_t . Hence

$$\begin{aligned}
595 \quad & T_t^*(dx^{i_1} \wedge \cdots \wedge dx^{i_k}) \\
& = dx^{i_1} \wedge \cdots \wedge dx^{i_k} + \left(\sum_{s=1}^k \partial_{x^j} a^{i_s} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^{i_k} \right) \Delta t \\
& \quad + \left(\sum_{s=1}^k \partial_{x^j} e_i^{i_s} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^{i_k} \right) \Delta \eta_i \\
& \quad + \left(\sum_{s < r} \partial_{x^j} e_i^{i_s} \partial_{x^l} e_i^{i_r} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^l \wedge \cdots \wedge dx^{i_k} \right) \Delta t \\
& \quad + o(\Delta t)
\end{aligned} \tag{B11}$$

600 Combining Eqs.(B8) and (B11), with application of Ito's lemma, all terms of order $o(\Delta t)$ are then removed, to obtain

$$\begin{aligned}
& T_t^* \theta = f(T_t(x)) T_t^*(dx^{i_1} \wedge \cdots \wedge dx^{i_k}) \\
& = \theta + \left\{ (\langle \nabla f, a \rangle + \frac{1}{2} e_i^\top H_f e_i) dx^{i_1} \wedge \cdots \wedge dx^{i_n} \right. \\
& \quad + \sum_{s=1}^k f \partial_{x^j} a^{i_s} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^{i_k} \\
& \quad + \left(\sum_{s < r} f \partial_{x^j} e_i^{i_s} \partial_{x^l} e_i^{i_r} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^l \wedge \cdots \wedge dx^{i_k} \right) \\
605 \quad & \left. + \left(\sum_{s=1}^k \langle \nabla f, e_i \rangle \partial_{x^j} e_i^{i_s} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^{i_k} \right) \right\} \Delta t \\
& \quad + \left\{ \langle \nabla f, e_i \rangle dx^{i_1} \wedge \cdots \wedge dx^{i_k} + \sum_{s=1}^k f \partial_{x^j} e_i^{i_s} dx^{i_1} \wedge \cdots \wedge dx^j \wedge \cdots \wedge dx^{i_k} \right\} \Delta \eta_i \\
& \quad + o(\Delta t).
\end{aligned} \tag{B12}$$

To simplify Eq.(B12), wedge algebra is applied and the high-order infinitesimal $o(\Delta t)$ is ignored. Accordingly, $T_t^* \theta$ is more compactly written as

$$610 \quad T_t^* \theta = \theta + \mathcal{M}(\theta) \Delta t + \mathcal{N}_i(\theta) \Delta \eta_i, \tag{B13}$$

for some differential k -forms $\mathcal{M}(\theta)$ and $\mathcal{N}_i(\theta)$.

Author contributions. The first author contributed the main idea and mathematical derivation, and wrote the first draft of the paper. The second author contributed all the physical interpretations of the derived equations, a much more detailed introduction, and a more detailed comparison of the proposed scheme with the LU equation. The third author contributed his insight and enthusiasm on this problem and worked on the writing of the draft. This work is done based on the fruitful discussion of all the three authors since the very early stage.

Competing interests. The authors declare no competing interests.

Acknowledgements. The authors would like to express their gratitude towards Wei Pan, Darryl Holm, Dan Crisan, Long Li, and Etienne Mémin for their patient explanation and insightful discussion. The research of YZ was supported by the ANR Melody project when he was a postdoc at Ifremer. The research VR is supported by the company SCALIAN DS and by France Relance through the MORAANE project.
620 The research BC is supported by ERC EU SYNERGY Project No. 856408-STUOD, and the the support of the ANR Melody project.

References

- Anderson, J. L.: An adaptive covariance inflation error correction algorithm for ensemble filters, *Tellus A: Dynamic Meteorology and Oceanography*, 59, 210 – 224, 2007.
- Berner, J., Shutts, G., Leutbecher, M., and Palmer, T.: A spectral stochastic kinetic energy backscatter scheme and its impact on flow-
625 dependent predictability in the ECMWF ensemble prediction system, *Journal of the Atmospheric Sciences*, 66, 603–626, 2009.
- Brenier, Y.: Polar factorization and monotone rearrangement of vector-valued functions, *Communications on Pure and Applied Mathematics*, 44, 375–417, 1991.
- Brzeźniak, Z., Capiński, M., and Flandoli, F.: Stochastic partial differential equations and turbulence, *Mathematical Models and Methods in Applied Sciences*, 1, 41–59, 1991.
- 630 Buizza, R., Miller, M., and Palmer, T.: Stochastic representation of model uncertainties in the ECMWF Ensemble Prediction System, *Quarterly Journal Royal Meteorological Society*, 125, 2887–2908, 1999.
- Chern, S. S., Chen, W. H., and Lam, K. S.: *Lectures on Differential Geometry*, WORLD SCIENTIFIC, <https://doi.org/10.1142/3812>, 1999.
- Flandoli, F.: The interaction between noise and transport mechanisms in PDEs, *Milan Journal of Mathematics*, 79, 543–560, 2011.
- Franzke, C., O’Kane, T., Berner, J., Williams, P., and Lucarini, V.: Stochastic climate theory and modeling, *Wiley Interdisciplinary Reviews: Climate Change*, 6, 63–78, 2015.
635
- Gottwald, G. and Melbourne, I.: Homogenization for deterministic maps and multiplicative noise, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 469, 2013.
- Gugole, F. and Franzke, C.: Numerical development and evaluation of an energy conserving conceptual stochastic climate model, *Mathematics of climate and weather forecasting*, in press, 219.
- 640 Harlim, J. and Majda, A. J.: Catastrophic filter divergence in filtering nonlinear dissipative systems, *Communications in Mathematical Sciences*, 8, 27–43, 2010.
- Hasselmann, K.: Stochastic climate models. Part I: theory, *Tellus*, 28, 473–485, 1976.
- Holm, D. D.: Variational principles for stochastic fluid dynamics, *Proceedings. Mathematical, Physical, and Engineering Sciences / The Royal Society*, 471, 2015.
- 645 Holm, D. D. and Luesink, E.: Stochastic Wave–Current Interaction in Thermal Shallow Water Dynamics, *Journal of Nonlinear Science*, 31, 2021.
- Houtekamer, P. and Mitchell, H. L.: A Sequential Ensemble Kalman Filter for Atmospheric Data Assimilation, *Monthly Weather Review*, 129, 123–137, 2001.
- Jain, A., Timofeyev, I., and Vanden-Eijnden, E.: Stochastic mode-reduction in models with conservative fast sub-systems, arXiv preprint
650 arXiv:1410.3004, 2014.
- Kotsuki, S., Ota, Y., and Miyoshi, T.: Adaptive covariance relaxation methods for ensemble data assimilation: Experiments in the real atmosphere, *Quarterly Journal of the Royal Meteorological Society*, 143, 2001–2015, 2017.
- Kunita, H.: *Stochastic flows and stochastic differential equations*, vol. 24, Cambridge university press, 1997.
- Leith, C.: Atmospheric predictability and two-dimensional turbulence, *Journal of the Atmospheric Sciences*, 28, 145–161, 1971.
- 655 Leon, A. B. D.: On the effect of stochastic Lie transport noise on fluid dynamic equations, Ph.D. thesis, Imperial College London, 2021.

- Leroux, S., Brankart, J.-M., Albert, A., Brodeau, L., Molines, J.-M., Jamet, Q., Le Sommer, J., Penduff, T., and Brasseur, P.: Ensemble quantification of short-term predictability of the ocean dynamics at kilometeric-scale resolution: A Western Mediterranean test-case, *Ocean Science Discussions*, pp. 1–36, 2022.
- 660 Leutbechner, M., Ollinaha, P., Lock, S.-J., Lang, S., Bechtold, P., Beljaars, A., Bozzo, A., Forbes, R., Haiden, T. Hogan, R., and Sandu, I.: Stochastic representations of model uncertainties in the IFS, in: *ECMWF/WWRP Workshop: Model Uncertainty*, ECMWF, Reading, 2016.
- Li, H., Kalnay, E., and Miyoshi, T.: Simultaneous estimation of covariance inflation and observation errors within an ensemble Kalman filter, *Quarterly Journal of the Royal Meteorological Society*, 135, 2009.
- Majda, A. J., Timofeyev, I., and VandenEijnden, E.: Models for stochastic climate prediction., *Proceedings of the National Academy of Sciences of the United States of America*, 96 26, 14 687–91, 1999.
- 665 Mémin, E.: Fluid flow dynamics under location uncertainty, *Geophysical & Astrophysical Fluid Dynamics*, 108, 119–146, 2014.
- Mikulevicius, R. and Rozovskii, B.: Stochastic Navier-Stokes Equations for Turbulent Flows, *SIAM J. Math. Anal.*, 35, 1250–1310, 2004.
- Miyoshi, T.: The Gaussian Approach to Adaptive Covariance Inflation and Its Implementation with the Local Ensemble Transform Kalman Filter, *Monthly Weather Review*, 139, 1519–1535, 2011.
- 670 Orszag, S.: Analytical theories of turbulence, *Journal of Fluid Mechanics*, 41, 363–386, 1970.
- Penland, C. and Sardeshmukh, P.: The optimal growth of tropical sea surface temperature anomalies, *Journal of climate*, 8, 1999–2024, 1995.
- Pope, S.: Lagrangian PDF methods for turbulent flows, *Annu. Rev. Fluid Mech.*, 26, 23–63, 1994.
- Poterjoy, J.: A Localized Particle Filter for High-Dimensional Nonlinear Systems, *Monthly Weather Review*, 144, 59–76, 2016.
- Raanes, P. N., Bocquet, M., and Carrassi, A.: Adaptive covariance inflation in the ensemble Kalman filter by Gaussian scale mixtures, 675 *Quarterly Journal of the Royal Meteorological Society*, 145, 53 – 75, 2019.
- Resseguier, V.: Mixing and fluid dynamics under location uncertainty, Ph.D. thesis, Université Rennes 1, 2017.
- Resseguier, V., Mémin, É., and Chapron, B.: Geophysical flows under location uncertainty, Part I Random transport and general models, *Geophysical & Astrophysical Fluid Dynamics*, 111, 149 – 176, 2016.
- Resseguier, V., Pan, W., and Fox-Kemper, B.: Data-driven versus self-similar parameterizations for stochastic advection by Lie transport and 680 location uncertainty, *Nonlinear Processes in Geophysics*, 27, 209–234, 2020.
- Resseguier, V., Li, L., Jouan, G., Dérian, P., Mémin, E., and Chapron, B.: New trends in ensemble forecast strategy: uncertainty quantification for coarse-grid computational fluid dynamics, *Archives of Computational Methods in Engineering*, 28, 215–261, 2021.
- Reynolds, C., Leutbecher, M., Batté, L., Chen, S., Christensen, H., Klasa, C., Pegion, P., Plant, B., Raynaud, L., Roberts, N., Sandu, I., Singleton, A., Sommer, M., Swinbank, R., Tennant, W., and Theis, S.: Reports from working group 3 : What are the pros/cons of existing 685 model uncertainty schemes and how should these be measured?, in: *ECMWF/WWRP Workshop: Model Uncertainty*, ECMWF, Reading, 2016.
- Sapsis, T. and Majda, A.: A statistically accurate modified quasilinear Gaussian closure for uncertainty quantification in turbulent dynamical systems, *Physica D: Nonlinear Phenomena*, 252, 34–45, 2013.
- Schlee, F. H., Standish, C. J., and Toda, N. F.: Divergence in the Kalman Filter, *AIAA Journal*, 5, 1114–1120, 1966.
- 690 Tandeo, P., Ailliot, P., Bocquet, M., Carrassi, A., Carrassi, A., Miyoshi, T., Pulido, M., Pulido, M., and Zhen, Y.: A Review of Innovation-Based Methods to Jointly Estimate Model and Observation Error Covariance Matrices in Ensemble Data Assimilation, *Monthly Weather Review*, 2020.

- Tibshirani, R. and Knight, K.: The Covariance Inflation Criterion for Adaptive Model Selection, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 61, 1999.
- 695 Warneford, E. S. and Dellar, P. J.: The quasi-geostrophic theory of the thermal shallow water equations, *Journal of Fluid Mechanics*, 723, 374 – 403, 2013.
- Ying, Y. and Zhang, F.: An adaptive covariance relaxation method for ensemble data assimilation, *Quarterly Journal of the Royal Meteorological Society*, 141, 2015.
- Zhen, Y. and Harlim, J.: Adaptive error covariances estimation methods for ensemble Kalman filters, *Journal of Computational Physics*, 294, 700 619–638, 2015.