

1 Title: **European native oyster reef ecosystems are universally Collapsed**

2

3 Authors: Philine S.E. zu Ermgassen^{1*}, Hannah McCormick², Alison Debney², José Fariñas-

4 Franco³, Celine Gamble², Chris Gillies⁴, Boze Hancock⁵, Ane T. Laugen⁶, Stéphane Pouvreau⁷,

5 Joanne Preston⁸, William G. Sanderson⁹, Åsa Strand¹⁰, Ruth H. Thurstan¹¹

6

7 ¹ Changing Oceans Group, School of Geosciences, University of Edinburgh, James Hutton Rd,

8 King's Buildings, Edinburgh EH9 3FE, UK. ; 0000-0002-3409-0644

9 philine.zu.ermgassen@ed.ac.uk *Corresponding Author

10 ² Zoological Society of London, ZSL London Zoo, Regent's Park, London, NW1 4RY, UK.

11 ³ Marine and Freshwater Research Centre and Department of Natural Resources and the

12 Environment. School of Science and Computing, Atlantic Technological University, Old

13 Dublin Road, Galway H91 T8NW, Ireland

14 ⁴ SeaGen Aquaculture, Newhaven, Victoria, Australia and TropWATER, James Cook

15 University, Townsville, Australia.

16 ⁵ Global Oceans Team, The Nature Conservancy, University of RI, USA, 02882

17 ⁶ Department of Natural Sciences, Centre for Coastal Research, University of Agder,

18 Postboks 422, NO-4604 Kristiansand, Norway

19 ⁷ LEMAR, Ifremer, 29840 Argenton en Landunvez, France

20 ⁸ Institute of Marine Sciences, University of Portsmouth, Ferry Road, Portsmouth, PO4 9LY,

21 UK

22 ⁹ Institute of Life and Earth Sciences, EGIS, Heriot-Watt University, Edinburgh, Scotland

23 EH14 4AS, UK

24 10 Department of Environmental Intelligence, IVL Swedish Environmental Research
25 Institute, Kristineberg 566, 451 78 Fiskebäckskil,
26 Sweden
27 11 Centre for Ecology and Conservation, University of Exeter, Cornwall, TR13 0JE, UK.; 0000-
28 0002-8045-1631

29

30

31 **Keywords**

32 *Ostrea edulis*, shellfish reef, threats, IUCN Ecosystem Red List, habitat restoration, historical
33 ecology, shifted baseline

34

35

36 **Abstract**

37 Oyster reefs are often referred to as the temperate functional equivalent of coral reefs. Yet
38 evidence for this analogy is lacking for the European native species *Ostrea edulis* and its
39 biogenic habitat. Recently assembled historical data provide a unique opportunity to
40 develop a robust definition for this ecosystem type, confirm that *O. edulis* are biogenic reef
41 builders, and assess its current conservation status. Today, *O. edulis* typically occur as
42 scattered individuals or, in a few locations, as dense clumps over a few m², however,
43 historically *O. edulis* reef ecosystems persisted at large scales. A key finding is that *O. edulis*
44 reef ecosystems should therefore be assessed at the >ha scale.

45

46 Using the IUCN Red list of Ecosystems Framework, we conclude the European native oyster
47 reef ecosystem type is Collapsed under three of five criteria (A: reduction in geographic
48 distribution, B: restricted geographic range, and D: disruption of biotic processes and
49 interactions). Criterion C (environmental degradation) was assessed as data deficient and
50 Criterion E (quantitative risk analysis) was not completed as the ecosystem was already
51 deemed collapsed.

52

53 Our assessment has far reaching implications for conservation policy and action, and shows
54 that the scale of current restoration efforts fall far short of what is necessary for ecosystem
55 recovery.

56

57

58 Introduction

59 Oyster reef ecosystems were once a widely distributed ecosystem type in temperate coastal
60 seas and estuaries globally (Beck et al., 2011; Gillies et al., 2020; zu Ermgassen et al., 2012,
61 Williams et al. in press, Thurstan et al. in review a, b). In historical accounts, oyster reefs
62 were described as the temperate equivalent of coral reefs (Hamm 1881), forming elevated
63 three-dimensional structures over large areas (Williams, 1837; zu Ermgassen et al., 2012). In
64 parts of the world, these reefs were so massive they represented a navigational hazard
65 (Hinke, 1916, Dumain 1832). Reef-building oyster species have a long history of human
66 exploitation, with evidence of extensive collection and consumption of oysters from
67 middens across Europe, Asia, Australia and the Americas illustrating that oysters have been
68 an important food and cultural resource in coastal communities for thousands of years
69 (Szabó & Amesbury 2011, Rick et al. 2016, Fariñas-Franco et al. 2018, Thurstan et al., 2020,
70 Astrup et al. 2021). Dramatic declines in the extent of oyster reefs were documented
71 globally following European colonisation and the industrial revolution (Alleway & Connell,
72 2015; Beck et al., 2011; Thurstan et al., 2013; zu Ermgassen et al., 2012, Thurstan et al.,
73 2020). Although pollution, harsh winters and changes to hydrological conditions were noted
74 in historical texts to have caused localised extinctions (Krøyer 1837, Royal Commission 1866,
75 Holmes 1927), the primary driver of loss of oyster reef ecosystems has been extraction by
76 fishing (Went 1961, Thurstan et al. in review a).

77

78 Today, ecosystem-forming oyster species including those in the genus of *Saccostrea*,
79 *Crassostrea* and *Ostrea species* still form high relief, complex reef structures in many
80 temperate estuaries across the globe (Bahr & Lanier, 1981; Hedgpeth, 1954; Gillies et al.
81 2017, Norgard et al. 2018), albeit over significantly smaller areas, and with a substantially

82 reduced habitat complexity relative to historical records (zu Ermgassen et al. 2012; Gillies et
83 al. 2018). This is, however, not the case in Europe, where (with the exception of a few
84 locations) the native oyster, *Ostrea edulis*, predominantly exists as scattered individuals,
85 occurring at densities rarely greater than 1 individual · m⁻² (Thorngren et al. 2019; Allison et
86 al. 2020; Pouvreau et al, 2023). Recognition of the degraded status of *O. edulis* populations
87 and the reef ecosystems they form is reflected in their many conservation designations,
88 with *O. edulis* and its habitat recognised as threatened and/or declining in Region II and
89 Region III (Greater North Sea and Celtic Sea respectively) under the OSPAR convention
90 (OSPAR Commission 2009), its recognition by some member states under the “Reefs”
91 feature of the Habitats Directive (European Council 1992), and its inclusion in some
92 Biodiversity Action Plans (e.g. UKBAP 1999). These designations are critically important for
93 the protection of *O. edulis* habitats and ecosystems.

94

95 The growing recognition of the degraded and yet ecologically important status of *O. edulis*
96 oyster reef ecosystems has resulted in increasing efforts to restore the habitat at numerous
97 locations across its native range (Preston et al. 2021; Pouvreau et al. 2023). Restoration
98 activities are set to increase both in scope and scale as the focus on restoration of degraded
99 terrestrial and marine habitats gains momentum at a national and international level
100 (United Nations General Assembly, 2020, EU Commission 2022). An improved description of
101 the ecosystem’s main physical and biological attributes prior to significant disturbance is
102 thus essential for developing a historical baseline against which current and future recovery
103 efforts can establish targets, assess progress and determine the efficacy of conservation
104 interventions including for the application of the EU Restoration Law.

105

106 Recent work by Thurstan et al. (in review a,b) documented the historical locations,
107 ecosystem characteristics and extents of *O. edulis* reef ecosystems in Europe. The broad
108 spatial scale and highly resolved nature of the data presents a unique historical record for
109 marine ecosystems, which provides a novel opportunity to visualise the form and extent of
110 *O. edulis* reef ecosystems prior to its widespread degradation. Thurstan et al. (in review a)
111 found that *O. edulis* reefs were historically widely distributed throughout coastal waters of
112 Europe and North Africa, as well as in the southern North Sea, to 80 m depth. They gathered
113 numerous descriptions of reefs extending over many ha or even km² and forming complex
114 structures with vertical relief “composed of several layers” (Levasseur 2006), with “oysters,
115 almost placed one on top of the other like stones, forming a wall” (Marsili 1715).
116 Additionally, they identified sources describing the rich benthic community associated with
117 this complex structured habitat. These descriptions all serve to elucidate the historical
118 extent and physical and biological characteristics of oyster reef ecosystems throughout
119 Europe, based primarily on a 1800-1930 baseline (Thurstan et al. in review b).

120

121 Here, we follow the IUCN Red List of Ecosystems Framework (Bland et al. 2017) to assess
122 the current status of the European native oyster reef ecosystem. Ecosystem red lists are one
123 of the headline indicators in the monitoring framework for the post-2020 Global Biodiversity
124 Framework (Kunming-Montreal Global Biodiversity Framework), and therefore play a critical
125 role in providing structured evidence to support policy development and decision making.
126 We develop a much needed definition of the European native oyster reef ecosystem type
127 based on historical descriptions of *O. edulis* reefs and analogous shellfish ecosystems
128 formed by *Ostrea* species in other regions. Our definition and Red List Assessment can be
129 used to guide national and Europe-wide conservation strategies, prioritise and monitor

130 restoration action, inform resource management, and to raise public awareness to support
131 management and protection policy.

132

133 **Methods**

134 Applying the IUCN Red List of Ecosystems assessment framework to an ecosystem requires
135 that the ecosystem in its functional and collapsed states are clearly defined, and that the
136 pathways to collapse (drivers of decline) are clearly identified. Once the definitions are
137 clear, the current status of the ecosystem can be assessed by applying those definitions to
138 current data. Assessment is undertaken by applying each of five criterion (A [reduction in
139 distribution], B [restricted distribution], C [environmental degradation], D [disruption of
140 biotic processes] and E [quantitative assessment of risk]), with the final classification equal
141 to the highest threat level identified (Bland et al. 2017).

142

143 *Development of ecosystem and ecosystem Collapse definitions*

144 Following the IUCN Red List of Ecosystems Framework (Bland et al. 2017), we developed a
145 comprehensive definition of the European native oyster ecosystem type created by *O.*
146 *edulis*, and the associated threshold of collapse. While definitions of oyster habitats exist
147 (e.g. OSPAR Commission 2009), comparison of modern definitions with historical sources
148 illustrate clearly that modern definitions describe a habitat in a degraded state, as opposed
149 to an ecosystem type (Table S1a-e, Figure S1). Modern definitions based on small habitat
150 patches, fail to adequately describe the full physical and biological attributes and key
151 processes of the ecosystem and are therefore ill equipped to be applied as ecological
152 baselines against which ecosystem condition can be assessed.

153

154 We reviewed existing definitions of shellfish ecosystems globally to identify important
155 ecosystem attributes representative of shellfish ecosystems irrespective of ecosystem-
156 forming species, including the IUCN Global Ecosystem Typology (Keith et al. 2020) and
157 “biogenic reef habitat” *sensu* Brown et al. (1997) (Table S2a-c). These included the structure
158 and form of reefs, the formation of biogenic structure created by oysters when occurring in
159 high densities, the contribution of dead shells to maintain a positive shell budget (Hemeon
160 et al. 2020, Solinger et al. 2022), and the spatial scale at which these ecosystems historically
161 functioned (Table S2a-c). Historical evidence of past *O. edulis* reef structure, spatial scale,
162 and functions were assessed (Table S2b-d). Oyster density is a key attribute of oyster reef
163 condition (zu Ermgassen et al. 2021; Pouvreau et al. 2021), but quantitative assessments of
164 densities were not available from the baseline period of assessment. Furthermore, habitat
165 descriptions from the principle period of documentary evidence (1800-1930) often
166 described reefs which were known to be overexploited (Möbius 1877, Krøyer 1837, Table
167 S1a-c, Figure S1). Our threshold densities for ecosystem assessment were therefore based
168 on cumulative evidence from descriptions of the ecosystem, catch rate information, and
169 quantitative information from related species in other geographies, as well as the current
170 understanding of *O. edulis* reef formation (Pouvreau et al. 2021, Table S2d-e).

171

172 Our definition of the collapsed European native oyster reef ecosystem and the threshold of
173 collapse was derived from relevant literature and expert knowledge on the pathways to
174 collapse.

175

176 *Collating baseline and current ecosystem data*

177

178 The risk of collapse of the European native oyster reef ecosystems was assessed using the
179 IUCN Red List of Ecosystems guidelines set out in Rodriguez et al. (2015) and Bland et al.
180 (2017). Significant declines in *O. edulis* reefs were observed in the 1800s (Thurstan et al.
181 2013, Royal Commission 1866), or earlier (Went 1961, Giovio 1524, Levasseur, 2006). In
182 contrast, data from the past 50 years is limited to biological records of species occurrence
183 (e.g. through the Global Biodiversity Information Facility), which may represent shell
184 remains as opposed to live individuals, catch data which are often challenging to
185 disaggregate from aquaculture production (e.g. FAO), or stock assessments from a few
186 geographically limited locations (e.g. Tully and Clarke 2012, Thorngren et al. 2019, Jenkin et
187 al. 2023, The Marine Institute and Bord Iascaigh Mhara 2023). Due to this lack of more
188 recent data, the IUCN Ecosystem Red Listing risk assessment was undertaken relative to a c.
189 1750 baseline using the historical data in Thurstan et al. (in review a).

190

191 Thurstan et al. (in review a, b) extracted historical data on the presence, condition and
192 function of *O. edulis* reefs from targeted searches of government records, nautical charts,
193 popular media, and scientific journals. Identified locations were further assessed for
194 whether there was high or low confidence that oysters were present in sufficient abundance
195 to be reef-forming based on descriptions, landings data and catch per unit effort
196 information. These data were primarily recorded between 1800-1930 (Thurstan et al. (in
197 review b), yet we consider them representative of a c. 1750s baseline for oyster reef
198 presence. It is recognised that, following exploitation, oyster reef ecosystem quality declines
199 before reef locations are extirpated (zu Ermgassen et al. 2012, Table S1a). Locations
200 recorded as oyster reefs between 1800-1930 therefore likely represent reefs that were
201 extant in 1750, even if their condition at the time of recording was degraded. Furthermore,

202 historical sources described discovering new oyster grounds until the late 19th century,
203 highlighting that some *O. edulis* reef habitats remained unexploited until this time period
204 (Thurstan et al. in review a). Only historical locations assigned a high confidence that *O.*
205 *edulis* were once present at densities at which they formed reefs were included in our
206 assessment. Areas in the southern Mediterranean with historical records of *O. edulis* were
207 therefore excluded from the analysis, as there is low confidence in the historical
208 documentary evidence that these populations were reef building, although it is not known
209 whether this lack of evidence is the result of an already shifted baseline, or the natural
210 ecological condition of *O. edulis* populations in the region (Thurstan et al. in review a).

211

212 To identify locations where *O. edulis* reef currently meets our ecosystem definition, data
213 describing the location and ecosystem attributes of remaining oyster habitats were
214 identified by using the Google search engine using the terms “oyster, *Ostrea*, COUNTRY” for
215 each country known to fall within the historical distribution of the European native oyster
216 (based on Thurstan et al. in review b). The searches were undertaken between September
217 2022 and June 2023. Where recent surveys were not identified for a country, or where data
218 were inconclusive, data was requested from local experts.

219

220 *Application of Red List criteria*

221 The collated historical and recent data were used to assess the risk of collapse in Criterion A
222 (reduction in geographic distribution), under sub criterion 3 (relative to a 1750 baseline) and
223 Criterion B (restricted geographic range), under sub criterion 1 and 2 (Area of Occupancy,
224 Extent of Occurrence). Spatial data representing the location of sites (historical and current)
225 meeting the *O. edulis* reef ecosystem definition, were processed using QGIS software

226 version 3.24 (QGIS Development Team). For Criteria A, the change in the extent was
227 assessed both by comparing the number of locations where *O. edulis* reef was recorded
228 historically and presently, and by comparing the described extents of *O. edulis* reef, where
229 such data were available historically (Thurstan et al. in review a and b). To determine the
230 Extent of Occurrence (Criterion B1), point locations of all high confidence historical *O. edulis*
231 reef occurrences (Thurstan et al. in review a,b), were used to draw a minimum convex
232 polygon using the Minimum Bounding Geometry tool in QGIS 3.24. To determine the Area
233 of Occupancy (Criterion B2), a grid layer of 10x10km squares was then overlaid, and all grid
234 squares that contained at least one historical oyster reef point were identified (Coordinate
235 Reference System used is ETRS89-extended / LAEA Europe).

236

237 Criterion C (environmental degradation) was assessed using shell substrate as a defining
238 non-living feature of oyster reefs, the loss of which “reduces the capacity of the ecosystem
239 to sustain its characteristic biota” (Bland et al. 2017). We sought to identify data from the
240 literature, suitable for assessing the current distribution and extent of shell substrate.

241 Criterion D assessed the risk of disruption of biotic processes, relative to a 1750 baseline
242 (sub criterion 3; Bland et al. 2017). Evidence for the ecological functions associated with *O.*
243 *edulis* reefs historically and recently were examined. Criteria E (quantitative risk analysis),
244 and Criteria B (subcriterion 1 and 2) require a thorough understanding of the existing and
245 future threats (pathways to collapse) facing the ecosystem being assessed. Expert opinion
246 and literature review were used to assess whether identified threats still impact, or have the
247 potential to impact, *O. edulis* reef condition.

248

249 **Results**

250

251 *Definition of the European native oyster reef ecosystem type*

252 European native oyster reefs are found from 20-42 ppt salinity, where the underlying
253 sediment is not overly mobile and current speeds are typically 0.05-0.45 m s⁻¹ (Pogoda et al.,
254 2023). European native oyster reefs have a rich, diverse, and distinct associated community,
255 supporting a higher species richness and abundance of species than surrounding
256 unstructured habitats (Kennon et al. 2023). While populations of *O. edulis* persist as non-
257 native species on the eastern coast of North America, its native range is restricted to Europe
258 from Norway to the African coast of the Mediterranean and into the Black Sea (Thurstan et
259 al. in review a).

260

261 European native oyster reefs can be defined as areas with high densities of multiple size
262 classes of *Ostrea* spp., on a shell dominated substrate (Table 1). The ecosystem contains
263 patches with high oyster density, often forming clumps of oysters and creating a complex
264 three-dimensional structure (Bodvin et al. 2011, Kennon et al. 2023, Thurstan et al. in
265 review a, b, Table S2b-d, Figure 1). Associated bivalve species, such as *O. stentina* (in the
266 Mediterranean) and *Mytilus edulis* also contribute to the reef structure (Möbius 1877), but
267 the primary ecosystem engineer is *O. edulis*. These patches may be interspersed with areas
268 of low structural complexity (Figure 1b) or other habitats, such as eelgrass beds or maerl
269 beds (Abancourt 1842, Marine Institute & Bord Iascaigh Mhara 2023).

270

271 The European native oyster ecosystem is biogenic, with previous generations forming the
272 underlying substrate on which the reef is built (Thurstan et al. in review a, Table S2b-d,
273 Table 1). While accurate mapping of *O. edulis* reefs historically is rare, the spatial scale at

274 which reefs historically persisted is best visualised through mapping undertaken on the
275 French oyster beds in the early 1900's (Joubin & Guérin-Ganivet 2009, Thustan et al. b,
276 Figure 2). While individual reefs within the ecosystem may predominantly form on the scale
277 of hectares (Figure 2a), multiple reefs persist within a wider ecosystem (Figure 2b). A
278 resilient ecosystem persists on a broader biogeographical scale, or metapopulation, even
279 when individual reefs within it are smothered or negatively impacted (e.g. Krøyer 1837). The
280 European native oyster reef ecosystem should therefore be assessed across a km² scale.

281

282 *Definition of European native oyster reef ecosystem Collapse*

283 The European native oyster reef ecosystem is considered collapsed at the point at which
284 there are no longer multiple size classes of oysters in the local population, gregarious
285 settlement leading to clumps are absent, and oysters do not contribute to biogenic
286 formation of three-dimensional structure at the hectare scale, assessed at the km² scale
287 (Table 1, Figure 1c). This results in a change from shell dominated substrate to sand, mud or
288 subtidal mixed sediment without significant (>25%) shell cover (Kasoar et al. 2015). The
289 associated community is therefore representative of the alternative underlying substrate,
290 which may be soft bottomed, low complexity habitat, cobble, or subtidal mixed sediments,
291 depending on the location. Where invasive species have moved in to occupy the niche
292 previously held by *O. edulis*, the associated community may instead reflect this shift. A
293 Collapsed European native oyster ecosystem does not span the expected depth range of the
294 species, nor the necessary spatial extent to classify as an ecosystem (>ha patches within km²
295 extents of European native oyster ecosystem). The collapsed European native oyster
296 ecosystem does not deliver ecosystem functions such as water filtration, nutrient cycling,
297 enhanced biodiversity, sediment stabilisation or shell production at significant scales.

298

299 *Pathways to collapse*

300 Oyster reef ecosystems are particularly sensitive to collapse, as oysters and their shells are
301 the preferred settlement substrate for oyster larvae (Rodriguez-Perez et al. 2019, Colsoul et
302 al. 2020). The removal of the biogenic habitat therefore disrupts or interrupts the life cycle
303 of the primary ecosystem engineer, *O. edulis*, and can tip the ecosystem into a state of
304 negative feedback (Figure 3). The loss of oysters or reduction in larval recruitment has
305 primarily occurred through the removal of oysters (i.e. fishing) (Thurstan et al. 2013).
306 Additionally, habitat disturbance from bottom towed gears (Ezgeta -Balić et al. 2021),
307 sedimentation (Sander et al. 2021), pollution (Helmer et al. 2019), invasive species (Drapkin
308 1963, Preston 2020a) and disease (Virvilis & Angelidis 2006, Culloty and Mulcahy 2007) all
309 play a role in the reduction or loss of oyster reefs in some locations (Helmer et al. 2019,
310 Pouvreau et al., 2023, Table S3). Finally, changes in salinity regime have also historically
311 played a role in the local extirpation (Krøyer 1837) or establishment (Collin 1884) of native
312 oyster populations.

313

314 **Applying the IUCN Ecosystem Red Listing criteria**

315

316 *Ecosystem Red listing Criterion A - reduction in distribution*

317 Criterion A considers the change in extent of the ecosystem type over time (Bland et al.
318 2017). No recent records of *O. edulis* persisting at higher densities over areas > ha were
319 identified throughout the native range of the European native oyster. As such, the extent of
320 the European native oyster reef ecosystem was deemed to have declined from being
321 present in 606 (10 km²) grid cells in c. 1750 (Figure 4b), to zero in the present day (Table S4).

322 In addition, while the spatial extent of oyster reef ecosystems was only documented from
323 317 of 1197 recorded historical locations, those locations encompassed a known reef area
324 of >1.7 million ha (Thurstan et al. in review a, b). There are no records of reef extent
325 meeting the ecosystem criteria of > ha (Table 1) in the present day (Table S4) and the
326 European native oyster reef ecosystem type is therefore deemed to be Collapsed under
327 category A3.

328

329 *Ecosystem Red listing Criterion B - restricted distribution*

330 Criterion B considers the current range of the ecosystem type (Bland et al. 2017). The
331 historical Extent of Occurrence was found to be 7,718,991 km² and the historical Area of
332 Occupancy was found to be 606 (10 km²) grid squares (Figure 4) . No current records of *O.*
333 *edulis* persisting at higher densities over areas > ha scales were identified (Table S4), which
334 qualifies the ecosystem type as Collapsed under sub criterion B1 and B2. Furthermore,
335 numerous threats were identified as still driving declines in *O. edulis* populations (Table S3).

336

337 *Ecosystem Red listing Criterion C - environmental degradation*

338 Criterion C considers the condition of abiotic attributes of the ecosystem which have a
339 defining role in ecological processes and/or the distribution of an ecosystem type (Bland et
340 al. 2017). The shell substrate underlying oyster reefs is a key abiotic attribute of the system.
341 While removal of living oysters by its nature removes the shell which would otherwise be
342 contributed to building the primary physical attribute of the ecosystem (Solinger et al.
343 2022), shell material can persist in the marine environment for millenia (Fariñas-Franco et
344 al. 2018, Sanders et al. 2021). This makes it challenging to assess the extent and condition of
345 areas of shell dominated substrate currently present in European seas and estuaries.

346 Although it can be unequivocally stated that the extent and condition of shell dominated
347 substrate has declined significantly over the past centuries as a result of destructive fishing
348 practices and increased sedimentation (Helmer et al. 2019, Sanders et al. 2021), it was not
349 possible to identify independent records of the extent and condition of oyster shell and shell
350 of species associated with oyster reef community (e.g. *Mytilus edulis*, Möbius 1877)
351 deposited on the seabed. While there are known locations where oyster and associated
352 bivalve shells are the dominant feature of the substrates (e.g. Todorova et al. 2009), there is
353 insufficient data reported to assess their extent and condition. This assessment therefore
354 deemed European native oyster reef ecosystem type to be data deficient under Criterion C.
355

356 *Ecosystem Red listing Criteria D - disruption of biotic processes*

357 Criterion D considers the degree to which biotic processes and interactions change within
358 the extent of the ecosystem. Oysters are both allogenic and autogenic ecosystem engineers,
359 substantially altering biotic processes and interactions both through their feeding activity
360 and their physical structure (Smyth and Roberts 2010, Kennon et al. 2023, Lee et al 2020,
361 2023). That *O. edulis* reef ecosystems have a distinct associated community is well
362 established (Krøyer 1837, Möbius 1877, summarised in Thurstan et al. in review a). While a
363 distinct associated community has also been recorded in remnant areas (Kennon et al.
364 2023), the loss of living oysters and their reef-associated community from the seafloor and
365 large-scale shift from structured reefs to sediments, has resulted in substantial changes to
366 the biotic community and biotic interactions (Reise 1982). The loss of trophic interactions is
367 perhaps most strikingly evidenced by the loss of oyster reefs as a key foraging ground for
368 shorebirds such as the oystercatcher (*Haematopus ostralegus*), which was described in 1801
369 as follows: “Oystercatcher, oyster thief, oyster collector.... The oystercatcher also swims, but

370 *is more likely to be seen walking along the beach. At low tide, it seems to be particularly*
371 *cheerful; then it runs around with a hooting sound, looking for its food, which consists*
372 *mainly of oysters. The bird knows how to break open the shells very skillfully, without hurting*
373 *its beak on the sharp edges. If they are closed too tightly, it hits them against a rock so that*
374 *they crack. If it can't find oysters, it will eat mussels, snails and other worms, even dead*
375 *animals.” (Lippold 1801, translated from German). Today, mussels, clams and worms*
376 *dominate the diet of *H. ostralegus* in Europe (Pol et al. 2009). *O. edulis* reefs historically*
377 *supported a range of ecological functions, however, modern data to quantify those*
378 *functions is lacking (zu Ermgassen et al. 2020). Overall, this assessment deemed the*
379 *European native oyster reef ecosystem type to be Collapsed under Criterion D.*

380

381 *Ecosystem Red listing Criterion E - quantitative assessment of risk*

382 Criterion E considers the probability of future ecosystem collapse. Threats to the existing
383 oyster populations in Europe were identified from the established pathways to collapse
384 (summarised in Figure 3, Table S3). It was, however, not possible to assess the probability of
385 future collapse, as the ecosystem type is already deemed collapsed (Criteria A, B and D). As
386 such, this assessment deemed Criterion E to be “Not Applicable” to the European native
387 oyster reef ecosystem type.

388

389 **IUCN Ecosystem Red Listing Assessment Outcome**

390 The overall threat ranking in the IUCN Ecosystem red listing assessment reflects the highest
391 risk ranking. In the case of the European native oyster reef ecosystem type, this threat
392 ranking is Collapsed (Table 2).

393

394 **Discussion**

395 That European native oyster reef ecosystems were assessed as Collapsed (Table 2) is a stark
396 finding, and should promote wider conversations about how much (or little) we know about
397 the status of our marine environments, and the subsequent implications of these
398 knowledge gaps for ocean policy and management. European native oyster reef ecosystems
399 historically covered millions of hectares of the European seafloor at a range of depths
400 (Thurstan et al. in review a,b), across which their reef structures, created by living and dead
401 shells, formed vertical relief and interstitial spaces that supported highly diverse, distinct
402 associated communities (Figure 1, Table S2b-d). These ecosystems would have provided
403 important ecosystem functions such as larval output, enhanced biodiversity, water
404 filtration, nutrient cycling, sediment stabilisation and enhanced productivity at multiple
405 trophic levels (Lippold 1801, Christianen et al. 2018, zu Ermgassen et al. 2020, Lee et al.
406 2020, 2023, Kennon et al. 2023). They formed reef systems, where individual reefs could be
407 many ha in size, with numerous reefs occurring across the system at the scale of several km²
408 (Figure 2). In contrast, today there are no known locations where reefs with high densities
409 of *O. edulis* are found at the scale of more than 0.1 ha in extent (Table S4).

410

411 Our assessment of the European native oyster reef ecosystem relative to a c. 1750 baseline
412 using the IUCN framework provides a deeper time dimension than existing assessments
413 which classify European native oyster habitats based on more recent data, as being
414 threatened and/or declining throughout much of their range (OSPAR Commission, 2009),
415 Endangered (Mediterranean infralittoral oyster beds, European Environment Agency 2022)
416 or Critically Endangered (*Ostrea edulis* beds on Atlantic shallow sublittoral muddy mixed
417 sediments, EU Red List of habitats, Gubbay et al. 2016). In general, current definitions of *O.*

418 *edulis* habitats (e.g. OSPAR Commission, 2009, Cameron 2022) reflect a significantly
419 degraded ecological state, a “shifted baseline”, relative to the historically described
420 ecosystem (Thurstan et al. in review a, Table S1a,b, S2a). This is because declines in the
421 condition of *O. edulis* reef ecosystems were already being documented by the early 1700s
422 (Pontoppidan 1769, Krøyer 1837, Brehm 1872, Levasseur 2006, Table S1a, Figure S1), prior
423 to scientific monitoring or commonly accepted historical baselines (e.g. Möbius 1877,
424 Krøyer 1837, Table S1b). This shifted baseline presents a challenge for oyster restoration
425 both in policy and in practice. For example, *O. edulis* reefs were not included as biogenic
426 reefs during the process of developing UK marine SACs, as it was not believed they were
427 capable of forming reefs (Holt et al. 1998). In developing the definition of the *O. edulis* reef
428 ecosystem type, however, we illustrated that there was substantial historical evidence that
429 criteria for being considered a “biogenic reef habitat” *sensu* Brown et al. (1997), were met
430 by oyster reefs historically (Table S2b). Our ecosystem type definition can also serve to
431 inform an understanding of *O. edulis* reef “reference ecosystem” attributes (Gann et al
432 2019), the minimum population size, area or density needed for the ecosystem to recover,
433 all of which have been identified as a critical knowledge gap (Preston et al. 2020b, zu
434 Ermgassen et al. 2020, McAfee et al. 2021). While the definition developed here does not
435 provide a quantitative answer to this important question, it does illustrate that restoration
436 projects will have to be vastly scaled-up for ecosystem scale recovery to be achieved.

437

438 Our assessment was undertaken at a time when governments and NGOs are seeking to
439 address the important issue of scaling up ecological restoration efforts (as exemplified by
440 the UN Decade on Ecosystem Restoration and EU Nature Restoration Law). While European
441 native oyster reef restoration efforts have been pilot-scale to date, there are increasing

442 efforts to scale up, both in the nearshore and offshore (Preston et al. 2020b, zu Ermgassen
443 et al. 2021). Our findings and the developed ecosystem definition highlight how critical
444 these efforts are. In the past, high levels of ecosystem resilience to disturbances such as
445 harsh winters, sedimentation and predation was possible because of the large scale,
446 variable depth range and high abundance at which oysters were found (Thurstan et al. in
447 review a). European native oyster ecosystems no longer exist at a scale capable of providing
448 ecosystem resilience or function (Table S4), with each of these drivers of decline now
449 considered to be a significant threat at individual locations (Pouvreau et al., 2023, Helmer et
450 al. 2019, Table S3).

451

452 Despite the Collapsed status of the European native oyster reef ecosystem, the benefits
453 associated with the recovery of shellfish reefs, even at a smaller spatial scale, should not be
454 understated. In particular in locations where oyster populations are protected from harvest,
455 remnant *O. edulis* populations can build three dimensional complex habitats (Bodvin et al.
456 2011, Smyth et al. 2020, Pouvreau et al. 2023), and to support a diverse epibiotic
457 community and distinct associated community (Smyth and Roberts 2010, Kennon et al.
458 2023). Smaller scale habitat restoration efforts are a key stepping stone to larger scale
459 ecosystem restoration (zu Ermgassen et al. 2016), ultimately leading to a tipping point
460 where recovery is self-sustaining.

461

462 That the European native oyster reef ecosystem type is Collapsed is an important indicator
463 of the intensely degraded status of European marine benthic ecosystems, not only due to
464 the scale of loss of ecosystem function that their decline represents, but also because they
465 may be a proxy for other sensitive, less commercially important and therefore less well

466 historically documented ecosystems. The current Collapsed state of the European native
467 oyster reef ecosystem is therefore a powerful warning that the state of the European seas is
468 more dire than commonly acknowledged when limiting our assessments to more recent
469 baselines. This evidence should be taken into consideration when planning the long term
470 recovery of these highly impacted waters, for example when developing or applying the EU
471 Restoration Law.

472

473 **Acknowledgements**

474 This work was funded by the Flotilla Foundation. RHT was funded by the Convex Seascape
475 Survey, and the COST Action MAF-WORLD CA20102, supported by COST (European
476 Cooperation in Science and Technology), www.cost.eu. Figure 1 was produced by Maria
477 Eggertsen, funded by the European Research Council (ERC) under the European Union’s
478 Horizon 2020 research and innovation programme (grant agreement No 856488 – ERC
479 Synergy project “SEACHANGE: Quantifying the impact of major cultural transitions on
480 marine ecosystem functioning and biodiversity”). The authors would like to thank the Native
481 Oyster Restoration Alliance Historical Ecology working group, who’s earlier work identifying
482 historical locations of oyster reefs helped lay the foundations for this assessment, and Prof
483 Aydin and Dr Yildiz for their local knowledge.

484

485 **References- 80 max**

486 Abancourt (1842), Archives du SHD de Vincennes, MAR/CC5/623, Pêche des huîtres,
487 mémoires et documents divers, Rapport d’Abancourt au préfet maritime, 18 octobre
488 1842.

489 Alleway, H. K., & Connell, S. D. (2015). Loss of an ecological baseline through the
490 eradication of oyster reefs from coastal ecosystems and human memory.
491 *Conservation Biology*, 29, 795–804. <https://doi.org/10.1111/cobi.12452>

492 Allison, S., Hardy, M., Hayward, K., Cameron, T. C. & Underwood, G.J.C. (2020).
493 Strongholds of *Ostrea edulis* populations in estuaries in Essex, SE England and their
494 association with traditional oyster aquaculture: evidence to support a MPA
495 designation. *Journal of the Marine Biological Association of the United Kingdom*, 100,
496 27–36.

497 Astrup, P.M., Benjamin, J., Stankiewicz, F., Woo, K., McCarthy, J., Wiseman, C.,
498 Baggaley, P., Jerbić, K., Fowler, M., Skriver, C. & Bailey, G. (2021). A drowned
499 Mesolithic shell midden complex at Hjarnø Vesterhoved, Denmark and its wider
500 significance. *Quaternary Science Reviews*, 258, 106854.

501 Bahr, L.M., & Lanier, W.P. (1981). *The ecology of intertidal oyster reefs of the South*
502 *Atlantic coast: A community profile* (FWS/OBS-81/15; p. 105).

503 Beck, M. W., Brumbaugh, R. D., Airoidi, L., Carranza, A., Coen, L. D., Crawford, C.,
504 Defeo, O., Edgar, G. J., Hancock, B., Kay, M., Lenihan, H. S., Luckenbach, M. W.,
505 Toropova, C. L., Zhang, G., & Guo, X. (2011). Oyster reefs at risk and
506 recommendations for conservation, restoration and management. *BioScience*, 61(2),
507 107–116.

508 Bland, L. M., Keith, D., Miller, R. M., Murray, N.J., & Rodríguez, J. P. (2017).
509 *Guidelines for the application of IUCN Red List of Ecosystems categories and criteria*
510 [Resource]. [https://www.iucn.org/resources/publication/guidelines-application-](https://www.iucn.org/resources/publication/guidelines-application-iucn-red-list-ecosystems-categories-and-criteria)
511 [iucn-red-list-ecosystems-categories-and-criteria](https://www.iucn.org/resources/publication/guidelines-application-iucn-red-list-ecosystems-categories-and-criteria)

512 Bodvin, T., Mortensen, S., Jelmert, A., Strand, Ø. & Grefsrud, E.G. (2011): Utredning
513 av europeisk flatøsters *Ostrea edulis* L. – Kunnskapsoversikt med forslag til
514 handlingsplan. DN-utredning 10-2011.
515 [http://www.miljødirektoratet.no/old/dirnat/attachment/2857/DN-utredning-10-](http://www.miljødirektoratet.no/old/dirnat/attachment/2857/DN-utredning-10-2011.pdf)
516 [2011.pdf](http://www.miljødirektoratet.no/old/dirnat/attachment/2857/DN-utredning-10-2011.pdf)

517 Brown AE, Burn AJ, Hopkins JJ, Way SF (1997) Joint Nature Conservation Committee.
518 The Habitats Directive: selection of Special Areas of Conservation in the UK. JNCC
519 Report No. 270. p.291. Peterborough, UK.

520 Brehm (1872) *Illustriertes Tierleben für Volk und Schule*, bearbeitet von Freidrich
521 Schrödler, Verlag des Bibliographischen Instituts.

522 Cameron, T. 2022. *Defining Oyster Beds in the Blackwater Estuary*. NECR411. Natural
523 England.

524 Collin, J. S. (1884). *Om Limfjordens tidligere og nuværende marine fauna, med særligt*
525 *hensyn til bløddyraunaen*. Danmark Ministeriet for kirke- og undervisningsvæsenet.
526 Kjøbenhavn : Gyldendal.

527 Colsoul, B., Pouvreau, S., Di Poi, C., Pouil, S., Merk, V., Peter, C., Boersma, M. &
528 Pogoda, B. (2020). Addressing critical limitations of oyster (*Ostrea edulis*)
529 restoration: Identification of nature-based substrates for hatchery production and
530 recruitment in the field. *Aquatic Conservation: Marine and Freshwater Ecosystems*,
531 30, 2101–2115

532 Culloty, S. C. & Mulcahy, M.F. (2007). *Bonamia ostreae in the native oyster Ostrea*
533 *edulis*. A review. Marine Environment and Health Series.

534 Drapkin, E. I. (1963). Effect of *Rapana bezear* Linne on the Black Sea fauna. *Doklady*
535 *Akademiia nauk SSSR 151 700–703 Translation Biological Science Section, 151, 964-*
536 *966.*

537 Dumain, L. (1832) Survey of the Coast of Louisiana [1806-1807, communicated to the
538 Senate 1811]. Class IV, Commerce and Navigation. In: *Document No. 158 in the*
539 *American State Papers. Vol. 7. (Eds. W. Lowrie, M. St. Clair Clarke), Gales and Seaton,*
540 *Washington, pp. 839-842.*

541 Ezgeta -Balić, D., Vrgoč, N., Isajlović, I., Medvešek, D., Vujević, A., Despalatović, M. &
542 Cvitković, I. (2021). Comparison of beam trawl catch, by-catch and discard in fishing
543 and non-fishing areas – a case study from the northern Adriatic Sea. *Mediterranean*
544 *Marine Science, 22, 108–120.*

545 EU Commission (2022) Proposal for a regulation of the European Parliament and of
546 the Council on nature restoration Accessed on November 14th 2023 [https://eur-](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A52022PC0304)
547 [lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A52022PC0304](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A52022PC0304)

548 European Council Directive (1992) Council Directive 92/43/EEC
549 <http://data.europa.eu/eli/dir/1992/43/2013-07-01> (accessed October 2023)

550 European Environment Agency (2022) European Red List of Habitats - enhanced by
551 EEA, May 2022. [https://www.eea.europa.eu/en/datahub/datahubitem-](https://www.eea.europa.eu/en/datahub/datahubitem-view/de2276d8-e295-4cd7-89c9-88812065db87)
552 [view/de2276d8-e295-4cd7-89c9-88812065db87](https://www.eea.europa.eu/en/datahub/datahubitem-view/de2276d8-e295-4cd7-89c9-88812065db87) [Accessed December 10th 2023]

553 Fariñas-Franco, J.M., Allcock, A.L., Roberts, D. (2018) Protection alone may not
554 promote natural recovery of biogenic habitats of high biodiversity damaged by
555 mobile fishing gears, *Marine Environmental Research 135, 18–28.*
556 [https://doi.org/10.1016/j.marenvres.2018.01.009.](https://doi.org/10.1016/j.marenvres.2018.01.009)

557 Fariñas-Franco, J.M., Pearce, B., Mair, J.M., Harries, D.B., MacPherson, R.C., Porter,
558 J.S., Reimer, P.J. & Sanderson, W.G. (2018). Missing native oyster (*Ostrea edulis* L.)
559 beds in a European Marine Protected Area: Should there be widespread restorative
560 management? *Biological Conservation*, 221, 293–311

561 Gann, G.D., McDonald, T., Walder, B., Aronson, J., Nelson, C.R., Jonson, J., Hallett,
562 J.G., Eisenberg, C., Guariguata, M.R., Liu, J., Hua, F., Echeverría, C., Gonzales, E.,
563 Shaw, N., Decler, K. & Dixon, K.W. (2019). *International principles and standards for*
564 *the practice of ecological restoration*. Society for Ecological Restoration, Washington,
565 D.C. 20005 U.S.A.

566 Gillies, C.L., McLeod, I.M., Alleway, H.K., Cook, P., Crawford, C., Creighton, C.,
567 Diggles, B., Ford, J., Hamer, P., Heller-Wagner, G. and Lebrault, E. (2018) Australian
568 shellfish ecosystems: Past distribution, current status and future direction. *PLoS One*,
569 13(2), p.e0190914.

570 Gillies, C. L., Castine, S., Alleway, H. K., Crawford, C., Fitzsimons, J. A., Hancock, B.,
571 Koch, P., McAfee, D., McLeod, I. M., & zu Ermgassen, P. S. E. (2020). Conservation
572 status of oyster reef ecosystem of Southern and Eastern Australia. *Global Ecology*
573 *and Conservation*, e00988. <https://doi.org/10.1016/j.gecco.2020.e00988>

574 Giovio P. (1524) *Libro de' pesci romani*. Translation by C. Zancaruolo, Venice 1560.
575 First edition in latin language was printed in Rome in 1524 and 1527, titled *De*
576 *piscibus marinis lacustribus fluviatilibus item de testaceis ac salsamentis liber*.

577 Ritter von Hamm, W. (1881) *Gesammelte kleine Schriften von Dr. Wilhelm Ritter*
578 *von Hamm*.

579 Hemeon, K.M., Ashton-Alcox, K.A., Powell, E.N., Pace, S.M., Poussard, L.M., Solinger,
580 L.K. & Soniat, T.M. (2020). Novel shell stock–recruitment models for *Crassostrea*

581 *virginica* as a function of regional shell effective surface area, a missing link for
582 sustainable management. *Journal of Shellfish Research*, 39, 633–654.

583 Hinke, W. J. (1916) Report of the Journey of Francis Louis Michel from Berne,
584 Switzerland, to Virginia, October 2, 1701-December 1, 1702. Part II. *The Virginia*
585 *Magazine of History and Biography* 24, 113-141.

586 Hedgpeth, J. W. (1954). Bottom communities of the Gulf of Mexico. *Fishery Bulletin*
587 *U.S. Fish and Wildlife Service*, 55, 203–216.

588 Holmes, H.B. (1927). *An investigation of saw dust pollution in relation to oysters in*
589 *Yaquina Bay*. Department of Commerce, Bureau of Fisheries, California

590 Jenkin, A., Trundle, C., Daniels, C., Sturgeon, S. & Street, K. (2023) Fal oyster survey
591 summary report 2023. p13. Cornwall Inshore Fisheries and Conservation Authority.

592 Joubin L, Guérin-Ganivet J (2009). Cartes des gisements de coquilles comestibles des
593 côtes de France. Ifremer. [https://doi.org/10.12770/c502ed30-0d7d-11de-a4b1-](https://doi.org/10.12770/c502ed30-0d7d-11de-a4b1-000086f6a603)
594 [000086f6a603](https://doi.org/10.12770/c502ed30-0d7d-11de-a4b1-000086f6a603)

595 Kasoar, T., zu Ermgassen, P.S.E., Carranza, A., Hancock, B. & Spalding, M. (2015).
596 New opportunities for conservation of a threatened biogenic habitat: a worldwide
597 assessment of knowledge on bivalve-reef representation in marine and coastal
598 Ramsar Sites. *Marine and Freshwater Research*, 66, 981–988.

599 Kennon, N.A., Robertson-Jones, A., Jemmett, S., Hugh-Jones, T., Bell, M.C. &
600 Sanderson, W.G., 2023. Rotational fishing enables biodiversity recovery and provides
601 a model for oyster (*Ostrea edulis*) habitat restoration. *PLoS One*, 18(3), p.e0283345.

602 Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. and Kingsford, R.T. (eds.) (2020). The IUCN
603 Global Ecosystem Typology 2.0: Descriptive profiles for biomes and ecosystem
604 functional groups. Gland, Switzerland: IUCN.

605 Lee, H.Z., Davies, I.M., Baxter, J.M., Diele, K. & Sanderson, W.G., 2020. Missing the
606 full story: First estimates of carbon deposition rates for the European flat oyster,
607 *Ostrea edulis*. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 30(11),
608 pp.2076-2086.

609 Lee, H.Z., Davies, I.M., Baxter, J.M., Diele, K. & Sanderson, W.G., 2023. A blue carbon
610 model for the European flat oyster (*Ostrea edulis*) and its application in
611 environmental restoration. *Aquatic Conservation: Marine and Freshwater*
612 *Ecosystems*.

613 Levasseur, O., 2006. Histoire de l'huître en Bretagne, Skol Vreizh, Morlaix. 83 p.

614 Lippold, G.H.C., Funke, C.P., (1810) Neues Natur- und Kunstlexicon enthaltend die
615 wichtigsten und gemeinnützigsten Gegenstände aus der Naturgeschichte,
616 Naturlehre, Chemie und Technologie.

617 McAfee, D., Costanza, R. & Connell, S.D. (2021). Valuing marine restoration beyond
618 the 'too small and too expensive.' *Trends in Ecology & Evolution*, 36, 968–971

619 Marine Institute & Bord Iascaigh Mhara (2023). Shellfish Stocks and Fisheries Review
620 2022: An assessment of selected stocks. Marine Institute., Galway, Ireland

621 Mortensen, S., Børgwald, M., Strohmeier, T., Strand, Ø., Larsson, S.B., Laugen, A.T.,
622 Reamon, M.C., & Marcussen, J.B. (2023). Kartlegging av østersbestander i Rogaland i
623 2021 og 2023 - innsamling av data som grunnlag for bestandsforvaltning. Rapport fra
624 havforskningen 2023-55 ISSN: 1893-4536. 22 s.
625 <https://www.hi.no/hi/nettrapporter/rapport-fra-havforskningen-2023-55>

626 Möbius, K.A. (1877). *Die Auster und die Austernwirthschaft*. Wiegandy, Hempel & Parey,
627 Berlin.

628 Norgard, TC, Bigg, MI, MacConnachie, SEM, Finney, JL and Gillespie, GE (2018) Index site
629 surveys for Olympia oysters (*Ostrea lurida* Carpenter, 1864) in British Columbia – 2009
630 to 2017. Can. Tech. Rep. Fish. Aquat. Sci. 3153: viii + 88 p.

631 Philpots, J. R. (1891). *Oysters and all about them* (Vol. 2). John Richardson and Co.

632 Pogoda, B., Hausen, T., Rothe, M., Bakker, F., Hauser, S., Colsoul, B., Dureuil, M.,
633 Krause, J., Heinicke, K., Pusch, C., Eisenbarth, S., Kreutle, A., Peter, C., & Pesch, R.
634 (2023). Come, tell me how you live: Habitat suitability analysis for *Ostrea edulis*
635 restoration. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 33(7), 678–
636 695. <https://doi.org/10.1002/aqc.3928>

637 Pol, M. van de, Ens, B.J., Oosterbeek, K., Brouwer, L., Verhulst, S., Tinbergen, J.M.,
638 Rutten, A.L. & Jong, M.D. (2009). Oystercatchers' bill shapes as a proxy for diet
639 specialization: More differentiation than meets the eye. *Ardea*, 97, 335–347.

640 Pontoppidan E (1769) Den Danske Atlas. København, Dannemark.

641 Pouvreau, S., Cochet, H., Bargat, F., Petton, S., Le Roy, V., Guillet, T., & Potet, M.
642 (2021). Current distribution of the residual flat oysters beds (*Ostrea edulis*) along the
643 west coast of France . SEANOE . <https://doi.org/10.17882/79821>

644 Pouvreau S., Lapègue S., Arzul I., Boudry P. (2023). Fifty years of research to counter
645 the decline of the European flat oyster (*Ostrea edulis*): a review of French
646 achievements and prospects for the restoration of remaining beds and revival of
647 aquaculture production. *Aquatic Living Resources*, 36, 13 (18p.).
648 <https://doi.org/10.1051/alr/2023006>

649 Preston J, Fabra M., Johnson E., Helmer L., Hendy I. (2020a) Interactions of larval
650 dynamics and substrate preference have ecological significance for benthic
651 biodiversity and *Ostrea edulis* Linnaeus, 1758 in the presence of *Crepidula fornicata*.

652 Journal of Aquatic Conservation: Marine and Freshwater Ecosystems. DOI:
653 10.1002/aqc.3446.

654 Preston J., Gamble, C., Debney, A., Helmer, L., Hancock, B. and zu Ermgassen, P.S.E.
655 (eds) (2020b). *European Native Oyster Habitat Restoration Handbook*. The Zoological
656 Society of London, UK., London, UK.

657 Reise, K. (1982). Long-term changes in the macrobenthic invertebrate fauna of the
658 Wadden Sea: are polychaetes about to take over? *Netherlands Journal of Sea
659 Research*, 16, 29–36

660 Rick, T.C., Reeder-Myers, L.A., Hofman, C.A., Breitburg, D., Lockwood, R., Henkes, G.,
661 Kellogg, L., Lowery, D., Luckenbach, M.W., Mann, R., Ogburn, M.B., Southworth, M.,
662 Wah, J., Wesson, J. & Hines, A.H. (2016). Millennial-scale sustainability of the
663 Chesapeake Bay Native American oyster fishery. *Proceedings of the National
664 Academy of Sciences*, 113, 6568–6573.

665 Rodriguez, J. P., Keith, D. A., Rodriguez-Clark, K. M., Murray, N. J., Nicholson, E.,
666 Regan, T. J., Miller, R. M., Barrow, E. G., Bland, L. M., Boe, K., Brooks, T. M., Oliveira-
667 Miranda, M. A., Spalding, M., & Wit, P. (2015). A practical guide to the application of
668 the IUCN Red List of Ecosystems criteria. *Philosophical Transactions of the Royal
669 Society B: Biological Sciences*, 370(1662). <https://doi.org/10.1098/rstb.2014.0003>

670 Rodriguez-Perez, A., James, M., Donnan, D.W., Henry, T.B., Møller, L.F. & Sanderson,
671 W.G., (2019). Conservation and restoration of a keystone species: Understanding the
672 settlement preferences of the European oyster (*Ostrea edulis*). *Marine Pollution
673 Bulletin*, 138, pp.312-321.

674 Royal Commission (1866) Report of the Commissioners appointed to inquire into the
675 sea fisheries of the United Kingdom, with appendix and minutes of evidence.
676 London: Eyre and Spottiswoode.

677 Sander, L., Hass, H.C., Michaelis, R., Groß, C., Hausen, T. & Pogoda, B. (2021). The
678 late Holocene demise of a sublittoral oyster bed in the North Sea. *PLoS One*, 16,
679 e0242208.

680 Smyth, D. & Roberts, D. (2010). The European oyster (*Ostrea edulis*) and its epibiotic
681 succession. *Hydrobiologia*, 655, 25–36.

682 Solinger, L.K., Ashton-Alcox, K.A., Powell, E.N., Hemeon, K.M., Pace, S.M., Soniat, T.M. &
683 Poussard, L.M. (2022). Oysters beget shell and vice versa: generating management
684 goals for live oysters and the associated reef to promote maximum sustainable yield
685 of *Crassostrea virginica*. *Canadian Journal of Fisheries and Aquatic Sciences*, 79,
686 1241–1254.

687 Szabó, K., Amesbury, J.R. 2011 Molluscs in a world of islands: The use of shellfish as
688 a food resource in the tropical island Asia-Pacific region, *Quaternary International*,
689 239,8-18, <https://www.sciencedirect.com/science/article/pii/S1040618211001285>

690 Thorngren, L., Bergström, P., Holthuis, T.D. & Lindegarth, M. (2019). Assessment of
691 the population of *Ostrea edulis* in Sweden: A marginal population of significance?
692 *Ecology and Evolution*, 00:1-12.

693 Thurstan, R.H., McCormick, H., Preston, J., Ashton, E.C., Bennema, F.P., Bratoš
694 Cetinić, A., Brown, J.H., Cameron, T.C., da Costa, F., Donnan, D., Ewers, C.,
695 Fortibuoni, T., Galimany, E., Giovanardi, O., Grancher, R., Grech, D., Hayden-Hughes,
696 M., Helmer, L., Jensen, K.T., Juanes, J.A., Latchford, J., Moore, A.B.M., Moutopoulos,
697 D.K., Nielsen, P., von Nordheim, H., Ondiviela, B., Peter, C., Pogoda, B., Poulsen, B.,

698 Pouvreau, S., Roberts, C.M., Scherer, C., Smaal, A.C., Smyth, D., Strand, Å,
699 Theodorou, J.A., & zu Ermgassen, P.S.E. (in review a). The world was our oyster:
700 Records reveal the vast historical extent of European oyster reef ecosystems.
701 <https://doi.org/10.32942/X20W43>

702 Thurstan, R.H., McCormick, H., Preston, J., Ashton, E.C., Bennema, F.P., Bratoš
703 Cetinić, A., Brown, J.H., Cameron, T.C., da Costa, F., Donnan, D., Ewers, C.,
704 Fortibuoni, T., Galimany, E., Giovanardi, O., Grancher, R., Grech, D., Hayden-Hughes,
705 M., Helmer, L., Jensen, K.T., Juanes, J.A., Latchford, J., Moore, A.B.M., Moutopoulos,
706 D.K., Nielsen, P., von Nordheim, H., Ondiviela, B., Peter, C., Pogoda, B., Poulsen, B.,
707 Pouvreau, S., Scherer, C., Smaal, A.C., Smyth, D., Strand, Å, Theodorou, J.A., & zu
708 Ermgassen, P.S.E. (in review b). Historical dataset details the distribution, extent and
709 form of lost *Ostrea edulis* reef ecosystems. <https://doi.org/10.32942/X28C99>

710 Thurstan, R. H., Diggles, B. K., Gillies, C. L., Strong, M. K., Kerkhove, R., Buckley, S. M.,
711 King, R. A., Smythe, V., Heller-Wagner, G., Weeks, R., Palin, F., & McLeod, I. (2020).
712 Charting two centuries of transformation in a coastal social-ecological system: A
713 mixed methods approach. *Global Environmental Change*, *61*, 102058.
714 <https://doi.org/10.1016/j.gloenvcha.2020.102058>

715 Thurstan, R. H., Hawkins, J. P., Raby, L., & Roberts, C. M. (2013). Oyster (*Ostrea*
716 *edulis*) extirpation and ecosystem transformation in the Firth of Forth, Scotland.
717 *Journal for Nature Conservation*, *21*(5), 253–261.
718 <https://doi.org/10.1016/j.jnc.2013.01.004>

719 Todorova, V., Micu, D. & Klisurov, L. (2009). Unique oyster reefs discovered in the
720 Bulgarian Black Sea. *Comptes Rendus de l'Académie bulgare des Sciences*, *62*, 871–
721 874

722 Tully, O., Clarke, S. (2012). The Status and Management of Oyster (*Ostrea edulis*) in
723 Ireland. Irish Fisheries Investigations No. 24. The Marine Institute, Rinville,
724 Oranmore, Ireland.

725 United Nations General Assembly. (2020a). *Decade on Ecosystem Restoration*.
726 Available at: <https://www.decadeonrestoration.org> [Accessed 25/01/22]

727 UKBAP, (1999). Native oyster (*Ostrea edulis*). Species Action Plan. In UK Biodiversity
728 Group. Tranche 2 Action Plans. Vol.V. Maritime Species and Habitats. English Nature
729 for the UK Biodiversity Group, Peterborough.

730 Virvilis, C. & Angelidis, P. (2006). Presence of the parasite *Marteilia* sp. in the flat
731 oyster (*Ostrea edulis* L) in Greece. *Aquaculture*, 259, 1–5

732 Went, Arthur E. J. (1961) Historical notes on the oyster fisheries of Ireland.
733 *Proceedings of the Royal Irish Academy. Section C: Archaeology, Celtic Studies,*
734 *History, Linguistics, Literature*, 62, 195–223. JSTOR,
735 <http://www.jstor.org/stable/25505107>.

736 Williams, J. L. (1837). *The Territory of Florida: Or Sketches of the Topography, Civil*
737 *and Natural History, of the Country, the Climate, and the Indian Tribes, from the First*
738 *Discovery to the Present Time*. A. T. Goodrich.

739 Williams, J.M., Lau, S.C.Y., Thomas, M., Tsao, C.S., & Russell, B.D. (in press). A long
740 history: social-ecological systems as drivers of oyster reef loss in the Pearl River Delta
741 and the broader Asian region. *Oceanography and Marine Biology: An Annual Review*,
742 volume 62.

743 zu Ermgassen, P.S.E., Bos, O., Debney, A., Gamble, C., Glover, A., Pogoda, B.,
744 Pouvreau, S., Sanderson, W., Smyth, D. & Preston, J. (eds) (2021). *European Native*

745 *Oyster Habitat Restoration Monitoring Handbook*. The Zoological Society of London,
746 UK., London, UK.

747 zu Ermgassen, P. S. E., Spalding, M. D., Blake, B., Coen, L. D., Dumbauld, B., Geiger,
748 S., Grabowski, J. H., Grizzle, R., Luckenbach, M., McGraw, K. A., Rodney, B., Ruesink,
749 J. L., Powers, S. P., & Brumbaugh, R. D. (2012). Historical ecology with real numbers:
750 Past and present extent and biomass of an imperilled estuarine ecosystem.
751 *Proceedings of the Royal Society B*, 279, 3393–3400.
752 <https://doi.org/10.1098/rspb.2012.0313>

753 zu Ermgassen, P., Hancock, B., DeAngelis, B., Greene, J., Schuster, E., Spalding, M. &
754 Brumbaugh, R.D. (2016). *Setting objectives for oyster habitat restoration using*
755 *ecosystem services: A manager's guide*. The Nature Conservancy, Arlington VA.

756 zu Ermgassen, P.S.E., Thurstan, R.H., Corrales, J., Alleway, H., Carranza, A., Dankers,
757 N., DeAngelis, B., Hancock, B., Kent, F., McLeod, I., Pogoda, B., Liu, Q. & Sanderson,
758 W.G. (2020). The benefits of bivalve reef restoration: A global synthesis of
759 underrepresented species. *Aquatic Conservation: Marine and Freshwater*
760 *Ecosystems*, 30, 2050–2065.

761

762

763

764 **Figures**

765 Figure 1. Artist's impression of a European native oyster reef ecosystem, based on historical
766 descriptions of associated species and habitat forms from Thurstan et al. in review a. Panel
767 A illustrates high density and relief oyster reef, which may on a larger scale be interspersed
768 with patches or lower complexity habitat (Panel B). Panel C illustrates the degraded habitat
769 structure now representative of oyster habitats. Artist: Maria Eggertsen.

770

771 Figure 2. A) Histogram of reported sizes of oyster reefs from the historical (<1910) literature.
772 B) An example map illustrating one of the French charts from which oyster reef extent was
773 extracted, digitised from *Joubin and Guérin-Ganivet 1910*. The chart illustrates how reefs in the
774 Bay of Cancale at various stages of degradation due to overfishing were mapped at the > ha
775 scale, but were distributed over many km² of the bay.

776

777 Figure 3. Pathways to collapse as identified by literature review and expert opinion.

778

779 Figure 4. Past (c.1750) Extent of Occurrence (A) and Area of Occupancy (B) of the European
780 native oyster (*Ostrea edulis*) reef ecosystem, based on locations identified by Thurstan et al.
781 in review a,b as having high confidence that oyster reef was historically present.

782 Coordinate Reference System: ETRS89-extended / LAEA Europe.

783

784

785

786

787 **Supporting Material**

788 **Supporting materials S1:** Summary of evidence of an early (<1800) shifted baseline in *Ostrea edulis*
789 reef condition.

790

791 **Supporting materials S2** Summary of the evidence underpinning the definition of the European
792 native oyster reef ecosystem.


793

794 **Supporting materials S3.** Summary of evidence of ongoing threats to *Ostrea edulis* populations in
795 Europe. Threats listed relate to those illustrated in Figure 3 within the main manuscript.

796

797 **Supporting materials S4** Summary of literature relating to the current presence of *Ostrea edulis* reef
798 ecosystems in coastal nations within the native range of the species.


799



800

801 Figure 1. Artist's impression of a European native oyster reef ecosystem, based on historical
802 descriptions of associated species and habitat forms from Thurstan et al. in review a. Panel
803 A illustrates high density and relief oyster reef, which may on a larger scale be interspersed
804 with patches or lower complexity habitat (Panel B). Panel C illustrates the degraded habitat
805 structure now representative of oyster habitats. Artist: Maria Eggertsen.

806



807

808 Figure 2. A) Histogram of reported sizes of oyster reefs from the historical (<1910) literature.

809 B) An example map illustrating one of the French charts from which oyster reef extent was

810 extracted, digitized from Joubin and Guérin-Ganivet 1910. The chart illustrates how reefs in


811 the Bay of Cancale at various stages of degradation due to overfishing were mapped at the >

812 ha scale, but were distributed over many km² of the bay.

813

814

815



816

817 Figure 3. Pathways to collapse as identified by literature review and expert opinion. Colours

818 indicate which component of the pathway to collapse is affected by each driver listed. Blue


819 indicates that the driver results in loss of substrate, green indicates that the driver results in

820 reduced larval survival and purple indicates the driver results in loss of adult oysters. Solid

821 circles indicate a unidirectional negative impact, whereas dashed circles indicate that the

822 effect may be positive or negative (see Table S3 for examples).

823



824

825 Figure 4. Past (c.1750) Extent of Occurrence (A) and Area of Occupancy (B) of the European
 826 native oyster (*Ostrea edulis*) reef ecosystem, based on locations identified by Thurstan et al.
 827 in review a,b as having high confidence that oyster reef was historically present.

828 Coordinate Reference System: ETRS89-extended / LAEA Europe.

829

830

831 **Tables**

832 Table 1 Proposed reef attributes (physical form and functional features) of the European
 833 native oyster reef ecosystem. Adapted from Gillies et al. (2020) to represent habitats built
 834 by *O. edulis*. This table aims to aid the delineation of reef ecosystems versus alternative
 835 ecosystems with oyster populations.

Attribute	Fully functional reef ecosystems	Partially functional reef ecosystems	Oyster populations within alternate ecosystems	References
1. Oyster density and size frequency	>20 live oysters m ⁻² representing multiple size classes	5-20 oysters m ⁻² representing multiple size classes	<5 oysters m ⁻² multiple size classes may not be represented	Pouvreau et al. 2021
2. Shell cover	>25 % cover		<25% cover	Kasoar et al. 2015 Kennon et al. 2023
3. Shell budget and reef height	Increasing or stable spatial extent and/or height.		Little or no evidence of shell substrate	Hemeon et al. 2020, Solinger et al. 2022
4. Patch size and number	Multiple patches of reef (> 5m ²), which may be separated by a few m to cover an area > 1h	Multiple patches of reef (> 5m ²), which may be separated by a few m to cover an area < 1h	Few or no patches of oyster reef	Krøyer 1837, Joubin and Guérin-Ganivet 2009

836

837 Table 2 IUCN Ecosystem Red Listing Assessment Outcomes, where CO = collapsed, DD =
 838 Data Deficient and NA= Not Applicable. The overall threat ranking is based on the highest
 839 risk ranking.

Criterion	A: Reduction in extent*	B: Restricted geographic distribution §	C: Environmental degradation* 0	D: Disruption of biotic processes	E: Quantitative analysis±	Overall threat ranking
1	DD	CO	DD	DD	NA	CO
2	NA	CO	NA	NA	NA	
3	CO	CO	DD	DD	NA	

840 Sub-criteria: *1= Past 50 years, 2= Next 50 years, 3=Since 1750; §1= Extent of Occurrence,

841 2= Area of Occupancy, 3= # threat locations; ±1= ≤50% in 50 years, 2= ≤20% in 50 years, 3=

842 ≤10% within 100 years.

Supplementary online materials

Title: European native oyster reef ecosystems are universally collapsed

Authors: Philine S.E. zu Ermgassen^{1*}, Hannah McCormick², Allison Debney², Jose M. Fariñas-Franco³, Celine Gamble², Chris Gillies⁴, Boze Hancock⁵, Ane T. Laugen⁶, Stéphane Pouvreau⁷, Joanne Preston⁸, William G. Sanderson⁹, Åsa Strand¹⁰, Ruth H. Thurstan¹¹

¹ Changing Oceans Group, School of Geosciences, University of Edinburgh, James Hutton Rd, King's Buildings, Edinburgh EH9 3FE, UK. ; 0000-0002-3409-0644 philine.zu.ermgassen@ed.ac.uk

² Zoological Society of London, ZSL London Zoo, Regent's Park, London, NW1 4RY, UK.

³ Marine and Freshwater Research Centre and Department of Natural Resources and the Environment. School of Science and Computing, Atlantic Technological University, Old Dublin Road, Galway H91 T8NW, Ireland

⁴ SeaGen Aquaculture, Newhaven, Victoria, Australia and TropWATER, James Cook University, Townsville, Australia.

⁵ Global Oceans Team, The Nature Conservancy, University of RI, USA, 02882

⁶ Department of Natural Sciences, Centre for Coastal Research, University of Agder, Postboks 422, NO-4604 Kristiansand, Norway

⁷ LEMAR, Ifremer, 29840 Argenton en Landunvez, France

⁸ Institute of Marine Sciences, University of Portsmouth, Ferry Road, Portsmouth, PO4 9LY, UK

⁹ Institute of Life and Earth Sciences, EGIS, Heriot-Watt University, Edinburgh, Scotland EH14 4AS, UK

¹⁰ Department of Environmental Intelligence, IVL Swedish Environmental Research Institute, Kristineberg 566, 451 78 Fiskebäckskil, Sweden

¹¹ Centre for Ecology and Conservation, University of Exeter, Cornwall, TR13 0JE, UK.; 0000-0002-8045-1631

Supporting materials S1 Summary of evidence of an early (<1800) shifted baseline in *Ostrea edulis* reef condition.

Table S1a: Summary of evidence that data on *Ostrea edulis* habitat form and function from the mid-1800s-early 1900s represents a shifted baseline relative to pristine or fully functioning *Ostrea edulis* reef ecosystems, i.e. that the condition of described beds, in particular those which provide insight into form and function, such as Möbius 1877 and Krøyer 1837, are describing oyster reefs which have been overfished and impacted for over a century.

Year of observation	Quotation	Location	Reference
1703	<p>"The tenant who, at the beginning of the last century, was in possession of the banks, petitioned (1700) to either be released from the lease or to receive a reduction in the expense. When, however, on further investigation of the matter, the conclusion was reached that he had destroyed the banks by fishing them too hard, he was deprived of them (1703), to let them rest until the expiry of the lease term (1706); but equally to pay the stipulated lease fee."</p>	Wadden Sea	Krøyer (1837)
1769	<p>"They are then 10 to 12 fathoms of water in the open sea, with the scrapers provided for the purpose, under full sail, cut up, taken out of the shoals, or put into the bath, whereupon the shoals are again lowered into the sea, or are then carried away until they have had as much as they desire. In such a boat, 3 or 4 scrapers can be towed, or if a dozen boat are set out, they can, in a hurry, scrape up 20 or 40,000 oysters. But as this oyster bed never rests, or is always abused in summer and winter, and also in the unused months, in a bad or disorderly manner, even by farmers far away and other strangers, it is already very ruined, and will soon be destroyed, unless it is put under proper inspection, that nothing is harvested for 2 or 3 years, and then only at certain times. which protection could best be put into effect, if the inhabitants of Skagen alone became entitled to this oyster catch."</p> <p>"On the Slesvig banks in 12 hrs in good conditions, 3-4000 oysters, or Aalbaekkerne which use bigger boats 5-5600 oysters in one day."</p> <p><u>In contrast</u> Kroyer (1837): "it is generally believed that the banks have improved since the Peace. In the first years after the war a boat could at most scrape a few hundred oysters a day; now, however, more than double that."</p>	Skagen, Denmark	Pontoppidan (1769)
~1830	<p>"In the Wash, about fifty years ago, were enormous oyster beds; one extending nearly the whole length of the Wash and continuing outside about 50 miles. One bed in particular, which was discovered about forty years ago, being (as the fishermen state) a fathom and a half deep, with</p>	The Wash, England	Harding (1882)

	nothing but oysters. Now everything is changed; the oysters on these beds are nearly exhausted, there not having been a fall of spat for a great number of years..."		
1849	"The oysters, when he was a boy, were as deep as the Town Hall is high. The bank was two miles long by half a mile wide" ... "Attributes the failure to the French working eight or ten miles to the southward of Brighton. This is the ground where there was a great oyster bed which has been dredged out. The oysters fell off first and the French tore away the oysters and made a trawl ground of it. It is 30 years since the oysters were torn away."	Brighton / English Channel	Buckland & Walpole (1879)
1870	"In the year 1870 a small oyster bed was discovered at the mouth of the Thames, north east from Whitstable. It was about 18 metres long by 6 metres broad. Forty-eight hours later 75 boats were there, close alongside of one another, fishing up the oysters.... Upon every old oyster which was taken were found only from nine to ten young ones of different ages. This bed had never been previously disturbed, and the oysters were accordingly found in their natural condition."	Margate Sands, England	Möbius (1883)
1871	"I remember when the boats could go out and dredge at Milford Haven, round by the Stack Rock, and each boat would get from 1000 to 1200 oysters in three or four hours. The last catch I took, I took 16 boats catch for a week; they had moderately fine weather during the week, and the largest catch was 600 oysters each, there were three men to a boat, and all of them did not bring in 5,000 in the gross, and that was starvation to the men."	Milford Haven, Wales	Anon (1876)
1874	"OUT TO THE OYSTER BEDS. We were, on Thursday last, one of a party of about a dozen persons who went out to the Oyster beds recently discovered off Douglas Bay. The swift sailing yacht Lizzie was at our disposal, and, with the fair wind which prevailed, she, in considerably under an hour, took us out to the scene of the destructive operations of the thirty boats which we found playing havoc with the bivalves. Anyone looking at these vessels from the shore would almost fancy that they were anchored in line, head on to the land, but we found that this was a delusion to which the distance lent a semblance of reality. The vessels were crossing and re-crossing the paths of each other in every possible way, and the men appeared to work with a degree of vigour which lent strength to the assertion made by one of them to us, that in a fortnight hence there would not be an oyster left on the bank. That they are already getting scarce is apparent from the fact that in the dredges large quantities of stones are now brought up, indicating that already the bed has been broken up to the bottom. Each boat has out, so far as we could see, four dredges, so that in all there were about 120 of these destructive engines at work on the bank. The dredges are lifted and emptied about every fifteen or twenty minutes. Say that this operation is performed three times an hour, it follows that there are 300 hauls per hour, or 3,000 in a working day of ten hours, made from the bank. Of course	Douglas, Isle of Man	Isle of Man Times (1874)

	<p>we have no means of knowing the exact extent of the oyster bed, but it is very evident that, no matter how extensive it may be, if these boats are permitted to "rag at it" (as the local phrase is) without intermission, in season and out of season, the total destruction of the bank is only a question of a very limited time. The operations of the boats extended over a space of about a mile and a-half to two miles in length, and about half a mile or so in width; so that we may presume that the bank is about that in dimensions. The vessels engaged are cutter-rigged, and appeared to be handy, smart boats, for, notwithstanding the deterring influence of the dredges out on the weather side, they went through the water at a good speed. The depth of water at the bank is from 30 to 35 fathoms. Out of the thirty boats engaged in the fishery, there was only one Manx boat (a Ramsey smack) that we saw...."</p>		
<1875	<p>"I will read a very short extract from my report to the Board of Trade [on an inquiry at Poole]... It is dated the 19 of June 1875: 'Some years ago the fishery was very productive. I had witnesses before me who agreed in stating that they had taken 2000 or 3000, or even 5000 oysters a day. The witnesses before me were also agreed that 500 or 600 oysters a day constituted now a good catch; that this number could only be taken at the very commencement of the season, and that the take rapidly fell off.... There is no reason to suppose that the small stock of oysters is attributable to any failure of spat. On the contrary, it is clear, from the age of the few oysters found, that a certain amount of spat must have fallen in each of the two last years. The fishermen themselves admit that the oysters are over-dredged; and I have no doubt whatever that the gradual failure of the oyster fishery in Poole Harbour is due to over-dredging."</p>	Poole Harbour, England	Anon (1876)
<1872	<p>"It took 20 boats seven years to dredge away these oysters. There is a fathom more water on the bed now than when they began to dredge. The oysters were thick on that bed and they used to spat. There is no dredging on it now as there are no more oysters to dredge. One boat has got 30,000 oysters a week..."</p>	Ramsey, Isle of Man	Buckland & Walpole (1879)
1879	<p>"There are oysters, in patches, in the bay. They are in lumps, 8 or 10 together. They would be 4 to 6 inches long. Nobody fishes for them here. There has been trawling for many years here by strange boats. The local trawling has increased of late years. The inshore ground has only recently been worked..."</p>	Aberystwyth Bay, Wales	Buckland & Walpole (1879)
<1906	<p>"It is certain that in the past, in each river of the Bay of Quiberon, the oyster bed was continuous and that in the past it was linked to the large natural bed of the open sea [...]. The oysters, in the most favourable conditions, rest on a hard soil, formed of old shells which, when packed and mixed with mud, form a solid ground. The oysters are sometimes isolated, sometimes attached to each other to form more or less large clumps."</p>	Bay of Quiberon	Joubin (1907)

Table S1b. Descriptions by the natural historian Karl Möbius are frequently referred to as a historical baseline, however, descriptions by Möbius (1883) and even by Krøyer (1837), who write some decades earlier, provide a description of impacted and long exploited oyster reefs off the coast of Denmark and Germany.

Year of observation	Quotation	Location	Reference
1877	<p>“In no place upon the seaflats do oysters grow upon rocky bottom. They grow best where there is a substratum of old oyster and other shells. The most of them lie singly, and they are seldom found growing together in clumps of masses. The wide-spread notion that they are found growing firmly attached to the sea-bottom, and piled upon one another, layer upon layer, is accordingly false. Upon the best of the Schleswig-Holstein beds the dredge must drag over a surface of from 1 to 3 square metres, and often over a greater distance, in order to secure a single full grown oyster.”</p>	Schleswig-Holstein, Germany	Möbius (1883)
1877	<p>“The living oysters do not lie in thick masses, stuck to and on one another on the banks, rather mostly more than one meter from one another apart, so that on a meter square of bank area mostly less than one adult oyster will be caught. And yet smaller is the count of half grown oysters. On the largest and most fruitful Bank Huntje, were in ten surveys undertaken between 1730-1852 on average for every 1000 grown oyster only 484 half grown caught, and on most other banks relatively fewer half grown.”</p>	Wadden Sea	Möbius (1883)

1837	<p>“If the sea-bed consists of solid rock, or of loose stones, some oysters are attached to the projections of the rock, or to the individual stones, but many also lie loose on the bottom. Where this consists of clay, sand, or silt, all the oysters must naturally lie loose, except where some are grouped in irregular clusters of three, four, or five individuals. More than five to six I have never seen united; and it is also evident that, if the oysters lay in many layers one above another, and, as a natural consequence, grew together in great masses, the underlings would be hindered not only in their development, but also in opening their shells, and consequently die in a shorter or longer time. The easternmost banks are so far from forming outcrops on the sea-bed that they usually lie in or on the edge of the deeper trenches in the sea-bed.”</p>	Wadden Sea	Krøyer (1837)
------	---	------------	---------------




Figure S1. Oysters caught during semi-quantitative dredge surveys of known oyster banks in the Wadden Sea from 1709-1830, illustrating that fisheries had overexploited oyster reefs in the region prior to published descriptions of the ecology of oyster reefs (published by Krøyer 1837 and Möbius 1877). Krøyer

(1837) provided an in depth review of the status of the Wadden sea oyster banks in the 1830s, drawing on survey data and reports from 1700 onwards. While "oyster banks" continued to be discovered during this time period, nine of the 44 banks known in 1730 were already considered "fished to nothing".

Supporting materials S2 Summary of the evidence underpinning the definition of the European native oyster reef ecosystem.

Table S2a: Examples of current definitions of habitats with mention of *O. edulis*. It is worth noting that *O. edulis* is listed as a feature of “Atlantic littoral *Mytilus edulis* beds on mixed substrata”, “Assemblages of Mediterranean euryhaline and/or eurythermal lagoon biocenosis on sand”, and “Assemblages of the euryhaline and/or eurythermal lagoon biocenosis on mud” (EUNIS, 2022). *O. edulis* not among the species listed for “Bivalve reefs in the Atlantic circalittoral zone” or “Bivalve reefs in the Atlantic littoral zone” (EUNIS, 2022). *O. edulis* not explicitly mentioned nor excluded from: “Infralittoral biogenic habitat”, “Black Sea infralittoral biogenic habitat” (EUNIS, 2022).

Related policy or framework	Attribute defined	Description	Reference
OSPAR	Oyster beds	“ <i>Ostrea edulis</i> occurring at densities of 5 or more per m ² on shallow mostly sheltered sediments (typically 0–10m depth, but occasionally down to 30m). There may be considerable quantities of dead oyster shell making up a substantial portion of the substratum.”	OSPAR Commission (2009)
Ramsar	Bivalve reef	“Bivalve reef consists of large areas of biogenic habitat, dominated by living bivalves where the complex structure of hard shells supports a distinct community that is persistent through time. Expanding on this general definition: ‘large areas’ typically consist of multiple patches, at least some of which are larger than 5 m ² ; ‘dominated’ means at least 25% cover of live shell matter across that space – non-living shell (cultch) may further add to habitat structure and to continuity over time, but without new growth they are unlikely to persist; a ‘distinct community’ is one that supports species and interactions that are rare or absent in surrounding communities; and ‘persistent through time’ describes communities that are likely to remain over decadal timescales or longer.”	Koasar et al. (2015)
EUNIS	Bivalve reefs in the Atlantic infralittoral zone	Infralittoral reefs in the Atlantic formed by bivalves such as <i>Limaria hians</i> and <i>Ostrea edulis</i>	EUNIS (2022)
EUNIS	<i>Ostrea edulis</i> beds on Atlantic infralittoral muddy mixed sediment	Dense beds of the oyster <i>Ostrea edulis</i> can occur on muddy fine sand or sandy mud mixed sediments. There may be considerable quantities of dead oyster shell making up a substantial portion of the substratum. The clumps of dead shells and oysters can support large numbers of <i>Ascidella aspersa</i> and <i>Ascidella scabra</i> . Sponges such as <i>Halichondria bowerbanki</i> may also be present. Several conspicuously large polychaetes, such as <i>Chaetopterus variopedatus</i> and terebellids, as well as additional suspension-feeding polychaetes such as <i>Myxicola infundibulum</i> and <i>Sabella pavonina</i> may be important in distinguishing this biotope, whilst the <i>Opisthobranch Philine</i>	EUNIS (2022)

	aperta may also be frequent in some areas. A turf of seaweeds such as <i>Plocamium cartilagineum</i> , <i>Nitophyllum punctatum</i> and <i>Spyridia filamentosa</i> may also be present. This biotope description may need expansion to account for oyster beds in England.		
EUNIS	Polychaete worm reefs in the Black sea infralittoral zone	Biogenic circalittoral reefs formed by a variety of polychaete worms. In more sheltered and freshwater-influenced environments the non-native serpulid tubeworm <i>Ficopomatus enigmaticus</i> is the most common reef building species. In moderately exposed environments reefs formed by the serpulid <i>Vermiliopsis infundibulum</i> are present. Finally, on lower infralittoral rock serpulids form massive reefs in collaboration with bivalves (i.e. <i>Ostrea edulis</i> , <i>Mytilus galloprovincialis</i>). These reefs are an important component of the Black Sea ecosystem and are characterised by high biodiversity.	EUNIS (2022)

Table S2b. Based on the current status of *O. edulis* habitats in Europe, it has previously been called into doubt whether *O. edulis* is capable of forming biogenic habitat. For example, *O. edulis* reefs were explicitly not considered as biogenic reefs during the UK marine SACs Project (Holt et al. 1998), because in their current highly impacted state, and the remaining habitats left, it was “doubtful if natural beds would qualify as reefs”. Here we provide a summary of the historical evidence that *O. edulis* reefs fulfilled the attributes of “biogenic reef habitat” *sensu* Brown et al. (1997), prior to their widespread degradation by human activities.

Biogenic reef habitat attributes	Evidence from <i>O. edulis</i> reef	Reference
Biogenic reef must “create a substratum which is reasonably discrete and substantially different to the underlying or surrounding substratum ” (Brown et al. 1997)	<p>“The oyster banks of Wicklow have become hard like a rock, as is generally believed for want of dredging. The more the banks are dredged, the more oysters breed. It would do the banks great good to be broken up by a heavy dredge worked from a large smack.”</p> <p>“These great oyster banks are situated on patches in the North Sea, especially off the Dutch coast. The trawlers carefully avoid these beds as the heavy clumps tear the nets.”</p> <p>“The trawlers avoid this “rough ground” as they call it, as much as possible; but when they do by accident get on to it, the oysters are so numerous that they fill up the trawl net and nearly bring up the vessel”</p> <p>... “The dredgermen who go out fishing in the North Sea (as I found out in my inquiries at Yarmouth) come across every now and then an</p>	<p>Irish Fisheries (1836)</p> <p>Buckland and Walpole (1879)</p> <p>Buckland (1875)</p>

	<p>enormous tract of oyster ground, which tears their nets all away to pieces, and for that reason they get away from it.”</p> <p>“It is certain that in the past, in each river of the Bay of Quiberon, the oyster bed was continuous and that in the past it was linked to the large natural bed of the open sea [...]. The oysters, in the most favourable conditions, rest on a hard soil, formed of old shells which, when packed and mixed with mud, form a solid ground. The oysters are sometimes isolated, sometimes attached to each other to form more or less large clumps.”</p> <p>“It was reported that large oyster beds had been discovered off New Quay, but strong dredges would be required to open them.”</p> <p>[In Bay of Saint Brieuc, North Brittany] “The Parliament of Brittany issued a decree on 16 October 1784, because the Saint Brieuc bed was almost completely exhausted: “In many places where it was formerly composed of several layers, only mud is currently being removed””.</p> <p>The seafloor is filled with oysters, almost placed one on top of the other like stones, forming a wall.”</p> <p>“In no place upon the seafats do oysters grow upon rocky bottom. They grow best where there is a substratum of old oyster and other shells.</p> <p>“The oysters, when he was a boy, were as deep as the Town Hall is high. The bank was two miles long by half a mile wide”</p> <p>The oyster reefs found by us are constructed mainly of <i>Ostrea edulis</i> shells, with calcareous tubes of serpulid polychaetes also present as cementing material. They represent erect biogenic structures with a distinguishing irregular, branching or netted shape with serrated margins attaining 7 m height, 30–50 m length and 10 m width</p> <p>“Trawlers have lately found oyster beds in the North Sea. The huge ‘Skelling Bank’ off Heligoland consists of numerous patches of oysters; other great oyster beds have been found off the Dutch coast by trawlers. A trawl has a beam which spreads about 40 feet, and especially in very deep water, where dredges will not work nicely, it picks up a great many oysters...”</p>	<p>Joubin (1907)</p> <p>The Aberystwith Observer (1897)</p> <p>Levasseur (2006)</p> <p>Marsili (1715)</p> <p>Möbius (1883)</p> <p>Buckland and Walpole (1879)</p> <p>Todorova et al. (2009)</p> <p>Philpots (1891)</p>
<p>Biogenic reef “unit should be ... somewhat raised” (Brown et al. 1997)</p>		
<p>Biogenic reef “unit should be substantial in size (generally of the order of a metre or two across as a minimum)” (Brown et al. 1997)</p>		

	<p>In the year 1870 a small oyster bed was discovered at the mouth of the Thames, north east from Whitstable. It was about 18 metres long by 6 metres broad. Forty-eight hours later 75 boats were there, close alongside of one another, fishing up the oysters.... Upon every old oyster which was taken were found only from nine to ten young ones of different ages. This bed had never been previously disturbed, and the oysters were accordingly found in their natural condition.</p> <p>“Over the Schleswig-Holstein seaflds there exist 50 oyster beds of very different sizes. The largest is not far from 2 km long, but the greater number are shorter than this. Their breadth is much less than their length, which is in the same direction as the channels along the slopes of which they lie....”</p> <p>Each boat has out, so far as we could see, four dredges, so that in all there were about 120 of these destructive engines at work on the bank. The dredges are lifted and emptied about every fifteen or twenty minutes. Say that this operation is performed three times an hour, it follows that there are 300 hauls per hour, or 3,000 in a working day of ten hours, made from the bank. Of course we have no means of knowing the exact extent of the oyster bed, but it is very evident that, no matter how extensive it may be, if these boats are permitted to "rag at it" (as the local phrase is) without intermission, in season and out of season, the total destruction of the bank is only a question of a very limited time. The operations of the boats extended over a space of about a mile and a-half to two miles in length, and about half a mile or so in width; so that we may presume that the bank is about that in dimensions.</p>	<p>Möbius (1883)</p> <p>Isle of Man Times (1874)</p>
--	--	--

Table S2c. The IUCN Ecosystem Red Listing process requires that the developed ecosystem definition is cross-referenced to relevant ecological classifications such as the IUCN Global Ecosystem Typology (Keith et al. 2020). Here we provide a summary of evidence that *O. edulis* reefs historically fulfilled the ecological trait criteria of forming “Shellfish beds and reefs”, as defined by the IUCN Global Ecosystem Typology (Keith et al. 2020).

Ecological trait	Evidence from <i>O. edulis</i> reefs	Reference
High productivity and moderate diversity, heterotrophic energy.	The oysters, which, as we have seen above, is itself so worthless, and consequently peaceful, animal, is given at the price of a great many	Krøyer (1837)

	<p>enemies. It is this circumstance which makes the oyster-banks so interesting to a zoologist; one can probably always be sure ... of drawing from the sea between the oysters a host of other animals, whose amusement is undoubtedly largely derived from the oyster. Such animals are various species of crabs, starfish, snails, worms, &c. Although the oyster by its shell should appear to be amply protected against these enemies, it will be evident from what follows that even the adult oyster does not always find a reassuring shelter in the armour with which nature has equipped it; still less can this be the case with the very infant. So great is the fertility of the oyster, that I alone find on our banks a long greater number of young oysters. When the full-grown oyster is drawn up from the depths, its shell is generally found lined with calcareous or membranous tubes, which serve as a dwelling-place for various worms (serpularae or worm-tubes, &c.); with balans, anomia, chitonians, ascidians, and similar immobile or unwieldy animals. The more of these there are, the more loopholes are formed (and especially if two or three oysters are together) for numerous small crabs (<i>Porcellana longicornis</i> and <i>Galatea strigosa</i>) of size from that of a grain of manure to that of a large pea, for polynids, ophids, &c. ; oyster young, however, are comparatively rarely seen; I suppose because they are for the most part devoured by these many small but very large predators."</p>	
<p>Structural complexity from shell aggregations.</p>	<p>In the year 1870 a small oyster bed was discovered at the mouth of the Thames, north east from Whitstable. It was about 18 metres long by 6 metres broad. Forty-eight hours later 75 boats were there, close alongside of one another, fishing up the oysters.... Upon every old oyster which was taken were found only from nine to ten young ones of different ages. This bed had never been previously disturbed, and the oysters were accordingly found in their natural condition.</p> <p>"There are oysters, in patches, in the bay. They are in lumps, 8 or 10 together.</p> <p>"The seed is like a viscus or glue, which immediately attaches itself to trees and stones and, in the absence of these, often to the oysters themselves, so that three or four can still be attached to an oyster."</p>	<p>Buckland and Walpole (1879)</p> <p>Oedmann (1743)</p>

	<p>“In June, when oysters are most scarce, they reproduce on the bottom of the sea, by leaving out their eggs or spawn, which looks like a drop of glue, and attaches itself to everything in the sea, often to oysters themselves.”</p>	Wilmsen (1831)
<p>Dominated by sessile filter-feeders, secondary deposit-feeders.</p>	<p>“Generally the [oyster] net is allowed to drag from five to ten minutes... and the entire contents of the bag emptied upon the deck. This mass consists of old oyster shells, mussels of various kinds, living oysters, snails, crabs, worms, starfish, sea-urchins, polyps, sponges, and seaweeds, which are generally mixed up with sand and mud... Despite these manifold cleansings [by fishermen for market] many oysters when they are exposed for sale are covered with dead and living animals, and the peculiar odour which oysters have when carried into the interior arises from the death and decay of the organic materials upon the outside of the shells...”</p>	Möbius (1883)
	<p>Even the shells of the living oysters are inhabited. Barnacles (<i>Balanus crenatus</i>) [...] often cover the entire surface of one of the valves. Frequently the shells are bedecked with yellowish tassels a span or more in length, each of which is a community of thousands of small gelatinous bryozoa (<i>Alcyonidium gelatinosum</i>), or they are overgrown by a yellowish sponge (<i>Halichondria panicea</i>) [...] Upon many beds the oysters are covered with thick clumps of sand which are composed of the tubes of small worms (<i>Sabellaria anglica</i>). These tubes, called 'sand-rolls' resemble organ-pipes, and are formed from grains of sand cemented into shape by means of slime from the skin of the worm [...]. Upon certain beds near the south point of the island of Sylt [...] there lives upon the oyster shells a species of tube-worm (<i>Pomatoceros triquetra</i>) [...] The shells of many oysters upon these beds also carry what are called 'sea hands' (<i>Alcyonium digitatum</i>) which are white or yellow communities of polyps of the size and shape of a clumsy glove. Often the oyster shells are also covered over with a brownish, clod-like mass, which consists of branched polyps (<i>Eudendrium rameum</i> and <i>Sertularia pumila</i>) or they may be covered with tassels of yellow stems which are nearly a finger long and have at their distal ends reddish polyp-heads</p>	Möbius (1883)

	<p>(<i>Tubularia indivisa</i>). Among these polyps, and extending out beyond them, are longer stems, which bear light yellow or brown polyp-cups (<i>Sertularia argentea</i>) [...]. I once took off and counted, one by one, all the animals living upon two oysters. Upon one I found 104 and upon the other 221 animals of three different species [...]. Soles [...] stone-picks, and stingrays [...] are abundant upon the oyster banks.</p> <p>in addition to the “fern” (<i>Sertularia abietina</i>), which is not disfiguring, there are a good many soft worm tubes (<i>Terehella</i>, or the like), sea-squirts (<i>Ciona intestinalis</i>), and small shell-fish (<i>Crenella diseors</i>) to be cleaned off.</p> <p>“These shells are very often covered with productions of the coral kind: they are frequently loaded also with small muscles and multitudes of worms, but only on the convex side, which appears to be the upper one, so that the animal rests on the flat side. It frequently happens that both shells are quite pierced through, and gnawed by worms in the same manner as old wood.”</p> <p>Very often one finds corallianic plants on the oyster shells; often they are also covered with mussels and other plants.</p>	<p>Holt (1902)</p> <p>Beckmann (1800)</p> <p>Lippold and Funke (1810)</p>
--	--	---

Table S2d. Summary of relevant historical catch rates and habitat descriptions, depicting the reef-building nature and high densities of *O. edulis* encountered in *O. edulis* reef ecosystems historically. For further analysis, see Thurstan et al. in review a,b.

Evidence from <i>O. edulis</i> reef	Reference
<p>Fishermen from Holland and Germany dredge for oysters here, especially during the months of August, September, and October, and often catch, at a single drag of the dredge, as many as 1,000 oysters. Sometimes great bunches of oysters growing attached to one another are gathered into the net.</p>	<p>Möbius (1883)</p>
<p>“In Ballycroy Bay, and the Sound of Bulls-mouth, three thousand oysters may be taken in a day, with a dredge. They are often sold for 3d per hundred.... There are several natural oyster beds in Broadhaven and Blacksod bays, and in Achil Sound; they are open to the public and dredged.”</p>	<p>Irish Fisheries (1836)</p>
<p>“I remember when the boats could go out and dredge at Milford Haven, round by the Stack Rock, and each boat would get from 1000 to 1200 oysters in three or four hours. The last catch I took, I took 16 boats catch for a week; they had moderately fine weather during the week, and the largest catch was 600 oysters each, there</p>	<p>Anon (1876)</p>

<p>were three men to a boat, and all of them did not bring in 5,000 in the gross, and that was starvation to the men."</p>	
<p>"I will read a very short extract from my report to the Board of Trade [on an inquiry at Poole]... It is dated the 19 of June 1875: 'Some years ago the fishery was very productive. I had witnesses before me who agreed in stating that they had taken 2000 or 3000, or even 5000 oysters a day. The witnesses before me were also agreed that 500 or 600 oysters a day constituted now a good catch; that this number could only be taken at the very commencement of the season, and that the take rapidly fell off.... There is no reason to suppose that the small stock of oysters is attributable to any failure of spat. On the contrary, it is clear, from the age of the few oysters found, that a certain amount of spat must have fallen in each of the two last years. The fishermen themselves admit that the oysters are over-dredged; and I have no doubt whatever that the gradual failure of the oyster fishery in Poole Harbour is due to over-dredging."</p>	<p>Anon (1876)</p>
<p>"Clew Bay abounds with oysters, where they are taken in large quantities, (considering the wretched small dredges with which they are fished for,) out of an open boat, rowed by two men, and a third holding the dredge rope. They seldom catch more than a thousand a day, as they find it difficult to dispose of them, even at a moderate price... the expense of sending them up [to Dublin] by carriers runs away with any profit that the fishermen would derive from a good market, and they now seldom fish for them unless they are bespoken."</p>	<p>Brabazon (1848)</p>
<p>"It took 20 boats seven years to dredge away these oysters. There is a fathom more water on the bed now than when they began to dredge. The oysters were thick on that bed and they used to spat. There is no dredging on it now as there are no more oysters to dredge. One boat has got 30,000 oysters a week..."</p>	<p>Buckland and Walpole (1876)</p>
<p>"Helligsø - 'one of the oldest banks in the whole Limfjord. It was initially extremely rich, so that even in one day, about 14,000 oysters were fished of a single boat despite the fishermen's inferior exercise scraping... The bank decreased year by year in fullness, but was still in the spring of 1868 so rich that a single boat in one day scraped about 5000 stkr. Now it gives so little that the scraping can't pay off... has been abandoned."</p>	<p>Collin (1871)</p>
<p>From the locality the bed extends not less than 60 miles in the North-West direction, where they lie very thick; 1200 have been caught here in the space of four hours by the trawl net. Towing by steam power, the whole space of ground appears almost inexhaustible, at all events it will take a great number of years to exhaust it. The water being deep, improved dredges are required, and steam winches to heave them up with. If enterprising gentlemen were to form a company, and have a few steamers of about 70 tons built, with wells in them, and fitted with steam winches, it is highly probably it would pay well... The vessels, I should estimate, would bring from 35 to 50 thousand [oysters] per week, and that ought to pay well. Already small sailing vessels have been getting 20 thousand per week, without the aid of steam power. ... Grimsby has now twenty oyster vessels."</p>	<p>Olsen (1885)</p>

“The period of the Cancale Fishery is known as “la Caravane” [...]. The 1909 “Caravane” involved 6 trips of 360 boats each, manned by 2500 men. From 10 April to 24 April, fishing took place for 38 hours and 45 minutes. The number of oysters caught was 16 million.”	Joubin (1910)
“In Auray River (near the bay of Quiberon), in 1885, 150 boats, manned by 1648 men, have dredged 4.2 million oysters within 4 days...”	Joubin (1907)
There were then very few boats in the Bay, from six to seven thousand oysters were often got in one day with only one dredge, but when larger boats from Jersey with superior tackle came this became a small haul.	The Cambrian News and Merionethshire Standard (1889)
“Until the last fishing season which lasted 6 months & 10 days, this industry (Bay of Brest fishery) came to the aid of 576 fishermen on 144 boats & that 14 million oysters were sold.”	Archives du Service Historique de la Défense de Vincennes (1849)
“It is by far the best stocked bed on the coast, and may, I think, be considered in a satisfactory condition, since the catch comprised oysters of all ages	Holt (1902)
“An old oyster often has twenty small ones attached to it”	Beckmann (1800)

Table S2e. Reported densities of reef building oysters from geographies outside or Europe. *Ostrea* spp. densities reflect densities in current exploited and protected areas. Reported *Crassostrea virginica* densities are provided for context. *C. virginica* are known reef builders, yet as a result of over-exploitation, mean densities recorded in the 1880s in Virginia, USA were similar to those reported in degraded *O. edulis* habitats today.

Species	Location	Mean density range (Ind m ⁻²)	Reference
<i>Ostrea angasi</i> (exploited)	Australia	17-67	Jones and Gardener 2016
<i>Ostrea lurida</i> (protected)	British Columbia	21-507 (across the 9 of 14 sites sampled, with a mean density of >20 Ind m ⁻²)	Norgard et al. 2018
<i>Crassostrea virginica</i> (exploited)	Texas	24-67	Moore and Danglede 1917
<i>Crassostrea virginica</i> (over exploited)	Virginia	0.07-0.57	Winslow 1881

Table S3. Summary of evidence of ongoing threats to *Ostrea edulis* populations in Europe. Threats listed relate to those illustrated in Figure 3 within the main manuscript.

Threat	Summary of evidence for current trend/future threat
Over harvesting	<p>Over harvesting is the leading cause for the decline in <i>O. edulis</i> historically. There are very few wild fisheries still active in Europe, as most stocks were fished to economic extinction by the 1930s (Thurstan et al. 2013). Many of the remaining fisheries are suffering from poor recruitment and declining catches (Southern IFCA 2019, Kent and Essex IFCA 2022, Marine Institute & Bord lascaigh Mhara 2023). Although the reasons for these recent declines are diverse and include, for example, the introduction of invasive species and disease, fishing remains a pressure on these declining populations, as reflected by fisheries restrictions to aid in the recovery of the population (Southern IFCA 2019, Kent and Essex IFCA 2022, Marine Institute & Bord lascaigh Mhara 2023).</p> <p>Illegal and unregulated harvesting is also an ongoing threat to native oyster populations and their recovery. <i>O. edulis</i> stocks in Strangford Lough, Northern Ireland recovered to >1 million individuals in 2007 but declined to 650,000 by 2005 (Smyth et al. 2009). This decline was linked to illegal and unregulated harvesting (Smyth et al. 2009), which is supported by the continued observation that <i>O. edulis</i> populations remain suppressed in areas which are not policed. In contrast, well protected areas are maintaining their population (Smyth et al. 2023).</p>
Poor water quality (Eutrophication)	<p>The <i>O. edulis</i> population in the Mar Menor, Spain, was estimated at ~135 million in 1989 (Garcia Garcia et al 1989), but has declined to levels at which there is no evidence of recruitment (Ruiz et al. 2020).</p>
Pollution (TBT)	<p>TBT impacted <i>O. edulis</i> populations (Thain and Waldock 1986), but this threat has been declining since the banning of TBT in the 1980s (Crouch estuary, UK; Rees et al. 2001).</p>
Climate change	<p>Temperatures above 28°C are detrimental to <i>O. edulis</i>, and temperatures above 36°C are fatal (Eymann et al. 2020), this may have implications for part of the species range, in particular in the eastern Mediterranean, under a warming climate (Sakalli 2017) as well as lagoonal areas. For example, water temperatures in the Mar Menor (Spain) reached 31°C in July 2023 (McGeer 2023). At lower temperatures, populations may be impacted by a temperature-mediated skewed sex ratio, with a study on two UK populations finding that higher temperatures resulted in a male-skewed sex ratio, which may impact population viability (Eagling et al. 2018). Furthermore, there may be interactions between climate change and other threats, such as invasive species. For example, rising temperatures will differentially affect <i>C. gigas</i> and <i>O. edulis</i>, and it is as yet unknown what its impact will be on the balance in population growth between the two species (Stechele et al. 2022).</p>
Invasive species (<i>Crassostrea gigas</i>)	<p>There are many examples of <i>C. gigas</i> and <i>O. edulis</i> coexisting, either within the same water body, but at differing depths (Staglicic et al. 2020), or within mixed reefs (Christianen et al. 2018, Thorngren et al. 2019). Given the currently limited presence of <i>O. edulis</i> in European coasts and seas, it is challenging to draw conclusions on whether interactions between these two species are positive for <i>O. edulis</i>, or negative. In areas of substrate limitation, <i>C.</i></p>

	<p><i>gigas</i> has been observed to provide an important source of substrate (Christianen et al. 2018), however, <i>C. gigas</i> has also been observed to consume the larvae of <i>O. edulis</i>, as well as having a high overlap in feeding traits, which could indicate that competitive interactions will become more important when <i>O. edulis</i> numbers increase (Ezgeta-Balic et al. 2020). In marine management policies, the precautionary principle is therefore applied and further introduction of <i>C. gigas</i> is not recommended (Ezgeta-Balic et al. 2020). <i>C. gigas</i> is expanding its range across Europe and will therefore impact increasing numbers of sites where <i>O. edulis</i> was historically present (Anglès d'Auriac et al. 2017).</p>
<p>Invasive species (<i>Rapana venosa</i>)</p>	<p>Introduced to the Black Sea in the 1940s, the predatory gastropod <i>Rapana venosa</i>, had consumed almost all oysters on the large Gudaota oyster bank by the 1950s (Zolotarev & Terentyev 2012). This species continues to present a significant issue to the recovery of shellfish populations, with impacts on shellfish species throughout the Black Sea (Janssen et al. 2014) and in the northern Adriatic, where it was first recorded in 1973 (Savini & Occhipinti-Ambrogi 2006).</p>
<p>Invasive species (<i>Crepidula fornicata</i>)</p>	<p><i>C. fornicata</i> can dominate substrates it invades and change the nature of the substrate to one of anoxic mud and making it less suitable for <i>O. edulis</i> settlement. This is especially problematic as their niches strongly overlap (Blanchard et al 2008). There are also direct interactions with <i>O. edulis</i>. Where substrate is limiting <i>C. fornicata</i> may outcompete <i>O. edulis</i> for settlement substrate (Preston et al. 2020). Furthermore, <i>C. fornicata</i> larvae may outcompete <i>O. edulis</i> larvae for food (Blanchard et al. 2008, Preston 2020). Finally, also of relevance to the ecosystem red listing, where <i>C. fornicata</i> is abundant, it can result in reduced biodiversity associated with <i>O. edulis</i> habitats, relative to when <i>C. fornicata</i> is absent (Lown et al. 2021).</p>
<p>Disease (<i>Marteilia refringens</i>)</p>	<p><i>Marteilia refringens</i> was first recorded in Europe in 1968, where it caused high mortality (Grizel et al. 1974). <i>Martelia</i> spp. was implicated in the extinction of <i>O. edulis</i> from Gulf of Thessaloniki, Greece (Virvilis and Anglidis 2006)</p>
<p>Disease (<i>Bonamia ostreae</i>)</p>	<p><i>Bonamia ostreae</i> is an invasive haplosporidan parasite that has been a leading cause of mortality in populations of <i>O. edulis</i>, since its introduction into Europe in the late 1970s (Grizel et al. 1985), causing over 90% mortality in 4+ ages oysters in naïve populations (Culloty & Mulcahy 2006). The disease caused by <i>B. ostreae</i> has subsequently been implicated in the decline and economic extinction of numerous remnant <i>O. edulis</i> populations, and the disease is considered a potentially limiting factor to restoration efforts in some locations (Laing et al. 2005). Recently, genetic markers which may indicate resistance or tolerance to <i>B. ostreae</i> have been identified (Sambade et al. 2022), and many restoration efforts are now seeking to work with resistant or tolerant individuals in their active restoration efforts (Kamermaans et al. 2023). The effectiveness of the markers in response to a <i>B. ostreae</i> challenge have yet to be tested, but there is evidence that populations with a longer exposure history to <i>B. ostreae</i> or slow-growth life histories have better survival in the presence of the parasite (Culloty et al. 2008; Egerton et al. 2020).</p>

Salinity changes	<p><i>O. edulis</i> is found in European waters with salinities ranging from c.20 ppt to 42ppt (Davis & Ansell 1962, Ruiz et al. 2020). Historically there are examples of increases in salinity resulting in the establishment and growth of <i>O. edulis</i> populations in the Limfjord, Denmark (Collin 1884). Similarly, within the Roskilde fjord and along the central eastern coast of Jutland (around Horsensfjord), oysters were previously recorded in areas where salinities are now too low for viable populations (Krøyer 1837, Rasmussen et al. 2007).</p>
High sedimentation	<p>Avoiding high sedimentation rates is a critical factor in site selection for <i>O. edulis</i> restoration efforts (Hughes et al. 2022, Kamermaans et al. 2022). Fishers in some locations have traditionally sought to combat the effects of sedimentation through the practice of “harrowing”, during which gear is towed over oyster habitats to remove sediment prior to larval settlement (Bromley et al. 2016).</p> <p>While increased sedimentation was not a reported driver of decline in the documentary historical literature (Thurstan et al. in review b), it has been implicated as a potential driver of decline in late Holocene reefs (Sander et al. 2021). It is widely accepted that deforestation during numerous periods from the 1000 onwards resulted in increased siltation of European coastal ecosystems (Poirier et al. 2011). With improved land use management, river sediment load in Europe may decrease in the future (Bakker et al. 2008).</p>
Destructive fishing methods/capital dredging	<p>Bottom towed gears have been recognized as a threat to <i>O. edulis</i> populations since the early 1800s (e.g. Krøyer 1837 and references within Table S3). Towed gears disturb the habitat, flatten the structure, and re-suspend sediments (Watling and Norse 1998, Oberle et al. 2016). As a result, bottom towed gears are known to reduce the diversity and complexity of seabed habitats (Hiddink et al. 2019, Pitcher et al. 2022). Much of the European shallow seas is trawled 1-10 times a year, and many inshore waterways are dredged (Eigaard et al. 2017). There are, however, areas where bottom towed gears are banned (e.g. Milford Haven, Wales; Sweden, Norway in waters <60 m depth), and where recovery of <i>O. edulis</i> habitat is not precluded.</p>

Table S4

Summary of literature relating to the current presence of *Ostrea edulis* reef ecosystems in coastal nations within the native range of the species.

Country	Any m ² areas with >20 oysters m ⁻² ?	Any > 1 ha areas with >20 oysters m ⁻²	Details
Albania	no	no	<i>O. edulis</i> and <i>O. stentina</i> present in species checklist. (Dhora 2009) No further mentions identified.
Algeria	no	no	<i>O. edulis</i> present, but not "strong presence" in north-western Algeria (Husseini and Talet 2019).
Belgium	no	no	No <i>O. edulis</i> collected from the scuba operated video footage on the Westhinder sandbank (Houziaux et al. 2007).
Bosnia and Herzegovina	no	no	Neum Bay, individuals of <i>O. edulis</i> also encountered when collecting mussels (Trozic-Borovac et al. 2022).
Croatia	no	no	<i>O. edulis</i> common, but not dominating locally (Zavodnik and Kovacic 2000). Max densities from survey <2 oysters m ⁻² . Coexistence with <i>C. gigas</i> (Staglicic et al. 2020). Listed as present in Novigrad and Karin Seas, but no further mention (Kruschel et al. 2011).
Cyprus	no	no	<i>O. edulis</i> present in species list (Fischer 1997).
Denmark	no	no	No recent records of <i>O. edulis</i> from the Wadden Sea. Max <i>O. edulis</i> biomass in the Limfjord is given as >0.25kg m ⁻² over small area, with most areas 0.01-0.05kg m ⁻² (Fomsgaard and Petersen 2015).
Egypt	no	no	<i>O. edulis</i> present in Nile lagoons, but not a dominant species (Bernasconi and Stanley 1994). Few <i>O. edulis</i> individuals (28-30 total) found at two sites within Burullus Lagoon (Orabi et al. 2018).
England	no	no	Some areas within the oyster beds have <i>O. edulis</i> at >5 m ⁻² , but average in Blackwater c. 3m ⁻² (Allison 2018); Fal catch rates c. 1-5kg hr ⁻¹ (Sturgeon et al. 2022). In Solent; <i>Ostrea edulis</i> stock assessment (Southern Inshore Fisheries and Conservation Authority 2019) only 33% of tows were positive (contained an oyster). Of these positive tows, the average CPUE (kg/m/hr/>70mm) was 3.29 with a range of 1.27 - 5.18. The <i>Ostrea edulis</i> population had declined by 96% in the last two decades (1999-2019; Helmer et al 2019). The threshold CPUE to reinstate a fishery is 15 oysters m ⁻² hr ⁻¹ .

France	yes	no	Patches with <i>O. edulis</i> at >20 m ² , but not over extended area (< 1000 m ²). Many places in Brittany were density is less than 1 oyster m ⁻² . See database on Pouvreau et al. (2021) https://www.seanoe.org/data/00686/79821/
Georgia	no	no	Large bank of <i>O. edulis</i> extirpated by <i>Rapana venosa</i> in the 1950s (Zolotarev and Terentyev 2012).
Germany	no	no	<i>O. edulis</i> extirpated in Germany (Gerken and Schmidt 2014)
Greece	no	no	<i>O. edulis</i> beds in the Gulf of Thessaloniki extinct due to <i>Martelia</i> (Virvilis and Angelidid 2006).
Ireland and Northern Ireland	yes	no	Lough Foyle- <i>O. edulis</i> densities <4 m ⁻² even in the best areas (Loughs Agency 2020). Area of beds in Inner Tralee Bay 4km ² , <i>O. edulis</i> density range 0-50 m ⁻² , but most areas <1 m ⁻² . In other sites in Ireland (Kilkieran Bay) densities did not exceed 5 m ⁻² (Tully and Clarke 2012, Marine Institute & Bord Iascaigh Mhara, 2021). In Strangford Lough, in 2004 some areas settled with >30 m ⁻² (Smyth et al. 2009), but there was subsequent decline to low numbers (Smyth et al. 2020).
Israel	no	no	<i>O. edulis</i> reared from introduced spat (Shpigel 1989); Present in species list (Fischer 1997).
Italy	no	no	<i>O. edulis</i> on 90 % of shells of <i>Pinna nobilis</i> , but <i>Pinna nobilis</i> at c. 0.1 m ⁻² in Sardinian estuaries (Addis et al. 2009). <i>O. edulis</i> on species check list for all areas barr the southern and middle Adriatic (Renda et al. 2022).
Lebanon	no	no	Listed as present after 1950, but not in records prior to 1950 (Crocetta et al. 2013).
Libya	no	no	Single shell from beach (Hera and Harris 2015). Oysters listed as present (Bek-Benghazi et al. 2020). Eleven <i>O. edulis</i> found at two sites in Regata (Abushaala et al. 2014).
Malta	no	no	No wild <i>O. edulis</i> populations known (Agius et L. 1978).
Montenegro	no	no	No survey of wild <i>O. edulis</i> , but <i>O. edulis</i> imported from Mali Ston (Croatia) for growth trials (Joksimovic et al. 2011 and Cataudella et al. 2005); <i>O. edulis</i> mentioned in check list as found on hard substrate (Petovic 2018).
Morocco	no	no	Only few oysters sampled from Morocco. All were <i>O. stentina</i> (Lapegue et al. 2006).
Norway	yes	no	Several <i>O. edulis</i> habitat areas along the Norwegian coast with patches with >50 m ² (Bodvin 2011; AT Laugen et al, unpublished data). Quantitative data on extent are lacking but known patches are generally small (Mortensen et al 2023; Ane Timenes Laugen Pers Comm).
Scotland	yes	no	Scattered individual <i>O. edulis</i> found around Shetland (Shelmerdine and Leslie 2009) and several areas around the Scottish west coast (Sanderson unpubl. data). Three

			areas with high density (>5 m ²) and populations measurable in the thousands, but are limited in extent to 10s m ² (Sanderson unpubl. data). Loch Ryan is the only known remnant population measurable in millions and where densities are often >5 m ² (Ramday et al. 2024). In Loch Ryan densities can reach c. 20 oysters m ⁻² in limited areas (10s m ² , Sanderson unpubl. data).
Slovenia	no	no	Present as epifauna on <i>Cladocora caespitosa</i> reefs (Piacco et al. 2014). Listed as a fouling species in the harbour of Piran (Ferrario et al. 2018).
Spain	no	no	Live <i>O. edulis</i> in 4% of samples from the river mouth (Rio Ulla) and no other sites. Not habitat forming (Cadee 1968). Population of 300,000 <i>O. edulis</i> found in natural bed in San Cibrán, but no habitat description given (Ruiz et al. 1992). No samples in extensive species list from Bay of Algeciras (van Aartsen et al. 1984); <i>O. edulis</i> beds >1 km in length but max <i>O. edulis</i> density <3 m ² and at A Caleira (Ortigueira) <i>O. edulis</i> bed c. 300m in length with some patches with oysters <5 m ² (Iglesias et al. 2013). Mar Manor population too low for natural recovery (Ruiz et al. 2020)
Sweden	yes	no	Numerous oyster beds near Resö, with the largest covering 26,000 m ² . The beds are mixed with <i>C. gigas</i> . Mean native oyster density ranges from 0.4-12.9 m ² , although patches with c. 20 oysters m ² exist (Holthuis 2022).
Syria	no	no	No recent evidence available in the literature or OBIS.
The Netherlands	no	no	Mixed shellfish reef in the Dutch Voordelta, <i>O. edulis</i> 6.8 +0.6 m ² in patches within the reef. <i>C. gigas</i> and <i>Mytilus edulis</i> more abundant. (Christiansen et al. 2018)
Tunisia	no	no	<i>O. edulis</i> rare in south eastern Tunisia (Ktari-Chakron and Azouz 1971). <i>O. edulis</i> listed as very common at one site, but "Oysters (<i>Ostrea edulis</i> var. tarentina) never form exploitable beds." (Lubet and Azouz 1969). 68 oysters from Bizert Lagoon and the Gulf of Hammamet in northern Tunisia- mix of <i>C. gigas</i> and <i>O. stentina</i> (Dridi et al. 2008). Six <i>O. edulis</i> and seven <i>O. stentia</i> specimens found across 28 locations in Skhira Bay (Boudaya et al. 2019)
Turkey	no	no	31 tonnes <i>O. edulis</i> harvested from the wild nationally in 2006. (Yildiz et al. 2011); Oysters present in the Marmara Sea (Sütcuoglu and Korun 2011); <i>O. edulis</i> almost disappeared from Gemlik Bay. The Istanbul Strait, a hard-bottom fauna has been replaced by a soft bottom fauna of polychaetes (öztürk and öztürk 1996). <i>O. edulis</i> not habitat forming in Turkish waters (Pers. comm. Dr. Aydin 2021)

Wales	no	no	Milford Haven intertidal density mean 0.03 \pm 0.01 s.e. oysters m ⁻² , with a max of 0.2 oysters m ⁻² (zu Ermgassen 2017). Subtidal density at various sites ranged from 0.05-0.17 m ⁻² (Lock 2017).
-------	----	----	--

References

- Abushaala, N.M., Shaibi, T., Howaege, H.M., (2014). Molluscan fauna of hard substrate along the coastal zone of western Libya. *International Journal of Bioassays*, 3, 3211-3217.
- Addis, P., Secci, M. & Cau, A. (2009). Density, size structure, shell orientation and epibiontic colonization of the fan mussel *Pinna nobilis* L. 1758 (Mollusca: Bivalvia) in three contrasting habitats in an estuarine area of Sardinia (W Mediterranean). *Scientia Marina*, 73, 143–152.
- Agius, C., Jaccarini, V. & Ritz, D.A. (1978). Growth trials of *Crassostrea gigas* and *Ostrea edulis* in inshore waters of Malta (Central Mediterranean). *Aquaculture*, 15, 195–218.
- Allison, S., Hardy, M., Hayward, K., Cameron, T.C. & Underwood, G.J.C. (2020). Strongholds of *Ostrea edulis* populations in estuaries in Essex, SE England and their association with traditional oyster aquaculture: evidence to support a MPA designation. *Journal of the Marine Biological Association of the United Kingdom*, 100, 27–36.
- Anglès d’Auriac, M.B., Rinde, E., Norling, P., Lapègue, S., Staalstrøm, A., Hjermmann, D.Ø. & Thaulow, J. (2017). Rapid expansion of the invasive oyster *Crassostrea gigas* at its northern distribution limit in Europe: Naturally dispersed or introduced? *PLOS ONE*, 12, e0177481.
- Anon (1876) Select Committee to inquire into Reasons for Scarcity of Oysters and Effect of Measures adopted after Report of Royal Commission on Sea Fisheries, 1866. Report, Proceedings, Minutes of Evidence, Appendix, Index.
- Archives du Service Historique de la Défense de Vincennes (1849) Lettre du Préfet Maritime de Brest au Ministre de la Marine et des Colonies précisant la fermeture du dragage des huîtres plates et leur vente, n°544, 26 avril 1849. CC5/623.
- Bakker, M.M., Govers, G., van Doorn, A., Quetier, F., Chouvardas, D. & Rounsevell, M. (2008). The response of soil erosion and sediment export to land-use change in four areas of Europe: The importance of landscape pattern. *Geomorphology*, Human and climatic impacts on fluvial and hillslope morphology, 98, 213–226.
- Beckmann (1800) Observations respecting oysters, and the places where found. *Philosophical Magazine and Journal* 6, 233-240. London: Alex. Tilloch.
- Bernasconi, M.P. & Stanley, D.J. (1994). Molluscan biofacies and their environmental implications, Nile Delta Lagoons, Egypt. *Journal of Coastal Research*, 10, 440–465.
- Bek-Benghazi, N., Al-Mgoushi, A., Hadoud, D., Shakman, E. (2020) Marine Mollusca of the Libyan waters, the southern Mediterranean Sea. J. Black Sea/Mediterranean Environment 26,263-285.
- Blanchard, M. (2009). Recent expansion of the slipper limpet population (*Crepidula fornicata*) in the Bay of Mont-Saint-Michel (Western Channel, France). *Aquat. Living Resour.*, 22, 11–19.
- Blanchard, M., Pechenik, J.A., Giudicelli, E., Connan, J.-P. & Robert, R. (2008). Competition for food in the larvae of two marine molluscs, *Crepidula fornicata* and *Crassostrea gigas*. *Aquat. Living Resour.*, 21, 197–205.

- Bodvin T, Mortensen, S., Helmet, A., Øyvind Strand, H.I. & Grefsrud, E.G. (2011) Utredning av europeisk flatøsters *Ostrea edulis* L. – Kunnskapsoversikt med forslag til handlingsplan. *DN-utredning 10-2011*. 19p.
- Boudaya, L., Mosbahi, N., Dauvin, J.-C. & Neifar, L. (2019). Structure of the benthic macrofauna of an anthropogenic influenced area: Skhira Bay (Gulf of Gabès, central Mediterranean Sea). *Environmental Science and Pollution Research*, 26, 13522–13538.
- Brabazon W (1848) The deep sea and coast fisheries of Ireland - with suggestions for the working of a fishing company. James McGlashan, Dublin.
- Bromley, C., McGonigle, C., Ashton, E.C. & Roberts, D. (2015). Restoring degraded European native oyster, *Ostrea edulis*, habitat: is there a case for harrowing? *Hydrobiologia*, 768, 151–165.
- Brown AE, Burn AJ, Hopkins JJ, Way SF (1997) Joint Nature Conservation Committee. The Habitats Directive: selection of Special Areas of Conservation in the UK. JNCC Report No. 270. p.291. Peterborough, UK.
- Buckland FT (1875) Fisheries (Norfolk): report on the fisheries of Norfolk, especially crabs, lobsters, herrings, and the Broads. London: Her Majesty's Stationery Office.
- Buckland FT, Walpole S (1879) Commissioners for Sea Fisheries of England and Wales. London, HMSO.
- Cadée, G., 1968. Molluscan biocoenoses and thanatocoenoses in the Ria de Arosa, Galicia, Spain. *Zoologische Verhandlungen*, 95(1), pp.1-121.
- Cataudella, S.; Massa, F.; Crosetti, D. (eds.) Interactions between aquaculture and capture fisheries: a methodological perspective. Studies and Reviews. General Fisheries Commission for the Mediterranean. No. 78. Rome, FAO. 2005. 229p.*
- Christianen, M.J.A., Lengkeek, W., Bergsma, J.H., Coolen, J.W.P., Didderen, K., Dorenbosch, M., Driessen, F.M.F., Kamermaans, P., Reuchlin-Hugenholtz, E., Sas, H., Smaal, A., van den Wijngaard, K.A. & van der Have, T.M. (2018). Return of the native facilitated by the invasive? Population composition, substrate preferences and epibenthic species richness of a recently discovered shellfish reef with native European flat oysters (*Ostrea edulis*) in the North Sea. *Marine Biology Research*, 14, 590–597.
- Collin, J (1871) Tidsskrift for populære Fremstillinger af Naturvidenskaben. Kort over Østersbankerne i Limfjorden. Denmark, København.
- Collin, J.S. & Denmark. Ministeriet for kirke- og undervisningsvæsenet. (1884). *Om Limfjordens tidligere og nuværende marine fauna, med særligt hensyn til bløddyraunaen*. Kjøbenhavn : Gyldendal.
- Crocetta, F., Bitar, G., Zibrowius, H. & Oliverio, M. (2013). Biogeographical homogeneity in the eastern Mediterranean Sea. II. Temporal variation in Lebanese bivalve biota. *Aquatic Biology*, 19, 75–84.
- Culloty, S.. C. & Mulcahy, M.F. (2007). *Bonamia ostreae* in the native oyster *Ostrea edulis*. A review. Marine Environment and Health Series.
- Culloty, S.C., Cronin, M.A. & Mulcahy, M.F. (2008). Potential resistance of a number of populations of the oyster *Ostrea edulis* to the parasite *Bonamia ostreae*. *Aquaculture*, 237, 41–58.
- Davis, H.C. & Ansell, A.D. (1962). Survival and Growth of Larvae of the European Oyster, *O. edulis*, at Lowered Salinities. *Biological Bulletin*, 122, 33–39.
- Dhora, D. (2009) Mollusks of Albania *Arch. Biol. Sci.*, Belgrade, 61 (3), 537-553 DOI:10.2298/ABS0903547D
- Dridi, S., Romdhane, M.S., Heurtebise, S., Cafsi, E.M., Boudry, P. and Lapegue, S., 2008. Genetic characterisation of oyster populations along the north-eastern coast of Tunisia. *African Journal of Marine Science*, 30(3), pp.489-495.

- Eagling, L.E., Ashton, E.C., Jensen, A.C., Sigwart, J.D., Murray, D. & Roberts, D. (2018). Spatial and temporal differences in gonad development, sex ratios and reproductive output, influence the sustainability of exploited populations of the European oyster, *Ostrea edulis*. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 28, 270–281.
- Egerton, S., Lynch, S.A., Prado-Alvarez, M., Flannery, G., Brennan, E., Hugh-Jones, T., Hugh-Jones, D. & Culloty, S.C. (2020). A naïve population of European oyster *Ostrea edulis* with reduced susceptibility to the pathogen *Bonamia ostreae*: are S-strategy life traits providing protection? *Integrative and Comparative Biology*.
- EUNIS (2022) EUNIS marine habitat classification review (tabular)- version 1, 2022. <https://www.eea.europa.eu/en/datahub/datahubitem-view/6388330ea-90e6-4e41-81ea-e70f25ae7117> Accessed December 5th 2023.
- Ezgeta-Balić, D., Šantić, D., Šegvić-Bubić, T., Bojanić, N., Bužanić, M., Vidjak, O., Varezić, D.B., Stagličić, N., Kundid, P., Peharda, M., Žužul, I., Grubišić, L. & Briski, E. (2020). Competitive feeding interactions between native *Ostrea edulis* and non-native *Crassostrea gigas* with implications of introducing *C. gigas* into commercial aquaculture in the eastern Adriatic Sea. *Marine Environmental Research*, 160, 105051.
- Fomsgaard, K. & Petersen, J.K. (2015) Pleje af østersbestanden i Limfjorden. DTU Aqua-rapport nr. 291-2015. Institut for Akvatiske Ressourcer, Danmarks Tekniske Universitet. 31 pp.
- Ferrario, J., Bogi, C., Cardeccia, A., Langeneck, J., Marchini, A., Ulman, A., & Occhipinti Ambrogi, A. (2018). Fouling community in the harbour of Piran (Slovenia). *Biologia Marina Mediterranea*, 25, 147-151.
- Fischer, W. (1997) Ein Beitrag zur marinen Molluskenfauna Griechenlands. Die Bivalven der Drepano Beach (Igoumenitsa, Epirus). *Club Conchylia Informationen* 29,57-60.
- García García, B., Perez Camacho, A., Abellan Martinez, E. and Garcia Alcazar, A. (1989). Stock assessment and distribution of the European flat oyster, *Ostrea edulis* L., in the Mar Menor (Murcia, SE of Spain). *Aquaculture: A Biotechnology in Progress* (eds, N. De Pauw, E. Jaspers, H. Ackefors and N. Wilkins) European Aquaculture Society, 1, pp. 307-317.
- Gerken, J. & Schmidt, A. (2014). Aktueller Status der Europäischen Auster (*Ostrea edulis*) und Möglichkeiten einer Wiederansiedlung in der deutschen Nordsee. Bundesamt für Naturschutz, Bonn-Bad Godesberg.
- Grizel, H., Comps, M., Bonami, J.R., Cousserans, F., Duthoit, J.L., Le Pennec, M.A. (1974). Recherches sur l'agent de la maladie de la glande digestive de *Ostrea edulis* Linné. *Sci. Pêche, Bull. Inst. Pêches Marit.* 240, 7–30.
- Grizel, H. (1985). *Etude des récentes épizooties de l'huître plate Ostrea edulis Linné et de leur impact sur l'ostréiculture bretonne*(Doctoral dissertation, Université des sciences et techniques du Languedoc).
- Harding CW (1882) Inquiries concerning the propagation of American smelt and shad, and notes on the fisheries of the Wash in England. Bulletin of the United States Fish Commission 1, 428-429
- Helmer, L., Farrell, P., Hendy, I., Harding, S., Robertson, M. & Preston, J. (2019). Active management is required to turn the tide for depleted *Ostrea edulis* stocks from the effects of overfishing, disease and invasive species. *PeerJ*, 7:e6431, <https://doi.org/10.7717/peerj.6431>.
- Héra, Z. & Haris, A. (2015) Marine molluscs from Marsa El Brega, Libya (Mollusca: Gastropod and Bivalvia). *Natura Somogyiensis* 27, 37-44. DOI:10.24394/NatSom.2015.27.37

- Hiddink, J.G., Kaiser, M.J., Sciberras, M., McConnaughey, R.A., Mazor, T., Hilborn, R., Collie, J.S., Pitcher, C.R., Parma, A.M., Suuronen, P., Rijnsdorp, A.D. & Jennings, S. (2020). Selection of indicators for assessing and managing the impacts of bottom trawling on seabed habitats. *Journal of Applied Ecology*, 57, 1199–1209.
- Holt, T.J., Rees, E.I., Hawkins, S.J. & Seed, R. (1998). *Biogenic Reefs volume IX. An overview of dynamic and sensitivity characteristics for conservation management of marine SACs. Scottish Association for Marine Science (UK Marine SACs Project)*.
- Holt EWL (1902) The public oyster bed on the coasts of Counties Wicklow and Wexford. Report on the sea and inland fisheries of Ireland for 1901, Part II, Scientific Investigations. Department of Agriculture and Technical Instruction for Ireland.
- Holthuis, T., (2022). Inventering av ostronbankar runt Resö. Överby Naturvård på uppdrag av boende på Resö. [Survey of oyster beds around Resö. Survey performed by Överby Naturvård on request by residents at Resö.] 12 pp.
- Houziaux J.-S., Haelters J. & Kerckhof, F. (2007) Annex 4: Biodiversity science: a case study from Belgian marine waters. *ICES SGBIODIV Report 2007* 18-25.
- Hughes, A., Bonačić, K., Cameron, T., Collins, K., da Costa, F., Debney, A., van Duren, L., Elzinga, J., Fariñas-Franco, J.M., Gamble, C., Helmer, L., Holbrook, Z., Holden, E., Knight, K., Murphy, J.A.J., Pogoda, B., Pouvreau, S., Preston, J., Reid, A., Reuchlin-Hugenholtz, E., Sanderson, W.G., Smyth, D., Stechele, B., Strand, Å., Theodorou, J.A., Uttley, M., Wray, B. & zu Ermgassen, P.S.E. (n.d.). Site selection for European native oyster (*Ostrea edulis*) habitat restoration projects: An expert-derived consensus. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 33, 721-736.
- Hussein, K. S. & Talet, L. B. (2019) A preliminary inventory of biodiversity and benthic habitats of “Plane” Island (Paloma) in Oran Bay, north western Algeria (western Mediterranean). *J. Black Sea/Mediterranean Environment* 25:49-72.
- Iglesias Estepa, D., García Alves, L., Manuel Parada Encisa, J., No Couto, E. & Miranda Bamio, M., (2013) D12. GIS on oyster beds including information on the different aspects assessed within the project in relation to natural oyster beds. *Oysterecover final report*.
- Isle of Man Times. June 27 1874, p4. Manx National Heritage, Isle of Man.
- Irish Fisheries (1836) First Report of the Commissioners of Inquiry into the State of the Irish Fisheries, with the minutes of evidence and the appendix. HMSO, Dublin.
- Janssen, R., Knudsen, S., Todorova, V. & Hoşgör, A.G. (2014). Managing Rapana in the Black Sea: Stakeholder workshops on both sides. *Ocean & Coastal Management*, 87, 75–87.
- Jones H., Gardner C. 2016. Small Bivalve Survey. Assessment and Stock Status Update: 2016 *Ostrea angasi*- Georges Bay *Venerupis largillierti* – Northern Zone, Georges Bay. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Joubin L. (1907) Gisements des mollusques comestibles des côtes de France: la région d’Auray (Morbihan). Bulletin de l’Institut Océanographique (Monaco), n°89. 55 p.
- Joubin L (1910) Études sur les gisements de mollusques comestibles des Côtes de France: la baie de Cancale. Bulletin de l’Institut Océanographique, No.174, 30 Juin 1910.
- Kamermaans, P., Blanco, A., Dalen, P. van, Engelsma, M., Bakker, N., Jacobs, P., Dubbeldam, M., Sambade, I.M., Vera, M. & Martinez, P. (2023). Bonamia-free flat oyster (*Ostrea edulis* L.) seed for restoration projects: non-destructive screening of broodstock, hatchery production and test for Bonamia-tolerance. *Aquat. Living Resour.*, 36, 11.

- Kammarmans, P., Perdon, J. & Bergsma, J. (2022). *Sediment suitability of Frisian Front search areas for European flat oyster (Ostrea edulis) restoration* (No. C021/22). Wageningen Marine Research, Yerseke. Joksimović, A., Mandić, S., Ilić, Z., Đurović, M., Đerić, N. Potential of commercial farming of the European Flat Oyster, *Ostrea edulis*, in the Kotor Bay (Orahovac). V International conference "Aquaculture & Fishery" - Conference Proceedings 242-247.
- Kasoar, T., zu Ermgassen, P.S.E., Carranza, A., Hancock, B. & Spalding, M. (2015). New opportunities for conservation of a threatened biogenic habitat: a worldwide assessment of knowledge on bivalve-reef representation in marine and coastal Ramsar Sites. *Marine and Freshwater Research*, 66, 981–988.
- Keith, D.A., Ferrer-Paris, J.R., Nicholson, E. and Kingsford, R.T. (eds.) (2020). The IUCN Global Ecosystem Typology 2.0: Descriptive profiles for biomes and ecosystem functional groups. Gland, Switzerland: IUCN.
- Kent and Essex Inshore Fisheries and Conservation Authority (2022) Update on native oyster research in Kent and Essex. <https://www.kentandsex-ifca.gov.uk/website-content/agenda-item-b5-native-oysters-1668506822.pdf> [Accessed December 14th 2023]
- Kruschel, C., Schultz, S. T., & Stiefel, J. (2011). High diversity and sensitivity to coastal development of fish and invertebrates of the Novigrad and Karin seas: A metacommunity study in the northern Adriatic, Croatia. In *Annales: Series Historia Naturalis* (Vol. 21, No. 1, p. 11). Scientific and Research Center of the Republic of Slovenia.
- Krøyer H (1837). De danske østersbanker - et bidrag til kundskab om Danmarks Fiskerier. København.
- Ktari-Chakroun, F. and Azouz, A., (1971). Les fonds chabutables de la région sud-est de la Tunisie (golfe de Gabès). Bulletin de l'Institut National Scientifique et Technique d'Océanographie et de Pêche. 2,5-47.
- Laing, I., Walker, P. & Areal, F. (2005). *A feasibility study of native oyster (Ostrea edulis) stock regeneration in the United Kingdom*. Native Oyster Stock Regeneration - A Review of Biological, Technical and Economic Feasibility. CEFAS.
- Lallias, D., Tarris, N., Boudry, P., Bonhomme, F. & Lapegue, S. (2006). Variance of reproductive success in the European flat oyster *Ostrea edulis* assessed by microsatellite-based parentage analyses in a natural population and in hatchery. 98 Annual Meeting National Shellfisheries Association, Monterey, CA (USA), 26-30 Mar 2006. <https://archimer.ifremer.fr/doc/00000/3334/>.
- Levasseur O (2006) Histoire de l'huître en Bretagne. Skol vreizh 58, 83 p.
- Lippold GHC; Funke CP (1810) Neues Natur- und Kunstlexicon enthaltend die wichtigsten und gemeinnützigsten Gegenstände aus der Naturgeschichte, Naturlehre, Chemie und Technologie.
- Lock, K. (2017). *Native oyster, Ostrea edulis, Milford Haven Waterway Survey report 2016*. Seasearch.
- Lown, A.E., Hepburn, L.J., Heywood, J.L. & Cameron, T.C. (2021). European native oysters and associated species richness in the presence of non-native species in a southern North Sea estuary complex. *Conservation Science and Practice*, 3, e361.
- Lubet, P. and Azouz, A., 1969. Etude des fonds chabutables du Golfe de Tunis. Bulletin de l'Institut National Scientifique et Technique d'Océanographie et de Pêche. 3, 87-112.
- Marine Institute & Bord lascaigh Mhara (2022). Shellfish Stocks and Fisheries Review 2021: An assessment of selected stocks. Marine Institute., Galway, Ireland

- Marine Institute & Bord Iascaigh Mhara (2023). Shellfish Stocks and Fisheries Review 2022: An assessment of selected stocks. Marine Institute., Galway, Ireland
- Marsili F (1715) Giornali e memorie per la ricognizione della spiaggia pontificie dell'Adriatico (Journals and Memoirs for the Reconnaissance of the Papal Beaches of the Adriatic). MS 72 F 4 J, Biblioteca Universitaria di Bologna.
- McGeer, C. (2023) In hot water: Mar Menor records historic 31C. EuroWeekly News. <https://euroweeklynews.com/2023/07/21/mar-menor-records-historic-31-degree-july-2023/> [Accessed December 14th 2023]
- Moore, H.F. & Danglede, E. (1915). Condition and extent of the natural oyster bed and barren bottoms of Lavaca Bay, Texas. In: *Report of the U.S. Commission of Fisheries for 1914*. pp. 1–45.
- Mortensen, S., Bøggwald, M., Strohmeier, T., Strand, Ø., Larsson, S.B., Laugen, A.T., Reamon, M.C., Marcussen, J.B. (2023). Kartlegging av østersbestander i Rogaland i 2021 og 2023 - innsamling av data som grunnlag for bestandsforvaltning. Rapport fra havforsknigen 2023-55 ISSN: 1893-4536. 22 s. <https://www.hi.no/hi/nettrapporter/rapport-fra-havforsknigen-2023-55>
- Möbius K (1883) The oyster and oyster culture. Berlin, Hempel & Parey. 1877. Translated by HJ Rice, pp. 683-747.
- Norgard, TC, Bigg, MI, MacConnachie, SEM, Finney, JL and Gillespie, GE (2018) Index site surveys for Olympia oysters (*Ostrea lurida* Carpenter, 1864) in British Columbia – 2009 to 2017. Can. Tech. Rep. Fish. Aquat. Sci. 3153: viii + 88 p.
- Oberle, F.K.J., Storlazzi, C.D., Hanebuth, T.J.J. (2016) What a drag: Quantifying the global impact of chronic bottom trawling on continental shelf sediment. *Journal of Marine Systems* 159, 109-119.
- Oedmann J (1743) Nachrichten von Austern, nach dem Verlangen der Königl. Akademie der Wissenschaften im 2. Quartale 1743 von Johan Oedmann pages 118-120 in Der Königl. Schwedischen Akademie der Wissenschaften Abhandlungen aus der Naturlehre, Haushaltungskunft und Mechanik, auf das Jahr 1744. Sechster Band. Georg Christan Grund, Hamburg, 1751.
- Olsen OT (1885) The fisherman's practical navigator, Second Edition. Imray & Son, London.Orabi, O.H., El-Badry, A.E.-M.A. & Alhejoj, I. (2018). Response of mollusk assemblages to environmental conditions: A case study from the Burullus Lagoon, Egypt. *Journal of the Geological Society of India*, 91, 583–588.
- OSPAR Commission (2009) Background document for *Ostrea edulis* and *Ostrea edulis* beds. 22 pages.
- Öztürk, B. & Öztürk, A.A. (1996). On the biology of the Turkish straits system. *Bulletin de l'Institut oceanographique, Monaco, n° special 17 (1996)*, 205–221.
- Petović, S., 2018. Additions to the checklist of the malacofauna of the Boka Kotorska Bay (south-east Adriatic Sea). *Studia Marina*, 31(1), pp.23-36.
- Philpots JR (1891) Oysters and all about them. John Richardson and Co., London.
- Pitacco, V., Orlando-Bonaca, M., Mavric, B., & Lipej, L. (2014). Macrofauna associated with a bank of *Cladocora caespitosa* (Anthozoa, Scleractinia) in the Gulf of Trieste (northern Adriatic) /Macrofauna associata ad un banco di *Cladocora caespitosa* (Anthozoa, Scleractinia) nel Golfo di Trieste (Adriatico settentrionale). In *Annales: Series Historia Naturalis* (Vol. 24, No. 1, p. 1). Scientific and Research Center of the Republic of Slovenia.
- Pitcher, C.R., Hiddink, J.G., Jennings, S., Collie, J., Parma, A.M., Amoroso, R., Mazor, T., Sciberras, M., McConnaughey, R.A., Rijnsdorp, A.D., Kaiser, M.J., Suuronen, P. & Hilborn, R. (2022). Trawl impacts on the relative status of biotic communities of seabed sedimentary habitats in 24 regions worldwide. *Proceedings of the National Academy of Sciences*, 119, e2109449119.

- Poirier, C., Chaumillon, E. & Arnaud, F. (2011). Siltation of river-influenced coastal environments: Respective impact of late Holocene land use and high-frequency climate changes. *Marine Geology*, 290, 51–62.
- Pontoppidan E (1769) Den Danske Atlas. København, Danmark.
- Pouvreau S., Cochet H., Bargat F., Petton S., Le Roy V., Guillet T., Potet M. (2021). Current distribution of the residual flat oysters beds (*Ostrea edulis*) along the west coast of France . SEANOE . <https://doi.org/10.17882/79821>
- Preston, J., Fabra, M., Helmer, L., Johnson, E., Harris-Scott, E. & Hendy, I.W. (2020). Interactions of larval dynamics and substrate preference have ecological significance for benthic biodiversity and *Ostrea edulis* Linnaeus, 1758 in the presence of *Crepidula fornicata*. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 30, 2133–2149.
- Ramday G., Kennon N.A., Cooper-Young E., Hanley N., Hugh-Jones, T., Janson M., Mair J.McD., Robertson Jones A., Waring K., Weber S. & Sanderson W.G (2024). The oyster population and scope for environmental enhancement in Loch Ryan. Confidential report to LROFC Ltd and Solway Firth Partnership. 18pp.
- Rasmussen, P., Strand Petersen, K., & Ryves, D. B. (2007). Environmental change in Danish marine waters during the Roman Warm Period inferred from mollusc data. *GEUS Bulletin*, 13, 21–24. <https://doi.org/10.34194/geusb.v13.4967>
- Rees, H.L., Waldock, R., Matthiessen, P. & Pendle, M.A. (2001). Improvements in the epifauna of the Crouch Estuary (United Kingdom) following a decline in TBT concentrations. *Marine Pollution Bulletin*, 42, 137–144.
- Renda, W., Amati, B., Bogi, C., Bonomolo, G., Capua, D., Dell'Angelo, B., & et al. (2022). The new Checklist of the Italian Fauna: marine Mollusca. Biogeographia. *The Journal of Integrative Biogeography*, 37.
- Ruiz, C., Martínez, D., Mosquera, G., Abad, M. & Sánchez, J.L. (1992). Seasonal variations in condition, reproductive activity and biochemical composition of the flat oyster, *Ostrea edulis*, from San Cibrao (Galicia, Spain). *Marine Biology*, 112, 67–74.
- Ruiz, J.M, Albentosa, M., Aldeguer, B., Álvarez-Rogel, J., Antón, J., Belando, M.D., Bernardeau, J., Campillo, J.A., Domínguez, J.F., Ferrera, I., Fraile-Nuez, E., García, R., Gómez-Ballesteros, M., Gómez, F., González-Barberá, G., Gómez-Jakobsen, F., León, V.M., López-Pascual, C., Marín-Guirao, L., Martínez-Gómez, C., Mercado, J.M.m Nebot, E., Ramos, A., Rubio, E., Santos, J., Santos, F., Vázquez-Luis, M., Yebra, L. (2020) Informe de evolución y estado actual del Mar Menor en relación al proceso de eutrofización y sus causas. Informe de asesoramiento técnico del Instituto Español de Oceanografía (IEO). 165p.
- Sakalli, A., (2017) Sea surface temperature change in the Mediterranean Sea under climate change: A linear model for simulation of the sea surface temperature up to 2100. *Applied Ecology and Environmental Research* 15: 707-716.
- Sambade, I.M., Casanova, A., Blanco, A., Gundappa, M.K., Bean, T.P., Macqueen, D.J., Houston, R.D., Villalba, A., Vera, M., Kamermans, P. & Martínez, P. (2022). A single genomic region involving a putative chromosome rearrangement in flat oyster (*Ostrea edulis*) is associated with differential host resilience to the parasite *Bonamia ostreae*. *Evolutionary Applications*, 15, 1408–1422.
- Sander, L., Hass, H.C., Michaelis, R., Groß, C., Hausen, T. & Pogoda, B. (2021). The late Holocene demise of a sublittoral oyster bed in the North Sea. *PLOS ONE*, 16, e0242208.
- Savini, D. & Occhipinti-Ambrogi, A. (2006). Consumption rates and prey preference of the invasive gastropod *Rapana venosa* in the Northern Adriatic Sea. *Helgol Mar Res*, 60, 153–159.

- Shelmerdine, R.L. & Leslie, B. (2009). *Restocking of the native oyster, Ostrea edulis, in Shetland: habitat identification study*. Scottish Natural Heritage Commissioned Report No. 396.
- Shpigel M. (1989) Gametogenesis of the European Flat Oyster (*Ostrea edulis*) and Pacific Oyster (*Crassostrea gigas*) in Warm Water in Israel. *Aquaculture* 80, 343-349.
- Smyth, D.M., Horne, N.S., Ronayne, E., Millar, R.V., Joyce, P.W.S., Hayden-Hughes, M. & Kregting, L. (2020). Wild gregarious settlements of *Ostrea edulis* in a semi-enclosed sea lough: a case study for unassisted restoration. *Restoration Ecology*, 28, 645–654
- Smyth, D., Millar, R., Clements, A., McIvenny, H. & Hayden-Hughes, M. (2023). Population dynamics of the European native oyster in a Marine Conservation Zone exposed to unregulated harvesting. *Aquat. Living Resour.*, 36, 3.
- Smyth, D., Roberts, D. & Browne, L. (2009). Impacts of unregulated harvesting on a recovering stock of native oysters (*Ostrea edulis*). *Marine Pollution Bulletin*, 58, 916–922.
- Southern Inshore Fisheries and Conservation Authority (2019) Solent oyster stock assessment. Summer 2019. 13p. Available at: <https://secure.toolkitfiles.co.uk/clients/25364/sitedata/Redesign/Solent-Dredge-Fisheries/Solent-Oyster-Survey-2019.pdf>
- Stagličić, N., Šegvić-Bubić, T., Ežgeta-Balić, D., Bojanić Varezić, D., Grubišić, L., Žuvić, L., Lin, Y. & Briski, E. (2020). Distribution patterns of two co-existing oyster species in the northern Adriatic Sea: The native European flat oyster *Ostrea edulis* and the non-native Pacific oyster *Magallana gigas*. *Ecological Indicators*, 113, 106233.
- Stechele, B., Maar, M., Wijsman, J., Van der Zande, D., Degraer, S., Bossier, P. & Nevejan, N. (2022). Comparing life history traits and tolerance to changing environments of two oyster species (*Ostrea edulis* and *Crassostrea gigas*) through Dynamic Energy Budget theory. *Conservation Physiology*, 10.
- Sturgeon, S., Street, K., Trundle, C., and Jenkin, A. 2021. Fal Oyster Fishery 2020-2021 Season Permit Statistics Report. Cornwall Inshore Fisheries and Conservation Authority (Cornwall IFCA), Hayle.
- Sütçüoğlu, N. & Korun, J. (2011). Detection of natural bacterial biota of European flat oyster (*Ostrea edulis*, L. 1758) from the northern Sea of Marmara. *Journal of Aquatic Sciences and Engineering*, 26, 13–26.
- Thain, J.E. & Waldock, M.J. (1986). The impact of Tributyl Tin (TBT) antifouling paints on molluscan fisheries. *Water Science and Technology*, 18, 193–202.
- The Aberystwith Observer, 24 June 1897 p4. The National Library of Wales.
- The Cambrian News and Merionethshire Standard, 15 March 1889, p8. The National Library of Wales.
- Thorngren, L., Bergström, P., Holthuis, T.D. & Lindegarth, M. (2019). Assessment of the population of *Ostrea edulis* in Sweden: A marginal population of significance? *Ecology and Evolution* 9, 13877–13888.
- Thurstan, R.H., McCormick, H., Preston, J., Ashton, E.C., Bennema, F.P., Bratoš Cetinić, A., Brown, J.H., Cameron, T.C., da Costa, F., Donnan, D., Ewers, C., Fortibuoni, T., Galimany, E., Giovanardi, O., Grancher, R., Grech, D., Hayden-Hughes, M., Helmer, L., Jensen, K.T., Juanes, J.A., Latchford, J., Moore, A.B.M., Moutopoulos, D.K., Nielsen, P., von Nordheim, H., Ondiviela, B., Peter, C., Pogoda, B., Poulsen, B., Pouvreau, S., Roberts, C.M., Scherer, C., Smaal, A.C., Smyth, D., Strand, Å, Theodorou, J.A., zu Ermgassen, P.S.E. (in review a) The world was our oyster: Records reveal the vast historical extent of European oyster reef ecosystems. <https://doi.org/10.32942/X20W43>

- Thurstan, R.H., McCormick, H., Preston, J., Ashton, E.C., Bennema, F.P., Bratoš Cetinić, A., Brown, J.H., Cameron, T.C., da Costa, F., Donnan, D., Ewers, C., Fortibuoni, T., Galimany, E., Giovanardi, O., Grancher, R., Grech, D., Hayden-Hughes, M., Helmer, L., Jensen, K.T., Juanes, J.A., Latchford, J., Moore, A.B.M., Moutopoulos, D.K., Nielsen, P., von Nordheim, H., Ondiviela, B., Peter, C., Pogoda, B., Poulsen, B., Pouvreau, S., Scherer, C., Smaal, A.C., Smyth, D., Strand, Å., Theodorou, J.A., zu Ermgassen, P.S.E. (in review b) Historical dataset details the distribution, extent and form of lost *Ostrea edulis* reef ecosystems. <https://doi.org/10.32942/X28C99>
- Todorova V, Micu D, Klisurov L (2009) Unique oyster reefs discovered in the Bulgarian Black Sea. *Hydrobiologie* 62, 871- 874.
- Trožić-Borovac S, Ičanović J, Imamović A, Bijedić A, Čelebić M. 2022. Ecological state of the environment and phenotypic characteristics of Mussel *Mytilus Galloprovincialis* Lamarck, 1873 in the Neum Bay (Bosnia and Herzegovina). *Veterinaria*, 71(2), 205-218.
- Tully, O. & Clarke, S. (2012). *The status and management of oyster (Ostrea edulis) in Ireland*. Irish Fisheries Investigations. Marine Institute.
- van Aartsen, J.J., Menkhorst, H.P.M.G. and Gittenberger, E., 1984. The marine Mollusca of the Bay of Algeciras, Spain, with general notes on Mitrella, Marginellidae and Turridae. *Basteria*, 2, pp.3-134.
- Virvilis, C. & Angelidis, P. (2006). Presence of the parasite *Marteilia* sp. in the flat oyster (*Ostrea edulis* L) in Greece. *Aquaculture*, 259, 1–5.
- Watling, L. & Norse, E.A. (1998). Disturbance of the seabed by mobile fishing gear: A comparison to forest clearcutting. *Conservation Biology*, 12, 1180–1197.
- Wilmsen FP (1831) Handbuch der Naturgeschichte für die Jugend und ihre Lehrer. Zweiter Band. Druck und Verlag von C.F. Amelang, Berlin.
- Winslow, F. (1881). Report on the oyster beds of the James River, Virginia, and of Tangier and Pocomoke Sounds, Maryland and Virginia. *Annual Report of the United States Coast and Geodetic Survey*, Appendix 11, 269–353
- Yildiz, H., Berber, S., Acarli, S. & Vural, P. (2011). Seasonal variation in the condition index, meat yield and biochemical composition of the flat oyster *Ostrea edulis* (Linnaeus, 1758) from the Dardanelles, Turkey. *Italian Journal of Animal Science*, 10, e5.
- Zavodnik, D., Kovačić, M., (2000) Index of marine fauna in Rijeka bay (Adriatic Sea, Croatia). *Natura Croatica* 9,297-379. *ISSN 1330-0520*
- Zolotarev, P.N. & Terentyev, A.S. (2012). Changes in the macrobenthic communities of the Gudauta Oyster Bank. *Oceanology*, 52, 231–237.
- zu Ermgassen, P.S.E. (2017) Milford Haven Native Oyster Regeneration Project - Stage One (current status and practicalities). A report commissioned by West Wales Shellfishermans Association Ltd. supported by the Milford Haven Native Oyster Regeneration Management Group. pp54.