

SUPPLEMENTARY MATERIAL

BYCATCH COMPOSITION

Table S1. Bycatch composition for beach seining (BS) and bottom trawling (BT) (Nunes & Rosa 1998) in municipality of Lucena, state of Paraíba, northeast Brazil. TAXA: order, family, and species, classification based on Nelson et al. (2016). (* data previously published by Passarone et al. (2019); sp. specimens identified at family level; sp.1, sp.2 (...) new species in process of description by other authors); N: number of individuals; N (%): percentage of individuals; IUCN: Level of species extinction risk according to the categories proposed by IUCN and assessed by ICMBio [data deficient (DD), least concern (LC), near threatened (NT), not assessed (NA)]; FMFG: feeding mode functional group classification, by Elliott et al. (2007) [detritivore (DV), herbivore (HV), omnivore (OV), piscivore (PV), zoobenthivore (ZB), zooplanktivore (ZP)]; EUFG: estuary use functional group classification, by Elliott et al. (2007) [marine migrant (MM), marine straggler (MS), estuarine (ES)]; REF.: references (x information not available in the literature).

TAXA	BS	BT	IUCN	FMFG	EUFG	REF.
	N (N %)	N (N%)				
TORPEDINIFORMES						
Narcinidae						
<i>Narcine brasiliensis</i> (Olfers, 1831)		1 (0.02%)	LC	ZB	MS	Bornatowski et al. 2006, Vasconcelos-Filho & Oliveira 1999
PRISTIFORMES						
Rhinobatidae						
<i>Pseudobatos percellens</i> (Walbaum, 1792)	1 (< 0.01%)		DD	ZB	MS	Shibuya 2005, Passos et al. 2013
MYLIOBATIFORMES						
Dasyatidae						
<i>Hypanus guttatus</i> (Bloch and Schneider, 1801)	3 (0.01%)		LC	ZB	MS	Gianeti 2011, Vasconcelos-Filho & Oliveira 1999
Gymnuridae						
<i>Gymnura micrura</i> (Bloch and Schneider, 1801)	5 (0.02%)		NT	PV	MS	Yokota et al. 2013, McEachran 1990
Urotrygonidae						
<i>Urotrygon microphthalmum</i> Delsman, 1941		1 (0.02%)	LC	ZB	MS	Santander-Neto 2015, FAO 2016
Myliobatidae						
<i>Aetobatus narinari</i> (Euphrasen, 1790)	1 (< 0.01%)		DD	ZB	MS	Schluessel et al. 2010, Tagliafico et al. 2012
ANGUILLIFORMES						
Muraenidae						
<i>Gymnothorax funebris</i> Ranzani, 1839	1 (< 0.01%)		DD	ZB	MS	Randall 1967, Vasconcelos-Filho & Oliveira 1999
Ophichthidae						
<i>Ophichthus cylindroideus</i> (Ranzani, 1839)	1 (< 0.01%)	1 (0.02%)	LC	x	MM	Favero et al. 2019

Table S1. Continuation.

Moringuidae						
<i>Neoconger mucronatus</i> Girard, 1858	1 (< 0.01%)		LC	ZB	x	Mikkola 2016
CLUPEIFORMES						
Pristigasteridae						
<i>Chirocentrodon bleekermanus</i> (Poey, 1867)*	2903 (9.36%)	215 (3.56%)	LC	ZP	MS	Corrêa et al. 2005, Passos et al. 2013
<i>Odontognathus mucronatus</i> Lacepède, 1800*	149 (0.48%)	372 (6.17%)	LC	ZP	MM	Santos et al. 2011, Araújo 2012
<i>Pellona harroweri</i> (Fowler, 1917)*	3943 (12.72%)	1288 (21.36%)	LC	ZP	MS	Muto et al. 2008, Ribeiro et al. 2019
Engraulidae						
<i>Anchoa filifera</i> (Fowler, 1915)*	114 (0.37%)		LC	ZP	MM	Passos et al. 2013, Vasconcelos-Filho & Oliveira 1999
<i>Anchoa marinii</i> Hildebrand, 1943		23 (0.38%)		ZP	MM	Ribeiro et al. 2019
<i>Anchoa</i> sp.	7 (0.02%)		-	-	-	
<i>Anchoa spinifer</i> (Valenciennes, 1848)*	55 (0.18%)	4 (0.07%)	LC	PV	MM	Nizinski 2002, Vasconcelos-Filho & Oliveira 1999
<i>Anchovia clupeioides</i> (Swainson, 1839)*	172 (0.55%)	182 (3.02%)	LC	ZP	MM	De Paiva et al. 2008, Vasconcelos-Filho & Oliveira 1999
<i>Cetengraulis edentulus</i> (Cuvier, 1829)*	4960 (16%)	30 (0.50%)	LC	ZP	MS	De Paiva et al. 2008, Vasconcelos-Filho & Oliveira 1999
<i>Engraulis anchoita</i> Hubbs and Marini, 1935		12 (0.20%)		ZP	MS	Ribeiro et al. 2019
<i>Lycengraulis grossidens</i> (Spix and Agassiz, 1829)*	603 (1.95%)	84 (1.39%)	LC	PV	MM	Bortoluzzi et al. 2006, Mai & Vieira 2013
Clupeidae						
<i>Brevoortia pectinata</i> (Jenyns, 1842)		13 (0.22%)		ZP	ES	Ribeiro et al. 2019
<i>Harengula clupeiola</i> (Cuvier, 1829)*	213 (0.69%)		LC	ZP	MS	De Paiva et al. 2008, Vasconcelos-Filho & Oliveira 1999
<i>Harengula</i> sp.	2 (0.01%)		-	-	-	
<i>Lile piquitinga</i> (Schreiner, Miranda and Ribeiro, 1903)	25 (0.08%)		LC	ZP	MS	Sales 2015, Vasconcelos-Filho & Oliveira 1999
<i>Opisthonema oglinum</i> (Lesueur, 1818)*	6364 (20.53%)	10 (0.17%)	LC	ZP	MS	Vasconcelos-Filho 1979, Vasconcelos-Filho & Oliveira 1999
<i>Rhinosardinia bahiensis</i> (Steindachner, 1879)	26 (0.08%)		LC	ZP	ES	Clark & Pessanha 2015

Table S1. Continuation.

<i>Rhinosardinia</i> sp.	3 (0.01%)		-	-	-	
SILURIFORMES						
Ariidae						
<i>Ariidae</i> sp.	28 (0.09%)		-	-	-	
<i>Aspistor luniscutis</i> (Valenciennes, 1840)	14 (0.05%)		LC	PV	MS	Bessa et al. 2012
<i>Aspistor quadriscutis</i> (Valenciennes, 1840)*	48 (0.15%)	1 (0.02%)	LC	PV	MM	Almeida 2018, de Andrade-Tubino et al. 2008
<i>Bagre bagre</i> (Linnaeus, 1766)*	59 (0.19%)	7 (0.12%)	NT	PV	MM	Pinheiro-Sousa et al. 2015, Mourão et al. 2014
<i>Bagre marinus</i> (Mitchill, 1815)*	199 (0.64%)	30 (0.50%)	DD	ZB	MM	Mendoza-Carranza 2003, Segura-Bertolini & Mendoza-Carranza 2013
<i>Cathorops</i> sp.	21 (0.07%)	99 (1.64%)	-	-	-	
<i>Cathorops spixii</i> (Agassiz, 1829)*	152 (0.49%)		LC	ZB	ES	Barletta & Blaber 2007
<i>Cathorops agassizii</i> Eigenmann and Eigenmann, 1888)	120 (0.39%)		LC	ZB	ES	Barletta & Blaber 2007
<i>Notarius grandicassis</i> (Valenciennes, 1840)	17 (0.05%)	11 (0.18%)	LC	ZB	MM	Mendes & Barthem 2010, Marцениuk et al. 2017
<i>Sciades herzbergii</i> (Bloch, 1794)	22 (0.07%)		LC	ZB	ES	Giarrizzo & Saint-Paul 2008, Vasconcelos-Filho & Oliveira 1999
MUGILIFORMES						
Mugilidae						
<i>Mugil curema</i> Valenciennes, 1836	1 (< 0.01%)		DD	ZB	MM	Medeiros et al. 2017, Vasconcelos-Filho & Oliveira 1999
<i>Mugil curvidens</i> Valenciennes, 1836	1 (< 0.01%)		DD	DV	MM	Favero et al. 2019
<i>Mugil</i> sp.	14 (0.05%)		-	-	-	
BLENNIIFORMES						
Dactyloscopidae						
<i>Dactyloscopus crossotus</i> Starks, 1913		6 (0.10%)	LC	ZB	MM	Schwamborn 2004, Favero et al. 2019
ATHERINIFORMES						
Atherinopsidae						

Table S1. Continuation.

<i>Atherinella blackburni</i> (Schultz, 1949)	4 (0.01%)		LC	ZP	MM	Gonzalez & Vaske-Júnior 2017, Favero et al. 2019
<i>Atherinella brasiliensis</i> (Quoy and Gaimard, 1825)	1 (< 0.01%)		LC	OV	ES	Favero et al. 2019
BELONIFORMES						
Hemiramphidae						
<i>Hyporhamphus unifasciatus</i> (Ranzani, 1841)	2 (0.01%)		NT	HV	MM	Favero et al. 2019
Belonidae						
<i>Strongylura marina</i> (Walbaum, 1792)	16 (0.05%)		LC	PV	MM	Ribeiro et al. 2019
CARANGIFORMES						
Carangidae						
<i>Carangoides bartholomaei</i> (Cuvier, 1833)	1 (< 0.01%)		LC	PV	MM	De Paiva et al. 2008, Favero et al. 2019
<i>Caranx crysos</i> (Mitchill, 1815)	1 (< 0.01%)		LC	PV	MM	Sley et al 2009, Barletta & Blaber 2007
<i>Caranx hippos</i> (Linnaeus, 1766)	12 (0.04%)		LC	PV	MS	Gonzalez et al. 2021, Vasconcelos-Filho & Oliveira 1999
<i>Caranx latus</i> Agassiz, 1831	3 (0.01%)		LC	PV	MS	Gonzalez et al. 2021, Vasconcelos-Filho & Oliveira 1999
<i>Caranx</i> sp.	1 (< 0.01%)		-	-	-	
<i>Chloroscombrus chrysurus</i> (Linnaeus, 1766)*	356 (1.15%)	22 (0.36%)	LC	ZB	MS	Oliveira-Silva & Lopes 2002, Vasconcelos-Filho & Oliveira 1999
<i>Hemicaranx amblyrhynchus</i> (Cuvier, 1833)	1 (< 0.01%)		LC	ZB	MM	Ribeiro et al. 2019
<i>Oligoplites palometa</i> (Cuvier, 1832)	2 (0.01%)	1 (0.02%)	LC	PV	MM	Ribeiro et al. 2019
<i>Oligoplites saurus</i> (Bloch and Schneider, 1801)	1 (< 0.01%)		LC	PV	MM	Ribeiro et al. 2019, Vasconcelos-Filho & Oliveira 1999
<i>Pseudocaranx dentex</i> (Bloch and Schneider, 1801)	6 (0.02%)		LC	ZB	MM	Sazima 1998, Rowling & Raines 2000
<i>Selene brownii</i> (Cuvier, 1816)*	37 (0.12%)		LC	ZB	MM	Bomfim 2014, Lima et al. 2021
<i>Selene setapinnis</i> (Mitchill, 1815)	3 (0.01%)	8 (0.13%)	LC	PV	MM	Ribeiro et al. 2019
<i>Selene vomer</i> (Linnaeus, 1758)*	90 (0.29%)	5 (0.08%)	LC	PV	MS	Daros 2014, Vasconcelos-Filho & Oliveira 1999
<i>Trachinotus carolinus</i> (Linnaeus, 1766)	12 (0.04%)		LC	ZB	MM	Denadai et al. 2013

Table S1. Continuation.

<i>Trachinotus falcatus</i> (Linnaeus, 1758)	2 (0.01%)		LC	ZB	MS	Höfling et al. 2000, Vasconselos-Filho & Oliveira 1999
<i>Trachinotus goodei</i> Jordan and Evermann, 1896	1 (< 0.01%)		LC	ZB	MS	Stefanoni 2008, Araújo et al. 2016
<i>Trachinotus marginatus</i> Cuvier, 1832	1 (< 0.01%)		LC	x	MS	Passos et al. 2013
<i>Trachinotus</i> sp.	2 (0.01%)		-	-	-	
ISTIOPHORIFORMES						
Sphyraenidae						
<i>Sphyraena guachancho</i> Cuvier, 1829	16 (0.05%)	8 (0.13%)	LC	PV	MM	Akadjé et al. 2013, Reis-Filho et al. 2010
PLEURONECTIFORMES						
Paralichthyidae						
<i>Citharichthys</i> sp.	1 (< 0.01%)		-	-	-	
<i>Citharichthys spilopterus</i> Günther, 1862	8 (0.03%)		LC	ZB	MM	Vasconselos-Filho et al. 2010, Vasconselos-Filho & Oliveira 1999
<i>Paralichthys brasiliensis</i> (Ranzani, 1842)	6 (0.02%)		LC	ZB	MM	Anderson et al. 2015, Vasconselos-Filho & Oliveira 1999
Achiridae						
<i>Achirus achirus</i> (Linnaeus, 1758)	7 (0.02%)		LC	ZB	ES	Mourão et al. 2014, Vasconselos-Filho & Oliveira 1999
<i>Achirus declivis</i> Chabanaud, 1940	21 (0.07%)	5 (0.08%)	LC	ZB	ES	Ribeiro et al. 2019, Vasconselos-Filho & Oliveira 1999
<i>Achirus lineatus</i> (Linnaeus, 1758)	1 (< 0.01%)	2 (0.03%)	LC	ZB	ES	Vasconselos-Filho et al. 2010, Vasconselos-Filho & Oliveira 1999
<i>Trinectes paulistanus</i> (Miranda, 1915)	20 (0.06%)		LC	ZB	MM	Contente et al. 2009, Favero et al. 2019
Cynoglossidae						
<i>Symphurus plagusia</i> (Bloch and Schneider, 1801)		3 (0.05%)	LC	ZB	MM	Vasconselos-Filho et al. 2010, Favero et al. 2019
<i>Symphurus tessellatus</i> (Quoy and Gaimard, 1824)	29 (0.09%)	10 (0.17%)	LC	ZB	MM	Chaves & Serenato 1998, Favero et al. 2019
SYNGNATHIFORMES						
Dactylopteridae						
<i>Dactylopterus volitans</i> (Linnaeus, 1758)	6 (0.02%)		LC	ZB	MM	Guedes et al. 2015, Favero et al. 2019

Table S1. Continuation.

SCOMBRIFORMES						
Trichiuridae						
<i>Trichiurus lepturus</i> Linnaeus, 1758*	1154 (3.72%)	21 (0.35%)	LC	PV	MM	Vasconcelos-Filho et al. 2010, Vasconcelos-Filho & Oliveira 1999
Scombridae						
<i>Scomberomorus brasiliensis</i> Collette, Russo and Zavala-Camin, 1978	11 (0.04%)		LC	PV	MS	Mourão et al. 2014, Vasconcelos-Filho & Oliveira 1999
<i>Scomberomorus cavalla</i> (Cuvier, 1829)	5 (0.02%)		LC	PV	MS	Fonteles-Filho 1988, Medeiros et al. 2018
Stromateidae						
<i>Peprilus paru</i> (Linnaeus, 1758)	1 (< 0.01%)		LC	PV	MS	Bowman et al. 2000, Passos et al. 2013
PERCIFORMES						
Centropomidae						
<i>Centropomus parallelus</i> Poey, 1860	3 (0.01%)		LC	ZB	MM	Lira et al. 2017, Vasconcelos-Filho & Oliveira 1999
<i>Centropomus ensiferus</i> Poey, 1860	1 (< 0.01%)		LC	PV	MM	Rueda & Defeo 2003
Gerreidae						
<i>Diapterus auratus</i> Ranzani, 1842	2 (0.01%)		LC	ZB	MM	Ribeiro et al. 2019, Vasconcelos-Filho & Oliveira 1999
<i>Diapterus rhombeus</i> (Cuvier, 1829)	9 (0.03%)		LC	ZB	MM	Ribeiro et al. 2019, Vasconcelos-Filho & Oliveira 1999
<i>Eucinostomus argenteus</i> Baird and Girard, 1855		10 (0.17%)	LC	ZB	MM	Leão 2016, Vasconcelos-Filho & Oliveira 1999
<i>Eucinostomus gula</i> (Quoy and Gaimard, 1824)	10 (0.03%)	4 (0.07%)	LC	ZB	MM	Vasconcelos-Filho et al. 2010, Vasconcelos-Filho & Oliveira 1999
<i>Eucinostomus melanopterus</i> (Bleeker, 1863)	1 (< 0.01%)		LC	ZP	MM	Araújo et al. 2016, Vasconcelos-Filho & Oliveira 1999
Mullidae						
<i>Pseudupeneus maculatus</i> (Bloch, 1793)	4 (0.01%)		LC	ZB	MS	Soares et al. 2020, Vasconcelos-Filho & Oliveira 1999
Serranidae						
<i>Rypticus randalli</i> Courtenay, 1967	2 (0.01%)	2 (0.03%)	LC	PV	MS	Medeiros et al. 2017, Ribeiro et al. 2019

Table S1. Continuation.

Haemulidae						
<i>Conodon nobilis</i> (Linnaeus, 1758)*	155 (0.5%)	28 (0.46%)	LC	ZB	MS	Lira et al. 2019, Vasconcelos-Filho & Oliveira 1999
<i>Genyatremus luteus</i> (Bloch, 1790)	8 (0.03%)	1 (0.02%)	LC	ZB	MS	Barletta & Blaber 2007, Vasconcelos-Filho & Oliveira 1999
<i>Haemulopsis corvinaeformis</i> (Steindachner, 1868)*	520 (1.68%)	426 (7.06%)	LC	ZB	MS	Silva et al. 2018, Vasconcelos-Filho & Oliveira 1999
<i>Orthopristis ruber</i> (Cuvier, 1830)	1 (< 0.01%)	2 (0.03%)	LC	ZB	MS	Almeida-Silva et al. 2015, Ribeiro et al. 2019
Lutjanidae						
<i>Lutjanus jocu</i> (Bloch and Schneider, 1801)	1 (< 0.01%)		NT	ZB	MS	Monteiro et al. 2009, Vasconcelos-Filho & Oliveira 1999
Polynemidae						
<i>Polydactylus virginicus</i> (Linnaeus, 1758)*	916 (2.95%)	48 (0.80%)	LC	ZB	MM	Barletta & Blaber 2007, Vasconcelos-Filho & Oliveira 1999
SCORPAENIFORMES						
Triglidae						
<i>Prionotus punctatus</i> (Bloch, 1793)	3 (0.01%)		LC	ZB	MS	Ribeiro et al. 2019, Vasconcelos-Filho & Oliveira 1999
MORONIFORMES						
Ephippidae						
<i>Chaetodipterus faber</i> (Broussonet, 1782)*	48 (0.15%)	2 (0.03%)	LC	HV	MM	Medeiros et al. 2017, Vasconcelos-Filho & Oliveira 1999
ACANTHURIFORMES						
Sciaenidae						
<i>Bairdiella ronchus</i> (Cuvier, 1830)	1 (< 0.01%)	2 (0.03%)	LC	ZB	MM	Medeiros et al. 2017, Vasconcelos-Filho & Oliveira 1999
<i>Cynoscion jamaicensis</i> (Vaillant and Bocourt, 1883)	2 (0.01%)	15 (0.25%)	LC	PV	MM	Rondineli et al. 2007, Ribeiro et al. 2019
<i>Cynoscion leiarchus</i> (Cuvier, 1830)	3 (0.01%)	1 (0.02%)	LC	PV	MM	Ribeiro et al. 2019, Vasconcelos-Filho & Oliveira 1999
<i>Cynoscion</i> sp.	1 (< 0.01%)		-	-	-	
<i>Cynoscion microlepidotus</i> (Cuvier, 1830)		8 (0.13%)	LC	ZB	ES	Araújo et al. 2016, Reis-Filho et al. 2010
<i>Cynoscion virescens</i> (Cuvier, 1830)	10 (0.03%)	27 (0.45%)	LC	ZB	MM	FAO 2016

Table S1. Continuation.

<i>Isopisthus parvipinnis</i> (Cuvier, 1830)*	2181 (7.04%)	311 (5.16%)	LC	PV	MM	Lira et al. 2013a, Silva-Júnior et al. 2015
<i>Larimus breviceps</i> Cuvier, 1830*	987 (3.18%)	677 (11.23%)	LC	ZP	MM	Santos et al. 2022, 2016
<i>Macrodon ancylodon</i> (Bloch and Schneider, 1801)*	102 (0.33%)	12 (0.20%)	LC	ZB	MM	Bezerra-Figueiredo et al. 2014, Mourão et al. 2014
<i>Menticirrhus americanus</i> (Linnaeus, 1758)*	82 (0.26%)	22 (0.36%)	DD	ZB	MM	Lira et al. 2013b, Freitas et al. 2011
<i>Menticirrhus littoralis</i> (Holbrook, 1847)	1 (< 0.01%)	4 (0.07%)	DD	ZB	MM	Braun & Fontoura 2004, Ribeiro et al. 2019
<i>Micropogonias furnieri</i> (Desmarest, 1823)	14 (0.05%)		LC	ZB	MM	Mendoza-Carranza & Vieira 2008, Vasconcelos-Filho & Oliveira 1999
<i>Nebris microps</i> Cuvier, 1830	28 (0.09%)	15 (0.25%)	LC	ZB	ES	Mourão et al. 2014, Xavier et al. 2012
<i>Ophioscion punctatissimus</i> Meek and Hildebrand, 1925*	87 (0.28%)	256 (4.24%)	DD	ZB	MS	Zahorcsak et al. 2000, Passos et al. 2013
<i>Ophioscion</i> sp.	65 (0.21%)					
<i>Ophioscion</i> sp.1	32 (0.1%)		-	-	-	
<i>Ophioscion</i> sp.2	4 (0.01%)		-	-	-	
<i>Ophioscion</i> sp.3	99 (0.32%)		-	-	-	
<i>Paralonchurus brasiliensis</i> (Steindachner, 1875)	16 (0.05%)	29 (0.48%)	LC	ZB	MM	Robert et al. 2007
<i>Pogonias cromis</i> (Linnaeus, 1766)		1 (0.02%)	NE	ZB	MM	Blasina et al. 2010, Ribeiro et al. 2019
<i>Stellifer brasiliensis</i> (Schultz, 1945)*	129 (0.42%)	170 (2.82%)	LC	ZB	MM	Dantas et al. 2015, Vasconcelos-Filho & Oliveira 1999
<i>Stellifer microps</i> (Steindachner, 1864)*	783 (2.53%)		LC	ZB	ES	Barletta & Blaber 2007
<i>Stellifer rastrifer</i> (Jordan, 1889)*	1609 (5.19%)	433 (7.18%)	LC	ZB	MM	Barletta & Blaber 2007, Vasconcelos-Filho & Oliveira 1999
<i>Stellifer</i> sp.	11 (0.04%)	387 (6.42%)	-	-	-	
<i>Stellifer</i> sp.1	166 (0.54%)		-	-	-	
<i>Stellifer stellifer</i> (Bloch, 1790)*	740 (2.39%)	662 (10.98%)	LC	ZB	ES	Dantas et al. 2015, Barletta et al. 2008
<i>Umbrina coroides</i> Cuvier, 1830	2 (0.01%)		LC	ZB	MS	Zahorcsak et al. 2000, Ribeiro et al. 2019
SPARIFORMES						
Lobotidae						

Table S1. Continuation.

<i>Lobotes surinamensis</i> (Bloch, 1790)	1 (< 0.01%)		LC	ZB	MS	Franks et al. 2003, Vasconselos-Filho & Oliveira 1999
TETRAODONTIFORMES						
Ostraciidae						
<i>Acanthostracion quadricornis</i> (Linnaeus, 1758)		1 (0.02%)	LC	ZB	MS	FAO 2016, De Paiva 2013
Tetraodontidae						
<i>Lagocephalus laevigatus</i> (Linnaeus, 1766)	40 (0.13%)		LC	HV	MM	Denadai et al. 2012, De Andrade et al. 2016
<i>Sphoeroides testudineus</i> (Linnaeus, 1758)	27 (0.09%)		DD	ZB	ES	Araújo 2012, Reis-Filho et al. 2010
<i>Sphoeroides greeleyi</i> Gilbert, 1900	2 (0.01%)		LC	ZB	MM	Lima 2014, Schultz et al. 2002
Diodontidae						
<i>Chilomycterus antennatus</i> (Cuvier, 1816)	1 (< 0.01%)		NA	ZB	MS	FAO 2016
<i>Chilomycterus spinosus</i> (Linnaeus, 1758)	1 (< 0.01%)		LC	ZB	MS	Almeida-Silva et al. 2015, Vasconselos-Filho & Oliveira 1999
<i>Diodon holocanthus</i> Linnaeus, 1758	1 (< 0.01%)		LC	ZB	MS	López-Peralta et al. 2002, Vasconselos-Filho & Oliveira 1999

REFERENCES

AKADJE C, KONAN JK, DIABY M, N'DA K & LELOC'H F. 2013. Diet of the barracuda *Sphyraena guachancho* in Côte d'Ivoire (Equatorial Eastern Atlantic Ocean). *Cybium* 37: 285-293.

ALMEIDA RDS. 2018. Distribuição espacial das guildas tróficas de peixes estuarinos no litoral amazônico brasileiro. *UFMA*, 44 p.

ALMEIDA-SILVA PH, TUBINO RA, ZAMBRANO LC, HUNDER DA, GARRITANO SR & MONTEIRO-NETO C. 2015. Trophic ecology and food consumption of fishes in a hypersaline tropical lagoon. *J Fish Biol* 86: 1781-1795.

ANDERSON AB, CARVALHO-FILHO A, MORAIS RA, NUNES LT, QUIMBAYO JP & FLOETER SR. 2015. Brazilian tropical fishes in their southern limit of distribution: checklist of Santa Catarina's rocky reef ichthyofauna, remarks and new records. *Check List* 11: 1688.

ARAÚJO FG, DE AZEVEDO MCC & GUEDES APP. 2016. Inter-decadal changes in fish communities of a tropical bay in southeastern Brazil. *Reg Stud Mar Sci* 3: 107-118.

ARAUJO PRV. 2012. Variação espaço-temporal e ecologia trófica de *Sphoeroides testudineus* (Linnaeus, 1758) e *Colomesus psittacus* (Bloch & Schneider, 1801)

(Actinopterygii, Tetraodontidae) no estuário do rio Mamanguape, Paraíba, Brasil. *UEPB*, 55 p.

BARLETTA M, AMARAL CS, CORRÊA MFM, GUEBERT F, DANTAS DV, LORENZI L & SAINT-PAUL U. 2008. Factors affecting seasonal variations in demersal fish assemblages at an ecocline in a tropical-subtropical estuary. *J Fish Biol* 73: 1314-1336.

BARLETTA M & BLABER S. 2007. Comparison of fish assemblages and guilds in tropical habitats of the Embley (Indo-west Pacific) and caete (Western Atlantic) estuaries. *Bull Mar Sci* 80: 647-680.

BESSA E, FERNANDEZ WS, CRISTINA A, ARCURI D, TURRA A & AEROPORTO J. 2012. Life history of three catfish species (Siluriformes: Ariidae) from southeastern Brazil. *Biota Neotrop* 12: 74-83.

BEZERRA-FIGUEIREDO M, CARVALHO NETA RNF, SILVA-NUNES JL & DA SILVA DE ALMEIDA Z. 2014. Feeding habits of *Macrodon ancylodon* (Actinopterygii, Sciaenidae) in northeast, Brazil. *Rev Biol Mar Oceanogr* 49: 559-566.

BLASINA GE, BARBINI SA & DÍAZ DE ASTARLOA JM. 2010. Trophic ecology of the black drum, *Pogonias cromis* (Sciaenidae), in Mar Chiquita coastal lagoon (Argentina). *J Appl Ichthyol* 26: 528-534.

- BOMFIM AC. 2014. Bioecologia da ictiofauna marinha descartada pelo arrasto camaroeiro em praias da Baía Potiguar, Brasil. UFRN, 115 p.
- BORNATOWSKI H, ABILHOA V & FREITAS MO. 2006. Sobre a alimentação de *Narcine brasiliensis* na baía de Ubatuba-enseada, São Francisco do Sul, Santa Catarina, Brasil. Est Biol 28: 57-60.
- BORTOLUZZI T, ASCHENBRENNER AC, CAMILA R, ROOS DC, LEPKOSKI ED, MARTINS JA, MAYKOL G, QUEROL E & QUEROL MV. 2006. Hábito alimentar da Sardinha Prata, *Lycengraulis grossidens* (Spix & Agassiz, 1829), (Pisces, Engraulidae), Rio Uruguai Médio, Sudoeste do Rio Grande do Sul, Brasil. Biod Pamp 4: 11-23.
- BOWMAN RE, STILLWELL CE, MICHAELS WL & GROSSLEIN MD. 2000. National Oceanic and Atmospheric Administration National Marine Fisheries Service Food of Northwest Atlantic Fishes and Two Common Species of Squid. NOAA TM, 155 p.
- BRAUN AS & FONTOURA NF. 2004. Reproductive biology of *Menticirrhus littoralis* in southern Brazil (Actinopterygii: Perciformes: Sciaenidae). Neotrop Ichthyol 2: 31-36.
- CHAVES PTC & SERENATO A. 1998. Diversidade de dietas na assembléia de linguados (Teleostei, Pleuronectiformes) do manguezal da Baía de Guaratuba, Paraná, Brasil. Rev Bras Oceanogr 46: 61-68.
- CLARK FJK & PESSANHA ALM. 2015. Diet and ontogenetic shift in habitat use by *Rhinosardinia bahiensis* in a tropical semi-arid estuary, north-eastern Brazil. J Mar Biol Assoc U K 95: 175-183.
- CONTENTE RF, STEFANONI MF & SPACH HL. 2009. Growth rates of four Hawaiian deep slope fishes: a comparison of methods for estimating age and growth from otolith microincrement widths, 63-69 p.
- CORRÊA CE, CHAVES PDT & BITTENCOURT PR. 2005. Biology of *Chirocentrodon bleekermanus* (Poey, 1867) (Clupeiformes: Pristigasteridae) in a Continental Shelf Region of Southern Brazil. Braz Arch Biol Technol 48: 419-427.
- DANTAS DV, BARLETTA M & COSTA MF. 2015. Feeding ecology and seasonal diet overlap between *Stellifer brasiliensis* and *Stellifer stellifer* in a tropical estuarine ecocline. J Fish Biol 86: 707-733.
- DAROS FALM. 2014. Estudo da ictiofauna costeira no litoral do Paraná e adjacências através de censos visuais e microquímica de otólitos. UFPR, 106 p.
- DE ANDRADE AC, SANTOS SR, VERANI JR & VIANNA M. 2016. Guild composition and habitat use by Tetraodontiformes (Teleostei, Acanthopterygii) in a south-western Atlantic tropical estuary. J Mar Biol Assoc U K 96: 1251-1264.
- DE ANDRADE-TUBINO MF, RIBEIRO ALR & VIANNA M. 2008. Organização espaço-temporal das ictiocenoses demersais nos ecossistemas estuarinos Brasileiros: Uma Síntese. Oecol Bras 12: 640-661.
- DENADAI MR, BORGES-SANTOS F, BESSA E, SILVA-FERNANDEZ W & TURRA A. 2013. Population Biology and Diet of the Pompano *Trachinotus carolinus* (Perciformes: Carangidae) in Caraguatatuba Bay, Southeastern Brazil. J Mar Biol Oceanogr 2: 2.
- DENADAI MR, SANTOS FB, BESSA E, BERNARDES LP & TURRA A. 2012. Population biology and diet of the puffer fish *Lagocephalus laevigatus* (Tetraodontiformes: Tetraodontidae) in Caraguatatuba Bay, south-eastern Brazil. Mar Biol Assoc U K 2: 407-412.
- DE PAIVA ACG, CHAVES PT & ARAÚJO ME. 2013. Distribution of estuarine fish fauna along coast of Brazil. Trop Oceanogr 2: 1-39.
- DE PAIVA ACG, TARSO P, CHAVES C & ARAÚJO ME. 2008. Estrutura e organização trófica da ictiofauna de águas rasas em um estuário tropical. Rev Bras Zool 25: 647-661.
- ELLIOT M, WHITFIELD AK, POTTER IC, BLABER SJM, CYRUS DP, NORDLIE FG & HARRISON TD. 2007. The guild approach to categorizing estuarine fish assemblages: a global review. Journal Compilation 8: 241-268.
- FAO - FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS. 2016. The living marine resources of the Western Central Atlantic Norfolk. Species identification guide for fishery purposes and American Society of Ichthyologists and Herpetologists special publication, n. 5. Virginia, USA, p. 426-427.
- FAVERO FLT, ARAUJO IMS & SEVERI W. 2019. Structure of the fish assemblage and functional guilds in the estuary of Maracáipe, Northeastern coast of Brazil. Bol Inst Pesca 45: 417.
- FONTELES-FILHO AA. 1988. Sinopse de informações sobre a cavala, *Scomberomorus cavalla*, e a serra, *S. brasiliensis*, no estado do Ceará, Brasil. Arq Cienc Mar 27: 21-48.
- FRANKS JS, VANDERKOOY KE & GARBER NM. 2003. Diet of Tripletail, *Lobotes surinamensis*, from Mississippi Coastal Waters. Gulf and Caribb Res 15: 27-32.
- FREITAS MO, HALUCH CF, ABILHOA V, CORRÊA MFM & HOSTIM-SILVA M. 2011. Estrutura populacional e biologia reprodutiva de *Menticirrhus americanus* (Linnaeus, 1758) (Teleostei, Sciaenidae) na baía de Ubatuba-enseada, Santa Catarina, Brasil. Biotemas 24: 47-59.
- GIANETI MD. 2011. Reprodução, alimentação, idade e crescimento de *Dasyatis guttata* (Bloch & Schneider, 1801) (Elasmobranchii; Dasyatidae) na região de Caiçara

do Norte - RN, 131 p. <https://pdfs.semanticscholar.org/0762/91025c8fe37ac813951f330aa26265a066b1.pdf>.

GIARRIZZO T & SAINT-PAUL U. 2008. Ontogenetic and seasonal shifts in the diet of the pemecou sea catfish *Sciades herzbergii* (Siluriformes: Ariidae), from a macrotidal mangrove creek in the Curuçá estuary, Northern Brazil. *Rev Biol Trop* 56: 861-873.

GONZALEZ JG ET AL. 2021. Trophic ecology of the juveniles of two jack species (*Caranx latus* and *C. hippos*) in contrasted tropical estuaries. *Estuarine, Coast Shelf Sci* 55: 107370.

GONZALEZ JG & VASKE-JÚNIOR T. 2017. Feeding ecology of the beach silverside *Atherinella blackburni* (atherinopsidae) in a tropical sandy beach, Southeastern Brazil. *Braz J Oceanogr* 65: 346-355.

GUEDES APP, ARAÚJO FG, PESSANHA ALM & MILAGRE RR. 2015. Partitioning of the feeding niche along spatial, seasonal and size dimensions by the fish community in a tropical Bay in Southeastern Brazil. *Mar Ecol* 36: 38-56.

HÖFLING JC, FERREIRA LI, DE OLIVEIRA MP, PAIVA-FILHO AM & PRADO A. 2000. Alimentação de peixes da família clupeidae do complexo estuarino lagunar de cananéia, São Paulo, Brasil. *Bioikos* 14: 12-20.

LEÃO GN. 2016. Aspectos da biologia de *Eucinostomus argenteus* Baird e Girard, 1855, gerreidae, capturado no canal de Santa Cruz - Pernambuco. UFRPE, 73 p. <http://www.tede2.ufrpe.br:8080/tede/bitstream/tede2/7087/2/George%20do%20Nascimento%20Leao.pdf>.

LIMA CSDS. 2014. Partição trófica de dois tetraodontidae simpátricos no estuário negativo do rio tubarão, Macau-RN, Brasil. UFPB.

LIMA RRSS ET AL. 2021. Fish bycatch caught by shrimp fisheries in Western Tropical Atlantic. 1 ed, Recife: CEPE, 206 p.

LIRA AS, LUCENA-FRÉDOU F, VIANA AP, EDUARDO LN & FRÉDOU T. 2017. Feeding ecology of *Centropomus undecimalis* (Bloch, 1792) and *Centropomus parallelus* (Poey, 1860) in two tropical estuaries in Northeastern Brazil. *Pan- Am J Aquat Sci* 12: 123-135.

LIRA AS, VIANA AP, EDUARDO LN, LUCENA-FRÉDOU F & FRÉDOU T. 2019. Population structure, size at first sexual maturity, and feeding ecology of *Conodon nobilis* (Actinopterygii: Perciformes: Haemulidae) from the coasts of Pernambuco, north-eastern Brazil. *Acta Ichthyol Piscat* 49: 389-398.

LIRA AS, VIANA AP & LUCENA-FRÉDOU F. 2013a. Determinação do hábito alimentar de *isophisthus parvipinis* (Curvier,

1830) (Actinopterygii: Sciaenidae) na região de Barra de Sirinhaém PE. XVIII CONBEP, 807-811 p.

LIRA AS, VIANA AP, NOLÉ LE, LUCENA-FRÉDOU F & FRÉDOU T. 2013b. Hábito alimentar do *Menticirrhus americanus* (Linnaeus, 1758) (Actinopterygii: Sciaenidae) na região de Barra de Sirinhaém PE. XVIII CONBEP, 801-805 p.

LÓPEZ-PERALTA RH, ARCILA T & ARCILA CT. 2002. Diet Composition of Fish Species from the Southern Continental Shelf of Colombia. *WorldFish Center Quarterly* 25: 23-29.

MAI ACG & VIEIRA JP. 2013. Review and consideration on habitat use, distribution and life history of *Lycengraulis grossidens* (Agassiz, 1829) (Actinopterygii, Clupeiformes, Engraulididae). *Biota Neotrop* 13: 121-130.

MARCENIUK AP, SICCHA-RAMIREZ R, BARTHEM RB & WOSIACKI WB. 2017. Redescription of *Notarius grandicassis* and *Notarius parmocassis* (Siluriformes; Ariidae), with insights into morphological plasticity and evidence of incipient speciation. *Syst Biodivers* 15: 274-289.

MCEACHRAN JD & SÉRET B. 1990. Gymnuridae. In: QUERO ET AL. (Eds). Check-list of the fishes of the eastern tropical Atlantic (CLOFETA), Paris: JNICT, Lisbon; SEI, UNESCO, p. 64-66.

MEDEIROS APM, XAVIER JHA, DA SILVA MB, AIRES-SOUZA L & ROSA IM DE L. 2018. Distribution patterns of the fish assemblage in the Mamanguape River Estuary, North-eastern Brazil. *Mar Biol Res* 14: 524-536.

MEDEIROS APM, XAVIER JHA & ROSA IML. 2017. Diet and trophic organization of the fish assemblage from the Mamanguape River Estuary, Brazil. *Lat Am J Aquat Res* 45: 879-890.

MENDES FL & BARTHEM RB. 2010. Hábitos alimentares de bagres marinhos (siluriformes: ariidae) do estuário amazônico. *Amazonia: Ci & Desenv* 5: 153-166.

MENDOZA-CARRANZA M. 2003. The feeding habits of gafftopsail catfish *Bagre marinus* (Ariidae) in Paraiso Coast, Tabasco, Mexico. *Hidrobiol* 13: 119-126.

MENDOZA-CARRANZA M & VIEIRA J. 2008. Whitemouth croaker *Micropogonias furnieri* (Desmarest, 1823) feeding strategies across four southern Brazilian estuaries. *Aquat Ecol* 42: 83-93.

MIKKOLA H (Ed). 2016. Fisheries and Aquaculture in the Modern World. Croatia INTECH d.o.o, 216 p. <http://dx.doi.org/10.5772/61558>.

MONTEIRO DP, GIARRIZZO T & ISAAC V. 2009. Feeding ecology of juvenile dog snapper *Lutjanus jocu* (Bloch and Shneider, 1801) (Lutjanidae) in intertidal mangrove

- creeks in Curuçá estuary (Northern Brazil). *Braz Arch Biol Technol* 52: 1421-1430.
- MOURÃO KRM, FERREIRA V & LUCENA-FRÉDOU F. 2014. Composition of functional ecological guilds of the fish fauna of the internal sector of the amazon estuary, Pará, Brazil. *An Acad Bras Cienc* 86: 1783-1800.
- MUTO EY, MALFARA DT, COELHO LI & SOARES LSH. 2008. Alimentação das sardinhas *Pellona harroweri* (Fowler, 1919) e *Chirocentrodon bleekermanus* (Poey, 1867), na região costeira de Santos, Estado de São Paulo. *Oceanografia e mudanças globais São Paulo: Inst Oceanogr* 2004: 287-302.
- NELSON JS, GRANDE TC & WILSON MVH. 2016. *Fishes of the World: Fifth Edition*. Canada, 1-707 p.
- NIZINSKI MS & MUNROE TA. 2002. Engraulidae: Anchovies. In: Carpenter KE (Ed), *FAO Species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Atlantic, Rome* 2: 764-794.
- NUNES CRR & ROSA RS. 1998. Composição e distribuição da ictiofauna acompanhante em arrastos de camarão na costa da Paraíba, Brasil. *Trab Oceanogr UFPE* 26: 67-83.
- OLIVEIRA-SILVA TJ & LOPES PRD. 2002. Notas sobre a alimentação e morfologia do aparelho digestivo de *Chloroscombrus chrysurus* (Linnaeus, 1766) (Actinopterygii, Carangidae) na Praia de Ponta da Ilha (Ilha de Itaparica, Bahia). *Rev Bras Zool* 4: 179-192.
- PASSARONE R, APARECIDO KC, EDUARDO LN, LIRA AS, SILVA LVS, JUSTINO AKS, CRAVEIRO C, SILVA ES & LUCENA-FRÉDOU F. 2019. Ecological and conservation aspects of bycatch fishes: An evaluation of shrimp fisheries impacts in Northeastern Brazil. *Braz J Oceanogr* 67: 1-10.
- PASSOS AC, CONTENTE RF, ABBATE-PAULO FV, SPACH HL, VILAR CC, JOYEUX JC, CARTAGENA BFC & FAVARO LF. 2013. Analysis of Fish Assemblages in Sectors Along a Salinity Gradient Based on Species, Families and Functional Groups. *Braz J Oceanogr* 61: 251-264.
- PINHEIRO-SOUSA DB, DA SILVA NK, PIOSKI NM, ROCHA ACG, CARVALHO-NETA RNF & ALMEIDA ZS. 2015. Aspectos alimentares e reprodutivos de *Bagre bagre* (Pisces, Ariidae) em um estuário da ilha de São Luís, Maranhão, Brasil. *Rev Bras Engenh Pesc* 8: 1-12.
- RANDALL JE. 1967. Food Habits of Reef Fishes of the West Indies. *Stud Trop Oceanogr* 5: 665-847.
- REIS-FILHO JA, NUNES JACC & FERREIRA A. 2010. Estuarine ichthyofauna of the Paraguaçu River, Todos os Santos Bay, Bahia, Brazil. *Biota Neotrop* 10: 301-311.
- RIBEIRO GC, CATTANI AP, HOSTIM-SILVA M, CLEZAR L, PASSOS AC, SOETH M, CARDOSO OR & SPACH HL. 2019. Marine ichthyofauna of Santa Catarina Island, Southern Brazil: checklist with comments on the species. *Biota Neotrop* 19: 1-12.
- ROBERT MDC, MICHELS-SOUZA MA & CHAVES PDT. 2007. Biology of *Paralonchurus brasiliensis* (Steindachner) (Teleostei, Sciaenidae) in Paran coast, Brazil. *Rev Bras Zool* 24: 191-198.
- RONDINELI GR, BRAGA FMS, TUTUI SLS & BASTOS GCC. 2007. Dieta de *Menticirrhus americanus* (Linnaeus, 1758) e *Cynoscion jamaicensis* (Vaillant e Bocourt, 1883) (Pisces, Sciaenidae) no sudeste do Brasil, Estado de São Paulo. *Bol Inst Pesca* 33: 221-228.
- ROWLING KR & RAINES LP. 2000. Description of the biology and an assessment of the fishery for silver trevally *Pseudocaranx dentex* off New South Wales. Cronulla, Australia: NSW, Fisheries Research Institute, 70 p.
- RUEDA M & DEFEO O. 2003. Spatial structure of fish assemblages in a tropical estuarine lagoon: Combining multivariate and geostatistical techniques. *J Exp Mar Bio Ecol* 296: 93-112.
- SALES NS. 2015. Influncia da salinidade na distribuio e dieta da ictiofauna em um esturio hipersalino. Campina Grande – PB. UFPB, 129 p.
- SANTANDER-NETO J. 2015. Dinmica populacional da raia *Urotrygon microphthalmum DELSMAN*, 1941 no nordeste do Brasil. UFRPE, 142 p.
- SANTOS LN, BROTTTO DS & ZALMON IR. 2011. Assessing artificial reefs for fisheries management: a 10-year assessment off Northern coast of Rio de Janeiro. CRC Press, 25-140 p.
- SANTOS LV, VASCONCELOS-FILHO JE, LIRA AS, SOARES A, EDUARDO LN, PASSARONE R, LE'-LOC'H F & LUCENA-FREDOU F. 2022. Trophic ecology and ecomorphology of the shorthead drum, *Larimus breviceps* (Acanthuriformes: Sciaenidae), from the northeastern Brazil. *Thalassas* 38: 1-11.
- SANTOS MN, ROCHA GRA & FREIRE KMF. 2016. Diet composition for three sciaenids caught off northeastern Brazil. *Rev Biol Mar Oceanogr* 51: 493-504.
- SAZIMA I. 1998. Field evidence for suspension feeding in *Pseudocaranx dentex*, with comments on ram filtering in other jacks (Carangidae). *Environ Biol Fishes* 53: 225-229.
- SCHLUESSEL V, BENNETT MB & COLLIN SP. 2010. Diet and reproduction in the white-spotted eagle ray *Aetobatus narinari* from Queensland, Australia and the Penghu Islands, Taiwan. *Mar Freshw Res* 61: 1278.

- SCHULTZ YD, FAVARO LF & SPACH HL. 2002. Aspectos reprodutivos de *Sphoeroides greeleyi* (Gilbert), Pisces, Osteichthyes, Tetraodontidae, da gamboa do Bagaçu, Baía de Paranaguá, Paraná, Brasil. Rev Bras Zool 19: 65-76.
- SCHWAMBORN SHL. 2004. Dinâmica e organização trófica de assembléias de peixes associadas aos prados de capim marinho (*Halodule wrightii*) de Itamaracá, Pernambuco. UFPE, 306 p.
- SEGURA-BERTTOLINI EC & MENDOZA-CARRANZA M. 2013. Importance of male gafftopsail catfish, *Bagre marinus* (Pisces: Ariidae), in the reproductive process. Cienc Mar 39: 29-39.
- SHIBUYA ARRS & SOARES MC. 2005. Note on the diet of the guitarfish, *Rhinobatos percellens* (Walbaun, 1792) (ELASMOGRANCHII: RHINOBATIDAE) from the coast of Paraíba, Brazil. Acta Biol Leopold 27: 63-64.
- SILVA JD, BARLETTA M, LIMA AR & FERREIRA GV. 2018. Use of resources and microplastic contamination throughout the life cycle of grunts (Haemulidae) in a tropical estuary. Environ Pollut 242: 1010-1021.
- SILVA-JÚNIOR CAB, VIANA AP, LUCENA-FRÉDOU F & FRÉDOU T. 2015. Aspects of the reproductive biology and characterization of Sciaenidae captured as bycatch in the prawn trawling in the northeastern Brazil. Acta Sci Biol Sci 37: 1.
- SLEY A, JARBOUI O, GHORBEL M & BOUAIN A. 2009. Food and feeding habits of *Caranx crysos* from the Gulf of Gabès (Tunisia). Mar Biol Assoc U K 89: 2009.
- SOARES A, LIRA AS, GONZALEZ JG, EDUARDO LN, LUCENA-FRÉDOU F, LE F, FERREIRA BP & FRÉDOU T. 2020. Feeding habits and population aspects of the spotted goatfish, *Pseudupeneus maculatus* (Perciformes: Mullidae), on the continental shelf of northeast Brazil. Sci Mar 84: 119-131.
- STEFANONI MF. 2008. Ictiofauna e ecologia trófica de peixes em ambientes paraiais da ilha das peças, Complexo estuarino de Paranguá, Paraná. UFPR, 143 p.
- TAGLIAFICO A, RAGO N, RANGEL S & MENDOZA J. 2012. Exploitation and reproduction of the spotted eagle ray (*Aetobatus narinari*) in the Los Frailes Archipelago, Venezuela. Fish Bull 110: 307-316.
- VASCONCELOS-FILHO AL. 1979. Estudo ecológico da região de Itamaracá, Pernambuco. Brasil. IV. Alimentação de Sardinha Bandeira, *Opisthonema oglinum*. Trab Oceanog UFPE 14: 105-116.
- VASCONCELOS-FILHO AL, NEUMANN-LEITÃO S, ESKINAZI-LEÇA E & OLIVEIRA AME. 2010. Hábitos alimentares de peixes consumidores secundários do Canal de Santa Cruz, Pernambuco, Brasil. Trop Oceanogr 38: 59-66.
- VASCONCELOS-FILHO AL & OLIVEIRA AME. 1999. Composição e ecologia da ictiofauna do Canal de Santa Cruz (Itamaracá - PE, Brasil). Trab Oceanog UFPE 27: 101-113.
- XAVIER JHA, CORDEIRO CAMM, TENÓRIO GD, DINIZ AF, PAULO EPN, ROSA RS & ROSA IL. 2012. Fish assemblage of the mamanguape environmental protection area, NE Brazil: Abundance, composition and microhabitat availability along the mangrove-reef gradient. Neotr Ichthyol 10: 109-122.
- YOKOTA L, GOITEIN R, GIANETI MD & LESSA RTP. 2013. Diet and feeding strategy of smooth butterfly ray *Gymnura micrura* in northeastern Brazil. J Appl Ichthyol 29: 1325-1329.
- ZAHORCSAK P, SILVANO RAM & SAZIMA I. 2000. Feeding biology of a guild of benthivorous fishes in a sandy shore on south-eastern Brazilian coast. Rev Bras Biol 60: 511-518.