

REVIEW OPEN ACCESS

Unraveling the Complexity of the N_e/N_c Ratio for Conservation of Large and Widespread Pelagic Fish Species: Current Status and Challenges

Chrystelle Delord^{1,2}  | Sophie Arnaud-Haond²  | Agostino Leone^{2,3,4}  | Jonathan Rolland⁵ | Natacha Nikolic^{5,6,7} 

¹UMR248 MARBEC, Univ. Montpellier, Ifremer, IRD, CNRS, La Réunion, France | ²UMR248 MARBEC, Univ. Montpellier, Ifremer, IRD, CNRS, Sète, France | ³Department of Earth and Marine Sciences (DiSTeM), University of Palermo, Palermo, Italy | ⁴National Biodiversity Future Center, Palermo, Italy | ⁵Centre de Recherche Sur la Biodiversité et l'Environnement (CRBE), Université de Toulouse, CNRS, IRD, Toulouse INP, Université Toulouse 3 – Paul Sabatier (UT3), Toulouse, France | ⁶Université de Pau et des Pays de l'Adour, INRAE, AQUA, ECOBIOP, Sain-Pée-sur-Nivelle, France | ⁷ARBRE – Agence de Recherche Pour la Biodiversité à La Réunion, Saint-Gilles, France

Correspondence: Natacha Nikolic (natachanikolic@hotmail.com)**Received:** 30 March 2024 | **Revised:** 9 September 2024 | **Accepted:** 13 September 2024**Funding:** This work was supported by Institut Français de Recherche pour l'Exploitation de la Mer.**Keywords:** conservation genetics | ecological genetics | population genetics – theoretical

ABSTRACT

Estimating and understanding the ratio between effective population size (N_e) and census population size (N_c) are pivotal in the conservation of large marine pelagic fish species, including bony fish such as tunas and cartilaginous fish such as sharks, given the challenges associated with obtaining accurate estimates of their abundance. The difficulties inherent in capturing and monitoring these species in vast and dynamic marine environments often make direct estimation of their population size challenging. By focusing on N_e , it is conceivable in certain cases to approximate census size once the N_e/N_c ratio is known, although this ratio can vary and does not always increase linearly, as it is influenced by various ecological and evolutionary factors. Thus, this ratio presents challenges and complexities in the context of pelagic species conservation. To delve deeper into these challenges, firstly, we recall the diverse types of effective population sizes, including contemporary and historical sizes, and their implications in conservation biology. Secondly, we outline current knowledge about the influence of life history traits on the N_e/N_c ratio in the light of examples drawn from large and abundant pelagic fish species. Despite efforts to document an increasing number of marine species using recent technologies and statistical methods, establishing general rules to predict N_e/N_c remains elusive, necessitating further research and investment. Finally, we recall statistical challenges in relating N_e and N_c emphasizing the necessity of aligning temporal and spatial scales. This last part discusses the roles of generation and reproductive cycle effective population sizes to predict genetic erosion and guiding management strategies. Collectively, these sections underscore the multifaceted nature of effective population size estimation, crucial for preserving genetic diversity and ensuring the long-term viability of populations. By navigating statistical and theoretical complexities, and addressing methodological challenges, scientists should be able to advance our understanding of the N_e/N_c ratio.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Evolutionary Applications* published by John Wiley & Sons Ltd.

1 | Introduction: Population Abundance and Effective Population Size in Conservation Genetics

As aptly enumerated by Swenson et al. (2024), abundance estimates and trends serve as essential metrics in conservation and resource management, playing key roles in assessing conservation status (Wilson, Kendall, and Possingham 2011), gauging the impacts of threats or recovery efforts (Jennings 2000; Ward-Paige et al. 2012), and determining quotas (such as allowable biological catch and annual catch limits in fisheries) for managed populations of both target and non-target species.

This paper focuses on large marine pelagic fish populations and the many logistical and conceptual challenges that researchers face when estimating their population abundance.

The first set of challenges is logistic: marine pelagic fish are known for their extensive movements across the three dimensions of the ocean. Marine pelagic fish populations may be unevenly distributed due to environmental factors such as water temperature, salinity, reproductive period, seasonal migrations and prey availability, complicating the establishment of representative sampling methods. Abundance estimates are typically based on fisheries data, which are expected to reflect the values of fishery stocks and can be poorly documented. The ocean is an immense and often difficult-to-access environment, making comprehensive surveys of pelagic fish census population size (N_c) costly in terms of time, resources, and logistics. Estimating the effective population size (N_e) of marine pelagic fish populations can be an alternative or a complementary measure for several reasons. First, N_e estimation techniques may be less invasive than traditional fish capture and counting methods and can be estimated based on the modeling of genetic information that allows addressing a larger spatial scale with a lower number of individuals. This can be particularly important for threatened by-catch: for example, scraping the skin of blue sharks with a pole when they are on the line and releasing them, as was done in a recent project (Nikolic et al. 2022). Second, N_e and its temporal fluctuations inform about a population's vulnerability to potential environmental disturbances and demographic trends, even for populations with high abundance such as large marine

pelagic species (e.g., Waples et al. 2018). In fact, both N_e and N_c are pivotal factors in influencing the capacity of populations to withstand extinction risks arising from demographic, environmental, or genetic stochastic events (Palstra and Fraser 2012). These events encompass sporadic recruitment failures, inbreeding depression, or declines in genetic diversity when populations are small (Soulé 1987; Boyce 1992; Frankham et al. 2003 by Palstra and Fraser 2012). Consequently, they are of paramount importance to conservation geneticists (Leroy et al. 2017). N_e is not a direct measure of abundance, but rather a crucial metric in population genetics, reflecting the health and viability of a population. It can also relate to demographic factors affecting abundance. A small N_e reflects a substantial degree of genetic erosion experienced by a population, thus serving as an indicator of its adaptive potential (Figure 1). Between two populations with equal N_c , the population with a smaller N_e is more likely to experience diminished genetic diversity, with a reduction of its adaptive potential compared to the one with a higher N_e (Ellegren and Galtier 2016).

For conservation purposes, it is important to distinguish the different types of population abundance and systematically provide a precise definition. First, various metrics have been employed to estimate the actual population size (N) also called the census population size (N_c). N_c may also be named N_A , N_B , N_T (Frankham 1995), and N_a (Palstra and Fraser 2012). These metrics undercover slightly contrasted quantities, including juveniles and senescent individuals or only breeding adults for example. It is not always straightforward to navigate and provide accurate definitions for each metric, as publications often contain confusion or inexact correspondences. For instance, when examining the papers of Frankham (1995) and then subsequently those of Palstra and Fraser (2012), which are supposed to reiterate Frankham's definition of N_c in terms of N_A , it becomes apparent that the definitions are not consistent. For example, Frankham (1995) uses the notation N_A to define abundance as the number of adults (N_A = breeding + senescent adults), whereas Palstra and Fraser (2012), citing Frankham (1995), define it as the annual census population size which is "The number of reproductively mature individuals in a population that may reproduce and hence contribute to the cohort of individuals




FIGURE 1 | Schematic representation of the total abundance of a population on the left (census size— N_c) and its effective population size (N_e) on the right, with intrapopulation genetic variability depicted by blue and orange colored points in equal relative proportions. N_e reflects the size of an ideal, homogeneous population (on the right) that would experience the same level of genetic drift as the census-size population (on the left). In scenarios of variable individual contributions to reproduction, the N_e of a population is generally lower than its total abundance N_c . Here, genetic variation represented by the orange color is more likely to be eliminated from the population (genetic erosion) as it is less frequent in proportion compared to the genetic variation represented by the blue color.

born in that year” named in that publication N_a instead of N_A . Waples (2024a) recommends using N_c as the number of adults alive at a given time to facilitate comparisons across species and emphasizes the importance of specifying whether juveniles are included or not. By synthesizing theoretical considerations and existing literature, we present a summary of various metrics in Table 1. Similarly, as conservation practitioners often focus on the number of reproducing adults (noted as N_B in Frankham (1995)), it is also important to avoid any confusion with the effective number of breeders N_b , that we will mention in the next section, and which is more akin to N_e . In population genetics, N_b is similar to a short-term N_e , scale of one reproductive cycle (instead of one generation), often used for specific periods such as a single breeding season for a given cohort (Waples and Teel 1990). The choice between the various N_c depends on the organisms being considered (Frankham 1995).

The second main challenge is conceptual and stems from the various definitions of N_e in the literature that have further contributed to confusion (Waples 2022). Initially, N_e is a genetic and evolutionary concept introduced by Sewall Wright in 1931 and 1938 as a theoretical number of individuals effectively contributing to the transmission of genetic diversity to the next generation, within an idealized population. It is defined in reference to the Wright-Fisher idealized population (1922, 1930, 1931), a hypothetical finite population with simplified characteristics (detailed below) where genetic drift is the sole operative factor, and the dynamics of allelic and genotypic frequencies across generations depends solely on the population census size (Wang, Santiago, and Caballero 2016). Hence, the Wright-Fisher model is commonly used to study the distribution of alleles in a population over time in the absence of evolutionary forces other than genetic drift. The idealized population under the Wright-Fisher model should not be confused with that of Hardy-Weinberg, for the latter examines allele frequency equilibrium in infinite populations under otherwise similar idealized conditions as the ones from Wright-Fisher. The Wright-Fisher ideal population is more precisely described as an ideal population where each generation consists of a fixed number of individuals reproducing randomly, and each generation is the result of a random sampling of individuals from the previous generation (non-overlapping generations), where changes in allele frequency are due to chance (genetic drift). So, each generation results from a binomial or multinomial (in the case of more than two alleles) sampling of gene copies from a pool of gametes, to which all individuals contribute equally. In such a population, the number of offspring produced by an individual follows a Poisson distribution. Since the 1960s (e.g., Kimura and Crow 1963), theories have emerged to predict the N_e under various inheritance modes and demographic scenarios. Subsequently, in the 1990s and 2000s (e.g., Schwartz, Tallmon, and Luikart 1999; Beaumont 2003), methodological advances have facilitated the estimation of the historical N_e of natural populations, owing to the rapid progress in molecular techniques and computational tools. Another manuscript dealing with this same special issue (Delord et al. 2024) is dedicated to these families of estimates. Considering these seminal definitions, it becomes evident that N_e cannot be simply equated to the notion of “Effective population size is the number of individuals that contribute offspring to the next generation.” Waples elucidates this concept exceptionally well in his 2022 publication. The number of parents contributing at least

one offspring (N_p) can be smaller or larger than N_e . In practice, the N_p/N_e ratio varies based on factors such as the sex ratio, the mean and variance in offspring number, and the relevance of inbreeding (Waples 2022). At this juncture, N_e can be defined as a concept in evolutionary genetics referring to a number of individuals and we will delve deeper into these distinctions in the forthcoming dedicated section.

In species with very large N_c such as marine fish (Hedrick 2005; Hedgecock and Pudovkin 2011), N_e often significantly lags behind N_c ($N_e \ll N_c$, Frankham 1995). Studies of marine species have reported estimates of N_e/N_c ratios ranging from 10^{-3} to 10^{-8} (Hauser and Carvalho 2008), implying N_e and N_c can, and apparently often, differ by several orders of magnitude.

For management and stock assessment purposes, it is essential to evaluate abundance trends to propose fishing quotas and ensure that ongoing exploitation is sustainable. Estimates based on fishing catch (e.g., catch-per-unit-effort modeling and may concern the number of individuals or the weight (biomass)) and biological (e.g., age-structured models) data are usually very difficult to collect and at best incomplete, hampering reliable N_e estimates. Alternative indirect methods for estimating abundance and demographic trends, such as tools based on the study of genetic variability, such as the N_e and N_e/N_c ratio, are thus increasingly used to facilitate the estimate of N_e . Moreover, the N_e/N_c ratio is an indicator of the extent of genetic variation expected in a population (Hedrick 2005). N_e , N_c and the N_e/N_c ratio thus provide complementary information about the conservation status of a population (Hoban et al. 2020, 2024), yet their estimation also presents significant challenges. Those challenges may arise from specific methodological issues related to chosen estimation methods, comprehensively covered in detail by several reviews (Wang and Caballero 1999; Wang 2005; Wang, Santiago, and Caballero 2016; Waples 2016a) and analytical assessments (e.g., Waples, Larson, and Waples 2016; Gilbert and Whitlock 2015). In addition, the direct comparison of N_e and N_c of a population is not a trivial task and requires considerable caution to be biologically meaningful (Palstra and Fraser 2012). The purpose of this paper is to provide scientists and conservation practitioners with a comprehensive review of the critical issues affecting the assessment of a population’s “abundance” (N_c) and effective population size. We illustrate some of these issues with examples from large-bodied, widely distributed marine species of high commercial or emblematic value with significant conservation concerns.

2 | Different Types of Effective Population Sizes: Spatial, Temporal, and Evolutionary Consequences

In population genetics, the precise definition of N_e depends on the spatial and temporal scale considered, and there are actually several types of N_e , all with specific implications in conservation biology. Firstly, it is crucial to determine the temporal scale for calculating N_e . We typically distinguish contemporary N_e (“short-term” or “contemporary effective population size”) from historical N_e (“long-term” or “historical effective population size”), although these timeframes may partially overlap and form a continuum depending on the methods used (Nadachowska-Brzyska, Konczal, and Babik 2022).

TABLE 1 | Definition and notation of various metrics of census and effective population size used in literature and mentioned in the main text, with example studies.

Metrics	Definitions and notations	Example(s)	Outline of existing estimation methods
Census size N_x	Total census size (N_T): all individuals (juveniles + adults) in the population Number of adults (N_A): number of adult individuals (including both breeders and senescent) in the population Annual census population size (N_a): number of sexually mature, i.e., potential breeder individuals contributing to the cohort of individuals born in that year Number of breeding adults (N_B): number of sexually mature adults including sterile individuals and non-breeding helpers	Frankham (1995) Frankham (1995), Waples (2024a) In Palstra and Fraser (2012), different from N_A in Frankham (1995) Frankham (1995) Ferchaud et al. (2016), Waples et al. (2018)	Direct estimation using classical approaches like counting, capture-mark recapture (CMR), and telemetry Indirect estimation using molecular markers: e.g., close-kin mark recapture (CKMR; Bravington, Skaug, and Anderson 2016), CRE (Creel–Rosenblatt Estimator; Creel and Rosenblatt 2013), the Moment estimator (Hettiarachige and Huggins 2018), and the genetic-based Capture–Mark–Recapture (g-CMR; Müller, Mercker, and Brün 2020). For more details see also Larroque and Balkenhol (2023) Deduction from the N_e/N_c ratio if N_e is known
Effective size N_e	Variance effective size ($N_e V$): reflects contemporary effective size, especially the rate of changes in allele frequencies across generations		Direct estimation using demographic information when available, e.g., Hill (1972) for a diploid and dioecious species: $N_e = \frac{4N_1 * T}{Vk + 2}$ N_1 being the cohort size at birth, T the generation time and V/k the lifetime variance in reproductive success across individuals from the cohort Many other equations were proposed to deal with various reproduction strategies (e.g., Wang, Santiago, and Caballero 2016), populations fluctuating in abundance (e.g., Engen, Lande, and Saether 2005) or structured populations (e.g., Wang and Caballero 1999; Gomez-Uchida et al. 2013; Ryman, Laikre, and Hössjer 2019), some of them also requiring some genetic information Deduction from the N_e/N_c ratio if N_e is known and under certain conditions (e.g., the AgeNe method for a population of constant age-structure and vital rates; Waples, Do, and Choquet 2011) Indirect estimation using molecular markers , e.g., following the variation in allele frequencies between samples collected at several successive time periods (the temporal method, e.g., Jorde and Ryman 1995)
	Inbreeding effective size ($N_e I$): reflects contemporary effective size, especially quantifying the rate of inbreeding (increase in homozygosity of alleles that are identical by descent)	Wright (1931) Crow (1954) is the first distinguishing $N_e I$ (originally defined by Wright) and $N_e V$	Deduction from the N_e/N_c ratio if N_e is known and under certain conditions (e.g., the AgeNe method for a population of constant age-structure and vital rates; Waples, Do, and Choquet 2011) Indirect estimation using molecular markers: e.g., the sibship method (e.g., Wang 2009; Akita 2020a, 2020b)

TABLE 1 (Continued)

Metrics	Definitions and notations	Example(s)	Outline of existing estimation methods
	Linkage-disequilibrium effective size (N_e LD): generally used to estimate contemporary effective size using independent, unlinked molecular markers, but could also reflect effective size from more or less distant past time periods, depending on the physical linkage and the recombination rates between linked loci. When physical linkage among loci is known it can be used to reconstruct N_e trajectories	Waples, Larson, and Waples (2016) Santiago et al. (2020)	Sometimes considered analogous to N_eV or N_eI (Waples 2024a) but whose similarity to N_eV and N_eI actually remains unclear (Wang, Santiago, and Caballero 2016; Ryman, Laikre, and Hössjer 2019) Indirect estimation using molecular markers , as described by Hill (1981) and Waples (2006), with a recent review by Waples (2024b)
	Coalescent effective size (N_e CO): typically reflects long-term effective size and is associated with the overall genetic diversity (θ) of the focal populations	Kingman's coalescent (see also Sjödin et al. 2005 and Wakeley 2020)	The value θ (theta) is a composite parameter representing the product of the effective population size (N_e) and the mutation rate (μ). In population genetics, $\theta = 4 N_e \mu$ is used to describe the genetic diversity and evolutionary history of a population, accounting for both the effective population size and the mutation rate Indirect estimation using molecular markers and the allele frequency spectra of one or more populations (e.g., Gutenkunst et al. 2009; Gattepaille, Günther, and Jakobsson 2016)
	Effective number of breeders (N_b): reflects contemporary effective size per reproductive cycle, as opposed to generational effective population size	Crow and Denniston (1988) (see also Waples and Teel 1990; Waples, Do, and Choquet 2011; Waples and Antao 2014)	Direct estimation using demographic information when available, e.g., N_b can be calculated for each sex (female $-f$, and male $-m$) (Crow and Denniston 1988): $N_b = \frac{kN-2}{k-1+\frac{2}{k}}$ and combined to obtain overall N_b per year (Wright 1938): $\frac{1}{N_b} = \frac{1}{4N_{\text{ef}}} + \frac{1}{4N_{\text{fm}}}$ With V_k the lifetime variance in reproductive success and $\frac{V_k}{k}$ the “index of variability” by Crow and
	Additive variance effective population size (N_{eAV}): reflects contemporary effective size	Hössjer, Laikre, and Ryman (2016)	Morton (1955) (Waples and Antao 2014). N is the number of individuals alive at any given time that have reached age at maturity. N_f and N_m is the number of adult females and males, respectively Deduction from the N_b/N_e ratio if N_e is known and under certain conditions (e.g., the AgeNe method for a population of constant age-structure and vital rates; Waples, Do, and Choquet 2011)
			Analytical approach for diploid organisms

Short-term N_e is relative to the present time or the last few generations of the sampled population. It reflects mostly the influence of the population's life cycle and demographic processes operating on recent time scales. It informs on the future prospects of a population and the risk of genetic erosion it may face in the near future (Wang, Santiago, and Caballero 2016). The most common contemporary N_e estimates used are those based on the loss of genetic diversity, specifically the loss of heterozygosity, through inbreeding (inbreeding N_e) and genetic drift (variance N_e) (Husemann et al. 2016). First distinguished by Crow (1954), they both refer to evolutionary processes occurring in one or a few recent generations (hence they are often referred to as “contemporary N_e ”) (Waples 2022). The variance N_e informs on the extent of genetic changes that can be observed within a given population over ecological time (Jorde and Ryman 1995, 2007). Even a small variance in N_e implies strong allelic frequency variations over time and thus a significant risk of genetic erosion (reduction in adaptive potential) through genetic drift.

Long-term N_e generally focuses on the composite parameter $\theta = 4N_e\mu$, where μ represents the mutation rate (Watterson 1975 by Waples 2022). Long-term N_e is relative to a population's evolutionary trajectory and reflects the combined influence of major evolutionary events, such as a bottleneck, founder effect or divergence events, occurring over a more or less ancient geological period (e.g., Pleistocene) on its current genetic diversity (Wang 2005).

Some methods (or combination of methods) can be used to estimate N_e at the present time, but also from various time periods in the past, hence capturing N_e trajectories through time (e.g., Santiago et al. 2020). In a conservation context, contemporary N_e is intuitively more meaningful, as it provides an expectation for future drift, rather than explaining how diversity got lost in the past. However, it is useful to compare it to an estimate of historical effective population size to replace the current health status of a population or stock in its evolutionary trajectory, and detect, for example, recent disturbances corresponding to increased anthropogenic pressures. Here, we can introduce the so-called “coalescence effective population size” (often denoted N_{eCO}), a long-term effective population size related to the average number of generations for a genealogy of genes to converge to its most recent common ancestor.

For contemporary N_e , we can introduce here the necessary distinction between the effective population size per generation or generational N_e and the effective population size per reproductive cycle (i.e., the effective number of breeders N_b). They differ notably in the case of populations with overlapping generations (in contrast to an idealized theoretical population). Generational N_e informs on past and recent evolutionary processes influencing a population's genetic diversity and its future. N_b , on the other hand, informs on the theoretical number of individuals contributing to the transmission of genetic information at a specific time step for a given cohort of individuals. It can thus be estimated for each reproductive cycle if information on several successive cohorts is available. N_b may be easier to estimate than generational effective population

size and allows for understanding specific processes occurring at the scale of the reproductive cycle, such as sexual selection or density-dependence (Waples and Antao 2014). Genetically based parentage methods are routinely used and are much easier to apply to a single season than to an entire adult lifetime (Waples and Antao 2014). The N_b parameter thus always represents a contemporary effective population size and can be estimated using the standard discrete-generation formula for inbreeding effective size with separate sexes (Crow and Denniston 1988) in the software AGENE (Waples, Do, and Choquet 2011; Waples and Antao 2014).

Contemporary N_e is typically linked to rates of change in allele frequency change or increase in inbreeding, while long-term N_e is associated with nucleotide diversity (Waples 2022). In practical applications, it is probably more beneficial to define N_e using demographic parameters such as the number of potential parents (N), and the mean (μ_k) and variance (σ_k^2) in offspring number per parent (k) (Waples 2022):

$$\text{Inbreeding } N_e = \frac{\mu_k N - 1}{\mu_k - 1 + \frac{\sigma_k^2}{\mu_k}} \quad (1)$$

$$\text{Variance } N_e = \frac{\mu_k (2N - 1)}{2 \left[1 + \frac{\sigma_k^2}{\mu_k} \right]} \quad (2)$$

The “variance effective population size” (often denoted N_{eV}) is generally contrasted with the “inbreeding effective population size” (often denoted N_{eI}) (Table 1). While the former reflects the effects of genetic drift on the degree of allelic frequency variation over time, the latter reflects the increase in homozygosity that can occur in a population when the occurrence of mating between related individuals increases, following a decline in its abundance, for example, N_{eV} and N_{eI} may thus be identical or significantly differ within the same population. These values indeed reflect slightly different time periods, with N_{eV} describing an effective population size at the present time while N_{eI} describes the effective population size of the parental generation (Trask et al. 2017). Similarly, these values may differ if allelic frequency variations occur at a faster rate than mating between related individuals and the increase in the rate of homozygosity within the studied population.

Moreover, N_{eV} and N_{eI} will often differ in the case of structured populations (Wang and Caballero 1999), that is in any scenario other than the ideal case where all subpopulations are of equal size and exchange equal, constant, and symmetrical gene flow. In this last case, once migration-drift equilibrium is reached, the N_{eV} and N_{eI} magnitudes at the overall metapopulation scale (Meta- N_e) are equivalent and converge towards a unique value (the so-called “eigenvalue” effective population size, often denoted N_{eig} , Ewens 1982; Wang and Caballero 1999; Ryman, Laikre, and Hössjer 2019). Furthermore, the sum of local N_{eV} yields this Meta- N_e value, which then gradually converges towards N_{eig} . At the local scale of each subpopulation, N_{eV} values can remain stable over time while N_{eI} values gradually approach the global Meta- N_e value (Ryman, Laikre, and Hössjer 2019).

In nature, the classical scenario is that of subpopulations differing in abundance and connected by variable or asymmetric gene flow. The situation is thus most often more complex, necessitating not only a clear distinction between the relative scope of N_{ev} and N_{el} but also between effective population size measures taken at the local (subpopulation) or global (Meta- N_e) scale, the latter no longer being calculable as a simple sum of local N_{ev} values (e.g., Gomez-Uchida et al. 2013). Wang and Caballero (1999) and Hössjer et al. (2015) developed mathematical models predicting the local effective population sizes of each subpopulation in different situations (e.g., monoicous or dioecious populations, haploid or diploid, with equal or variable local abundances). The estimate of local effective population sizes generally involves demographic parameters such as local variance in reproductive success but also genetic parameters indicating the presence of gene flow between subpopulations, such as the fixation index F_{ST} (higher when gene flow between subpopulations is of low intensity). Hössjer, Laikre, and Ryman (2016) optimize some of these models for the specific case of diploid species and compare N_{ev} and N_{el} magnitudes at the local and global scales (Meta- N_e).

Engen, Lande, and Saether (2005) propose a model predicting demographic effective population size (thus N_{ev}) from the N_e/N_c ratio in the case of fluctuating population sizes over time (expansion or decline). This model relies on Fisher's reproductive value concept (Fisher 1958) and on the assumed evolution of the frequency of a rare allele as a consequence of changes in age structure over time. Trask et al. (2017) apply this method to a population of Red-billed Choughs, also integrating two N_{el} effective population size estimation methods to compare them with Engen, Lande, and Saether's (2005) N_{ev} estimator. The latter can be calculated over a single generation or in the long term.

These different mathematical models that allow for direct estimation of effective population sizes are complex and rarely used on empirical datasets (but see Laikre et al. 2016; Trask et al. 2017), unlike indirect genetic estimation methods relying on linkage disequilibrium or allele frequency spectra. Nevertheless, they can be an interesting resource for the simulation of demographic scenarios and genomic data when the software eventually used to analyze or model data requires an approximate knowledge of the species. For example, when simulating a population structured by age, it is the N_c and demographic parameters (e.g., survival and fecundity rates) of each age class that can be set by the user (e.g., CKMRpop for forward-in-time simulation and tabulation of pairwise relationships in age-structured populations (Anderson and Dunham 2005; Anderson 2023); PySLiM mentions the concept of N_e as an emergent property of an individual-based simulation (see the link https://tskit.dev/pyslim/docs/latest/time_units.html?highlight=effective+size)). N_e can represent what can be called an "emergent property" of the simulated model. Its estimation will result from the demographic parameters set during simulation, which will notably influence the variance in reproductive success between simulated individuals, as well as from more stochastic variations related to how the chosen parameters are applied. For example, the method used to model genetic drift or the way random events are simulated can introduce variability in the N_e estimates. In such a case, it can be useful to use existing mathematical models first to (i) predict the theoretical N_e expected under a given simulation

scenario deemed as realistic for the organism targeted, and then (ii) to estimate the N_e from data, to check their match. Moreover, if one wishes to subsequently test the performance of indirect estimation methods of N_e based on genetic data, it may be useful to estimate several theoretical and observed N_e values (e.g., values of local and global N_e , short- or long-term) to identify the scenario (and associated parameters) that best optimizes the convergence between theoretical or data-based values.

From a more general perspective, it is essential to understand the different types of effective population sizes and the implications of their choice in theoretical and empirical contexts. An example provided by Hössjer, Laikre, and Ryman (2016) and taken up by Ryman, Laikre, and Hössjer (2019) is based on the concept of minimum viable population sizes (MVP, Shaffer 1981; Rai 2003; Reed et al. 2003) and specifically concerns the "50/500" rule also called Franklin-Soulé number (Soulé 1980; Franklin 1980). This rule states a homogeneous and isolated population must have a minimum N_e of 50 in the short term (contemporary time scale, 1 to few generations) and a minimum of 500 in the long term (historical time scale) in order to maintain a stable population size through time. According to this rule, the smallest value best fits an inbreeding effective population size, N_{ei} , and the largest an effective population size reflecting the loss of additive genetic variation (defined by Hössjer, Laikre, and Ryman 2016) denoted N_{eAV} (additive variance effective population size). The latter reflects the rate of loss of genetic variation due to the expression of phenotypic traits and is therefore directly related to the adaptive potential of an individual. Ryman, Laikre, and Hössjer (2019) show that different types of effective population sizes can vary greatly from each other within structured populations. They caution that some genetic tools available for estimating effective population size do not necessarily reflect N_{ei} or N_{eAV} and consequently cannot be used to verify the adequacy of an empirical population to the "50/500" rule. Moreover, the Franklin-Soulé number was based on livestock populations that have been purged by humans, and may tolerate higher levels of inbreeding compared to wild species (Lande 1995, see also Clarke et al. 2024). Considering that only approximately 10% of the spontaneous mutational variance is quasi-neutral (López and López-Fanjul 1993a, 1993b), or for maintaining variability over 10 generations, it has been proposed that the Franklin-Soulé number for wild species would need to be increased by a factor of 10, resulting in $N_e = 5000$ (Lande 1995).

3 | Qualitative Prediction of the N_e/N_c Ratio in Wild Populations: Influence of Life History and Demography

Understanding the influence of life history strategies and demography on N_e and N_e/N_c is crucial for elucidating how these factors shape population resilience and the maintenance of genetic diversity. Variabilities in life history traits and demographic characteristics can significantly impact N_e and N_e/N_c , thereby influencing the long-term viability of populations and their ability to adapt to environmental changes. Consequently, investigating this influence is essential for effective and sustainable management of wild populations. Significant work has been invested to identify general rules allowing estimating the N_e/N_c ratio based on a population biology and life history

traits (e.g., Hauser and Carvalho 2008; Waples et al. 2013; Wang, Santiago, and Caballero 2016; Waples 2024a). This, however, remains complex for wild populations (Palstra and Fraser 2012). On one hand, the reproductive strategies and life history traits of species influence their N_e by acting on the variance in reproductive success among individuals. For example, an imbalanced sex ratio can decrease the N_e , as can any factor that generates non-random mating among individuals (Nunney 1991, 1993, 1996), such as overlapping generations within age-structured populations. Species with very high fecundity and a large number of offspring (albeit with a high mortality rate at young life stages), such as “r-strategy” species—common among marine invertebrates—, often exhibit higher N_e and genetic diversity (Romiguier et al. 2014; Ellegren and Galtier 2016) than species with lower fecundity, producing fewer offspring (but more resistant, with lower mortality at young life stages), such as “K-strategy” species. However, beyond these species’ “typologies,” more demographic parameters will come into play. In iteroparous species (i.e., those that reproduce multiple times during their life such as large marine pelagic fish), N_e is influenced by life history traits such as longevity or age at maturity, which affect the variance in reproductive success (Lee, Engen, and Sæther 2011). Some of these life history traits may also influence the N_e/N_c ratio among species. Waples et al. (2013) show that three demographic parameters influence this ratio predictably: (1) adult lifespan, (2) age at maturity, and (3) and the coefficient of variation in age-specific fecundity. Indeed, a long adult longevity strengthens the influence of overlapping generations, reducing N_e . Furthermore, a long-lived species will have more numerous but also more spread out reproductive opportunities, with the risk that adult mortality between reproduction events period increases the variance in reproductive success among individuals and causes a decrease in N_e . This phenomenon is even more pronounced if mortality rates are higher in younger and intermediate age classes than in older age classes (i.e., species with type III survival curves, Pinder, Wiener, and Smith 1978), or if fecundity increases with age. In this case, the greatest contribution to reproduction will essentially be ensured by the small proportion of old adult individuals that have survived to the most fecund age classes, again leading to increased overall variability in reproductive success among individuals and a decrease in N_e . Fecundity parameters (including variations in fecundity among individuals of the same age, due to more or less favorable reproductive conditions, for example) as well as survival parameters, such as the annual adult mortality rate, thus have a significant influence on N_e and its relationship to total adult abundance (Waples 2016b). Barry, Broquet, and Gagnaire (2022) corroborate this finding through simulations, showing a pronounced decrease in N_e/N_c as adult lifespan increases. This trend is particularly acute under type III survivorship curves when fecundity increases with age, a common trait among marine fishes. The authors also reveal an inversely proportional relationship between adult longevity (i.e., reduction in the expected N_e/N_c ratio) and the observed genetic diversity in the genome of 16 species of marine fish in the Mediterranean Sea. The study of Barry, Broquet, and Gagnaire (2022) underlines the role of vital rates as critical drivers of N_e and genetic diversity levels within marine species in natural environments. In contrast, the rates of evolutionary processes are contingent upon N_e (Waples 2022).

Hill (1972) and Crow and Kimura (1970) mathematically summarize the relationship between N_e and the overall variance in reproductive success among individuals of the same cohort using the following equation, for a diploid species with separate sexes:

$$N_e = \frac{4N_1 * T}{V_k + 2} \quad (3)$$

where N_1 represents the total population size (or a cohort at birth), assumed constant over time, T represents generation time expressed as the average age of parents of a cohort at birth, and V_k represents the overall variance in reproductive success among individuals born in the same cohort.

Based on these theoretical and empirical findings, we could attempt to qualitatively predict the N_e/N_c ratio of a species of interest if (i) we can accurately know its demographic parameters or assign it to a taxon or life history trait typology for which robust and consistent estimates are already available, and if (ii) we know the order of magnitude of one of the two values N_e or N_c in order to deduce the second. For example, Waples, Do, and Choquet (2011) propose a method, implemented in the software AgeNe, allowing under certain assumptions the theoretical calculation of N_e and the N_e/N_c ratio using demographic parameter values of a population. In practice, however, for many species, it is rare to know this information accurately. Furthermore, life history traits are not the only parameters likely to influence, in reality, N_e and the N_e/N_c ratio (Waples 2024a).

Another complexity that remains unresolved but which will influence this ratio, as it has been demonstrated to impact N_e , pertains to organism size. Recent research (Lynch et al. 2023) has revealed that N_e decreases with increasing organism size. This phenomenon may be correlated with the inverse scaling of mutation rates with genome size (Lynch et al. 2016) and the inverse relationship between recombination rates and genome size (Lynch et al. 2011). However, the relationship between body size and genome size is complex (e.g., Glazier 2021) and requires further investigation to understand the underlying mechanisms. These points introduce an additional layer of complexity to the determination of N_e values, which are crucial for accurately estimating a parameter that aids in deriving N_c . Specifically, lower mutation rates in larger organisms could mean fewer genetic markers are available for estimating N_e , reducing the precision of such estimates. Furthermore, decreased recombination rates could lead to larger blocks of linked genes, which can bias estimates of genetic diversity and, consequently N_e . With lower recombination rates, linkage disequilibrium is higher because alleles within these larger blocks of linked genes are inherited together more often than by chance.

In marine species, for which very low N_e/N_c ratios have been reported by numerous empirical studies (Hauser and Carvalho 2008), different life history strategies and typologies can be observed. Elasmobranchs, which include species of rays and sharks such as the blue shark, *Prionace glauca*, exhibit highly diversified reproductive strategies and migration behaviors that can diversely influence on N_e (Domingues, Hilsdorf, and Gadig 2018). The correlation between N_e and N_c in the sandbar

shark, *Carcharhinus plumbeus*, the leopard shark, *Stegostoma fasciatum*, and the great white shark, *Carcharodon carcharias*, has suggested ratios close to 1 ($N_e \sim N_c$) (Andreotti et al. 2016; Dudgeon and Ovenden 2015; Portnoy et al. 2009). These long-lived species are characterized by a late maturity, a low fecundity remaining stable between age classes, and a low variance in reproductive success (Ovenden et al. 2016). However, other studies on species with similar life histories such as the blue shark (King et al. 2015) report lower N_e/N_c values. Furthermore, the study of genetic diversity and N_e in elasmobranchs remains underexplored to date, with several scientific works highlighting the need for its better integration within conservation programs (Ovenden et al. 2016; Domingues, Hilsdorf, and Gadig 2018).

Species of large pelagic bony fishes such as tuna, swordfish, and sailfish, are likely to exhibit strong variations in reproductive success due to their high abundance and fecundity, which often increases with age. Furthermore, bony fishes often follow a type III survival curve with a high birth rate and high juvenile mortality, meaning that mortality rates are higher in young (and intermediate) age classes than in older age classes, with many individuals dying after few reproductive opportunities. Hedgcock (1994) proposed that exceedingly low N_e/N_c ratios could result from a combination of prolific reproduction and elevated mortality among juveniles, leading to a significant variance in reproductive success substantially depressing N_e (Palstra and Ruzzante 2008). This concept has subsequently been completed and confirmed (Nunney 1996; Waples 2002; Hedrick 2005; Palstra and Ruzzante 2008). Thus, often in these species of type III, a large proportion of reproduction is ensured by a relatively small number of older and very fecund individuals. The sweepstakes reproductive success hypothesis (Hedgcock and Pudovkin 2011) also describes this phenomenon as one of the biological factors that may explain the very low N_e/N_c ratios reported in these species (e.g., Laconcha et al. (2015) for albacore tuna, Laurent and Planes (2007) for sardines). Furthermore, sex-ratio imbalances and temporal fluctuations in total abundance (linked, for example, to fishing exploitation) are also likely to influence the N_e and the N_e/N_c ratio (Pinsky and Palumbi 2014). Nevertheless, the commonly high total abundance for such species theoretically implies N_e ranging from a few hundred to several thousand (or tens of thousands) individuals (Pinsky and Palumbi 2014). Thus, despite supposed low N_e/N_c ratios, N_e itself can potentially be high, and its estimation can therefore be technically complex, due to challenges such as the need for large sample sizes (especially for highly abundant species) and potential bias from age structure. Waples (2016b) estimates that only extreme cases of sweepstakes reproductive success in marine teleosts could generate N_e/N_c values actually lower than 10^{-2} , a threshold well above the average of 10^{-3} to 10^{-8} reported above (see also Clarke et al. 2024). Moreover, many applications reporting low N_e and N_e/N_c ratios would more likely reflect, in reality, the chosen method for N_e estimation than a true biological reality (Hare et al. 2011; Bierne, Bonhomme, and Arnaud-Haond 2016; Waples 2016a; Irion et al. 2017). In southern bluefin tuna (*Thunnus maccoyii*), Waples et al. (2018) report a N_e/N_c ratio on the order of 10^{-1} , arguing that teleost life history traits do not systematically generate ratios as low as expected, despite the variability in reproductive success among individuals. Demographic parameters of teleosts may also vary between species. Thus, it cannot be excluded that even two species sharing

relatively similar strategies could exhibit contrasted N_e/N_c ratios if some of their demographic parameters differ (e.g., age at maturity, longevity, fecundity), as is the case with tunas (Juan-Jordá et al. 2013; Murua et al. 2017).

It is therefore not easy to predict the order of magnitude of N_e and the N_e/N_c ratio in large pelagic species based solely on broad categories of life history. In fact, although many estimates of the N_e/N_c ratio between species were reported using both theoretical approaches (Waples et al. 2013) and reviews of empirical case studies (e.g., Frankham 1995; Palstra and Ruzzante 2008; Buffalo 2021), it remains difficult to derive general rules to date (Palstra and Fraser 2012). The respective estimation and linking of these two quantities therefore require greater investments and the development of dedicated scientific programs. Detailed knowledge of the demographic parameters of each stock could provide more insightful predictions, but such data are often lacking. Moreover, beyond the inherent biological characteristics of species, the influence of other demographic factors, such as migration patterns, reproductive behavior, and environmental variability, also warrants consideration. For example, fluctuations in total population abundance (resulting, for example, from fishing exploitation, temporal fluctuation in recruitment, and environmental variability) or the geographical scale considered (in the context of populations structured into several subgroups of various connectivity patterns according to a metapopulation model) influence both N_e , and its relationship with N_c . The N_e/N_c ratio can also prove to be unstable over time depending on these factors (e.g., Baker et al. 2016). For example, fishing pressure exerted on a population can decrease N_e more rapidly than its N_e , causing a temporary increase in the N_e/N_c ratio (Allendorf et al. 2008). Repeated estimates over time are therefore necessary to better understand the potential vulnerability of a population or stock (Palstra and Fraser 2012). N_e and its relationship to N_c may also rely on the magnitude of N_c itself and some associated evolutionary parameters, such as recombination rates and linked selection, although consequent additional research should be carried on to better document these relationships (Ellegren and Galtier 2016; Buffalo 2021). Many other evolutionary factors like mutation, migration, and selection, as well as variations in recombination rate and linkage, can influence genetic variation (Waples 2022) and by consequence N_e and N_c (Waples 2024a). In populations with high abundance, the influence of selection can indeed outweigh that of genetic drift while in small population the random fluctuations in allele frequencies (genetic drift) mainly drives the fate of new mutations (Crow 1993; Lynch and Lande 1998).

In a general manner, documenting the N_e/N_c ratio in species of conservation interest is recognized as an essential path of future research in conservation genetics. To better understand the influence of life history traits on the N_e/N_c ratio, it is recommended to multiply estimations of both indices for various species and taxonomic groups (Hoban et al. 2024). Moreover, most methods for assessing population abundance (whether directly or indirectly using molecular markers) are relatively robust to low dispersal levels. In this respect, high cautiousness about statistical considerations and a good understanding of the spatial and temporal scope of N_e and N_c estimates will be essential and constitute the basis of the next sections of this summary.

4 | Challenges in Relating N_e and N_c

In this paper, we refrain from delving into specific statistical challenges and potential artifacts associated to each of the multiple methods to estimate N_e and/or N_c . Those are widely available from a number of excellent reviews (e.g., Hauser and Carvalho 2008; Bravington, Skaug, and Anderson 2016; Wang 2016; Waples 2016a, 2024a, 2024b; Waples, Larson, and Waples 2016; Waples, Waples, and Ward 2022) and also constitute the basis of a detailed review in the present issue (Delord et al. 2024).

Beyond the biological challenges summarized above, exploring the intricate relationship between N_e and N_c is also associated to significant statistical challenges, often revealing complexities that extend beyond initial assessments (Palstra and Fraser 2012). It requires sometimes treating differently the generations or cohorts. Some indirect estimation methods (i.e., based on the study of genetic diversity rather than demographic parameters (Wang 2016; Wang, Santiago, and Caballero 2016)) of N_e provide information integrating several generations, while estimates of N_c are generally one shot and applied to the current time. Palstra and Fraser (2012) and Waples (2024a) provide an overview of good practices when aligning N_e and N_c across recent temporal scales or generations. Waples (2024a) indicates that “short-term, single-generation estimates of N_e/N_c are the most meaningful and the best predictors of what to expect in the near future.” Waples et al. (2018) provide an example of cautious N_e/N_c estimation on the Southern Bluefin tuna, and Waples (2005) an example of the effort to match contemporary N_e to the appropriate time periods when computing the N_e/N_c ratio.

Relating N_e and N_c also implies being able to quantify correctly the degree of uncertainty around each of them, and the overall uncertainty associated with their ratio (Waples et al. 2018). However, the question of the best practices for constructing confidence intervals around estimates of N_e and N_c represents a field of research in itself (e.g., Hamilton, Tartakovsky, and Battocletti 2018; Waples, Larson, and Waples 2016; Jones, Ovenden, and Wang 2016). Finally, as mentioned in the Introduction, it is essential to clearly define and systematically report which N_c one aims to estimate. This quantity can indeed reflect the total population size, including immature individuals, or only adult and potentially reproductive individuals. The latter is recommended by Palstra and Fraser (2012) and by Waples (2024a), since it is the one that most directly influences N_e .

Thus, although many estimates of the N_e/N_c ratio between species are reported using both theoretical approaches (Waples et al. 2013) and reviews of empirical case studies (e.g., Frankham 1995; Palstra and Ruzzante 2008), deriving general rules for calculating the N_e/N_c ratio remains challenging (Palstra and Fraser 2012) due to the complexity and variability inherent in the biological factors influencing N_e and N_c in different species and contexts. The respective estimation of these two quantities therefore often requires greater investments and the development of dedicated scientific programs.

To directly access N_c (in terms of number of breeders), alternative methods (Table 1) have emerged in recent years such

as Close-kin mark-recapture (CKMR) (Bravington, Skaug, and Anderson 2016). This method exemplifies the requirements and challenges surrounding the development of N_c using genetic data. CKMR is a method based on the same principles as Capture-mark-recapture (CMR), where individuals are “marked” by their genotypes and considered “recaptured.” The number of parent–offspring (POP) relationships detected within a dataset, combined with the mathematical formulation of a demographic model based on vital population parameters (survival, fecundity, age at maturity, etc.), allows for the estimation of the most plausible total abundance (typically expressed as the number of breeding adult males and/or females). The CKMR model can generate robust estimates of population abundance when key assumptions such as annual reproduction and population size stability are met (Swenson et al. 2024). However, in cases of significant population declines or non-annual reproductive dynamics, a more complex CKMR model must be constructed to avoid biased parameter estimates (Swenson et al. 2024). Despite the need for further improvements to address large marine pelagic fish, the recent CKMR literature demonstrates both the willingness and the necessity to develop tools for assessing the abundance trends and CKMR emerges as a promising tool for integration into long-term monitoring programs (Swenson et al. 2024). Specific life-history traits, such as semelparity and parthenogenesis, render CKMR methods inapplicable (Bravington, Grewe, and Davies 2016). The CKMR approach can provide crucial parameters for stock assessment (Rodríguez-Rodríguez et al. 2022) and reduce uncertainty in these evaluations. For example, this method has been used to estimate the number of adult individuals of Southern bluefin tuna and to improve the definition of spawning stock biomass (Bravington, Grewe, and Davies 2014). Although we chose to detail CKMR in this paragraph due to its clear illustration of the principles and challenges of genetic N_c estimation, other methods such as CRE, the Moment estimator, and g-CMR (Table 1) also offer complementary approaches for estimating population abundance. CMR methods based on POPs demonstrated significant potential for informing research and management strategies in species conservation, but their results must be interpreted with caution particularly considering the critical role of spatial and temporal sampling intensity (Larroque and Balkenhol 2023). Davies, Bravington, and Thomson (2017) demonstrated that CKMR population estimates for Atlantic bluefin tuna (*Thunnus thynnus*) can exhibit significant bias if the spatial structure is not explicitly considered. Another study revealed, through simulations, a positive relationship between population density and the sampling effort required for unbiased estimates (Rosenblatt et al. 2023). Finally, the design of a sampling strategy hinges on understanding the life history (demographic data) and behavior (spatial data) of the species studied (Rosenblatt et al. 2023).

5 | Conclusion

In view of the challenges, even the impossibility, of determining individual numbers as is the case for large pelagic fish species, and the necessity to strive for minimally invasive technologies, molecular-based metrics offer undeniable advantages. Recent advances and increasing interest aim to address the challenges surrounding tools using molecular markers. This summary highlights the critical importance of deepening our

comprehension of N_e dynamics, particularly within the context of large pelagic fish species where the estimation of N_e and N_c is challenging and from which the estimated N_e/N_c ratio is submitted to large variation within and among taxonomic groups.

Estimating the N_e/N_c ratio is crucial for several reasons. Firstly, understanding N_e/N_c aids in assessing the long-term viability of species, especially in the face of environmental changes and anthropogenic threats. By doing so, it also provides valuable insights to inform conservation strategies aimed at preserving biodiversity, as accurate estimates of N_e/N_c are essential for implementing effective management practices, such as stock management and habitat restoration, to ensure the persistence of wild populations.

Despite its importance, accurately estimating the N_e/N_c ratio remains a challenge due to various technical and conceptual reasons. Technical challenges include the difficulty of gathering sufficient samples in large populations distributed across vast habitats, statistical limitations in estimating N_e and N_c , particularly in species with complex life histories and spatial dynamics but also the characteristics of genomes. Moreover, uncertainties in demographic parameters and the inherent stochasticity of genetic processes further complicate accurate estimation. Fundamentally, the failure to estimate N_e/N_c correctly stems from gaps in our understanding of the underlying biological mechanisms driving population dynamics, including factors such as reproductive strategies, migration patterns, and evolutionary processes.

To address these challenges, concerted efforts are needed to advance both theoretical frameworks and empirical methodologies for estimating N_e and N_c . This requires interdisciplinary collaboration among geneticists, ecologists, statisticians, and conservation biologists to develop innovative approaches that account for the complexities of natural populations. Additionally, investing in long-term monitoring programs and data collection efforts is crucial for improving the accuracy of N_e/N_c estimates across diverse taxonomic groups and ecosystems. Furthermore, integrating advances in genomic technologies and computational methods can enhance our ability to capture the intricacies of population dynamics and refine estimates of the N_e/N_c ratio.

In conclusion, by acknowledging the importance of estimating N_e/N_c , recognizing the challenges involved with long standing and new alternative methods, and proposing strategies for improvement, we can pave the road for more robust and reliable assessments of population health and genetic diversity. POPs-based methods are promising alternative methods, but they are challenging as they necessitate extensive and prolonged sampling efforts. By addressing all the challenges head-on, we can better inform conservation decisions and safeguard the future of wild populations and ecosystems.

Acknowledgments

This article is part of an editorial work within the POPSIZE project funded by the FEAMP—European Union. AL acknowledges the support of the NBFC to the University of Palermo funded by the PNRR, National Biodiversity Future Center—NBFC, Project ID CN00000033.

J.R. received funding from the Agence Nationale de la Recherche (CEBA: ANR-10-LABX-25-01; TULIP: ANR-10-LABX-0041, JCJC: ANR-23-CE02-0005-01). We are strongly grateful to Pierre-Alexandre Gagnaire (ISEM) for his constructive comments on the first draft of this publication. We would like to express our sincere gratitude to Special Issue Editor Joachim Mergeay for his thoughtful recommendations and constructive comments, which have significantly enhanced the quality and clarity of this article. We also thank the anonymous reviewers for their valuable feedback and suggestions, which have contributed to the improvement of our work.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The authors have nothing to report.

References

- Akita, T. 2020a. “Nearly Unbiased Estimator of Contemporary n_e Based on Kinship Relationships.” *Ecology and Evolution* 10: 10343–10352. <https://doi.org/10.1002/ece3.6421>.
- Akita, T. 2020b. “Nearly Unbiased Estimator of Contemporary Effective Mother Size Using Within-Cohort Maternal Sibling Pairs Incorporating Parental and Nonparental Reproductive Variations.” *Heredity* 124: 299–312. <https://doi.org/10.1038/s41437-019-0271-6>.
- Allendorf, F. W., P. R. England, G. Luikart, P. A. Ritchie, and N. Ryman. 2008. “Genetic Effects of Harvest on Wild Animal Populations.” *Trends in Ecology and Evolution* 23, no. 6: 327–337. <https://doi.org/10.1016/j.tree.2008.02.008>.
- Anderson, E. C. 2023. “CKMRpop: Forward-In-Time Simulation and Tallying of Pairwise Relationships.” R Package Version 0.1.3.9999.
- Anderson, E. C., and K. K. Dunham. 2005. “Spip 1.0: A Program for Simulating Pedigrees and Genetic Data in Age-Structured Populations.” *Molecular Ecology Notes* 5, no. 2: 459–461.
- Andreotti, S., M. Rutzen, S. van der Walt, et al. 2016. “An Integrated Mark-Recapture and Genetic Approach to Estimate the Population Size of White Sharks in South Africa.” *Marine Ecology Progress Series* 552: 241–253. <https://doi.org/10.3354/meps11744>.
- Baker, C. S., R. M. Hamner, M. Oremus, M. Stanley, P. Brown, and R. Constantine. 2016. *Estimating the Abundance and Effective Population Size of Māui's Dolphins Using Microsatellite Genotypes in 2010–11, With Retrospective Matching to 2001–07*, 1–44. Auckland: Department of Conservation.
- Barry, P., T. Broquet, and P. A. Gagnaire. 2022. “Age-Specific Survivorship and Fecundity Shape Genetic Diversity in Marine Fishes.” *Evolution Letters* 6, no. 1: 46–62. <https://doi.org/10.1002/evl3.265>.
- Beaumont, M. A. 2003. “Conservation Genetics.” In *Handbook of Statistical Genetics*, edited by D. J. Balding, M. Bishop, and C. Cannings, 2nd ed., 751–766. London, UK: Wiley.
- Bierne, N., F. Bonhomme, and S. Arnaud-Haond. 2016. “Dedicated Population Genomics for the Silent World: The Specific Questions of Marine Population Genetics.” *Current Zoology* 62, no. 6: 545–550. <https://doi.org/10.1093/cz/zow107>.
- Boyce, M. S. 1992. “Population Viability Analysis.” *Annual Review of Ecology and Systematics* 23: 481–506.
- Bravington, M., P. Grewe, and C. Davies. 2016. “Absolute Abundance of Southern Bluefin Tuna Estimated by Close-Kin Mark-Recapture.” *Nature Communications* 7: 13162. <https://doi.org/10.1038/ncomm513162>.

- Bravington, M. V., P. M. Grewe, and C. R. Davies. 2014. "Fishery-Independent Estimate of Spawning Biomass of Southern Bluefin Tuna Through Identification of Close-Kin Using Genetic Markers." FRDC Rep. CSIRO.
- Bravington, M. V., H. J. Skaug, and E. C. Anderson. 2016. "Close-Kin Mark-Recapture." *Statistical Science* 31: 259–274. <https://doi.org/10.1214/16-STS552>.
- Buffalo, V. 2021. "Quantifying the Relationship Between Genetic Diversity and Population Size Suggests Natural Selection Cannot Explain Lewontin's Paradox." *eLife* 10: e67509.
- Clarke, S. H., E. R. Lawrence, J. M. Matte, et al. 2024. "Global Assessment of Effective Population Sizes: Consistent Taxonomic Differences in Meeting the 50/500 Rule." *Molecular Ecology* 33: e17353. <https://doi.org/10.1111/mec.17353>.
- Creel, S., and E. Rosenblatt. 2013. "Using Pedigree Reconstruction to Estimate Population Size: Genotypes Are More Than Individually Unique Marks." *Ecology and Evolution* 3: 1294–1304. <https://doi.org/10.1002/ece3.538>.
- Crow, J. F. 1954. "Breeding Structure of Populations. II. Effective Population Number." In *Statistics and Mathematics in Biology*, edited by O. Kempthorne, T. Bancroft, J. Gowen, and J. Lush, 543–556. Ames (IA): Iowa State University Press.
- Crow, J. F. 1993. "Mutation, Mean Fitness, and Genetic Load." *Oxford Surveys in Evolutionary Biology* 9: 3–42.
- Crow, J. F., and C. Denniston. 1988. "Inbreeding and Variance Effective Population Numbers." *Evolution* 42: 482–495.
- Crow, J. F., and M. Kimura. 1970. *An Introduction to Population Genetics Theory*. New York: Harper & Row.
- Crow, J. F., and N. E. Morton. 1955. "Measurement of Gene Frequency Drift in Small Populations." *Evolution* 9: 202–214.
- Davies, C., M. Bravington, and R. Thomson. 2017. "Advice on Close-Kin Mark-Recapture for Estimating Abundance of Eastern Atlantic Bluefin Tuna: A Scoping Study." In *Atlantic-Wide Research Programme on Bluefin Tuna (ICCAT GBYP-PHASE 5) GBYP 07c/2015*. CSIRO Marine Laboratories, Castray Esplanade, Hobart, Tasmania, Australia: International Commission for the Conservation of Atlantic Tuna. https://www.iccat.int/GBYP/Docs/Close_Kin_Mark_Recapture_Phase_6_Scoping_Study.pdf.
- Delord, C., S. Arnaud-Haond, A. Leone, et al. 2024. "Advancements in Effective Population Size Estimation to Large Marine Populations: Integrating Literature Review and Methodological Application." *Evolutionary Applications*.
- Domingues, R. R., A. W. S. Hilsdorf, and O. B. F. Gadig. 2018. "The Importance of Considering Genetic Diversity in Shark and Ray Conservation Policies." *Conservation Genetics* 19, no. 3: 501–525. <https://doi.org/10.1007/s10592-017-1038-3>.
- Dudgeon, C. L., and J. R. Ovenden. 2015. "The Relationship Between Abundance and Genetic Effective Population Size in Elasmobranchs: An Example From the Globally Threatened Zebra Shark *Stegostoma fasciatum* Within Its Protected Range." *Conservation Genetics* 16, no. 6: 1443–1454. <https://doi.org/10.1007/s10592-015-0752-y>.
- Ellegren, H., and N. Galtier. 2016. "Determinants of Genetic Diversity." *Nature Reviews Genetics* 17, no. 7: 422–433. <https://doi.org/10.1038/nrg.2016.58>.
- Engen, S., R. Lande, and B. E. Saether. 2005. "Effective Size of a Fluctuating Age-Structured Population." *Genetics* 170, no. 2: 941–954. <https://doi.org/10.1534/genetics.104.028233>.
- Ewens, W. J. 1982. "On the Concept of the Effective Population Size." *Theoretical Population Biology* 21, no. 3: 373–378. [https://doi.org/10.1016/0040-5809\(82\)90024-7](https://doi.org/10.1016/0040-5809(82)90024-7).
- Ferchaud, A. L., C. Perrier, J. April, C. Hernandez, M. Dionne, and L. Bernatchez. 2016. "Making Sense of the Relationships Between N_e , N_b and N_c Towards Defining Conservation Thresholds in Atlantic Salmon (*Salmo salar*)." *Heredity* 117: 1–11. <https://doi.org/10.1038/hdy.2016.62>.
- Fisher, R. A. 1958. *The Genetical Theory of Natural Selection*. 2nd ed. New York, NY: Dover Publications.
- Frankham, R. 1995. "Effective Population Size/Adult Population Size Ratios in Wildlife: A Review." *Genetical Research Cambridge* 66, no. 2: 95–107. <https://doi.org/10.1017/S0016672300034455>.
- Franklin, I. R. 1980. "Evolutionary Change in Small Populations." In *Conservation Biology: An Evolutionary-Ecological Perspective*, edited by M. Soulé and B. Wilcox, 135–149. Sunderland, MA: Sinauer Associates.
- Gattepaille, L., T. Günther, and M. Jakobsson. 2016. "Inferring Past Effective Population Size From Distributions of Coalescent Times." *Genetics* 204, no. 3: 1191–1206. <https://doi.org/10.1534/genetics.115.185058>.
- Gilbert, K. J., and M. C. Whitlock. 2015. "Evaluating Methods for Estimating Local Effective Population Size With and Without Migration: Estimating n_e in the Presence of Migration." *Evolution* 69, no. 8: 2154–2166. <https://doi.org/10.1111/evo.12713>.
- Glazier, D. S. 2021. "Genome Size Covaries More Positively With Propagule Size Than Adult Size: New Insights Into an Old Problem." *Biology* 10, no. 4: 270. <https://doi.org/10.3390/biology10040270>.
- Gomez-Uchida, D., F. P. Palstra, T. W. Knight, and D. E. Ruzzante. 2013. "Contemporary Effective Population and Metapopulation Size (N_e and Meta- N_e): Comparison Among Three Salmonids Inhabiting a Fragmented System and Differing in Gene Flow and Its Asymmetries." *Ecology and Evolution* 3, no. 3: 569–580. <https://doi.org/10.1002/ece3.485>.
- Gutenkunst, R. N., R. D. Hernandez, S. H. Williamson, and C. D. Bustamante. 2009. "Inferring the Joint Demographic History of Multiple Populations From Multidimensional SNP Frequency Data." *PLoS Genetics* 5: e1000695.
- Hamilton, M. B., M. Tartakovsky, and A. Battocletti. 2018. "Speed-Ne: Software to Simulate and Estimate Genetic Effective Population Size (N_e) From Linkage Disequilibrium Observed in Single Samples." *Molecular Ecology Resources* 18: 714–728. <https://doi.org/10.1111/1755-0998.12759>.
- Hare, M. P., L. Nunney, M. K. Schwartz, et al. 2011. "Understanding and Estimating Effective Population Size for Practical Application in Marine Species Management." *Conservation Biology* 25, no. 3: 438–449. <https://doi.org/10.1111/j.1523-1739.2010.01637.x>.
- Hauser, L., and G. R. Carvalho. 2008. "Paradigm Shifts in Marine Fisheries Genetics: Ugly Hypotheses Slain by Beautiful Facts." *Fish and Fisheries* 9, no. 4: 333–362. <https://doi.org/10.1111/j.1467-2979.2008.00299.x>.
- Hedrick, D. 1994. "Does Variance in Reproductive Success Limit Effective Population Sizes of Marine Organisms?" In *Genetics and Evolution of Aquatic Organisms*, edited by A. R. Beaumont, 122–134. London, UK: Chapman & Hall.
- Hedrick, D., and A. H. Pudovkin. 2011. "Sweepstakes Reproductive Success in Highly Fecund Marine Fish and Shellfish: A Review and Commentary." *Bulletin of Marine Science* 87, no. 4: 971–1002. <https://doi.org/10.5343/bms.2010.1051>.
- Hedrick, P. 2005. "Large Variance in Reproductive Success and the N_e/N Ratio." *Evolution* 59, no. 7: 1596–1599. <https://doi.org/10.1111/j.0014-3820.2005.tb01809.x>.
- Hettiarachchi, C. K. H., and R. M. Huggins. 2018. "Inference From Single Occasion Capture Experiments Using Genetic Markers." *Biometrical Journal* 60: 463–479. <https://doi.org/10.1002/bimj.201700046>.
- Hill, W. G. 1972. "Effective Size of Population With Overlapping Generations." *Theoretical Population Biology* 3, no. 3: 278–289. [https://doi.org/10.1016/0040-5809\(72\)90004-4](https://doi.org/10.1016/0040-5809(72)90004-4).

- Hill, W. G. 1981. "Estimation of Effective Population-Size From Data on Linkage Disequilibrium." *Genetics Research* 38: 209–216.
- Hoban, S., M. Bruford, J. J. D'Urban, et al. 2020. "Genetic Diversity Targets and Indicators in the CBD Post-2020 Global Biodiversity Framework Must Be Improved." *Biological Conservation* 248: 108654.
- Hoban, S., J. M. da Silva, A. Hughes, et al. 2024. "Too Simple, Too Complex, or Just Right? Advantages, Challenges, and Guidance for Indicators of Genetic Diversity." *Bioscience* 74: biae006. <https://doi.org/10.1093/biosci/biae006>.
- Hössjer, O., L. Laikre, and N. Ryman. 2016. "Effective Sizes and Time to Migration–Drift Equilibrium in Geographically Subdivided Populations." *Theoretical Population Biology* 112: 139–156. <https://doi.org/10.1016/j.tpb.2016.09.001>.
- Hössjer, O., F. Olsson, L. Laikre, and N. Ryman. 2015. "Metapopulation Inbreeding Dynamics, Effective Size and Subpopulation Differentiation - A General Analytical Approach for Diploid Organisms." *Theoretical Population Biology* 102: 40–59.
- Husemann, M., F. E. Zachos, R. J. Paxton, and J. C. Habel. 2016. "Effective Population Size in Ecology and Evolution." *Heredity* 117, no. 4: 191–192. <https://doi.org/10.1038/hdy.2016.75>.
- Irion, D. T., L. R. Noble, A. A. Kock, et al. 2017. "Pessimistic Assessment of White Shark Population Status in South Africa: Comment on Andreotti Et al. (2016)." *Marine Ecology Progress Series* 577: 251–255.
- Jennings, S. 2000. "Patterns and Prediction of Population Recovery in Marine Reserves." *Reviews in Fish Biology and Fisheries* 10: 209–231.
- Jones, A. T., J. R. Ovenden, and Y.-G. Wang. 2016. "Improved Confidence Intervals for the Linkage Disequilibrium Method for Estimating Effective Population Size." *Heredity* 117: 217–223. <https://doi.org/10.1038/hdy.2016.19>.
- Jorde, P. E., and N. Ryman. 1995. "Temporal Allele Frequency Change and Estimation of Effective Size in Populations With Overlapping Generations." *Genetics* 139, no. 2: 1077–1090. <https://doi.org/10.1093/genetics/139.2.1077>.
- Jorde, P. E., and N. Ryman. 2007. "Unbiased Estimator for Genetic Drift and Effective Population Size." *Genetics* 177, no. 2: 927–935. <https://doi.org/10.1534/genetics.107.075481>.
- Juan-Jordá, M. J., I. Mosqueira, J. Freire, and N. K. Dulvy. 2013. "Life in 3-D: Life History Strategies in Tunas, Mackerels and Bonitos." *Reviews in Fish Biology and Fisheries* 23: 135–155. <https://doi.org/10.1007/s11160-012-9284-4>.
- Kimura, M., and J. F. Crow. 1963. "The Measurement of Effective Population Number." *Evolution* 17: 279–288.
- King, J. R., M. Wetklo, J. Supernaul, et al. 2015. "Genetic Analysis of Stock Structure of Blue Shark (*Prionace glauca*) in the North Pacific." *Fisheries Research* 172: 181–189. <https://doi.org/10.1016/j.fishres.2015.06.029>.
- Laconcha, U., M. Iriondo, H. Arrizabalaga, et al. 2015. "New Nuclear SNP Markers Unravel the Genetic Structure and Effective Population Size of Albacore Tuna (*Thunnus alalunga*)." *PLoS One* 10, no. 6: e0128247. <https://doi.org/10.1371/journal.pone.0128247>.
- Laikre, L., F. Olsson, E. Jansson, O. Hössjer, and N. Ryman. 2016. "Metapopulation Effective Size and Conservation Genetic Goals for the Fennoscandian Wolf (*Canis lupus*) Population." *Heredity* 117: 279–289. <https://doi.org/10.1038/hdy.2016.44>.
- Lande, R. 1995. "Mutation and Conservation." *Conservation Biology* 9: 782–791. <https://doi.org/10.1046/j.1523-1739.1995.09040782.x>.
- Larroque, J., and J. Balkenhol. 2023. "A Simulation-Based Evaluation of Methods for Estimating Census Population Size of Terrestrial Game Species From Genetically-Identified Parent-Offspring Pairs." *PeerJ* 11: e15151.
- Laurent, V., and S. Planes. 2007. "Effective Population Size Estimation on *Sardina pilchardus* in the Bay of Biscay Using a Temporal Genetic Approach." *Biological Journal of the Linnean Society* 90, no. 4: 591–602. <https://doi.org/10.1111/j.1095-8312.2007.00747.x>.
- Lee, A. M., S. Engen, and B.-E. Sæther. 2011. "The Influence of Persistent Individual Differences and Age at Maturity on Effective Population Size." *Proceedings of the Royal Society B* 278, no. 1722: 3303–3312. <https://doi.org/10.1098/rspb.2011.0283>.
- Leroy, G., E. L. Carroll, M. W. Bruford, et al. 2017. "Next-Generation Metrics for Monitoring Genetic Erosion Within Populations of Conservation Concern." *Evolutionary Applications* 11: 1066–1083. <https://doi.org/10.1111/eva.12564>.
- López, M. A., and C. López-Fanjul. 1993a. "Spontaneous Mutation for a Quantitative Trait in *Drosophila melanogaster*. I. Response to Artificial Selection." *Genetics Research* 61, no. 2: 107–116. <https://doi.org/10.1017/S0016672300031219>.
- López, M. A., and C. López-Fanjul. 1993b. "Spontaneous Mutation for a Quantitative Trait in *Drosophila melanogaster*. II. Distribution of Mutant Effects on the Trait and Fitness." *Genetics Research* 61, no. 2: 117–126. <https://doi.org/10.1017/S0016672300031220>.
- Lynch, M., M. Ackerman, J. F. Gout, et al. 2016. "Genetic Drift, Selection, and Evolution of the Mutation Rate." *Nature Reviews Genetics* 17: 704–714.
- Lynch, M., F. Ali, T. Lin, Y. Wang, J. Ni, and H. Long. 2023. "The Divergence of Mutation Rates and Spectra Across the Tree of Life." *EMBO Reports* 24, no. 10: e57561. <https://doi.org/10.15252/embr.202357561>.
- Lynch, M., L. M. Bobay, F. Catania, J. F. Gout, and M. Rho. 2011. "The Repatterning of Eukaryotic Genomes by Random Genetic Drift." *Annual Review of Genomics and Human Genetics* 12: 347–366. <https://doi.org/10.1146/annurev-genom-082410-101412>.
- Lynch, M., and R. Lande. 1998. "The Critical Effective Size for a Genetically Secure Population." *Animal Conservation* 1: 70–72. <https://doi.org/10.1111/j.1469-1795.1998.tb00229>.
- Müller, B., M. Mercker, and J. Brün. 2020. "Population Size Estimates Based on the Frequency of Genetically Assigned Parent–Offspring Pairs Within a Subsample." *Ecology and Evolution* 10: 6356–6363. <https://doi.org/10.1002/ece3.6365>.
- Murua, H., E. Rodriguez-Marin, J. D. Neilson, J. H. Farley, and M. J. Juan-Jordá. 2017. "Fast Versus Slow Growing Tuna Species: Age, Growth, and Implications for Population Dynamics and Fisheries Management." *Reviews in Fish Biology and Fisheries* 27: 733–773. <https://doi.org/10.1007/s11160-017-9474-1>.
- Nadachowska-Brzyska, K., M. Konczal, and W. Babik. 2022. "Navigating the Temporal Continuum of Effective Population Size." *Methods in Ecology and Evolution* 13, no. 1: 22–41. <https://doi.org/10.1111/2041-210X.13740>.
- Nikolic, N., F. Devloo-Delva, D. Bailleul, et al. 2022. "Stepping up to Genome Scan Allows Stock Differentiation in the Worldwide Distributed Blue Shark *Prionace Glauca*." *Molecular Ecology* 32: 1000–1019.
- Nunney, L. 1991. "The Influence of Age Structure and Fecundity on Effective Population Size." *Proceedings of the Royal Society B* 246, no. 1315: 71–76. <https://doi.org/10.1098/rspb.1991.0126>.
- Nunney, L. 1993. "The Influence of Mating System and Overlapping Generations on Effective Population Size." *Evolution* 47, no. 5: 1329–1341. <https://doi.org/10.1111/j.1558-5646.1993.tb02158.x>.
- Nunney, L. 1996. "The Influence of Variation in Female Fecundity on Effective Population Size." *Biological Journal of the Linnean Society* 59, no. 4: 411–425. <https://doi.org/10.1111/j.1095-8312.1996.tb01474.x>.

- Ovenden, J. R., G. M. Leigh, D. C. Blower, et al. 2016. "Can Estimates of Genetic Effective Population Size Contribute to Fisheries Stock Assessments?" *Journal of Fish Biology* 89, no. 6: 2505–2518. <https://doi.org/10.1111/jfb.13129>.
- Palstra, F. P., and D. J. Fraser. 2012. "Effective/Census Population Size Ratio Estimation: A Compendium and Appraisal." *Ecology and Evolution* 2, no. 9: 2357–2365. <https://doi.org/10.1002/ece3.329>.
- Palstra, F. P., and D. E. Ruzzante. 2008. "Genetic Estimates of Contemporary Effective Population Size: What Can They Tell Us About the Importance of Genetic Stochasticity for Wild Population Persistence?" *Molecular Ecology* 17, no. 5: 3428–3447. <https://doi.org/10.1111/j.1365-294X.2008.03842.x>.
- Pinder, J. E., J. G. Wiener, and M. H. Smith. 1978. "The Weibull Distribution: A New Method of Summarizing Survivorship Data." *Ecology* 59, no. 1: 175–179.
- Pinsky, M. L., and S. R. Palumbi. 2014. "Meta-Analysis Reveals Lower Genetic Diversity in Overfished Populations." *Molecular Ecology* 23, no. 1: 29–39. <https://doi.org/10.1111/mec.12509>.
- Portnoy, D. S., J. R. McDowell, C. T. McCandless, J. A. Musick, and J. E. Graves. 2009. "Effective Size Closely Approximates the Census Size in the Heavily Exploited Western Atlantic Population of the Sandbar Shark, *Carcharhinus plumbeus*." *Conservation Genetics* 10, no. 6: 1697–1705. <https://doi.org/10.1007/s10592-008-9771-2>.
- Rai, U. K. 2003. "Minimum Sizes for Viable Population and Conservation Biology." *Our Nature* 1, no. 1: 3–9. <https://doi.org/10.3126/on.v1i1.297>.
- Reed, D. H., J. J. O'Grady, B. W. Brook, J. D. Ballou, and R. Frankham. 2003. "Estimates of Minimum Viable Population Sizes for Vertebrates and Factors Influencing Those Estimates." *Biological Conservation* 113, no. 1: 23–34. [https://doi.org/10.1016/S0006-3207\(02\)00346-4](https://doi.org/10.1016/S0006-3207(02)00346-4).
- Rodríguez-Rodríguez, G., H. M. Ballesteros, E. Sánchez-Llamas, R. Bande, and R. F. Otero. 2022. "The State of the Art in Cost-Benefit of HTS Methods for Stock Assessment: An Overview." *Frontiers in Marine Science* 9: 1005534. <https://doi.org/10.3389/fmars.2022.1005534>.
- Romiguier, J., P. Gayral, M. Ballenghien, et al. 2014. "Comparative Population Genomics in Animals Uncovers the Determinants of Genetic Diversity." *Nature* 515, no. 7526: 261–263. <https://doi.org/10.1038/nature13685>.
- Rosenblatt, E., S. Creel, K. Gieder, J. Murdoch, and T. Donovan. 2023. "Advances in Wildlife Abundance Estimation Using Pedigree Reconstruction." *Ecology and Evolution* 13: e10650. <https://doi.org/10.1002/ece3.10650>.
- Ryman, N., L. Laikre, and O. Hössjer. 2019. "Do Estimates of Contemporary Effective Population Size Tell Us What We Want to Know?" *Molecular Ecology* 28, no. 8: 1904–1918. <https://doi.org/10.1111/mec.15027>.
- Santiago, E., I. Novo, A. F. Pardiñas, M. Saura, J. Wang, and A. Caballero. 2020. "Recent Demographic History Inferred by High-Resolution Analysis of Linkage Disequilibrium." *Molecular Biology and Evolution* 7, no. 12: 3642–3653. <https://doi.org/10.1093/molbev/msaa169>.
- Schwartz, M. K., D. A. Tallmon, and G. Luikart. 1999. "Using Genetics to Estimate the Size of Wild Populations: Many Methods, Much Potential, Uncertain Utility." *Animal Conservation* 2: 321–323.
- Shaffer, M. L. 1981. "Minimum Population Sizes for Species Conservation." *Bioscience* 31, no. 2: 131–134. <https://doi.org/10.2307/1308256>.
- Sjödín, P., I. Kaj, S. Krone, M. Lascoux, and M. Nordborg. 2005. "On the Meaning and Existence of an Effective Population Size." *Genetics* 169: 1061–1070.
- Soulé, M. E. 1980. "Thresholds for Survival: Maintaining Fitness and Evolutionary Potential." In *Conservation Biology: An Evolutionary-Ecological Perspective*, edited by M. E. Soulé and B. A. Wilcox, 151–169. Sunderland, MA: Sinauer.
- Soulé, M. E. 1987. *Viable Populations for Conservation*. Cambridge, UK: Cambridge University Press.
- Swenson, J. D., E. N. Brooks, D. Kacev, et al. 2024. "Accounting for Unobserved Population Dynamics and Aging Error in Close-Kin Mark-Recapture Assessments." *Ecology and Evolution* 14: e10854. <https://doi.org/10.1002/ece3.10854>.
- Trask, A. E., E. M. Signal, D. I. McCracken, S. B. Piertney, and J. M. Reid. 2017. "Estimating Demographic Contributions to Effective Population Size in an Age-Structured Wild Population Experiencing Environmental and Demographic Stochasticity." *Journal of Animal Ecology* 86, no. 5: 1082–1093. <https://doi.org/10.1111/1365-2656.12703>.
- Wakeley, J. 2020. "Developments in Coalescent Theory From Single Loci to Chromosomes." *Theoretical Population Biology* 133: 56–64. <https://doi.org/10.1016/j.tpb.2020.02.002>.
- Wang, J. 2005. "Estimation of Effective Population Sizes From Data on Genetic Markers." *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 360, no. 1459: 1395–1409. <https://doi.org/10.1098/rstb.2005.1682>.
- Wang, J. 2009. "A New Method for Estimating Effective Population Size From a Single Sample of Multilocus Genotypes." *Molecular Ecology* 18: 2148–2164.
- Wang, J. 2016. "A Comparison of Single-Sample Estimators of Effective Population Sizes From Genetic Marker Data." *Molecular Ecology* 25, no. 19: 4692–4711. <https://doi.org/10.1111/mec.13725>.
- Wang, J., and A. Caballero. 1999. "Developments in Predicting the Effective Size of Subdivided Populations." *Heredity* 82, no. 2: 212–226. <https://doi.org/10.1038/sj.hdy.6884670>.
- Wang, J., E. Santiago, and A. Caballero. 2016. "Prediction and Estimation of Effective Population Size." *Heredity* 117: 193–206. <https://doi.org/10.1038/hdy.2016.43>.
- Waples, R. K., W. A. Larson, and R. S. Waples. 2016. "Estimating Contemporary Effective Population Size in Non-Model Species Using Linkage Disequilibrium Across Thousands of Loci." *Heredity* 117, no. 4: 233–240. <https://doi.org/10.1038/hdy.2016.60>.
- Waples, R. S. 2002. "Evaluating the Effect of Stage-Specific Survivorship on the N_e/N Ratio." *Molecular Ecology* 11: 1029–1037.
- Waples, R. S. 2005. "Genetic Estimates of Contemporary Effective Population Size: To What Time Periods Do the Estimates Apply?" *Molecular Ecology* 14, no. 11: 3335–3352. <https://doi.org/10.1111/j.1365-294X.2005.02673.x>.
- Waples, R. S. 2006. "A Bias Correction for Estimates of Effective Population Size Based on Linkage Disequilibrium at Unlinked Gene Loci." *Conservation Genetics* 7: 167–184. <https://doi.org/10.1007/s10592-005-9100-y>.
- Waples, R. S. 2016a. "Tiny Estimates of the N_e/N Ratio in Marine Fishes: Are They Real?" *Journal of Fish Biology* 89, no. 6: 2479–2504. <https://doi.org/10.1111/jfb.13143>.
- Waples, R. S. 2016b. "Life-History Traits and Effective Population Size in Species With Overlapping Generations Revisited: The Importance of Adult Mortality." *Heredity* 117, no. 4: 241–250. <https://doi.org/10.1038/hdy.2016.29>.
- Waples, R. S. 2022. "What Is N_e , Anyway?" *Journal of Heredity* 113, no. 4: 371–379. <https://doi.org/10.1093/jhered/esac023>.
- Waples, R. S. 2024a. "The N_e/N Ratio in Applied Conservation." *Evolutionary Applications* 17: e13695. <https://doi.org/10.1111/eva.13695>.
- Waples, R. S. 2024b. "Practical Application of the Linkage Disequilibrium Method for Estimating Contemporary Effective Population Size: A Review." *Molecular Ecology Resources* 24: e13879. <https://doi.org/10.1111/1755-0998.13879>.

- Waples, R. S., and T. Antao. 2014. "Intermittent Breeding and Constraints on Litter Size: Consequences for Effective Population Size per Generation (N_e) and per Reproductive Cycle (N_b)." *Evolution* 68, no. 6: 1722–1734. <https://doi.org/10.1111/evo.12384>.
- Waples, R. S., C. Do, and J. Choquet. 2011. "Calculating N_e and N_e/N in Age-Structured Populations: A Hybrid Hill-Felsenstein Approach." *Ecology* 92, no. 7: 1513–1522.
- Waples, R. S., P. M. Grewe, M. W. Bravington, R. Hillary, and P. Feutry. 2018. "Robust Estimates of a High $n_e/$ Ratio in a Top Marine Predator, Southern Bluefin Tuna." *Science Advances* 4, no. 7: eaar775. <https://doi.org/10.1126/sciadv.aar7759>.
- Waples, R. S., G. Luikart, J. R. Faulkner, and D. A. Tallmon. 2013. "Simple Life History Traits Explain Key Effective Population Size Ratios Across Diverse Taxa." *Proceedings of the Royal Society B* 280, no. 1768: 20131339. <https://doi.org/10.1098/rspb.2013.1339>.
- Waples, R. S., and D. J. Teel. 1990. "Conservation Genetics of Pacific Salmon I. Temporal Changes in Allele Frequency." *Conservation Biology* 4: 144–156.
- Waples, R. S., R. K. Waples, and E. J. Ward. 2022. "Pseudoreplication in Genomic-Scale Data Sets." *Molecular Ecology Resources* 22: 503–518. <https://doi.org/10.1111/1755-0998.13482>.
- Ward-Paige, C. A., D. M. Keith, B. Worm, and H. K. Lotze. 2012. "Recovery Potential and Conservation Options for Elasmobranchs." *Journal of Fish Biology* 80: 1844–1869.
- Watterson, G. A. 1975. "On the Number of Segregating Sites in Genetical Models Without Recombination." *Theoretical Population Biology* 7: 256–276.
- Wilson, H. B., B. E. Kendall, and H. P. Possingham. 2011. "Variability in Population Abundance and the Classification of Extinction Risk." *Conservation Biology* 25: 747–757.
- Wright, S. 1931. "Evolution in Mendelian Populations." *Genetics* 16, no. 2: 97–159. <https://doi.org/10.1093/genetics/16.2.97>.
- Wright, S. 1938. "Size of a Population and Breeding Structure in Relation to Evolution." *Science* 87: 430–431.