

# The enigmatic coralligenous atolls of northern Corsica: possible scars from World War II

Marc VERLAQUE<sup>1\*</sup>, Charles-François BOUDOURESQUE<sup>1</sup>

<sup>1</sup>Aix Marseille University and Université de Toulon, CNRS, IRD, MIO (Mediterranean Institute of Oceanography), UM 110, 13009 Marseille, France.

\* Corresponding author: marclouis.verlaque@gmail.com

**Abstract.** In August 2011, during an oceanographic survey, a series of enigmatic coralligenous circles were discovered, between 105 and 125 m deep, off the coast of Cap Corse (Capicorsu, northern Corsica, Mediterranean Sea). Described under the name of coralligenous atolls, this new type of bio-concretion has attracted the attention of scientists and environment managers. Several prospecting surveys have been organized to unravel the mystery of these formations. Hitherto, only a natural and ancient origin (hydrothermal, biological, or hydrodynamic origin) has been considered. In the present paper, the alternative hypothesis of an anthropogenic and recent origin is proposed: the jettisoning of unused, generally unprimed, bombs by US aircraft squadrons based in Corsica, returning from their raids to the European continent, towards the end of World War II (1943-1945).

**Keywords:** Corsica, coralligenous atolls, bombing craters, jettison zone, World War II.

**Résumé.** Les énigmatiques atolls coralligènes du nord de la Corse : possibles cicatrices de la Seconde Guerre Mondiale. En août 2011, lors d'une campagne océanographique, un grand nombre d'énigmatiques cercles de coralligène ont été découverts, entre 105 et 125 m de profondeur, au large du Cap Corse (Capicorsu, nord de la Corse, Méditerranée). Décrits sous le nom d'atolls de coralligène, ce nouveau type de bio-concrétionnement a attiré l'attention des scientifiques et des gestionnaires de l'environnement. Plusieurs campagnes de prospection ont été organisées afin de percer le mystère de ces formations. Jusqu'à présent, seules des origines naturelles et anciennes (hydrothermale, biologique ou hydrodynamique) ont été envisagées. Nous proposons ici l'hypothèse opposée d'une origine récente et anthropique : le délestage des bombes non-utilisées et généralement non-amorcées, par les escadrilles américaines basées en Corse, au retour de leurs missions vers le continent européen, vers la fin de la Seconde Guerre Mondiale (1943-1945).

**Mots-clés:** Corse, atolls coralligènes, cratères de bombes, Seconde Guerre Mondiale, zone *jettison*.

## 1. Introduction

Over the last decades, the increased spatial resolution and accurate positioning capability of hydroacoustic instruments for seabed measurements, i.e., multibeam echosounder (MBES), sub-bottom profiler (SBP), and side-scan sonar (SSS), has led to the discovery of many different seabed structures. Among them, circular or elliptical structures have been described worldwide. Their origin can be either natural (i.e. physical, geological, or biological) or anthropogenic (i.e. dredging, dumping, explosives) (see the review by Diaz-Mendoza *et al.*, 2023).

On August 31, 2011, during the *CapCoral 2* oceanographic survey carried out by GIS Posidonie and the University of Corsica, a new type of coralligenous concretions was discovered, between 105 and 125 m depth, off the coast of Cap Corse (Capicorsu, northern Corsica, Mediterranean Sea) (Bonacorsi *et al.*, 2011, 2012a; Bonacorsi 2012). Circular in shape, these structures have been described under the name of *atolls coralligènes* (coralligenous atolls) (Bonacorsi *et al.*, 2012b). Since then, several multidisciplinary surveys have been dedicated to these bio-concretions to study them and attempt to understand the origin of their formation: *CoralCorse* (2013), *MedAtolls* (2014), *Gombessa 6 'Cap Corse'* (2021) and the most recent, organized by Andromede Oceanology (July 2023) (Bonacorsi *et al.*, 2013, 2014; Clabaut *et al.*, 2014; Deter *et al.*, 2022).

Since 2016, the area where the coralligenous atolls were discovered has been part of a marine park, the *Parc naturel marin du Cap Corse et de l'Agriate*.

To date, only a natural (hydrothermal, biological, or hydrodynamic) and ancient (i.e. several millennia) origin has been considered. Here, we propose and discuss an alternative hypothesis, that of an anthropogenic and recent origin: the jettisoning of unused bombs by US aircraft squadrons based in Corsica, returning from raids to southern Europe, towards the end of World War II.

Since 2016, the area where the coralligenous atolls were discovered has been part of a marine park, the *Parc naturel marin du Cap Corse et de l'Agriate*.

To date, only a natural (hydrothermal, biological, or hydrodynamic) and ancient (i.e. several millennia) origin has been considered. Here, we propose and discuss an alternative hypothesis, that of an anthropogenic and recent origin: the jettisoning of unused bombs by US aircraft squadrons based in Corsica, returning from raids to southern Europe, towards the end of World War II.

## **2. Characteristics of the coralligenous atolls**

Two fields of several hundred coralligenous atolls (more than a thousand according to Andromede Oceanology, <https://www.andromede-ocean.com>) are localized 22 and 31 km north of Cap Corse. These atolls have a diameter between 20 and 25 m (**Fig. 1a**). The best characterized atolls, according to Bonacorsi *et al.* (2012b, 2013), Clabaut *et al.* (2014), Pergent-Martini *et al.* (2014), Pergent *et al.* (2015), and Denter *et al.* (2022), consist of:

- a fragmented central dome (1 to 2 m in diameter and 0.2 to 0.5 m high), made up of coralligenous frameworks of different sizes (**Fig. 1b**),

- a halo of coastal detrital coarse sediments, ~10 m wide, with a high calcium carbonate content (85 to 92%), with a few scattered free-living rhodoliths (non-geniculate and unattached calcified rhodophytes) and numerous organic debris,
- a peripheral crown 1 to 3 m wide, simple, or split by a sandy strip, made up of coralligenous bank (*coralligène de plateau*), associated with a high density of free-living rhodoliths and numerous invertebrates (e.g. sponges, bryozoans, and echinoderms).

Other atolls lack a central dome. Seismic reflection surveys seem to indicate that the atolls are located on beds of coarse sediment, 1 to 2 m thick. No atolls have been observed on soft bottoms more than 3-4 m thick (Pergent-Martini *et al.*, 2014a, 2014b). A first <sup>14</sup>C dating of fragments of bio-concretions taken from the surface of a central dome gave an age of 6 200 years (Pergent-Martini *et al.*, oral communication, *Fête de la Science*, November 2021). Examination of photographs of thin sections of the bio-concretion (courtesy of Michèle Ferrandini) enabled us to determine the main coralligenous-building rhodophytes (*Lithophyllum* of the *stictiforme* group, *Mesophyllum expansum*, *Titanoderma* sp., *Neogoniolithon* sp., *Peyssonnelia* sp., *Polystrata* sp.).

Comparison of two images of the same dome taken in 2014 and 2021 showed an increase in crevices and sedimentation, but no change in the coverage of outstanding sessile species (Deter *et al.*, 2022). Examination of sonograms shows that the contiguous atolls do not merge but overlap (**Fig. 1c**). Their arrangement on the bottom is not uniform: atolls can form strings (**Fig. 1d**) or clusters (**Fig. 1e**).


The atolls are oases of species diversity with, on the central dome and the crown, a very rich benthic fauna (brachiopods, bryozoans, cnidarians, crustaceans, echinoderms, molluscs, sponges) (Pergent-Martini *et al.*, 2014a, 2014b; Pergent-Martini *et al.*, oral communication, *Fête de Science*, November 2021).

### **3. Characteristics of World War II bomb impact craters**

The comparison of aerial photographs of terrestrial areas bombed during World War II with Cap Corse sonograms enabled us to identify several analogies between bomb impact craters and coralligenous atolls, despite the smaller size of the former (usually, 6-12 m in diameter) (Capps Tunwell *et al.*, 2016; Dolejš *et al.*, 2020; Waga *et al.*, 2022):


- in a given area, the impact craters have approximately the same diameter (**Fig. 1f**);

- the arrangement in a string or in a cluster is common, probably reflecting different release conditions (altitude, operational mode) (**Fig. 1f** and **1g**);
- the contiguous craters overlap, reflecting the succession of impacts over time (**Fig. 1h**);
- some craters have a central dome of ejecta (**Fig. 1i**).



**Figure 1.** Coralligenous atolls and terrestrial bomb craters. **a:** example of concentration of coralligenous atolls off Cap Corse, on a mosaic of sonograms. **b:** atolls with a central dome. **c:** atoll overlapping. **d:** string of atolls. **e:** cluster of atolls. **f:** aerial view of a terrestrial bombed area from World War II. **g:** a string of terrestrial bomb craters. **h:** overlapping of terrestrial bomb craters. **i:** terrestrial bomb craters with a central dome. (**a-e** from Pergent-Martini *et al.*, 2014; **f, g, i** from © IGNF; **h** from Dolejš *et al.*, 2020, modified). Scale: 250 m (**a**), 25 m (**b-e, g-i**), 100 m (**f**).

The formation of a central dome of ejecta in reaction to an impact is a frequent phenomenon, and the coralligenous atolls with a double crown could result from induced circular waves. This is very well illustrated by a drop falling on the surface of water (**Fig. 2a**). Such features can also be observed in some meteorite impact craters on the surface of the Moon, planets and natural satellites (**Fig. 2b-c**).



**Figure 2.** a: impact of a drop on the surface of water with central ejecta and induced circular waves. b-c: the *Tycho* impact crater and the *Hesiodus A* concentric impact crater on the surface of the Moon (From Wikimedia commons).

#### 4. Hypotheses of natural and ancient events

Hitherto, only a natural and ancient origin (several millennia) has been considered for the formation of coralligenous atolls.

The first hypothesis is a hydrothermal origin (Bonacorsi *et al.*, 2012b). The expulsion of weakly buried gas in the sediment creates depressions on the seabed, called pockmarks. Discovered in all the world's oceans, their shape is usually circular or elliptical with elongation parallel to the direction of the bottom currents. Pockmarks can be grouped, aligned, linked, or isolated. They are often associated with sedimentary or tectonic structures (channels, faults, diapirs, synclines-anticlines) and preferentially develop when the sedimentary layer is thin (Garlan *et al.*, 2018). In the Mediterranean, environments in which deep water pockmarks preferentially occur are active mud volcanoes, vestigial mud volcanoes and corresponding fault systems, and most abundantly on active faults and submarine slumps (Dimitrov and Woodside, 2003).

The characteristics of Corsican coralligenous atolls are very different from those of pockmarks and hydrothermal sources (black or white smokers; Fouquet *et al.*, 1994). No current emissions of gases or fluids could be demonstrated in Corsican atoll areas (Deter *et al.*, 2021, 2022), which does not exclude the possibility that such emissions could have existed in the past. In southern Italy, for example at Ischia Island, the activity of volcanic underwater CO<sub>2</sub> vents is irregular over time (Foo *et al.*, 2018; Teixidó *et al.*, 2018).

Circular seabed structures can have a biological origin, e.g. fish nests (Kawase *et al.*, 2013), seagrass beds (Warwick, 2022) or shallow pits created by cetaceans (Nelson *et al.*, 1987), but in view of the depth and characteristics of coralligenous atolls, these origins can be ruled out. Another biological origin, dating back to a period when sea level was much lower than today, has been considered (Pergent-Martini *et al.*, 2014; Deter *et al.*, 2022). During the LGM (Last Glacial Maximum, ~20 000 years ago), the sea level was 120-135 m below the current sea level (Lichter *et al.*, 2010; Collina-Girard, 2012). The coralligenous atolls would have been formed under particular hydrodynamic conditions and at very shallow depths by photophilic calcified rhodophytes such as the *Neogoniolithon brassica-florida* reefs described in southern Tunisia (Denizot *et al.*, 1981; Pergent *et al.*, 2009; Langar *et al.*, 2011). However, *N. brassica-florida* reefs are thermophilic formations, today confined to the warmest regions of the Mediterranean (Boudouresque, 2003), and their presence in Corsica near the LGM is unlikely. Furthermore, no shallow formation similar to the coralligenous atolls has to date been discovered in the Mediterranean and the assemblage of calcified rhodophytes identified on the thin sections (see section 2) is characteristic of coralligenous concretions, therefore of a deep habitat.

The more or less large size and the totally fragmented state of the central domes has been attributed to a biological process of erosion. However, bioeroding organisms do not cause this type of dismantling of coralligenous formations. Sponges (*Cliona* spp.) and sea urchins (*Echinus* spp., *Sphaerechinus granularis*) erode the outcrop surface while bivalves (*Lithophaga lithophaga*, *Petricola* spp.) bore galleries (Sartoretto and Francour, 1997; Ballesteros, 2006; Boudouresque *et al.*, 2017; Gennaro *et al.*, 2020). Finally, a bottom current origin should be excluded because, at more than 100 m depth, currents are unidirectional rather than swirling.

## **5. A new hypothesis: a recent and human-driven event**

Based on **(i)** the similarities noted between the fields of coralligenous atolls and the terrestrial bombed areas of World War II, and **(ii)** the analysis of documents relating to the activity of the US aircraft squadrons, based in Corsica, towards the end of this war, we propose an alternative hypothesis to a natural and ancient event: the jettisoning of unused unprimed bombs towards the end of World War II (1943-1945), i.e. a recent and anthropogenic event.

### **5.1. Structural arguments**

The distribution of coralligenous atolls is similar to that of bomb craters. The clusters of atolls would correspond to a grouped or low altitude drop and the aligned atolls to a string drop at high altitude. In areas with high densities of atolls, most have exactly the same

diameter, which is difficult to explain by a natural cause but not at all with the hypothesis of bombs of the same calibre.

The wider diameter of the coralligenous atolls (20-25 m) compared to that of bomb craters (6-12 m) may be due to the characteristics of the marine environment (water, soft substrate) and the fact that unprimed bombs penetrate the substrate intact. The average speed of fall of World War II bombs (usually, 100 to 500 kg) was mainly a function of the altitude of release (e.g. 6 000 m: 300 m/s and 600 m: 150 m/s) (Ley and Schaefer, 1941). This speed must have been considerably reduced by the impact on the sea surface, but it would have had to remain sufficient to penetrate the soft substrate.

Neighbouring atolls do not merge, but overlap in a defined order with, at the end of the series, an entire atoll (which would be the last formed) as in the case of bomb impacts. The presence of a central dome and its fragmented state are consistent with impact craters with a central dome of ejecta. The presence or absence of the central dome and its variable size would be linked to the conditions of the impact (thickness of the sediment, and type, speed, and angle of penetration of the bomb). The <sup>14</sup>C dating of organisms taken from the surface of a dome, which indicates more than 6 000 years, is compatible with the ejecta of deep sedimentary layers. In a natural formation, the surface layers would be younger.

## **5.2. Historical arguments**

World War II ushered in the practice of massive aerial bombardments. When, due to counter-orders, weather (fog, clouds) or damage (enemy anti-aircraft defence), the planes were forced to return to their base without being able to complete their mission, it was either too dangerous for them to land with their bomb load, or the weight of the bombs would have increased their fuel consumption too much. It was therefore the rule, before landing at the departure airfield, to jettison unused bombs at sea, possibly quite close to the coast. The areas dedicated to this operation are referred to as 'jettison zones' (Wikipedia Contributors, 2023).

Numerous testimonies from World War II pilots show that these bomb dumps, following aborted missions, were very frequent (Granfield, 2011; Griggs, 2015). For example, on December 15, 1944, a single raid aborted due to fog led a squadron of 138 RAF Lancaster bombers to drop around 100 000 incendiary projectiles and nearly two hundred 'Cookies' (giant 1.8 metric ton bombs) in the English Channel (Wikipedia Contributors, 2023).

On October 4, 1943, Corsica was itself liberated from the Italian and then German occupation, thanks to an uprising by the local population, Corsican *maquisards*, the landing of Free French Forces

(the emblematic submarine *Casabianca*), as well as a change of heart among the Italian soldiers facing the Germans (Gambiez, 1973; Ferranti, 2013; Gregori, 2023).

The US army landed in Corsica in December 1943 and built 17 airfields for its hundreds of bombers and went so far as to give the name of an aircraft carrier to the island: the 'USS Corsica' (**Fig. 3a**) (Taddei, 2023). More than 2 000 bombers, fighters and reconnaissance planes from the USAAF (United States Army Air Forces), the RAF (Royal Air Force) and the French Air Force were thus positioned on the island. The main types of bombers used were the B17 Flying Fortress, B24 Liberators, B25 Mitchells, and B26 Marauders (**Fig. 3b-e**).

The famous French writer, poet, journalist and aviator Antoine de Saint-Exupery performed reconnaissance missions, from the Bastia-Puretta airfield (Corsica), to collect intelligence on German troop movements, in preparation for the Allied invasion of southern France. He disappeared while on one such reconnaissance mission, on July 31, 1944. The wreckage of his plane, a P38 Lightning, was discovered off the coast of Marseilles in 2000 (Vaudoit *et al.*, 2004; Pradel and Vanrell, 2008).

Each bomber pilot had to carry out at least 70 raids before being replaced. Until April 1945, several thousand raids were carried out on northern Italy and southern France. A strategic target was the Brenner Pass over which runs the railway line connecting Austria to Italy (Verona and Innsbruck are on this line), a line that was used to supply the German armies (**Fig. 3f**). From November 6, 1944, to April 25, 1945, the 57<sup>th</sup> Squadron of B25 Mitchells conducted 6 839 raids, lost 46 aircraft, and had 532 aircraft damaged (57th Bomb Wing Association, 2018; 57th Bombardment Wing, Twelfth Air Force, 2018).

Other important raids on Italy and southern France were carried out from Corsican airfields such as the bombing of Monte Cassino on February 15, 1944 (United States. Dept. of the Army, Office of Military History, 1947), of Nice and Saint-Laurent-du-Var on May 26, 1944. Corsica was also on the route of massive raids such as the one on Marseille on May 27, 1944, carried out from Brindisi. (Wikipedia Contributors, 2024).

During all bombing raids, human and material losses were enormous, and many bombers returned damaged and unable to carry out their mission. There are numerous testimonies of pilots forced to jettison their bombs at sea before returning to Corsica (Shepherd, 1996; Taddei, 2003; Shores *et al.*, 2018). These operations to offload unused and generally unprimed munitions required the designation of jettison zones off the Corsican coast, and




the two coralligenous atoll areas discovered north of Cap Corse are situated where one would expect them to be found: on the return route from raids on southern France and northern Italy (notably those on the Brenner Pass, **Fig. 3f**).



**Figure 3.** Corsica, playing the role of a giant Allied aircraft carrier during the Second World War. **a:** location of US military airfields in 1944 in Corsica (from Marcel, 2024). **b:** B17 Flying Fortress. **c:** B24 Liberator. **d:** B25 Mitchell. **e:** B26 Marauder (**b-e:** from Wikimedia commons). **f:** Location of the Brenner Pass and of the coralligenous atolls off Cap Corse, on the route of the Allied bomber squadrons (drawing © Marc Verlaque).

In the Baltic Sea where large quantities of conventional ammunition were dumped at the end and after World War II, many explosion craters were identified on the seabed (Kampmeier *et al.*, 2020; Papenmeier *et al.*, 2022; Díaz-Mendoza *et al.*, 2023) (**Fig. 4a**). In the *Posidonia oceanica* meadow of the east coast of Corsica, north of the Sulinzara airfield (between Sulinzara and U Cateraghju), approximatively north-south alignments of circular spots of dead matte have been observed (**Fig. 4b**) (Clabaut *et al.*, 2014). They may correspond to the Sulinzara (as Solenzara, **Fig. 3a**) jettison zone. Bomb impacts from World War II have also been described in *P. oceanica* meadows of Provence and French Riviera (Meinesz and Lefevre, 1984; Pergent-Martini, 1994; Pergent-Martini and Pergent, 1996; Pergent-Martini and Pasqualini, 2000) and Corsica (Clabaut *et al.*, 2014).

On the seabed of the Grande Rade of Toulon (France), Garlan *et al.* (2018) reported 661 high backscatter circular structures, with diameters ranging from 11 to 134 m, and very specific characteristics, which clearly differentiate them from other sedimentary, biological and geological figures encountered on the seabed (**Fig. 4c and 4d**).















**Figure 4. a:** explosion craters of around 25 m diameter in a post-war marine munition dumpsite (Baltic Sea; from Kampmeier *et al.*, 2020, modified). **b:** line of possible bomb impacts in the *Posidonia oceanica* meadow (East coast of Corsica, north of the Sulinzara airfield; courtesy of Gérard Pergent). **c-d:** circular seabed structures observed on multibeam echosounder images of the Grande Rade of Toulon, France (in red on the map; diameter: 11 to 134 m) (Source: Garlan *et al.* (2018). Reproduced and modified with permission from the Coastal Education and Research Foundation, Inc.). **e:** bathymetry images of a dumping areas. Insets show the typical torus shape of the structures of Class 1 (Southwestern Baltic Sea; from Diaz-Mendoza *et al.*, 2023, modified).

These circular structures are usually clearly separated but can sometimes partially overlap. They seem to be randomly arranged in terms of distribution and size, but sometimes they are arranged in a line; they are in this case series of circles with similar diameters and sharpness of outline. These sedimentary structures indicate that they have been generated by the explosion of bombs. The oldest ones most probably result from bombs dropped in 1940, and 1944 during the landing of the Allied forces.

In the Baltic Sea, Díaz-Mendoza *et al.* (2023) identified more than 3 000 circular seabed features (diameters between 6 and 77 m) corresponding to pockmarks, dumping spots and explosion craters (**Fig. 4e**). These authors propose a classification into six classes, which provides an overview of their formation mechanisms (**Table 1**). Coralligenous atolls correspond well to classes 1, 2 and 3, which all have an anthropogenic origin (i.e. detonation craters, bomb droppings and dumping of dredged materials). Like in Corsica, the circular elements of class 1 are often grouped, sometimes merged with edges of varying height and width, and often with a central elevation.

**Table 1.** Classification of circular seabed structures and schematic representation showing main features classes observed in the hydroacoustic data (Southwestern Baltic Sea; from Diaz-Mendoza *et al.*, 2023).

	Class	Description	Feature name / suspected origin	MBES /SSS Backscatter	SBP Profile
Anthropogenic features	1	Positive relief, ring-shaped features with high backscatter and in SBP internal chaotic reflectors. They are above mean seafloor depth and very often present internal elevations.	Dumping rings / Dumping of dredged material		
	2	Random high backscatter 'splotches', positive relief or slightly negative relief (depending on the hardness of the substrate).	Random dumping spots / Dumping of material		
	3	Flat areas of string of high backscatter rounded spots with no evidence of seismic acoustic anomalies.	Strings of dumping spots / Loss of dumping material		
	5	U-shaped craters without elevated rims.	Craters / ammunitions blasting		
Natural features	6	Craters without elevated rims and with gas related acoustic anomalies: acoustic turbidity, acoustic blanking, high-amplitude reflectors.	Pockmarks / Seepage activity		
Ambiguous	4	High backscatter ring-shaped features with elevated rims and vertical seismic acoustic anomalies, enhanced reflectors, velocity 'pull-up' reflectors, and in association with acoustic blanking.	No defined		

Backscatter intensity  Acoustic anomalies 

## 6. Conclusion

The various surveys carried out on the coralligenous atoll fields of Cap Corse have not yet collected any convincing geological,

hydrodynamic, biological, or current-driven elements to support the hypothesis of a natural and ancient origin.

During the most recent oceanographic survey (*Gombessa 6 Cap Corse*), the participants themselves once again highlighted the main feature that is difficult to reconcile with a natural origin: the very constant and very regular size of the atolls.


In contrast, the arguments, both structural and historical, in favour of a recent and anthropogenic origin of the coralligenous atolls, the jettisoning of unused and unprimed bombs during World War II, are numerous. It is a historical fact that, during this conflict, jettison zones were used around Corsica, and the possibility that the two fields of coralligenous atolls discovered off the coast of Cap Corse are the first two jettison zones to be located around the island deserves consideration. The data, particularly magnetometry data, collected during *Gombessa 6*, still being analyzed, will perhaps make it possible to confirm this hypothesis. If this were the case, it would be desirable, for security reasons, to seek to locate other possible jettison zones around Corsica.

Due to the originality of these formations, their high specific diversity, and a supposed ancient and natural origin, these atolls were considered, soon after being discovered, as natural monuments of high heritage value. Strong protection measures were therefore considered (e.g. ban on anchoring). One accident has already taken place in October 2018 with the collision between the ro-ro ferry *Ulysse* and the containership *CSL Virginia*, anchored 28 km north-west of the Cap Corse (Boudouresque *et al.*, 2019).

If their recent and anthropogenic origin were confirmed, would these potential protective measures become irrelevant? The answer is clearly no. Whatever the origin of the atolls, their large number and their raised structures (domes and crowns) on a sub-horizontal coastal detrital bottom create environmental conditions particularly favourable to the development of a coralligenous bank (*coralligène de plateau*) and the installation of a species-rich benthic flora and fauna, with high heritage value and high conservation priority (**Fig. 5**).

Even if their origin is not natural, these lush coralligenous formations deserve to be preserved. Furthermore, like other vestiges of World War II, such as the conning tower of the *Casabianca* submarine in Bastia (a replica of which is visible at Place Saint Nicolas) (Griffi and Preziosi, 1988; Wikipedia contributors, 2023) and the wreck of a B17 Flying Fortress in Calvi, 26 m depth (Barraqué *et al.*, 2009), the memorial aspect must also be considered.

Finally, independently of a concern for heritage conservation, if the hypothesis of a jettison zone is confirmed, a limitation of human activities (anchoring, trawling, drilling, dredging, installation of wind turbines and burying of submarine cables) should also be strongly recommended for safety reasons. Unexploded bomb dump areas are theoretically closed to navigation. For example, in the Channel/North Sea zone, there are at least three zones of this type: one near the English coast, one in the Thames estuary and the third in the Pas-de-Calais (Contributors to Wikipedia, 2023).



**Figure 5.** The recent and anthropogenic hypothesis of the formation of coralligenous atolls. **a-b:** diagrams of the formation of atolls with variable central dome in 1944. **c:** diagram of a coralligenous atoll with its rich benthic flora and fauna in 2024. (drawing © Marc Verlaque).

**Acknowledgements.** We thank Professors Christine Pergent-Martini and Gérard Pergent (University of Corsica Pasquale Paoli) for their comments and the sending of data collected during their oceanographic surveys, Professor Michèle Ferrandini (University of Corsica Pasquale Paoli) for sending photos of her thin sections of bio-concretions of coralligenous atolls, Alain Barcelo, Pr. Jacques Piazzola and Dr. Luigi Piazzoli for their suggestions and comments, the scientific journals (Continental Shelf Research – ELSEVIER, *Frontiers in Earth Science*, ISPRS International J. Geo-information, and *Journal of Coastal Research*) for permission to reproduce figures, and Michael Paul for proofreading this article.

## References

- 57th BOMB WING ASSOCIATION, 2018. *57th Bomb Wing Association Archives*. <https://57thbombwing.com/GalleryV/57thWingArchive>.
- 57th BOMBARDMENT WING, TWELFTH AIR FORCE, 2018. *Battle of the Brenner. World War II Operational Document*. Combined Arms Research Library Digital Library 113 p. <https://cgsc.contentdm.oclc.org/digital/collection/p4013coll8/search/searchterm/World%20War%20II%20Operational%20Documents/field/collec/mode/exact/conn/and>
- BALLESTEROS E., 2006. Mediterranean coralligenous assemblages: a synthesis of present knowledge. *Oceanogr. Mar. Biol.: Ann. Rev.*, 44: 123-195.
- BARRAQUÉ N., COLOMBINI H., DIRAIMONDO G., 2009. *La Corse. 100 plongées incontournables et nos autres coups de cœur. Tourisme patrimoine*. Éditions GAP, Challes-les-Eaux: 1-377.
- BONACORSI M., 2012. *Caractérisation des peuplements benthiques du Cap Corse (CapCoral 1 et 2)*. Thèse Doctorat Écologie marine, Université de Corse: 1-156.
- BONACORSI M., CLABAUT P., MARENGO M., PERGENT G., PERGENT-MARTINI C., 2011. *Cartographie des peuplements coralligènes du Cap Corse - Rapport de mission CAPCORAL 2, 27 août - 21 septembre 2011*. Contrat Agence des Aires Marines Protégées / GIS Posidonie : 1- 21.
- BONACORSI M., PERGENT-MARTINI C., CLABAUT P., PERGENT G., 2012a. *Cartographie des peuplements coralligènes du Cap Corse - Rapport CAPCORAL (CapCoral 1 et 2)*. Contrat Agence des Aires Marines Protégées / GIS Posidonie : 1- 120.
- BONACORSI M., PERGENT-MARTINI C., CLABAUT P., PERGENT G., 2012b. Coralligenous "atolls": discovery of a new morphotype in the Western Mediterranean Sea. *Comptes Rendus Biologies*, 335 (10-11) : 668-672.
- BONACORSI M., PERGENT-MARTINI C., CLABAUT P., DAMIER E., DANIEL B., BREAND N., MARENGO M., PERGENT G., 2013. Caractérisation des peuplements benthiques du Cap Corse. *Colloque Carhamb'ar, CARTographie des HAbitats Marins Benthiques: de l'Acquisition à la Restitution*, 2ème édition, Brest, 26 au 28 Mars 2013 : 84-88.
- BONACORSI M., ALAMI S., BREAND N., CLABAUT P., DANIEL B., PERGENT G., PERGENT-MARTINI C., 2014. Cartography of main coastal ecosystems (coralligenous and rhodolith beds) along the Corsican coasts. In: *Proceeding of the 2nd Mediterranean Symposium on the conservation of coralligenous & other calcareous bio-concretions* (Portoro, Slovenia, 29-30 October 2014), Langar H., Bouafif C., Ouerghi A. (eds.), RAC/SPA publ., Tunis: 37-42.
- BOUDOURESQUE C.F., 2003. The erosion of Mediterranean biodiversity. In: *The Mediterranean Sea: an overview of its present state and plans for future protection*, Rodríguez-Prieto C., Pardini G. (eds.), Servei de Publicacions de la Universitat de Girona: 53-112.
- BOUDOURESQUE C.F., BLANFUNE A., HARMELIN-VIVIEN M., PERSONNIC S., RUITTON S., THIBAUT T., VERLAQUE M., 2017. Where seaweed forests meet animal forests: the examples of macroalgae in coral reefs and the Mediterranean coralligenous ecosystem. In: *Marine animal forests*. Rossi S., Bramanti L., Gori.
- BOUDOURESQUE C.F., BLANFUNE A., MARTIN G., G., PERRET-BOUDOURESQUE M., TAUPIER-LETAGE I., 2019. The Virginia oil spill in Provence: a tale of inappropriate over-cleaning. *Rapp. Comm. internation. Mer Médit.*, 42: 100.
- CAPPS TUNWELL D., PASSMORE D.G., HARRISON S., 2016. Second World War bomb craters and the archaeology of Allied air attacks in the forests of the Normandie-Maine National Park, NW France. *J. Field Archaeol.*, 41 (3) : 312-330.
- CLABAUT P., AUGRIS C., PERGENT G., PERGENT-MARTINI C., PASQUALINI V., BONACORSI M., 2014. *Les fonds marins côtiers de Corse. Cartographie biomorphosédimentaire*. Éditions Quae, Versailles. 25 feuilles, échelle 1/20 000, livret d'accompagnement: 1-80.
- COLLINA-GIRARD J., 2012. *La Provence immergée. Plongées à Marseille et ses abords*. Éditions les Presses du Midi, Toulon: 1-282.

- CONTRIBUTORS TO WIKIPEDIA, 2023. *Munition immergée*, Wikipédia, l'encyclopédie libre, [https://fr.wikipedia.org/w/index.php?title=Munition\\_immerg%C3%A9e&oldid=210002434](https://fr.wikipedia.org/w/index.php?title=Munition_immerg%C3%A9e&oldid=210002434) (accessed on March 21, 2024).
- CONTRIBUTORS TO WIKIPEDIA, 2024. *Bombardement du 26 mai 1944*, Wikipédia, l'encyclopédie libre, 15 janvier 2024, 22:20 UTC, <[https://fr.wikipedia.org/w/index.php?title=Bombardement\\_du\\_26\\_mai\\_1944&oldid=211546003](https://fr.wikipedia.org/w/index.php?title=Bombardement_du_26_mai_1944&oldid=211546003)> [accessed on January24]
- DENIZOT M., GUELORGET O., MASSIEUX M., PERTHUISOT J.P., 1981. Une remarquable construction récifale à Mélobésiée dans une lagune sursalée du Sud-Est tunisien (la Bahiret el Biban). *Cryptog.-Algol.*, 2 (4): 253-266.
- DETER J., HOLON F., MOUILLOT D., HOCDE R., 2021. Gombessa 6 "Cap Corse" cruise: CTD profiles in Corsica, NW Mediterranean Sea, July 2021. *SEANOE*, <https://doi.org/10.17882/89637>.
- DETER J., BALLESTA L., MASSEY J.-L., GUILBERT A., HOLON F., DELARUELLE G., BLANDIN A., MARRE G., RAUBY T., MOUILLOT D., VELEZ L., DEJEAN T., DI IORIO L., HOCDE R., FURFARO G., JORRY S., BOUCHETTE F., FERRIER-PAGES C., CHARRIERE A., IZART A., FERRANDINI M., PLUQUET F., PERGENT G., PERGENT-MARTINI C., LUONGO G., DANOVARO G., BOISSERY P., CANCEMI M., 2022. State of knowledge on the deep coralligenous rings of Cape Corsica following the scientific expedition Gombessa 6 (2021). In: BOUAFIF C., OUERGHI A. (eds), 4th Mediterranean Symposium on the conservation of coralligenous & other calcareous bio-concretions, Genoa, Italy, 20-21 September 2022 SPA/RAC publ., Tunis: 45-50.
- DÍAZ-MENDOZA G. A., KRÄMER K., VON RÖNN G. A., SCHWARZER K., HEINRICH C., REIMERS H. C., WINTER C., 2023. Circular structures on the seabed: Differentiating between natural and anthropogenic origins-Examples from the Southwestern Baltic Sea. *Frontiers in Earth Science*, 11, 1170787.
- DIMITROV L., WOODSIDE J., 2003. Deep sea pockmark environments in the Eastern Mediterranean. *Mar. Geol.* 195: 263–276. doi:10.1016/S0025-3227(02)00692-8
- DOLEJŠ, M., PACINA, J., VESELY, M., BRÉTT, D., 2020. Aerial bombing crater identification: exploitation of precise digital terrain models. *ISPRS International J. Geo-information*, 9 (713):1-15.
- FERRANTI M., 2013. *Corse 1943: Les combattants de la liberté*. Albiana publ., Ajaccio: 1-170.
- FOO S.A., BYRNE M., RICEVUTO E., GAMBI M.C., 2018. The carbon dioxide vents of Ischia, Italy, a natural system to assess impacts of ocean acidification on marine ecosystems: an overview of research and comparisons with other vent systems. *Oceanogr. Mar. Biol.: Annual Rev.*, 56: 237-310.
- FOUQUET Y., AUZENDE J. M., BALLU V., BATIZA R., BIDEAU D., CORMIER M.H., GEISTDOERFER P., LAGABRIELLE Y., SINTON J., SPADEA P., 1994. Variabilité des manifestations hydrothermales actuelles le long d'une dorsale ultra rapide. Dorsale Est Pacifique entre 17° et 19° S (campagne NAUDUR). *C.R. Acad. Sci. Sér II*, 319 (11): 1399-1406.
- GAMBIEZ F., 1973. *Libération de la Corse*. Hachette Littérature publ., Paris: 1-318.
- GARLAN T., MATHIAS X., BRENON E., FAVRETTO-CRISTINI N., DESCHAMPS A., BEUCLER E., GUYOMARD P., MORIO O., 2018. Circular sedimentary figures of anthropic origin in a sediment stability context. *Journal of Coastal Research*, Special Issue 85: 411-415.
- GENNARO P., PIAZZI L., CECCHI E., MONTEFALCONE M., MORRI C., BIANCHI C.N., 2020. *Monitoraggio e valutazione dello stato ecologico dell'habitat a coralligeno. Il coralligeno du parete*. ISPRA publ., Roma: 1-64.
- GRANFIELD A., 2011. *Bombers over sand and snow: 205 Group RAF in World War II*, Pen & Sword Aviation Publ.: 1-384.
- GREGORI S., 2023. *Nouvelle histoire de la Résistance en Corse. Juillet 1940 – Septembre 1943. Forti saremu se saremu uniti*. Volumes I and II. Éditions Alain Piazzola, Ajaccio: 1-1034.
- GRIFFI T., PREZIOSI L., 1988. *Première mission en Corse occupée : avec le sous-marin Casabianca (décembre 1942-mars 1943)*. Éditions L'Harmattan, Paris: 1-190.
- GRIGGS A.L. (ed.), 2015. *Flying flak alley: personal accounts of World War II Bomber Crew Combat*. McFarland & Company publ., Jefferson: 1-267.
- KAMPMEIER M., VAN DER LEE E. M., WICHERT U., GREINERT J., 2020. Exploration of the munition dumpsite Kolberger Heide in Kiel Bay, Germany: Example for a standardised hydroacoustic and optic monitoring approach. *Continental Shelf Research*, 198, 104108: 1-21.
- KAWASE H., OKATA Y., ITO K., 2013. Role of huge geometric circular structures in the reproduction of a marine pufferfish. *Scientific reports*, 3(1): 1-5.
- LANGAR H., BESSIBES M., DJELLOULI A., PERGENT-MARTINI C., PERGENT G., 2011. The *Neogoniolithon brassica-florida* (Harvey) Setchell & LR Mason (1943) reef of Bahiret el Bibane Lagoon (Southeastern Tunisia). *J. Coast. Res.*, 27 (2): 394-398.

- LEY W, SCHAEFER H., 1941. How About Penetration Bombs? *United States Naval Institute Proceedings*, 67 (12): 466.  
<https://www.usni.org/magazines/proceedings/1941/december/how-about-penetration-bombs>
- LICHTER M., ZVIELY D., KLEIN M., SIVAN D., 2010. Sea-level changes in the Mediterranean: past, present and future – A review. *In: Seaweeds and their role in globally changing environments*. Israel A., Einav R., Seckbach J. (eds.), Springer publ., Dordrecht: 5-17.
- MARCEL, 2024. U.S.S. Corsica, *Avions légendaires*,  
<https://www.avionslegendaires.net/reportage/uss-corsica-lautre-ile-porte-avions-de-mediterranee>
- MEINESZ A., LEFEVRE J.R., 1984. Régénération d'un herbier à *Posidonia oceanica* quarante années après sa destruction par une bombe dans la rade de Villefranche (Alpes Maritimes). *In: International Workshop on Posidonia oceanica beds*, Boudouresque C.F., Jeudy de Grissac A., Olivier J. (eds.), GIS Posidonie publ., Marseille, 1: 39-44.
- NELSON C.H., JOHNSON K.R., BARBER J.H., 1987. Gray whale and walrus feeding excavation on the Bering shelf, Alaska. *Sediment. Petrol.* 57: 419-430. doi:10.1306/212F8B4D-2B24-11D7-8648000102C1865D
- PAPENMEIER S., DARR A., FELDENS P., 2022. Geomorphological data from detonation craters in the Fehmarn Belt, German Baltic Sea. *Data*, 7 (63): 1-10.  
<https://doi.org/10.3390/data7050063>
- PERGENT G., BESSIBES M., DJELLOULI A., EL ABED A., LANGAR H., MRABET R., PERGENT-MARTINI C., 2009. Le récif à *Neogoniolithon brassica-florida* de la lagune des Bibans (Tunisie). *Proceedings of the 1<sup>st</sup> symposium on the coralligenous and other calcareous bio-concretions of the Mediterranean Sea*, Tabarka, 15-16 January 2009, Pergent-Marini C., Bricchet M (eds.), RAC/SPA publ., Tunis: 117-122.
- PERGENT G., PERGENT-MARTINI C., CLABAUT P., BONACORSI M., SARTORETTO S., 2015. Les campagnes océanographiques en Corse: De la cartographie des habitats-clés du littoral à la découverte des atolls. Poster - *Colloque de Restitution Commission Nationale de la Flotte Côtière*, Talence, 11-12 juin 2015.
- PERGENT-MARTINI C., 1994. *Impact d'un rejet d'eaux usées urbaines sur l'herbier à Posidonia oceanica, avant et après la mise en service d'une station d'épuration*. Doctoral thesis, University of Corsica: 1-191.
- PERGENT-MARTINI C., PASQUALINI V., 2000. Seagrass population dynamics before and after the setting up of a wastewater treatment plant. *Biol. Mar. Medit.*, 7 (2): 405-408.
- PERGENT-MARTINI C., PERGENT G., 1996. Spatio-temporal dynamics of *Posidonia oceanica* beds near a sewage outfall (Mediterranean - France). *In: Seagrass biology: proceedings of an international workshop*, Kuo J., Phillips R.C., Walker D.I., Kirkman H. (eds.), Australia: 299-306.
- PERGENT-MARTINI C., ALAMI S., BONACORSI M., CLABAUT P., PERGENT G., 2014a. *Cartographie des peuplements benthiques du littoral de la Corse – Rapport CoralCorse*. Université de Corse & GIS Posidonie, Equipe EqEL: 1-79.
- PERGENT-MARTINI C., ALAMI S., BONACORSI M., CLABAUT P., DANIEL B., RUITTON, S., SARTORETTO S., PERGENT G., 2014b. New data concerning the coralligenous atolls of Cap Corse: an attempt to shed light on their origin. *Proceeding of the 2<sup>nd</sup> Mediterranean Symposium on the conservation of coralligenous & other calcareous bio-concretions* (Portoro, Slovenia, 29-30 October 2014), Langar H., Bouafif C., Ouerghi A. (eds.), RAC/SPA publ., Tunis: 129- 134.
- PRADEL J., VANRELL L., 2008. *St-Éxupéry. L'ultime secret. Enquête sur une disparition*. Éditions du Rocher, Monaco: 1-189.
- SARTORETTO S., FRANCOUR P., 1997. Quantification of bioerosion by *Sphaerechinus granularis* on 'coralligène' concretions of the western Mediterranean. *J. Mar. Biol. Ass. U.K.*, 77: 565-568.
- SHEPHERD D.W., 1996. *Of men and wings: the first 100 missions of the 449th Bombardment Group (January 1944 - July 1944), Fifteenth Air Force, World War II* – Based upon the Wartime Diary of C.A. Shepherd, 718th Squadron Norfield, Pub; First Edition: 1-343.
- SHORES C., MASSIMELLO G., GUEST R., OLYNYK F., BOCK W., THOMAS A., 2018. *A history of the Mediterranean Air War, 1940-1945. vol. 4: Sicily and Italy to the fall of Rome 14 May 1943 – 5 June 1944*, Casemate Publishers: 1-680.
- TADDEI D., 2003. *Décembre 1944 – Avril 1945, USS Corsica, l'île porte-avions*. Albania publ., Ajaccio, 233 p.
- TEIXIDÓ N., GAMBI M.C., PARRAVACINI V., KROEKER K., MICHELI F., VILLÉGER S., BALLESTEROS E., 2018. Functional biodiversity loss along natural CO<sub>2</sub> gradients. *Nature Communications*, 9 (5149): 1-9.
- UNITED STATES. DEPT. OF THE ARMY. OFFICE OF MILITARY HISTORY, 1947. The bombing of Cassino in: *United States Army in World War II.: The Mediterranean.*: 1-441.



- VAUDOIT H., CASTELLANO P., ROSENFELD A., 2004. *Saint-Ex, la fin du mystère*. Éditions Filipacchi – La Provence, Marseille : 1-224.
- WAGA J., SZYPUŁA B.J., FAJER M.J., 2022. Heritage of war: Analysis of bomb craters using Lidar (Kędzierzyn-Koźle, Poland). *Internat. J. Conserv. Sci.*, 13 (2): 593-608.
- WARWICK R.M., 2022. Seagrass 'fairy circles' on the Isles of Scilly. *J. Mar. Biol. Association U. K.*, 102(1-2): 63-68.
- WIKIPEDIA CONTRIBUTORS, 2023. *French submarine Casabianca (1935)*. Wikipedia, the free encyclopedia. Accessed on 19 April 2024.