



# Assimilation of radar freeboard and snow altimetry observations in the Arctic and Antarctic with a coupled ocean/sea ice modelling system

5 Alette Chenal<sup>1,2</sup>, Gilles Garric<sup>1</sup>, Charles-Emmanuel Testut<sup>1</sup>, Mathieu Hamon<sup>1</sup>, Giovanni Ruggiero<sup>1</sup>,  
Florent Garnier<sup>3</sup>, Pierre-Yves Le Traon<sup>1,4</sup>

<sup>1</sup> Mercator Océan International, Toulouse, 31400, France

<sup>2</sup> CNES, Toulouse, 31400, France

<sup>3</sup> Ministère de la Défense, Paris, 75007, France

<sup>4</sup> Ifremer, Plouzané, 29280, France

10

*Correspondence to:* Alette Chenal ([achenal@mercator-ocean.fr](mailto:achenal@mercator-ocean.fr)) and Gilles Garric ([ggarric@mercator-ocean.fr](mailto:ggarric@mercator-ocean.fr)).

**Abstract.** Sea ice and snow volume are essential variables for polar predictions, but operational systems still struggle to accurately capture their evolution. Satellite measurements now provide estimates of sea ice freeboard and snow depth. The combined assimilation of sea ice concentration (SIC), along-track altimetry radar freeboard data from Cryosat-2 and observations of snow depth from Cryosat-2 and SARAL is implemented in a multivariate approach in a global  $\frac{1}{4}^\circ$  ocean/sea ice coupled NEMO4.2/SI3 model. A multivariate experiment, performed on two full seasonal cycles 2017–2018, is compared to a free (no assimilation) and a SIC-only assimilation simulations. The multivariate technique increases the sea ice volume, even in the absence of freeboard and snow measurements during summer, and rapidly changes the spatial patterns of ice and snow thicknesses in both hemispheres, in accordance with the assimilated observations. The sea ice volume from the multivariate approach compares better with independent (not assimilated) estimates from IceSat-2 and CS2SMOS or SMOS in both hemispheres. The multivariate system performs better in the Arctic than in Antarctica where the ice and ocean separate analyses seem not designed to consider the strong interactions between upper oceanic layers and sea ice cover in the Southern Ocean and to prevent localised degradations. These results also confirm the importance of using variable snow and ice densities in a freeboard assimilation context. This study shows promising results for enhancing the capacity of assimilation systems to monitor the volume of sea ice and snow and paves the way for future satellite missions.

15  
20  
25

## 1 Introduction

In response to climate change, Arctic sea ice is continuing to decline and is regularly breaking historically low records, and, more recently, the entire year of 2023 showed the lowest sea ice extent in Antarctica ever seen in the satellite record (Gilbert and Holmes, 2024). October 2020 was the lowest end-of-summer sea ice volume since 2010 in the Arctic (Perovich et al., 2020). Given the rapid transformations affecting sea ice due to climate change, sea ice monitoring is of the utmost

30



importance. Assimilation techniques allow us to combine models and observations to improve our ability to monitor the ocean and sea ice state. Sea ice concentration (SIC) is currently assimilated in most sea ice data assimilation systems using different methods: nudging, Kalman filter variants, or 3DVAR variants (Uotila et al., 2019). However, one of the challenges in assimilating SIC is to extend the SIC information to other prognostic model variables such as sea ice thickness (SIT). Tietsche et al. (2013) concluded that in their Arctic model configuration, a proportional relationship between SIT and the SIC update was most effective for adjusting the modelled SIT. Massonnet et al. (2015) and Kimmritz et al. (2018) used the model covariances with a multivariate Ensemble Kalman Filter (EnKF) to update different sea ice variables, propagating the information from the observed SIC to the unobserved variables. Experiments have used EnKF or variations of this multivariate scheme with multidata frameworks: both SIC and SIT products have been assimilated in the Arctic (e.g. Cheng et al., 2023; Williams et al., 2023; Chen et al., 2024). The assimilation methods can vary, but the assimilated SIT products are usually thin SIT from the European Space Agency's (ESA) Soil Moisture and Ocean Salinity (SMOS) mission, thick SIT measured by the ESA satellite mission CryoSat-2 (CS2), with two processing techniques available (Ricker et al., 2014 or Kurtz and Harbeck, 2017), or an observational product that statistically combines information from the two (CS2SMOS, Ricker et al., 2017).

Xie et al. (2016) found that assimilating SMOS thin SIT data had significant benefits for SIC and SIT modelling in some regions near the ice edge. Mu et al. (2018) combined the use of both SMOS thin SIT and CS2 SIT product in their assimilation system and obtained better results than the observation-only CS2SMOS product, demonstrating the added value of the model dynamics. The assimilation of CS2SMOS merged product (Xie et al., 2018) reduced model biases compared to the assimilated data, and results were in better agreement with independent datasets, with no degradation of other sea ice variables. Fritzner et al. (2019) compared the assimilation of SIC combined separately with either CS2 SIT, SMOS SIT, or a snow thickness (SNT) dataset in a short simulation and concluded that CS2 SIT provides the best long-term model improvements compared to SMOS SIT. They also found that SNT assimilation had a weaker effect on the model than SIT assimilation. Other teams methods updated SIT in the Arctic with nudging (Fritzner et al., 2018; Blockley and Peterson, 2018; Balan-Sarojini et al., 2021), with ensemble optimal interpolation (Lee and Ham, 2022, 2023), and with an enthalpy-adjusting scheme to ensure a consistent update of all sea ice variables (Liu et al., 2024). These numerous studies highlight that sea ice assimilation remains an active and evolving research area. The absence of a clear consensus on the optimal method reflects the complexity of balancing model uncertainties, data availability, and computational efficiency to achieve the best possible agreement with observations.

Mu et al., (2020) and Cipollone et al., (2023) implemented multidata and multivariate sea ice assimilation in global configurations, but with Arctic-only CS2, SMOS, and CS2SMOS SIT products. They both found their experiments to agree with in-situ data. Luo et al., (2021) implemented a multivariate assimilation system in Antarctica and successfully assimilated SIC and SMOS SIT. They had to inflate their atmospheric ensemble forcing, even though it was unnecessary in a similar Arctic assimilation scheme, suggesting that differences in the impact of sea-ice data assimilation between the two poles. They stated that the implementation of Arctic sea-ice data assimilation cannot be simply extended to the Antarctic.



SIT can be retrieved from altimeter radar freeboard (RFB) measurements by using hydrostatic equilibrium and taking  
65 into account the height of the snow penetrated by the radar wave, a medium where the radar velocity is modified (Garnier et  
al., 2022). The sea water, ice and snow densities and the snow depth above the ice are required for the RFB-SIT conversion,  
and the assumptions made on these variables result in a significant uncertainty in the sea ice volume products (Kern et al.,  
2015; Kwok and Cunningham, 2015). The snow layer accounts for most of the uncertainty in the calculation of SIT from RFB  
(Garnier et al., 2021). The CS2 SIT products mentioned above use the Warren 99 (W99) snow climatology (Warren et al.,  
70 1999) or a modified version of it which is now known to be outdated and unreliable on most regions of the Arctic (Kern et al.,  
2015). Fiedler et al. (2022) is the first study to use the along-track CS2 RFB data in the Arctic, and to convert it into SIT using  
the modelled snow cover prior to the assimilation step. Their study results in a general improvement of the modelled SIT, with,  
in particular, a bias reduction in the Canadian Basin. This improvement extends into summer period, when no data is  
assimilated. However, they noted no substantial improvement in the Beaufort region due to a degradation of ice thicknesses  
75 below 1 m. Mignac et al. (2022) performed the same experiment, adding the SMOS SIT data to the along-tracks SIT computed  
from CS2 RFB and modelled snow, arguing that the SMOS SIT product performs better in thin ice areas of the Arctic. The  
thin SMOS ice assimilation was able to counteract the SIT overestimation that happens in the Arctic marginal seas when  
assimilating only CS2 products.

Other sources of uncertainty in the RFB-SIT conversion stems from the choice of ice and snow densities. The NEMO  
80 model uses constant snow and ice densities, whereas the observation products usually parametrize the ice density depending  
on the ice type (multi-year ice MYI, or first-year ice FYI, see Alexandrov et al., 2010) in the Arctic and on the season (see  
Kurtz and Markus, 2012) in the Antarctic. The choice of snow density varies in different SIT retrievals from RFB  
measurements, including options such as constant density, seasonally varying density, climatology-based density, or modelled  
density. Kern et al. (2015) stated the importance of having well calibrated density for the ice and they recommend using  
85 seasonally varying snow density instead of a constant. Positive model biases in sea ice volume compared to satellite altimetry  
estimates have been attributed mainly to ice density differences (Bocquet et al., 2024). New efforts are currently being made  
to get fresh measurements of sea ice densities: Jutila et al. (2022) measured ice densities on average higher than the values  
from Alexandrov et al. (2010) for both the FYI and MYI, resulting in 12.4 % and 16.7 % larger sea ice thickness values for  
FYI and MYI.

90 Knowing the large uncertainty associated with the sea ice volume products derived from RFB measurements, Sievers  
et al. (2023) directly assimilated the radar freeboard in the Arctic. In their assimilation scheme, they used a varying density for  
the ice, set as a function of the modelled salinity of the ice, and a linearly varying snow density depending on the season,  
following Mallett et al. (2020). The densities were not modified in the sea-ice model physics. They used the modelled snow to  
convert the freeboard to ice thickness and they updated sea ice concentration and sea ice thickness through data assimilation.  
95 They compared the resulting sea ice thickness with in-situ data, showing improvements in some regions of the Arctic and  
degradation in others, using a simulation without assimilation and another with assimilation of sea ice concentration only as  
references.



Our work is in line with that of Sievers et al. (2023). However, we decided to assimilate RFB together with snow thickness observations to update the snow in addition to the sea ice variables at a global scale, i.e. including the Arctic and Antarctica. Moreover, we kept a coherent parametrization between the assimilation scheme and the sea ice model, so we used the model fixed snow and ice densities.

In this study, we use the operational Kalman filter scheme deployed in the production of global reanalysis and forecast at Mercator Ocean to implement a multivariate sea ice assimilation scheme with sea ice concentration (SIC), sea ice volume (SIV) and snow volume (SNV). In contrast to the usual ice assimilation where the SIC model variable (univariate) is updated using SIC observations (monodata), this approach aims to assimilate along-track radar freeboard and altimetric snow depth observations in addition to the SIC observations (multidata) and to update SIC, SIV and SNV model variables (multivariate). We use the same assimilation method for the Arctic and Antarctic. We aim to provide first answers to the following scientific questions:

- Does the multivariate/multidata approach provide added value over the widespread univariate/monodata method? What are the impacts of using altimetric radar freeboard and altimetric snow observations in addition to the SSMIS SIC data?
- Are the current parametrizations in sea ice models sufficient for accurate assimilation of radar freeboard and snow measurements?
- What challenges arise when applying the same sea ice assimilation scheme to both the Arctic and Antarctic, given their differing physical environments and ice dynamics?

We describe the modelling and assimilation components, the data assimilated in the analysis system, and the experimental design in Section 2. Section 3 focuses on the performances of the assimilation setup while section 4 presents a comparison with independent satellite observations. Section 5 discusses the main results and conclusions are given in section 6.

## 2 Analysis system and experimental design

### 2.1 Modelling and assimilation system

#### 2.1.1 Global ice-ocean coupled model configuration

We use the ocean/sea ice coupled model Nucleus for European Modelling of the Ocean (NEMO) version 4.2 (Madec et al., 2022), coupled to the Sea Ice modelling Integrated Initiative (SI3, Vancoppenolle et al., 2023). Simulations are run on a ¼ degree tripolar horizontal grid (Madec and Imbard, 1996) with 75 oceanic vertical levels. The atmospheric forcing is the European Centre for Medium-Range Weather Forecasts (ECMWF) ERA5 atmospheric reanalysis (Hersbach et al., 2020) with a 1h frequency

The sea ice model SI3 describes the ice and snow behaviour with assumptions that for dynamics, ice is a non-newtonian 2D continuum, whereas for thermodynamics, it is a mushy layer covered by snow. Subgrid variability is represented through 11 sea ice thickness categories, with fixed boundaries. Global prognostic variables in SI3 are the sea ice velocity  $\mathbf{u}$  and its stress



130 tensor  $\sigma$ , and quantities computed in each thickness category: sea ice concentration, sea ice and snow volume per unit area, sea ice and snow enthalpy per unit area, and sea ice salt content. The model uses constant densities for the sea water, sea ice and snow with respective values of 1026, 917 and 330 kg/m<sup>3</sup>. Snow exclusively comes from the solid precipitations of the atmospheric forcing and disappears either by melting processes or by snow-ice conversion when the snow base gets below the sea level. In this study, we use the adaptative elastic-viscous plastic rheology and a parametrization to represent landfast sea ice. The ice model component is called every 3 ocean timesteps, that is, every 30 minutes.

### 135 2.1.2 Assimilation scheme

The assimilation system is the one used in the current near real time operational system (Lellouche et al., 2021). The 7-day assimilation cycle proceeds as follows: firstly, the model runs for the full cycle length for a ‘forecast’ trajectory, resulting in a forecast state. Observations available during the cycle time are loaded and processed as needed, with special care taken to define the observation errors. Using the forecast output and an observation operator, model variables are transformed into observation-equivalent variables that are consistent in space and time with the assimilated observations. Then, the analysis step produces 4D increments or model updates of the forecast trajectory. The increment depends on the innovation (observation minus model equivalent), weighted by the Kalman gain. We use a reduced-order Kalman filter derived from a singular evolutive extended Kalman (SEEK) filter (Brasseur and Verron, 2006; Lellouche et al., 2021). The Kalman gain is meant to balance the information from the model and the observations to get closer the real ocean and sea ice state: as such, it is based on the error covariance of the forecast and the observation errors. The model forecast error covariance is computed from a fixed ensemble of 4D ocean and ice state anomalies that vary seasonally.

The anomalies are computed from a long simulation without assimilation, using the same model configuration and parameters with respect to a running mean. Anomalies are computed on a reduced grid for the ocean (1 out of 2 points) and on a full grid for the sea ice. The increments at each model grid point are calculated independently in a local scheme, where a localization algorithm controls the spatial influence of observations. This approach helps to limit the impact of sampling noise on the increments. The last step of the assimilation cycle is the Incremental Analysis Update (IAU) that allows us to gradually introduce the analysis increments into the model (Benkiran and Greiner, 2008). The model runs a second time over the 7-day cycle for a ‘best’ trajectory; and at each timestep a tendency term is added to the model variables in the prognostic equations. The tendency term comes from the increment, modulated by a distribution function (Lellouche et al., 2013).

155 The ice and ocean analysis are separate, which means that ocean covariances are used for the ocean variables only, and the same applies for sea ice variables. The ocean analysis is multivariate and multidata, using sea level anomaly datasets from satellite altimetry (SEALEVEL\_GLO\_PHY\_L3\_NRT\_008\_044, 2023), sea surface temperature (SST) from OSTIA (Operational Sea Surface Temperature and Sea Ice Analysis, SST\_GLO\_SST\_L4\_NRT\_OBSERVATIONS\_010\_001, 2023), and temperature and salinity vertical profiles from in situ ARMOR and CORA-REP measurements (INSITU\_GLO\_PHYBGCWAV\_DISCRETE\_MYNRT\_013\_030, 2024). The ocean observations are not assimilated under the sea ice; except for the SST OSTIA because the product is calibrated with our assimilated SIC product and sets the under-



ice ocean surface temperature to the freezing point, which is consistent with the ice assimilation. Additionally, given the scarcity of in-situ data in the Southern polar ocean, no in-situ vertical profiles are assimilated below 60°S.

Two different methods are used for the ice assimilation: univariate and multivariate. They refer to the number of variables in the Kalman filter state vector, determining for which variables increments are calculated. In the univariate configuration, only a SIC increment is created, which means that only SIC observations can be assimilated. In the multivariate configuration, the state vector is made of sea ice concentration SIC, sea ice volume SIV, snow volume SNV, radar freeboard volume RFBV, and snow thickness SNT. This multivariate configuration allows us to assimilate a larger variety of data and to update the modelled ice accordingly. It is not required to use observational data on each of the state vector variables: when no data are given, the Kalman filter uses the model covariances to propagate the information from the observed variables to the unobserved ones. RFBV and SNT variables are included in the state vector due to the availability of observation datasets for these quantities. SIV and SNV are included because they are global prognostic variables of the ice model, essential for accurately describing the model state. SIC is included for both reasons.

The different variables updated in the sea ice assimilation cycle are listed in Table 1. The increments do not distinguish ice categories, they present total values aggregated over each grid cell. All increments are tempered by the IAU factor. The first updated model variable is the SIC. The analysis is created by adding the increment to the forecast:  $SIC_a = SIC_f + SIC_{inc}$ . Then, the total ice concentration is redistributed into each existing thickness category using a Gamma distribution, which adds most of the increment to the middle thickness categories and less to the extreme categories.

In the univariate system, all other updates are computed from this SIC increment: following Tietsche et al. (2013), the SIV is proportional to the sea ice concentration, with a constant varying depending on the hemisphere:  $h_{SH}^* = 1m$  and  $h_{NH}^* = 2m$ . The SNV increment is set to zero in the univariate method. In the multivariate method, SIV and SNV increments come directly from the Kalman filter algorithm. The algorithm updates the total ice and snow volumes for each grid cell, and then redistributes the updates to the individual ice categories. For the SIV, the algorithm adjusts the SIT in each category, starting with the thinnest ice. This prioritizes melting thinner ice first when the ice volume increment is negative. Changes are applied proportionally to the analysis SIC in each category, ensuring larger changes in categories with greater ice surface area. The SNV update accounts for the forecast SNT, analysis SIC, and SNV increment. When the SNV increment is zero, corrections are still applied, aiming at maintaining a constant SNV even under varying SIC conditions. Redistribution preserves the snow distribution across ice categories by adjusting the forecast SNT by the same ratio in each category. If the updated SNT exceeds a threshold defined as half the analysis SIT, it is capped to avoid unrealistic values. In such cases, the total snow volume may decrease compared to the forecast.

Then, the volumetric ice salinity and enthalpy are corrected in both methods by adjusting the previous ice salinity and enthalpy to the new ice volume  $SIV_a = SIV_f + SIV_{inc}$ . The volumetric snow enthalpy is also corrected following the same procedure. The updated volumetric ice salinity and enthalpy and the volumetric snow enthalpy are used to compute the ice salinity vertical profile, the salt mass content, and the snow and ice vertical temperature profile.

195



Updated variable	Univariate method	Multivariate method
SIC	Increment	Increment
SIV	$SIV_{inc} = h^* \times SIC_{inc}$	Increment
SNV	$SNV_{inc} = 0$	Increment
Volumetric ice salinity	Computed from $SIV_{inc}$ and forecast value.	
Volumetric ice enthalpy	Computed from $SIV_{inc}$ and forecast value.	
Volumetric snow enthalpy	No update	Computed from $SNV_{inc}$ and forecast value.

**Table 1: Variables updated during the assimilation cycle and their origin in both the univariate and multivariate systems.**

The different experiments presented in this paper show the evolution of the sea ice assimilation methods from a univariate and mono-data system, updating only SIC, to a multivariate and multidata setup.

## 2.2 Assimilated observing network

### 200 2.2.1 Sea ice concentration SSMIS

The observation data used for sea ice concentration (SIC) assimilation is the global daily reprocessed passive microwave dataset from the European Organization for the Exploitation of Meteorological Satellites (EUMETSAT) Ocean and Sea Ice Satellite Application Facility (OSISAF) OSI-450 (OSI SAF, 2022). Considering the large errors in satellite measurement in low SIC regions (Ivanova et al., 2015), we arbitrarily set to 0 the data values below 7.5%. Moreover, we only consider nominal data from the OSISAF algorithm, excluding data with coastal correction, interpolation, or climatology corrections. We use the spatial pattern of the “standard\_error” provided with the dataset to construct the observation error for the assimilation but we inflate linearly the error to obtain a maximum of 25% in the Arctic (same value as Lellouche et al., 2021) and 40% in Antarctica, and we set a minimum value of the error to 1%.

### 2.2.2 Radar freeboard RFB-LEGOS

210 The “laboratoire d’etudes en géophysique et océanographie spatiales” (LEGOS) scientists have used along tracks measurement from the CS2 satellite to create a freeboard dataset (Guerreiro et al., 2017; Laforge et al., 2021). Thanks to hydrostatic equilibrium, freeboard can provide sea ice thickness values using information of snow depth, and water, ice and snow densities. Altimetry measurements detect radar freeboard (RFB) due to the slower velocity of the radar wave when travelling through the snow (see equations in Bocquet et al., 2023). Radar freeboard values can be negative because of the term accounting for the radar speed reduction in the snow layer: it is not a real physical distance contrarily to ice freeboard.

215 We multiply the RFB values by the SSMIS data to assimilate radar freeboard volume per unit area (RFBV) in consistency with volumetric prognostic model quantities. We use the uncertainty provided with the dataset as the observation error,



constraining it to a range of 0.01 m to 5 m. The RFBV model equivalent is calculated from Bocquet et al. (2023) with constant sea water, sea ice and snow densities (Eq. 1).

$$220 \quad RFB = \frac{\rho_w - \rho_{ice}}{\rho_w} \cdot SIT - \left( \frac{\rho_{snow}}{\rho_w} + (1 + 0.00051\rho_{snow})^{1.5} - 1 \right) \cdot SNT = 0.106 \cdot SIT - 0.584 \cdot SNT \quad (1)$$

We use the LEGOS data because it provides concomitant RFB and snow data in both hemispheres. We assimilate two modes of CS2 instruments: the Synthetic Aperture Radar (SAR) for offshore regions and SAR Interferometric (SARin) for coastal areas. Due to potential truncation problems with the filtering of RFB measurements, and to be able to use the same method across different spatial resolutions of the configuration, we kept the full scales of SAR and SARin measurements. The data are only available during winter in both hemispheres. CS2 satellite tracks cover the entire ice domain of each hemisphere in about a month: during each assimilation cycle, important areas remain unobserved, especially at lower latitudes (Antarctica).

### 2.2.3 Snow thickness SNOW-KaKu

Snow thickness (SNT) data come from the KaKu LEGOS data (Garnier et al., 2021) and consist in the difference between CS2 Ku-band altimetric measurements, reflected by the ice, and SARAL Ka-band altimetric measurements, reflected by the snow. The data are provided in monthly gridded files, available in winter in each hemisphere. A temporal linear interpolation is applied to get SNT data at each weekly analysis. Due to SARAL orbital characteristics, no data are available for latitudes below 81.5°N. The observation error used in the analysis comes from the uncertainty supplied with the data, constrained to an arbitrary range of 0.01 m to 5 m. The snow data are assimilated as a thickness quantity, with the snow volume increment subsequently computed using the Kalman filter. It is important to note that the snow volume increment depends on all the assimilated data and reflects how well the volume correlates with them.

## 2.3 Experiments setup

Three experiments have been performed to assess the performance of the assimilation and the impact of the multivariate approach:

- FREE: experiment without any assimilation, used as a baseline of the model capacities;
- 240 • UNIVAR: experiment similar to the current operational system, using the previously described univariate SIC assimilation method;
- MULTIVAR: experiment with the multivariate assimilation scheme described previously, assimilating SIC, RFB and SNT observations, and updating the SIC, SIV and SNV model variables.

Characteristics of the three experiments are summarized in Table 2. All three experiments were conducted over two full annual cycles, 2017 and 2018, covering the period from 14/12/2016 to 26/03/2019. Initial conditions are based on the reanalysis GLORYS12V1 (Lellouche et al. (2021)).

Experiment name	Assimilated data	Analysis increments	Updated model variables
-----------------	------------------	---------------------	-------------------------





FREE	None	None	None
UNIVAR	SSMIS	SIC	SIC, SIV
MULTIVAR	SSMIS, RFB-LEGOS, SNOW-KaKu	SIC, SIV, SNV, RFBV, SNT	SIC, SIV, SNV

**Table 2: Experiments setup in terms of assimilated data, analysis increments and updated model quantities.**

### 3 Performances of the assimilation system


#### 250 3.1 Sea ice leads

We assess the experiments on their ability to correctly reproduce the amount of open waters within the sea ice extent, referred to “leads” hereafter. The area of sea ice leads offers valuable insights for predicting the Arctic sea ice extent (Zhang et al., 2018). The daily sea ice leads area timeseries are represented on Figure 1(a) in the Arctic and Figure 1(b) in the Antarctic. The sea ice leads content is computed by subtracting the sea ice area from the sea ice extent defined by cells where SIC > 15%.

255 We use two others different SIC datasets in order to quantify the spread among observations (Ivanova et al., 2015): the OSI-408 product (OSI SAF, 2017), derived from AMSR-2 satellite measurements and processed by the EUMETSAT OSISAF; and the CDR dataset (Meier et al., 2017; Peng et al., 2013) from the National Snow and Ice Data Center (NSIDC). All SIC data are interpolated on the polar stereographic SSMIS grid and use a consistent continental mask, ensuring the same area coverage.

260 As expected, the two assimilation experiments outperform the FREE experiment during summertime in terms of sea ice concentration coverage. In both hemispheres, FREE is not able to prevent excessive melting and shows a significant lack of sea ice, mainly in marginal areas, during July-October in Arctic and in January-April in Antarctica (not shown).

In the Arctic, the maximum lead surface area occurs in summer, more precisely at the beginning of the melting season. The daily surface area of leads peaks in July and then decreases with the retreat of the sea ice extent. The amount of leads remains constant from October to May in all the observations. In Antarctica, the lowest lead surface area is synchronous with the sea ice extent minimum in February-March. The observations then show an increase in leads area until its peak in 265 November–December, corresponding to the first third of the melting season. The southern observational datasets show strong agreement regarding the minimum lead surface; but diverge as the lead area increases. In both hemispheres, NSIDC and SSMIS observations respectively display the smallest and the largest amount of leads. The FREE experiment shows the smallest amount of leads remaining outside the range of the observations for most of the year in both hemispheres, and has a weaker 270 seasonal amplitude in the Arctic than the assimilated experiments and SSMIS and AMSR2 estimates, but comparable to NSIDC’s amplitude. The assimilation process rapidly and realistically increases the amount of leads in both the Arctic and Antarctic sea ice cover. The two assimilated experiments remain very close to the NSIDC leads area estimates during the northern hemisphere constant sea ice leads period, and they reproduce very well the rapid increase in lead surface area during spring. The UNIVAR experiment remains within the range of observational estimates throughout the year. The MULTIVAR 275 simulation exhibits the highest amount of leads during the peak period in July, even higher than the SSMIS observations.




**Figure 1: Daily time evolution of Arctic (a) and Antarctic (b) sea ice leads in millions of km<sup>2</sup> for SSMIS (black), AMSR2 (dashed black), NSIDC (dotted black) satellite data with the surface covered by them (shaded grey) and for FREE (blue), UNIVAR (green) and MULTIVAR (pink) experiments.**

280 In Antarctica, both the UNIVAR and MULTIVAR experiments have a consistently higher sea ice leads area than the FREE  
experiment and are thus in better agreement with the observations. They correctly reproduce the minimum leads area and its  
maximum, with the MULTIVAR experiment showing the highest amount of leads during the peak period in early December,  
still coherent with the SSMIS observations. However, during the second half of the increase in lead surface, the assimilated  
experiments show significant fluctuations that exceed the range of the observations. The fluctuations are linked to the  
285 occurrence of localized low-SIC and thin ice areas in the ice cover, called polynyas when they become open-water areas. Maps  
of the sea ice concentration in the assimilated observations and their difference to the experiments are shown on Figure 2 for  
both hemispheres. The well-known Weddell Sea “Maud-rise polynya” that appeared in winter 2017 (Jena et al., 2019) is not  
reproduced by the FREE experiment (Figure 2(b)). The UNIVAR and MULTIVAR experiments are able to reproduce this



290 polynya. However, in the assimilated simulations, the Maud-rise polynya begins to take shape from June 2017, earlier than in the observations, and the system struggle to keep an ocean uniformly covered in ice in the Weddell Sea. Other polynyas are present in few locations around the Antarctic: in the Amundsen Sea offshore of Pine Island Bay at 120°W in the UNIVAR and MULTIVAR simulations (Figure 2(b)), and near Iselin Bank at 180°E in the Ross Sea in the MULTIVAR simulation. These events appear repeatedly during the ice freezing period in 2017 and 2018.



295 **Figure 2: July 2018 in the Arctic (a) and September 2017 in the Antarctic (b) maps of the sea ice concentration, representing the observation SSMIS on the first column, and the difference between the experiences and the reference SSMIS observation on the following columns. The simulations are, in that order: FREE, UNIVAR and MULTIVAR. Root mean squared errors (RMS) are provided under each map.**

On the maps on Figure 2, sea ice concentration modelled by the UNIVAR simulation stands out and compares very well with the assimilated SSMIS dataset in the Arctic (RMSE of 0.04 in July 2018) and remains below the observation error in Antarctica (RMSE of 0.06 in September 2017). Despite leads metrics that moderately resemble SSMIS on average in the FREE experiment, its Arctic RMSE of 0.15 highlights inconsistencies in the modeled spatial patterns of sea ice concentration. Multivariate assimilation of RFB and SNT data reduces the Arctic SIC compared to SSMIS, mainly in the central Arctic. This lower SIC in central Arctic results in a RMSE of 0.18 for July 2018, the highest among the experiments. In that summer period, there are no RFB and SNT observations and the multivariate assimilation system creates the SIV and SNV increments from SIC observations and model covariances only. During the other months, the RMSE for the MULTIVAR simulation is lower, falling between the mean RMSEs of the UNIVAR and FREE simulations, which are 0.04 and 0.13, respectively. The Arctic

300  
305



mean RMSE of the UNIVAR and MULTIVAR simulations are similar in winter, but they differ in summer with the MULTIVAR simulation RMSE being 0.07 higher. In Antarctica, the FREE simulation presents mainly positive SIC biases in winter, particularly in the marginal ice zone (MIZ, defined by SIC values between 15% and 80%), and places the ice edge too far north compared to SSMIS observations (not shown) with mean RMSEs of 0.16 in September 2017 and 0.23 over the whole 2017-2018 months. The ice edge overestimation in the FREE experiment is corrected by the SIC assimilation in both UNIVAR and MULTIVAR simulations with comparable RMSEs of respectively 0.06 and 0.08 in September 2017 and the same values for the mean RMSEs over the whole 2017-2018 months.

In both hemispheres, the assimilation of SIC creates a higher presence of leads in the sea ice cover, in accordance with the SSMIS assimilated observations. The multivariate experiment alone even overestimates the quantity of leads during the seasonal maximum in the Arctic summertime. In the Antarctic, the two assimilated experiments reproduce the same variability and the occurrence of unobserved polynyas.

### 3.2 Snow volume

Figure 3(a) shows the probability density functions for snow thickness, radar freeboard using SAR mode, and radar freeboard using SARin mode, along with their model equivalents for the three experiments in the Arctic in April 2017. The SNOW-KaKu data in the Arctic present a zero-inflated bimodal, asymmetrical and positively skewed snow distribution with the first mode representing a snow thickness of 0 cm (no snow observed on the grid cell), and the second mode increasing in thickness as winter progresses and peaking at 13.6 cm in April 2017. The MULTIVAR snow distribution is very close to the Arctic SNOW-KaKu during winter (not shown) and matches perfectly in April. The UNIVAR and especially the FREE simulations accumulate excessive snow as winter progresses, leading to a positive bias by the end of the winter assimilation period as shown on Figure 3(a). The linear correlation (r-value) computed against the SNOW-KaKu observations in the Arctic results is consistently above 0.5 for MULTIVAR, peaking at 0.7 in December 2018 (not shown). The FREE and UNIVAR experiments exhibit systematic lower r-values, with the UNIVAR experiment having the lowest average correlation of 0.37. Compared to SNOW-KaKu estimates, the FREE and UNIVAR simulations present a spatially homogeneous overestimated snow thickness in Central Arctic and an underestimation in few areas such as north of the Canadian Archipelago, the east coast of Greenland, and in the Barents and Greenland seas (Figure 3(b)). This results in an excessive total snow volume of 1.24 Mkm<sup>3</sup> in the FREE experiment compared to that of 0.94 Mkm<sup>3</sup> estimated by SNOW-KaKu observations. In April 2017 (Figure 3 (b)), the MULTIVAR simulation represents closely both the SNOW-KaKu spatial pattern and the total snow volume amount with 0.91 Mkm<sup>3</sup>. This result is robust and remains valid for the other months of the year.





Figure 3: Top panels (a): Probability density functions (%) of the snow thickness, the radar freeboard SAR and radar freeboard SARin observations (dotted black) and their model equivalent for the FREE (blue), UNIVAR (green) and MULTIVAR (pink) experiments in the Arctic for April 2017. Middle (b), resp. bottom (c), row panels: snow volume per unit area [m<sup>3</sup>/m<sup>2</sup>], resp. radar freeboard volume per unit area, from SNOW-KaKu, resp. RFB LEGOS, (first column) and differences with FREE, UNIVAR and MULTIVAR experiments. Total snow and RFB volumes values and root mean squared difference (RMS) are provided under each map.

340



345 **Figure 4:** Top panels (a): Probability density functions (%) of the snow thickness, the radar freeboard SAR and radar freeboard  
 SARin observations (dotted black) and their model equivalent for the FREE (blue), UNIVAR (green) and MULTIVAR (pink)  
 experiments in the Antarctic for May and October 2017. Middle (b), resp. bottom (c), row panels: snow volume per unit area, resp.  
 radar freeboard volume per unit area, from SNOW-KaKu, resp. RFB LEGOS, (first column) and differences with FREE, UNIVAR  
 and MULTIVAR experiments in October 2017. Total snow and RFB volumes values and root mean squared difference (RMS) are  
 350 provided under each map.



In the Antarctic, the SNOW-KaKu data again exhibit a bimodal and positively skewed distribution, with a mode at 0.6 cm another at 11.6 cm in the first month of assimilation in May 2017 on Figure 4(a). As winter progresses, the second mode gets thicker and more frequent, peaking at 17.6 cm in October 2017. Among the simulation, the FREE experiment diverges the most from the observations, showing an increasing accumulation of snow as winter progresses, with a main mode 11.2 cm higher than the observed mode in October 2017. The UNIVAR and MULTIVAR experiments present lower snow thickness values compared to the observations during the whole 2017 and 2018 seasons, with main modes respectively 8.2 cm and 7.5 cm lower than the observed mode. The most significant snow positive biases in the FREE experiment are associated with thinner snow measurements in the SNOW-KaKu data, suggesting a thicker and more uniform snow cover, with a snow accumulation in the interior of the Weddell Sea, resulting in an excess of 1.06 million km<sup>3</sup> of snow compared to the SNOW-KaKu estimate (see Figure 4(b)). In comparison, the UNIVAR simulation presents a general thinner snow depth, maintaining however the accumulation in the Southwestern part of Weddell Sea. The MULTIVAR simulation has the weakest biases and is even able to reduce the high snow accumulation in the Weddell Sea present in the FREE simulation and to represent the thicker snow pattern measured in the SNOW-KaKu product downstream the Antarctica Peninsula. The biggest incoherence between the MULTIVAR simulation and the SNOW-KaKu observations is on the Pacific Ocean/Eastern Antarctic coastal sector, where the assimilated experiment does not reproduce the high snow thicknesses. The UNIVAR and MULTIVAR simulations have respectively 1.02 and 0.85 million km<sup>3</sup> less snow than SNOW-KaKu estimations in October 2017. The two simulations underestimate the SNOW-KaKu snow volume estimate for all the winter months of 2017 and 2018.

In both hemispheres, the MULTIVAR experiment consistently simulates snow depths closest to those used in the multivariate assimilation scheme. The assimilation of SNT is also able to rapidly modify the snow spatial distribution in accordance with the SNOW KaKu observations distribution. The agreement between the MULTIVAR experiment's snow thickness and the observations is higher in the Arctic than in the Antarctic.

### 3.3 Radar freeboard volume

The FREE simulation exhibits lower RFB values than the other experiments especially at the end of summer (not shown). FREE and UNIVAR have biases of respectively -6.6 cm and -7.9 cm in RFBV compared to the LEGOS observations in April 2017 (Figure 3(c)). The MULTIVAR simulation exhibits a very small bias of -0.5 cm in the assimilated region and a RMSE of 2.2 cm, below the observation error of both the SAR and SARin data. The largest differences compared to the LEGOS RFB estimates are located along the coasts around the Canadian Archipelago and to the east of Greenland, i.e. in SARin areas. The SARin data are assimilated with higher observation errors compared to SAR data, with mean values of 19.2 cm and 9.2 cm, respectively. The highest difference (> 40 cm) between MULTIVAR RFB values and LEGOS RFB estimates arises at the end of both 2017 and 2018 winters in the north of Greenland, an area where snow observations are not available. In summer, when no RFB observations are assimilated, the probability density function of the MULTIVAR RFB values remains more positively skewed than in other simulations. In November, when the observed data return after the summer break, the MULTIVAR experiment shows the lowest RMSE (2.6 cm) compared to the FREE (7.6 cm) and UNIVAR (8.3 cm)



385 experiment based on the 2017 and 2018 averages. However, the MULTIVAR simulation presents larger RFB biases in  
November, still below the mean observation error, than during the rest of the winter months when the errors relative to the  
RFB LEGOS dataset stay consistent.

LEGOS RFB measurements in the Antarctic present a similar gamma-type distribution as in the Arctic, with a  
decreasing SAR mode (from 3.3 cm to 0.5 cm) and increasing SARin mode (from 3.9 cm to 4.9 cm) between May and October  
390 2017 (Figure 4(a)). The simulations exhibit more uniform RFB values than in the Arctic with up to 20% of the RFB having  
the same value in the UNIVAR experiment in May 2017. The FREE, UNIVAR and MULTIVAR experiments have similar  
RFB SAR modes of respectively -0.4 cm, -1.0 cm and 0.3 cm in May 2017, lower than the observed SAR mode of 3.3 cm. As  
the season progresses, the FREE and UNIVAR simulations present an even more negative bias, with RFB modes respectively  
8 cm and 3 cm lower than the LEGOS RFB SAR mode in October 2017. A similar behaviour is shown for RFB SARin model  
395 equivalents, with the FREE and to a lesser extent the UNIVAR simulations frequently modelling negative RFB values that  
decrease as winter progresses. The MULTIVAR experiment is the only experiment to show a positively skewed distribution  
with positive modes in both SAR and SARin model equivalents throughout the duration of the simulation, aligning more  
closely to the LEGOS observations variability for the positive RFB values. The FREE and UNIVAR simulations display a  
general low bias in RFB all around the Antarctic (respectively -13.1 cm and -9.6 cm in average), with the most significant  
400 negative biases located in the two thicker RFB areas, indicating a more uniform RFB spatial distribution (Figure 4(c)). The  
MULTIVAR experiment has the lowest biases, -3.5 cm in average, and a RMSE of 4.47 cm. The FREE, UNIVAR and  
MULTIVAR simulations represent respectively 1.80, 1.33 and 0.48 million km<sup>3</sup> less RFBV than the LEGOS dataset. The  
underestimation of the southern RFB in the FREE simulation is likely due to the overestimation of the snow thickness in the  
Antarctic.

405 For all simulations and in both hemispheres, SAR measurements are in better agreement with the RFB model  
equivalent values compared to the SARin measurements. The MULTIVAR experiment shows the closest agreement with the  
observations among the simulations. The agreement between the RFB and SNV model equivalents from the MULTIVAR  
experiment and the observations is not as high in the Antarctic as in the Arctic.

## 4 Validation with independent datasets

### 410 4.1 Total freeboard: ICESat-2 data

Figure 5 presents the scatterplots between the monthly NSIDC ATL-20 gridded along-tracks total freeboard product  
measured by the ICESat-2 (Ice, Cloud and Land Elevation Satellite) ATLAS laser altimeter (Petty et al., 2023), and the total  
freeboard collocated in time and space for the LEGOS data and the FREE, UNIVAR and MULTIVAR experiments in the  
Arctic. The LEGOS total freeboard is made using LEGOS RFB and SNOW-KaKu data, and the constant water, ice and snow  
415 model densities. The MULTIVAR simulation and LEGOS data present similar linear correlation statistics (slopes and r-values)  
and MULTIVAR has better statistics than the FREE and UNIVAR experiments. The MULTIVAR simulation and the LEGOS






420 data have similar mean RMSE compared to ICESat-2 data (6.7 cm and 7.2 cm respectively). The MULTIVAR simulation and LEGOS data also display comparable mean total freeboard in January-February 2019, with values of 22.2 cm and 22.0 cm respectively, slightly thinner than the ICESat-2 estimate of 23.7 cm. The mean total freeboard for the FREE and UNIVAR experiments was found to be 19.4 cm and 15.0 cm, respectively, for the same period. The FREE and UNIVAR simulations consistently underestimate ICESat-2 total freeboard, especially in October 2018 with mean values of 9.2 cm and 6.6 cm respectively while the MULTIVAR experiment shows a mean value of 15.8 cm, aligning better with the mean total freeboard ICESat-2 estimate of 23.9 cm. The FREE experiment is not able to prevent excessive summer melting and exhibits unrealistic ice-free zones in October 2018. Higher statistical agreement in October 2018 for the MULTIVAR experiment shows that the data assimilation from the last winter positively impacts the simulation during the entire summer. However, compared to 425 ICESat-2, MULTIVAR still underestimates the thickness of the total freeboard at the end of Arctic summer. All the experiments exhibit correlations higher than 0.6 reflecting a general consistency with ICESat-2 total freeboard in terms of spatial distributions.


In Antarctica, simulated total freeboards show less agreement with ICESat-2 measurements compared to those in the 430 Arctic (Figure 6). All the experiments and the LEGOS estimations present a general more scattered plot in the south than in the north. In October 2018, the last month of the assimilation season in the southern hemisphere, the MULTIVAR total freeboard shows a greater variability than the FREE and UNIVAR total freeboard, in accordance however with the dispersion of the assimilated CS2 LEGOS RFB and SNOW-KaKu datasets. Both the MULTIVAR experiment and LEGOS data have a positive mean bias compared to the ICESat-2 data, of respectively +10.7 cm and +8.6 cm. The FREE simulation has a positive 435 bias cluster for thin total freeboard but underestimates the thicker freeboard values, resulting in a mean bias of +2.4 cm. The UNIVAR experiment is mostly underestimating ICESat-2 total freeboard values, with a mean bias of -11.9 cm. The melting season (January-February 2019) highlights the excessive thinning of the total freeboard in the simulations compared to the ICESat-2 data. The FREE experiment again has large unobserved ice-free zones with total freeboard values at 0 cm. The MULTIVAR experiment presents the highest total freeboard summer values among the experiments, with mean value of 19.6 440 cm (resp. 7.3 cm and 4.4 cm and for the FREE and UNIVAR experiments), still underestimating to a lesser extent the ICESat-2 mean values of 34.2 cm. The MULTIVAR simulation improves the concordance with ICESat-2 measurements with a systematic increase of the slopes in winter as in summer.

Assimilating radar freeboard and snow depth observations in the multivariate framework significantly reduces biases found with IceSat-2 total freeboard in both hemispheres. The agreement between modelled variables and IceSat-2 estimates is 445 stronger in the north than in the south.



450

**Figure 5:** Scatterplots of the monthly Arctic ICESat-2 total freeboard against FREE, UNIVAR, MULTIVAR experiments and LEGOS RFB/SND-KaKu data computed with model densities (black) for October 2018, beginning on the 14/10/2018 (experiments respectively in blue, green and pink; no LEGOS data), and for January-February 2019 (experiments respectively in orange, red and cyan). The  $x=y$  line (grey) and linear regressions for Oct 2018 (dotted black) and Jan-Feb 2019 (dashed black) are shown. Values of the linear slopes ( $s$ ) and the  $r$ -values ( $r$ ) are provided and all statistics are significant.



455

**Figure 6:** Idem Figure 5 but for Antarctica.

## 4.2 Sea ice volume


### 4.2.1 Total sea ice volume

The daily total ice volume values for each experiment are shown on Figure 7 (dotted lines). Figure 7 also presents the experiments collocated within the spatial coverage of the assimilated observations, which includes the central Arctic orbital gap and limited coverage of marginal seas (solid lines). Three different products are shown: (1) LEGOS<sub>og</sub>, the original SIV

460



LEGOS (Guerreiro et al., 2017), based on CS2 RFB and SNOW-KaKu measurements with varying snow and ice densities; (2) LEGOS\_mD, which uses the same measurements but applies constant snow and ice densities from the SI3 model; and only in the Arctic (3) CS2SMOS AWI, which combines SIV estimates from CS2 altimetric freeboard measurements of thicker ice and SMOS brightness temperature measurements of thinner ice (Ricker et al., 2017), using a modified W99 snow climatology and variable ice and snow densities.



470 **Figure 7: Time evolution of Arctic (a) and Antarctic (b) sea ice volume.** The daily values are presented for the simulations FREE (blue), UNIVAR (green) and MULTIVAR (pink), integrated over the whole hemisphere (dotted) and over the observation domain (plain lines). SIV observations used for comparison are computed over the LEGOS observation domain: LEGOS original SIT (LEGOS\_og, grey L in circles), SIT constructed from LEGOS observations of RFB and snow and the model constant ice and snow densities (LEGOS\_mD, black stars), and CS2SMOS AWI data in the Arctic (black dashes). The SIVOLU is computed using either SIC data provided by the supplier or the SIC OSISAF SSMIS data.



475 In the Arctic, the amount of sea ice remains consistently high throughout the entire simulation in the MULTIVAR  
experiment, resulting in sea ice maximums in average 13% and 48% higher than respectively the FREE and UNIVAR  
experiments. The FREE and UNIVAR simulations start each winter with a low sea ice volume compared to the observations.  
The MULTIVAR experiment presents systematically higher volume estimates and align better with CS2SMOS product in the  
beginning of November 2017 and 2018. The MULTIVAR SIV values increase rapidly during the first month of assimilation  
480 and follow closely the LEGOS\_mD observations. Even in summer, the MULTIVAR simulation maintains more ice volume  
in the Arctic than the other simulations. The UNIVAR simulation shows a particularly drastic decrease in its ice volume  
estimate relative to the FREE experiment and is consistently lower than all the observation products. On average over the  
entire simulation period, the UNIVAR experiment shows a decrease in sea ice volume of 23% while the MULTIVAR  
experiment shows a 21% increase compared to the FREE experiment. The assimilation of CS2 LEGOS RFB and SNOW-  
485 KaKu in the MULTIVAR experiment modifies the seasonal cycle of the sea ice volume estimates, with a maximum earlier  
than in the other simulations, and is more consistent with the observations.


In both hemispheres, the MULTIVAR experiment shows the largest sea ice volume, while UNIVAR has the smallest.  
Among the different products, LEGOS\_mD has the highest volume, followed by LEGOS\_og and –only in the Arctic–  
CS2SMOS. Notably, the products are highly sensitive to variations in snow and ice densities, with LEGOS\_mD showing in  
490 average respectively 1.48 million km<sup>3</sup>, resp. 5.6 million km<sup>3</sup>, more sea ice volume than the original LEGOS\_og in the Arctic,  
resp. the Antarctic.

#### 4.2.2 Comparison with SMOS satellite measurements

The CS2SMOS AWI product uses measurements from the SMOS satellite in addition to CS2 measurements. SMOS  
is known to have less uncertainties than CS2 on thin ice measurements (less than 1 m, Ricker et al., 2017). The LEGOS\_og  
495 displays a coherent sea ice thickness spatial distribution compared to the CS2SMOS product with the smallest RMSD (resp.  
mean difference) of 30 cm (resp. 5 cm, Figure 8). LEGOS\_mD presents a higher RMSD (resp. mean difference) of 39 cm  
(resp. 34 cm). The FREE simulation shows thinner ice than the CS2SMOS data in the central Arctic and on the east coast of  
Greenland, and thicker ice elsewhere. The UNIVAR simulation has a globally much thinner ice coverage with approximately  
half of its ice area covered by ice below 1 m thickness and the other half with ice between 1 m and 2 m height. The MULTIVAR  
500 experiment shows a higher ice volume compared to the other experiments, with a significant ice accumulation thicker than in  
the CS2SMOS product on the north of the Canadian Archipelago and Greenland. In that area of important deviation between  
CS2SMOS and MULTIVAR values, the assimilated SNOW-KaKu measurements are not available. In the LEGOS SIV  
observation domain, the simulations present a similar RMSD against the CS2SMOS product of 33 cm (FREE, MULTIVAR)  
and 31 cm (UNIVAR). The MULTIVAR modelled ice thickness has the same positive biases as the LEGOS\_mD product but  
505 keeps a thinner ice than the CS2SMOS data on the east coast of Greenland, similarly to the two other simulations. Outside of  
the LEGOS observations domain, the UNIVAR simulation shows the highest RMSD (65 cm) for the CS2SMOS SIT values



510 thicker than 1 m, while the FREE simulation has the highest RMSD (48 cm) for CS2SMOS SIT values thinner than 1 m among the three experiments. The RFB and snow assimilation in the MULTIVAR simulation corrects the FREE and UNIVAR underestimation of the ice thickness in the central Arctic region (RMSD of 38 cm) and presents fewer positive biases than the FREE simulation for the thin ice around the ice edge (RMSD of 27 cm).




515 **Figure 8: April 2017 sea ice volume maps in the Arctic for CS2SMOS dataset (reference) and its difference with the FREE, UNIVAR, and MULTIVAR experiments (first line) and the observations LEGOS\_og (original) and LEGOS\_mD (with model constant densities). Table: root mean square error (RMS) and mean difference (MD) between FREE, UNIVAR, MULTIVAR, LEGOS\_og, LEGOS\_mD and CS2SMOS data, calculated on the LEGOS zone and outside the LEGOS zone and for CS2SMOS sea ice thickness of less than or greater than 1m. The table colours highlight the values close to 0 (white) and the extremes (green for the RMS, and blue/red for the negative/positive MD).**

520 The LEGOS observations, considering both fixed and varying densities, present a very thick ice volume in the southern hemisphere (Figure 9). Similarly to the Arctic, the LEGOS\_mD shows thicker ice volumes than the LEGOS\_og data. Compared to SMOS data, both LEGOS estimates show a different ice field: the CS2 Antarctic ice thickness processed by the LEGOS is thicker with RMSE values of 78 cm (resp. 97 cm) for LEGOS\_og (resp. LEGOS\_mD) and the ice accumulations are measured on the northernmost part of the Weddell Sea with CS2 measurements, whereas SMOS satellite detects thick ice on the southernmost part of the Weddell Sea. The FREE and UNIVAR simulations have spatially homogeneous SIV distributions and similar RMSD compared to the SMOS data on the LEGOS domain (respectively 24 and 26 cm). The FREE 525 experiment has a consistent positive SIV bias compared to the SMOS dataset. Although most of the UNIVAR experiment's ice thickness is below 1 m, it underestimates SMOS ice thickness, except on areas close to the ice edge, where UNIVAR values align well with the SMOS measurements (mean difference of -2 cm). Compared to FREE and UNIVAR, the MULTIVAR simulation shows more important ice accumulations, in consistency with both LEGOS SIV data, and therefore has the highest



530 RMSE relative to the SMOS data on the LEGOS domain (38 cm). The MULTIVAR simulation does not reproduce the largest LEGOS SIV values and is therefore closer to the SMOS data than the LEGOS estimates. Outside the LEGOS domain, MULTIVAR corrects the positive bias noticed along the ice edge in the FREE simulation but degrades the performances of the UNIVAR simulation with a higher error (mean difference of 7 cm). The FREE simulation is the only experiment that does not reproduce correctly the Maud Rise polynya, which is seen in all observation products and in the two assimilated experiments.

535



540 **Figure 9: September 2017 sea ice volume maps in the Antarctic for the SMOS data (reference) and its difference to the FREE, UNIVAR, and MULTIVAR experiments (first line) and to the observations LEGOS\_og (original) and LEGOS\_mD (with model constant densities). Table: root mean square error (RMS) and mean difference (MD) between FREE, UNIVAR, MULTIVAR, LEGOS\_og, LEGOS\_mD and SMOS data, calculated on the LEGOS zone and outside the LEGOS zone. The table colours highlight the values close to 0 (white) and the extremes (green for the RMS, and blue/red for the negative/positive MD).**

In both hemispheres, for SIT < 1 m, using the multivariate assimilation scheme better aligns the modelled sea ice volume with the SMOS data, presenting a lower RMSE for the MULTIVAR experiment than the FREE experiment and the LEGOS\_mD data. However, the UNIVAR experiment shows more accurate sea ice volume estimates for thin ice than the MULTIVAR experiment when using SMOS measurements as a reference.

545



## 5 Discussion

### 5.1 Performances of the multivariate assimilation

The radar freeboard and snow thickness assimilation allows the multivariate assimilation experiment to correct the model biases against the assimilated datasets: the MULTIVAR simulation has the closest results to the RFB LEGOS and SNOW-KaKu products in both hemispheres. However, the comparison of the Antarctic snow and RFB equivalents shows less agreement with the assimilated observations than in the Arctic.

The univariate assimilation system only corrects the SIC variable and aims at keeping a constant SNV, with a dynamic threshold on the SNT. In the Antarctic, and to a lesser extent in the Arctic, the UNIVAR experiment displays a lower SNV compared to the FREE experiment. This result shows that our threshold is not appropriate in most of the Antarctic, and in some regions in the Arctic. Thanks to the snow assimilation, in the MULTIVAR simulation, the total volume of snow is adjusted but does not recover the total amount of observed snow in the Antarctic. A modification of the SNT threshold would improve the snow assimilation algorithm in that sense. Nevertheless, the SNOW-KaKu assimilation enables the simulations to reproduce the snow observations spatial distribution in both hemispheres. The snow cover completely melts in summer in each hemisphere and shows no long-term effect of the winter snow assimilation.

The MULTIVAR simulation shows higher RFB values than the FREE and UNIVAR simulations in both hemispheres, even in the absence of observations during the summer. However, a drift in the RFB equivalent is still observed during this season, leading to a negative bias in November/May, when the assimilation begins. This small negative bias suggests that the model's trajectory is below the observed values, a hypothesis supported by the significantly more pronounced bias observed in the FREE and UNIVAR simulations. In the Antarctic, the RFB underestimation is particularly significant: the FREE and UNIVAR simulations show mostly negative radar freeboard values due to an imbalance between their snow and ice thicknesses: their snow cover is too thick compared to their ice thickness. The initial state of ice and snow in the southern hemisphere found in the FREE experiment is much more different from the assimilated observations compared to the north. The multivariate assimilation process is then less effective in aligning the model with the observed data in the Antarctic than it is in the Arctic.

In both hemispheres, the MULTIVAR simulation produces RFB extremums that extend beyond the minimums and maximums observed in the FREE and UNIVAR simulations, and more closely align with the LEGOS observations. Despite this improvement, the MULTIVAR simulation does not capture the thickest and thinnest RFB LEGOS measurements. This discrepancy could be attributed to the spatial resolution mismatch between the observations and the model. Furthermore, it is important to recognize that the MULTIVAR simulation is not designed to replicate every extreme observation (such as a notably high SARin RFB of 4.3 m observed in October 2017 in Antarctica) as the assimilation scheme seeks to balance observational data with the model's physical constraints. Given the use of unfiltered RFB data in the assimilation, we do not expect the model to reproduce the exact observed values but rather a smoothed representation that respects the model's inherent dynamics.



The LEGOS observations are characterized by spatially significant data gaps in central Arctic and in the Canadian Archipelago. The MULTIVAR simulation smoothly assimilates the RFB and SNOW-KaKu data in these areas without any visible demarcations. Furthermore, due to the choice of parameters for the localisation algorithm in the assimilation scheme, the assimilated satellite tracks do not print on the modelled patterns. However, the largest RFB differences between the MULTIVAR experiment and the RFB LEGOS assimilated observations are located on the north of the Canadian Archipelago and Greenland, with an especially thin RFB in our simulation north of Greenland. No snow observations are available in this area, and the MULTIVAR presents thicker snow values than the FREE and UNIVAR simulations, suggesting either that in the absence of snow observations, an inaccurate modelled snow depth affects the RFB assimilation performance; or that the background error covariances matrix do not assess correctly the relationships between snow and radar freeboard variables.

When considering the sea ice volume, the experiments provide similar results in both hemispheres: the assimilation of SIC with the univariate method decreases the ice volume compared to the FREE simulation. The assimilation of RFB LEGOS and SNOW-KaKu creates the highest sea ice volume of all the simulations. The MULTIVAR experiment also displays a more accurate spatial distribution of the ice than the other experiments. The MULTIVAR modelled ice volume in the Arctic is very coherent with the LEGOS\_mD dataset in the Arctic, which is more consistent with our observation operator in terms of sea water, snow and sea ice densities. In the Antarctic, the modelled sea ice volume is consistently lower than the LEGOS\_mD product, probably due to lower model skills in representing sea ice in the Antarctic than in the Arctic (Massonnet et al., 2011) and more divergence between the modelled initial state and the assimilated observations, as discussed earlier.

## 5.2 Comparison with independent data

The IceSat-2 satellite measures the total freeboard through laser altimetry instruments, it is therefore completely independent from the radar altimetry-based LEGOS freeboard estimates. Previous section shows that assimilating LEGOS data reduces the errors in the simulations total freeboard estimates compared to ICESat-2 measurements. The comparison in the Antarctic also shows weaker correlations between IceSat-2 data and the experiments than in the Arctic. It should be emphasized that most of the comparisons made in the southern hemisphere with ICESat-2 data is done during summertime, without assimilation of radar freeboard and snow. The summer period of the southern ice is also known to be poorly represented by the models (e.g. Shu et al., 2020; Roach et al., 2020). In addition, the LEGOS data present less coherence with ICESat-2 compared to the Arctic. Nevertheless, the MULTIVAR simulation exhibits higher performance in terms of total freeboard compared to the other two simulations, particularly during the summer months. This demonstrates that the multivariate assimilation process induces changes in total freeboard that persist even when radar freeboard and snow are not assimilated.

Further comparison with in-situ independent observations in the Arctic show similar improvement with the multivariate assimilation system compared to the FREE and the UNIVAR experiments (see more details in Appendix A).

Sea ice thickness products obtained from brightness temperature measured by the SMOS satellite can be considered complementary to the altimetric ice products because they provide thin ice estimates (Kaleschke et al., 2024). In the Arctic, the CS2SMOS data shows thinner ice thicknesses than the LEGOS products (same as other CS2 products in Sallila et al., 2019)





but the observational datasets are still coherent (better spatial alignment and RMSD of the same order as the FREE simulation). In that hemisphere, differences between the simulations and the CS2SMOS data show a generally better agreement for the MULTIVAR simulation compared to the FREE and UNIVAR simulations. The predominant positive biases observed in the  
615 MULTIVAR simulation are consistent with the biases in the LEGOS\_mD product (i.e., north of the Canadian Archipelago and Greenland). However, Sallila et al., (2019) established that the CS2SMOS product tends to underestimate the thickness of thick ice in the Arctic when compared to in-situ measurements. Therefore, an overestimation of the CS2SMOS estimates is not an unexpected outcome for thicker ice. The CS2SMOS product estimates of thin ice, however, are in closer alignment with  
620 the in-situ Arctic measurements (Sallila et al., 2019). The more precise thin ice estimates from the UNIVAR experiment are compromised by the assimilation of CS2 data in the MULTIVAR experiment, when compared to the CS2SMOS values. It may be beneficial to increase the observation errors for the thicker RFB or in the marginal ice zone in order to reduce this degradation in comparison to the UNIVAR simulation.

In Antarctica, the SMOS product is restricted to ice with SIT < 1 m, and a similar situation as with the thin Arctic ice arises: the comparison with the SMOS Antarctic data shows a better agreement with the UNIVAR simulation. The  
625 MULTIVAR simulation predominantly overestimates the SMOS measurements, due to an overestimation of the assimilated LEGOS data compared to the SMOS estimates. The SMOS data however display a systematic underestimation of sea ice thickness in areas of ice divergence (Kaleschke et al., 2024); and the Antarctic sea ice shows generally divergent ice drifts (e.g. Petty et al., 2021). Moreover, the assimilated LEGOS data present little resemblance with the SMOS Antarctic measurements. However, the Southern Ocean lacks consistent in-situ data measurements of sea ice and snow to better evaluate satellite  
630 observations and models estimates. Here, The MULTIVAR simulation provides better statistics than the two other experiments against the IceSat-2 data thanks to the multivariate assimilation of LEGOS observation product, and it shows a better alignment with the SMOS data than the FREE simulation despite the assimilation of a LEGOS product that does not align with the SMOS data. The validation against these two independent datasets hence proves that the multivariate ice assimilation scheme in the Antarctic created an intermediate sea ice state between the LEGOS observations and the model.

### 635 **5.3 Ice and snow densities**

Sea ice thickness products obtained from CS2 radar altimetry measurements have significant uncertainties due to the assumptions made on values of snow thickness and ice and snow densities during the radar freeboard to ice thickness conversion (Kern et al., 2015; Kwok and Cunningham, 2015; Mallett et al., 2020; Garnier et al., 2021). Assimilating directly the radar freeboard allows us to control the origin of the uncertainties by using the rawest measurement possible and controlling  
640 all the assumptions made during the assimilation process. We decided to assimilate a satellite observed altimetry snow thickness, which uses the same radar altimetry techniques as the RFB product. Garnier et al. (2022) show that using coherent measurement techniques between the snow and freeboard datasets gives an accurate total freeboard value even when the snow-ice interface is biased.



The multivariate data assimilation proceeds for the RFB volume observations by constructing a model equivalent  
645 using the model SIV and SNV variables and the model fixed densities for water, ice and snow. The water density is nearly  
consistent in all the sea ice volume datasets, with values varying by only a few  $\text{kg/m}^3$ . However, the ice and snow density  
values vary a lot. The model's constant ice density is  $917 \text{ kg/m}^3$ , but the ice density in the Arctic depends on the ice age for  
LEGOS\_og and CS2SMOS with the values from Alexandrov et al. (2010) as extremums:  $882 \text{ kg/m}^3$  for the MYI and  $917$   
650  $\text{kg/m}^3$  for the FYI. Hence, assimilating radar freeboard and snow with the model constant ice density does affect mostly the  
MYI regions in the Arctic, which corresponds to the thicker ice regions that do not melt during summer, in the north of the  
Canadian Archipelago and Greenland. The difference of ice density results in an ice thickness 32% higher on MYI in the  
Arctic. The model constant snow density is  $330 \text{ kg/m}^3$ . Garnier et al. (2022) used a constant snow density of  $300 \text{ kg/m}^3$  in the  
Arctic for the LEGOS\_og product, with a consequently lower sea ice thickness than the model for equal RFB, snow thickness  
and ice density values. Densities in the observation products in the Antarctic are generally seasonally varying densities. The  
655 model's ice density exceeds that of the LEGOS\_og observations (mean:  $895 \text{ kg/m}^3$ ), with a particularly pronounced  
discrepancy in October. The model snow density is comparable on average to the LEGOS observation's snow densities in  
Antarctica but presents differences up to  $40 \text{ kg/m}^3$  for some winter months. This discrepancy between ice and snow densities  
brings additional variability in sea ice volume even when similar radar freeboard and snow measurements are used, as  
illustrated by the difference between the LEGOS\_og and LEGOS\_mD datasets. The constant densities parametrization in the  
660 model enhances the positive bias of the sea ice volume in the Arctic compared to the CS2SMOS product. In the experiments  
presented here, the uncertainties due to the densities are related to the RFB observation operator. Hence, these uncertainties  
increase the representation error in the analysis. Varying ice and/or snow densities are crucial features to be incorporated in  
the next version of the sea ice model: it would ensure a more accurate radar freeboard assimilation by lowering this  
representation error. One could for instance use the method from Zhang et al. (2022) to select the optimal freeboard-to-  
665 thickness conversion ratios values by fitting the resulting ice thickness to in-situ or airborne measurements. Moreover,  
implementing seasonally evolving densities in our model would increase the physical accuracy of the sea ice modelling. For  
instance, Mallett et al. (2020) offers a linear evolution of the snow density to account for the densification of the snow as  
winter passes. Sievers et al. (2023) use this relationship to implement a radar freeboard assimilation scheme with a varying  
snow density, but did not modify the density in the model physics.

#### 670 **5.4 Sea ice openings in Antarctica**

In both hemispheres, results showed that all assimilated experiments successfully corrected the biases of the FREE  
experiment with respect to the SIC variable. Univariate SIC assimilation provides the best performance for sea ice  
concentration as the covariances are not negatively affected by other quantities. The addition of freeboard radar and snow in a  
multivariate assimilation configuration takes the modelled SIC away from the OSISAF data during summer, i.e. when RFB  
675 and SNOW-KaKu data are unavailable and the SIV and SNT increments are built from model covariances. Summer remains



the most difficult season for systems to reproduce in both hemispheres. SIC passive microwave observations also have the greatest uncertainties during the melting season (Ivanova et al., 2014).

Sea-ice models using VP or EVP rheologies have been shown to reproduce the observed sea ice deformations only with high resolution horizontal grids (4.5 km grid spacing or lower, Wang et al., 2016; Spreen et al., 2017; Hutter et al., 2018).

680 Both assimilated experiments increased the amount of open water compared to the FREE experiment and increase the amount of sea ice leads on a coarser grid of  $\frac{1}{4}^\circ$ , i.e. grid cells of size between 10 km and 24 km in the Arctic. The multivariate experiment shows an even higher presence of open waters than the UNIVAR experiment during peak period in the boreal summer.

The assimilated experiments timeseries in the Antarctic display oscillations that are due to the occurrence of very localized low-SIC or open water areas, e.g. the so-called polynyas (Figure 1(b)). These openings only appear in the assimilated experiments. As none of these openings occur in the FREE experiment, the thick snow + ice layer likely insulates the ice and prevents melting from the ocean beneath. The occurrence of the Maud Rise Polynya in Sept-Oct 2017 (Jena et al., 2019) is reproduced by the UNIVAR experiment, but its size is underestimated (Fig 2b)). On the other hand, the size of this polynya is greatly overestimated by MULTIVAR and appears about 3 months in advance of the one observed by satellite. Furthermore, 690 the MULTIVAR (and UNIVAR to a lesser extent) experiments show the presence of other polynyas this winter 2017 and a few more during winter 2018. These events are the combination of a general reduction of snow and increase of ice freeboard with respect to the FREE simulation, but in specific areas where SIC or RFB observations show local minima. These reductions in the areas covered by ice finally expose the surface to the warm waters of the ocean. Once triggered, assimilation is no longer able to counteract the strong vertical instability and oceanic warming that prevent these openings from closing. However, some 695 of these activation zones correspond to fracture zones that have already been identified, either for reasons of atmospheric divergence (low pressure systems in Kwok et al., 2017) or linked to the local bathymetry (Reiser et al., 2019). These polynyas are the consequences of intense interactions between the ocean and the surface in our simulations on places where the equilibrium of model is very sensitive to any disturbance. Changes have been implemented in the assimilation system to mitigate the occurrence of these simulated polynyas (see paragraphs 2.1.2 and 2.2.1). The SST assimilation under the ice has 700 been activated to keep the surface waters close to the freezing point. Very few in situ profiles are available in the Southern Ocean, and some of them were radically changing the thermohaline properties of the ocean in a large area and over a long period of time, thus we did not activate the in-situ profile assimilation poleward  $60^\circ\text{S}$  to keep the modelled ocean stratification. We increased the maximum SIC observation error to 40% to moderate the intensity of sea ice assimilation in the Southern hemisphere. These modifications have reduced the likelihood of triggering polynyas in both UNIVAR and MULTIVAR 705 simulations.



## 6 Conclusion

This study presents the first implementation of a multivariate sea ice assimilation scheme in both the Arctic and Antarctica within a global  $\frac{1}{4}^\circ$  modelling and analysis system. This system, largely based on the Mercator operational system, already includes a multivariate ocean assimilation but currently only assimilates sea ice concentration (SIC). Our study  
710 enhances this capability by incorporating a multivariate ice assimilation approach, assimilating along tracks radar freeboard and snow depth jointly with sea ice concentration. By comparing simulations without assimilation, with univariate SIC assimilation, and with this innovative multivariate system, we assess the capabilities of the assimilation scheme. The univariate SIC assimilation method systematically decreases the ice volume compared to the FREE experiment and shows a thin ice bias compared to observations. The multivariate assimilation increases the sea ice volume in both hemispheres, enabling the  
715 modelled sea ice to converge on assimilated data sets. The spatial distribution of the sea ice and the snow is modified in accordance with the assimilated observations. Even in summer and in the observation's spatial holes, when no satellite altimetry observations are assimilated, the MULTIVAR experiment's ice variables are favorably modified by the multivariate ice assimilation. Moreover, the diagnosed freeboard from the multivariate system compares better with Iceat-2 independent observations in the Arctic and, to a lesser extent, in Antarctica.

The comparison with observations coming from SMOS satellite shows that the UNIVAR experiment agrees better  
720 with the more reliable SMOS sea ice volume estimates for thin ice (less than 1 m) than the MULTIVAR experiment. In the Antarctic, CS2 and SMOS sea ice volume estimates diverge, so assimilating CS2 radar freeboard takes the model results away from SMOS measurements. Increasing the error of altimetry measurements over marginal zones and thin ice surfaces or merging altimetry with SMOS estimates for ice are potential options in this multi-variate approach. Ultimately, the results of  
725 the assimilation scheme reflect a balance driven by our selection of assimilated observations: the simulation is restricted to an intermediate position between the assimilated data and the model's trajectory. Therefore, a degree of consistency between the assimilated and independent validation datasets is essential to effectively detect an improvement of the sea ice fields thanks to data assimilation techniques.

The multivariate assimilation system performs better in the Arctic than in the Antarctic, largely due to differences in  
730 the model's initial free state. In the southern hemisphere, the initial biases in the free simulation are larger than those in the northern part, making it more challenging for the assimilation to reconcile the model with observations. This highlights the critical role of the model's baseline state in a data assimilation system. Further, the significant differences in ice volume estimates due to the use of constant or non-constant densities show and confirm the importance of having a comprehensive modelled physics with observations measurements.

The results for the southern hemisphere also show the strong interactions with the oceanic surface layers in the life  
735 cycle of the sea ice cover. The choice of the assimilation parameters (analysis snow depth threshold, observation errors, localization radius) is still an ongoing work and further study in the assimilation methodology is needed to fully handle the strong coupled ocean/ice interactions at work in the Southern Ocean.



This multivariate assimilation system paves the way for the future integration of CIMR and CRISTAL satellite  
740 measurements in synergy into operational systems. The CRISTAL satellite, set for launch in 2028, will carry altimetry radar  
instruments equipped with both Ku-band and Ka-band radars, enabling simultaneous altimetry measurement of the air–snow  
and ice–snow interfaces. The CIMR satellite will measure the sea ice concentration with passive microwave imagers, allowing  
for sub-daily and high resolution (5 km) polar measurements.

745 *Data availability.* All the sea-ice reanalysis experiments are available on request. This study has been conducted using E.U.  
Copernicus Marine Service Product: Global Ocean Sea Ice Concentration Time Series REPROCESSED (OSI-SAF);  
<https://doi.org/10.48670/moi-00136>, available on [Global Ocean Sea Ice Concentration Time Series REPROCESSED \(OSI-SAF\) | Copernicus Marine Service](#). The LEGOS data (FBR, SNOW-KaKu and SIV LEGOS\_og) used in this study (doi  
10.6096/CTOH\_SEAICE\_2019\_12) were developed, validated by the CTOH/LEGOS, France and distributed by Aviso+:  
750 [Altimetry Sea Ice products from CTOH](#). ICESat-2 total freeboard was downloaded from <https://nsidc.org/data/atl20/versions/4>  
on the 06/06/2024, using the ‘monthly’ group of the netcdf files (Petty et al., 2023). SMOS Antarctic data was downloaded  
from [Tian-Kunze, X; Kaleschke, L \(2021\): SMOS-derived sea ice thickness in the Antarctic from 2010 to 2020 \(pangaea.de\)](#),  
version 3.2, last accessed on the 14/08/2024. The merging of CryoSat-2 and SMOS data (CS2SMOS) was funded by the ESA  
project SMOS & CryoSat-2 Sea Ice Data Product Processing and Dissemination Service and data from 01/12/2016 to  
755 27/03/2019 were obtained from <https://www.meereisportal.de> (grant: REKLIM-2013-04, Ricker et al., 2017). The data  
presented in the Appendix A consists in the BGEP ULS measurements, collected and made available by the Beaufort Gyre  
Exploration Program based at the Woods Hole Oceanographic Institution (<https://www2.whoi.edu/site/beaufortgyre/>) in  
collaboration with researchers from Fisheries and Oceans Canada at the Institute of Ocean Sciences; the ULS measurements  
in the Fram Strait are from the website <https://data.npolar.no> (Sumata et al., 2021); and the Operation IceBridge Quick Look  
760 measurements, available at <https://nsidc.org/data/nsidc-0708/versions/1> (Kurtz et al., 2016).

*Author contribution.* AC, GG and CET designed the analysis and the experiments and AC carried them out. AC and GG wrote  
the paper and CET, GR, MH and PYLT revised it. MH and GR helped with the experiments’ setup. FG provided the LEGOS  
datasets and shared valuable insights on the satellite altimetry observations. All named authors have participated in the present  
article and have brought contributions to the elaboration of its final version.

765 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements:* Special thanks to Sara Fleury, and the LEGOS team for providing the altimetric satellite observations  
(radar freeboard and snow depth data) and for their useful advice regarding these observations.

Thank you to Guillaume Samson, Clément Bricaud and Laurent Parent from Mercator Océan for technical assistance and  
support in conducting the experiments.



770 This study was supported by Mercator Ocean International (France) and Centre National des Etudes Spatiales (CNES, France) as part of the doctoral grant of AC.

Provision of datasets used within this study is also acknowledged: sea ice concentration products from OSI-SAF and NSIDC, ERA5 atmospheric reanalysis from ECMWF, ICESat-2 data from NASA, CS2SMOS and SMOS ice thickness from the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research (AWI), BGEP ice drafts from the Beaufort Gyre

775 Exploration Program based at the Woods Hole Oceanographic Institution, Fram Strait ice drafts from the Norwegian Polar Institute, Operation Ice Bridge Quick-Look ice thickness from the NASA National Snow and Ice Data Center.

The Scientific colour maps lipari and vik (Cramer, 2023) are used in this study to prevent visual distortion of the data and exclusion of readers with colour-vision deficiencies (Cramer et al., 2020).


780



## Appendix A. Comparison with in-situ measurements.

The in-situ data include Upward-Looking Sonar (ULS) moorings measurements in the Beaufort Sea, from the Beaufort Gyre Exploration Project (BGEP) with moorings A, B and D; and in the Fram Strait, from the Norwegian Polar Institute (NPI) (Sumata et al., 2021) with moorings F11, F12, F13 and F14. We also use airborne laser and radar altimeter measurements in the western Arctic from the Operation Ice Bridge Quick Look product (OiB-QL, Kurtz et al., 2016).

The ULS moorings are located in regions where the LEGOS data are fully available (both RFB and SNOW-KaKu). A distinction is made for OiB-QL measurements based on the availability of LEGOS data, highlighting the orbital hole that results from using SARAL-Altika measurements.




**Figure A1: Map of the Arctic and the different in-situ measurements used for validation of the simulations.**

785 BGEP ULS measurements, available all year long, are available for the whole duration of the simulations, and the NPI ULS data are available until August 2018. Airborne OiB-QL observations are collected only in spring, but they sample a variety of ice (MYI and FYI) and cover a significant area in the Arctic. OiB-QL measurements campaigns took place during 7 days in March 2017, 3 days in April 2017, 1 day in March 2018 and 6 days in April 2018. The comparison for all measurements is made at monthly frequency. The LEGOS values presented in this appendix are made from the LEGOS RFB data, the SNOW-KaKu data, and the model fixed densities (LEGOS\_mD).



790 **A.1 Beaufort Sea: BGEP ULS**




795 **Figure A2: Comparison of monthly average ice draft from LEGOS data, FREE, UNIVAR and MULTIVAR experiments within 200 km of the Beaufort Gyre Experiment Program ULS Moorings (Mooring A: triangle, Mooring B: circle, Mooring D: square) for the summer (empty symbols) and winter (solid symbols). The linear regression (dashed black line), slope (s) and r-value (r) are shown for each dataset. Methodology from Laxon et al. (2013).**

The Figure shows a remarkable agreement of ice drafts between BGEP data and all experiments. The LEGOS observations have less coherence with the BGEP ULS measurements than the experiments but still with very high statistics. The values that underestimate the BGEP measurements in all 3 experiments are mostly during summertime (empty markers). The MULTIVAR experiment exhibits less accuracy than the FREE and UNIVAR simulations, with more scattered values, inheriting the behaviour of assimilated LEGOS data. However, MULTIVAR ice drafts have higher correlation than those from LEGOS estimates and, further, the MULTIVAR experiment is able to keep the strong correlation obtained with the FREE ice draft values during summertime.

800

**A.2 Fram Strait: NPI ULS**



805 **Figure A3: Comparison of monthly average ice draft from LEGOS data, FREE, UNIVAR and MULTIVAR experiments within 200 km of the Norwegian Polar Institut (NPI) Fram Strait ULS Moorings for the summer (empty symbols) and winter (solid symbols). The slope (s) and r-value (r) are given for each dataset.**


The ULS ice draft measurements are thicker in the Fram Strait than in the Beaufort Sea. The LEGOS data is in general agreement with the NPI data but presents mostly thicker ice drafts than the ULS measurements. The FREE and UNIVAR ice





810 drafts consistently underestimate the ULS measurements, with very low slopes and r-values. Most of the summer values (empty markers) in these two experiments have ice drafts at 0 m. Assimilating LEGOS RFB and SNOW-KaKu results in higher ice drafts, especially in winter when the assimilation is effective. Large errors in the MULTIVAR experiment's summer ice drafts values compared to the NPI ULS measurements still remain in this region of the Fram Strait, where the ice front is highly variable.

### 815 A.3 Operation IceBridge QuickLook sea ice thickness



820 **Figure A4: Comparison of monthly average ice thickness from LEGOS data, FREE, UNIVAR and MULTIVAR experiments collocated with OiB-QL airborne measurements in the Arctic. Areas where LEGOS SNOW-KaKu and RFB measurements are available are respectively in black, blue (FREE), green (UNIVAR) and pink (MULTIVAR); otherwise orange (FREE), red (UNIVAR) and cyan (MULTIVAR) refer to regions where SNOW-KaKu data are not available. All ice thickness values are gridded onto a 0.4° latitude by 4° longitude Arctic grid, following the methodology of Tilling et al. (2018). The slope (s) and r-value (r) are given for each dataset.**

825 The LEGOS data and the OiB-QL ice thickness measurements are in general good agreement. The OiB-QL data presents a cluster of measurements between 1 and 2 m that is well reproduced by all experiments and by the LEGOS data. Thicker measurements from the OiB-QL 2017 and 2018 campaigns are underestimated by the FREE and UNIVAR experiments. These 2 experiments do not show ice thickness values higher than 4 m, whereas the OiB-QL measurements signal ice up to 6.6 m thick. The MULTIVAR simulation is able to reproduce thicker ice, resulting in a general better agreement with the OiB-SL measurements, in all regions: where all the assimilated data is available, and also where some or all of the assimilated data are missing. However, the MULTIVAR experiment's ice thickness values are very scattered, especially in the region where the

830 LEGOS data is not entirely available (no SNOW-KaKu poleward of 81.5°N; and no RFB LEGOS poleward of 88°N).



## References

- Alexandrov, V., Sandven, S., Wahlin, J., and Johannessen, O. M.: The relation between sea ice thickness and freeboard in the Arctic, *The Cryosphere*, 4, 373–380, <https://doi.org/10.5194/tc-4-373-2010>, 2010.
- 835 Balan-Sarajini, B., Tietsche, S., Mayer, M., Balmaseda, M., Zuo, H., de Rosnay, P., Stockdale, T., and Vitart, F.: Year-round impact of winter sea ice thickness observations on seasonal forecasts, *The Cryosphere*, 15, 325–344, <https://doi.org/10.5194/tc-15-325-2021>, 2021.
- Benkiran, M. and Greiner, E.: Impact of the Incremental Analysis Updates on a Real-Time System of the North Atlantic Ocean, <https://doi.org/10.1175/2008JTECHO537.1>, 2008.
- 840 Blockley, E. W. and Peterson, K. A.: Improving Met Office seasonal predictions of Arctic sea ice using assimilation of CryoSat-2 thickness, *The Cryosphere*, 12, 3419–3438, <https://doi.org/10.5194/tc-12-3419-2018>, 2018.
- Bocquet, M., Fleury, S., Piras, F., Rinne, E., Sallila, H., Garnier, F., and Rémy, F.: Arctic sea ice radar freeboard retrieval from the European Remote-Sensing Satellite (ERS-2) using altimetry: toward sea ice thickness observation from 1995 to 2021, *The Cryosphere*, 17, 3013–3039, <https://doi.org/10.5194/tc-17-3013-2023>, 2023.
- 845 Bocquet, M., Fleury, S., Rémy, F., and Piras, F.: Arctic and Antarctic sea ice thickness and volume changes from observations between 1994 and 2023, *Authorea*, 2024.
- Brasseur, P. and Verron, J.: The SEEK filter method for data assimilation in oceanography: a synthesis, *Ocean Dynamics*, 56, 650–661, <https://doi.org/10.1007/s10236-006-0080-3>, 2006.
- Chen, Y., Smith, P., Carrassi, A., Pasmans, I., Bertino, L., Bocquet, M., Finn, T. S., Rampal, P., and Dansereau, V.: 850 Multivariate state and parameter estimation with data assimilation applied to sea-ice models using a Maxwell elasto-brittle rheology, *The Cryosphere*, 18, 2381–2406, <https://doi.org/10.5194/tc-18-2381-2024>, 2024.
- Cheng, S., Chen, Y., Aydođdu, A., Bertino, L., Carrassi, A., Rampal, P., and Jones, C. K. R. T.: Arctic sea ice data assimilation combining an ensemble Kalman filter with a novel Lagrangian sea ice model for the winter 2019–2020, *The Cryosphere*, 17, 1735–1754, <https://doi.org/10.5194/tc-17-1735-2023>, 2023.
- 855 Cipollone, A., Banerjee, D. S., Iovino, D., Aydogdu, A., and Masina, S.: Bivariate sea-ice assimilation for global-ocean analysis–reanalysis, *Ocean Science*, 19, 1375–1392, <https://doi.org/10.5194/os-19-1375-2023>, 2023.
- Cramer, F.: Scientific colour maps, , <https://doi.org/10.5281/zenodo.8409685>, 2023.
- Cramer, F., Shephard, G. E., and Heron, P. J.: The misuse of colour in science communication, *Nat Commun*, 11, 5444, <https://doi.org/10.1038/s41467-020-19160-7>, 2020.
- 860 Fiedler, E. K., Martin, M. J., Blockley, E., Mignac, D., Fournier, N., Ridout, A., Shepherd, A., and Tilling, R.: Assimilation of sea ice thickness derived from CryoSat-2 along-track freeboard measurements into the Met Office’s Forecast Ocean Assimilation Model (FOAM), *The Cryosphere*, 16, 61–85, <https://doi.org/10.5194/tc-16-61-2022>, 2022.
- Fritzner, S., Graverson, R., Christensen, K. H., Rostosky, P., and Wang, K.: Impact of assimilating sea ice concentration, sea ice thickness and snow depth in a coupled ocean–sea ice modelling system, *The Cryosphere*, 13, 491–509, 865 <https://doi.org/10.5194/tc-13-491-2019>, 2019.



- Fritzner, S. M., Graverson, R. G., Wang, K., and Christensen, K. H.: Comparison between a multi-variate nudging method and the ensemble Kalman filter for sea-ice data assimilation, *Journal of Glaciology*, 64, 387–396, <https://doi.org/10.1017/jog.2018.33>, 2018.
- 870 Garnier, F., Fleury, S., Garric, G., Bouffard, J., Tsamados, M., Laforge, A., Bocquet, M., Fredensborg Hansen, R. M., and Remy, F.: Advances in altimetric snow depth estimates using bi-frequency SARAL and CryoSat-2 Ka–Ku measurements, *The Cryosphere*, 15, 5483–5512, <https://doi.org/10.5194/tc-15-5483-2021>, 2021.
- Garnier, F., Bocquet, M., Fleury, S., Bouffard, J., Tsamados, M., Remy, F., Garric, G., and Chenal, A.: Latest Altimetry-Based Sea Ice Freeboard and Volume Inter-Annual Variability in the Antarctic over 2003–2020, *Remote Sensing*, 14, 4741, <https://doi.org/10.3390/rs14194741>, 2022.
- 875 Gilbert, E. and Holmes, C.: 2023’s Antarctic sea ice extent is the lowest on record, *Weather*, 79, 46–51, <https://doi.org/10.1002/wea.4518>, 2024.
- Guerreiro, K., Fleury, S., Zakharova, E., Kouraev, A., Rémy, F., and Maisongrande, P.: Comparison of CryoSat-2 and ENVISAT radar freeboard over Arctic sea ice: toward an improved Envisat freeboard retrieval, *The Cryosphere*, 11, 2059–2073, <https://doi.org/10.5194/tc-11-2059-2017>, 2017.
- 880 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 885 Hutter, N., Losch, M., and Menemenlis, D.: Scaling Properties of Arctic Sea Ice Deformation in a High-Resolution Viscous-Plastic Sea Ice Model and in Satellite Observations, *Journal of Geophysical Research: Oceans*, 123, 672–687, <https://doi.org/10.1002/2017JC013119>, 2018.
- INSITU\_GLO\_PHYBGCWAV\_DISCRETE\_MYNRT\_013\_030: Global Ocean- In-Situ Near-Real-Time Observations, E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). [dataset], <https://doi.org/10.48670/moi-00036>, 2024.
- Ivanova, N., Johannessen, O. M., Pedersen, L. T., and Tonboe, R. T.: Retrieval of Arctic Sea Ice Parameters by Satellite Passive Microwave Sensors: A Comparison of Eleven Sea Ice Concentration Algorithms, *IEEE Transactions on Geoscience and Remote Sensing*, 52, 7233–7246, <https://doi.org/10.1109/TGRS.2014.2310136>, 2014.
- 895 Ivanova, N., Pedersen, L. T., Tonboe, R. T., Kern, S., Heygster, G., Lavergne, T., Sørensen, A., Saldo, R., Dybkjær, G., Brucker, L., and Shokr, M.: Inter-comparison and evaluation of sea ice algorithms: towards further identification of challenges and optimal approach using passive microwave observations, *The Cryosphere*, 9, 1797–1817, <https://doi.org/10.5194/tc-9-1797-2015>, 2015.
- 900 Jena, B., Ravichandran, M., and Turner, J.: Recent Reoccurrence of Large Open-Ocean Polynya on the Maud Rise Seamount, *Geophysical Research Letters*, 46, 4320–4329, <https://doi.org/10.1029/2018GL081482>, 2019.
- Jutila, A., Hendricks, S., Ricker, R., von Albedyll, L., Krumpfen, T., and Haas, C.: Retrieval and parameterisation of sea-ice bulk density from airborne multi-sensor measurements, *The Cryosphere*, 16, 259–275, <https://doi.org/10.5194/tc-16-259-2022>, 2022.



- 905 Kaleschke, L., Tian-Kunze, X., Hendricks, S., and Ricker, R.: SMOS-derived Antarctic thin sea ice thickness: data description and validation in the Weddell Sea, *Earth System Science Data*, 16, 3149–3170, <https://doi.org/10.5194/essd-16-3149-2024>, 2024.
- Kern, S., Khvorostovsky, K., Skourup, H., Rinne, E., Parsakhoo, Z. S., Djepa, V., Wadhams, P., and Sandven, S.: The impact of snow depth, snow density and ice density on sea ice thickness retrieval from satellite radar altimetry: results from the ESA-CCI Sea Ice ECV Project Round Robin Exercise, *The Cryosphere*, 9, 37–52, <https://doi.org/10.5194/tc-9-37-2015>, 2015.
- 910 Kimmritz, M., Counillon, F., Bitz, C. M., Massonnet, F., Bethke, I., and Gao, Y.: Optimising assimilation of sea ice concentration in an Earth system model with a multicategory sea ice model, *Tellus A: Dynamic Meteorology and Oceanography*, 70, 1–23, <https://doi.org/10.1080/16000870.2018.1435945>, 2018.
- Kurtz, N. and Harbeck, J.: CryoSat-2 Level 4 Sea Ice Elevation, Freeboard, and Thickness, Version 1, <https://doi.org/10.5067/96J00KIFDAS8>, 2017.
- 915 Kurtz, N., Studinger, M., Harbeck, J., Onana, V., and Yi, D.: IceBridge Sea Ice Freeboard, Snow Depth, and Thickness Quick Look (NSIDC-0708, Version 1), Boulder, Colorado, USA. NASA National Snow and Ice Data Center Distributed Archive Center [dataset], <https://doi.org/10.5067/GRIXZ91DE0L9>, 2016.
- Kurtz, N. T. and Markus, T.: Satellite observations of Antarctic sea ice thickness and volume, *Journal of Geophysical Research: Oceans*, 117, <https://doi.org/10.1029/2012JC008141>, 2012.
- 920 Kwok, R. and Cunningham, G. F.: Variability of Arctic sea ice thickness and volume from CryoSat-2, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373, 20140157, <https://doi.org/10.1098/rsta.2014.0157>, 2015.
- Kwok, R., Pang, S. S., and Kacimi, S.: Sea ice drift in the Southern Ocean: Regional patterns, variability, and trends, *Elementa: Science of the Anthropocene*, 5, 32, <https://doi.org/10.1525/elementa.226>, 2017.
- 925 Laforge, A., Fleury, S., Dinardo, S., Garnier, F., Remy, F., Benveniste, J., Bouffard, J., and Verley, J.: Toward improved sea ice freeboard observation with SAR altimetry using the physical retracker SAMOSA+, *Advances in Space Research*, 68, 732–745, <https://doi.org/10.1016/j.asr.2020.02.001>, 2021.
- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophysical Research Letters*, 40, 732–737, <https://doi.org/10.1002/grl.50193>, 2013.
- 930 Lee, J.-G. and Ham, Y.-G.: Satellite-Based Data Assimilation System for the Initialization of Arctic Sea Ice Concentration and Thickness Using CICE5, *Frontiers in Climate*, 4, 2022.
- Lee, J.-G. and Ham, Y.-G.: Impact of satellite thickness data assimilation on bias reduction in Arctic sea ice concentration, *npj Clim Atmos Sci*, 6, 1–12, <https://doi.org/10.1038/s41612-023-00402-6>, 2023.
- 935 Lellouche, J.-M., Le Galloudec, O., Drévilion, M., Régnier, C., Greiner, E., Garric, G., Ferry, N., Desportes, C., Testut, C.-E., Bricaud, C., Bourdallé-Badie, R., Tranchant, B., Benkiran, M., Drillet, Y., Daudin, A., and De Nicola, C.: Evaluation of global monitoring and forecasting systems at Mercator Océan, *Ocean Science*, 9, 57–81, <https://doi.org/10.5194/os-9-57-2013>, 2013.
- Lellouche, J.-M., Greiner, E., Bourdallé-Badie, R., Garric, G., Angélique, M., Drevillon, M., Bricaud, C., Hamon, M., Le Galloudec, O., Regnier, C., Candela, T., Testut, C.-E., Gasparin, F., Ruggiero, G., Benkiran, M., Drillet, Y., and Le Traon, P.-



- 940 Y.: The Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis, *Front. Earth Sci.*, 9, <https://doi.org/10.3389/feart.2021.698876>, 2021.
- Liu, X., Yao, J., Zhang, S., Wu, T., Chen, Z., Fang, Y., Chu, M., Yan, J., and Jie, W.: A Coordinated Sea-Ice Assimilation Scheme Jointly Using Sea-Ice Concentration and Thickness Observations With a Coupled Climate Model, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003608, <https://doi.org/10.1029/2023MS003608>, 2024.
- 945 Luo, H., Yang, Q., Mu, L., Tian-Kunze, X., Nerger, L., Mazloff, M., Kaleschke, L., and Chen, D.: DASSO: a data assimilation system for the Southern Ocean that utilizes both sea-ice concentration and thickness observations, *Journal of Glaciology*, 67, 1235–1240, <https://doi.org/10.1017/jog.2021.57>, 2021.
- Madec, G. and Imbard, M.: A global ocean mesh to overcome the North Pole singularity, *Climate Dynamics*, 12, 381–388, <https://doi.org/10.1007/BF00211684>, 1996.
- 950 Madec, G., Bourdallé-Badie, R., Chanut, J., Clementi, E., Coward, A., Ethé, C., Iovino, D., Lea, D., Lévy, C., Lovato, T., Martin, N., Masson, S., Mocavero, S., Rousset, C., Storkey, D., Müeller, S., Nurser, G., Bell, M., Samson, G., Mathiot, P., Mele, F., and Moulin, A.: NEMO ocean engine, <https://doi.org/10.5281/zenodo.6334656>, 2022.
- Mallett, R. D. C., Lawrence, I. R., Stroeve, J. C., Landy, J. C., and Tsamados, M.: Brief communication: Conventional assumptions involving the speed of radar waves in snow introduce systematic underestimates to sea ice thickness and seasonal growth rate estimates, *The Cryosphere*, 14, 251–260, <https://doi.org/10.5194/tc-14-251-2020>, 2020.
- 955 Massonnet, F., Fichet, T., Goosse, H., Vancoppenolle, M., Mathiot, P., and König Beatty, C.: On the influence of model physics on simulations of Arctic and Antarctic sea ice, *The Cryosphere*, 5, 687–699, <https://doi.org/10.5194/tc-5-687-2011>, 2011.
- Massonnet, F., Fichet, T., and Goosse, H.: Prospects for improved seasonal Arctic sea ice predictions from multivariate data assimilation, *Ocean Modelling*, 88, 16–25, <https://doi.org/10.1016/j.ocemod.2014.12.013>, 2015.
- 960 Meier, W. N., Fetterer, F., Savoie, M., Mallory, S., Duerr, R., and Stroeve, J.: NOAA/NSIDC Climate Data Record of Passive Microwave Sea Ice Concentration, Version 3, <https://doi.org/10.7265/N59P2ZTG>, 2017.
- Mignac, D., Martin, M., Fiedler, E., Blockley, E., and Fournier, N.: Improving the Met Office’s Forecast Ocean Assimilation Model (FOAM) with the assimilation of satellite-derived sea-ice thickness data from CryoSat-2 and SMOS in the Arctic, *Quarterly Journal of the Royal Meteorological Society*, 148, 1144–1167, <https://doi.org/10.1002/qj.4252>, 2022.
- 965 Mu, L., Yang, Q., Losch, M., Losa, S. N., Ricker, R., Nerger, L., and Liang, X.: Improving sea ice thickness estimates by assimilating CryoSat-2 and SMOS sea ice thickness data simultaneously, *Quarterly Journal of the Royal Meteorological Society*, 144, 529–538, <https://doi.org/10.1002/qj.3225>, 2018.
- Mu, L., Nerger, L., Tang, Q., Loza, S. N., Sidorenko, D., Wang, Q., Semmler, T., Zampieri, L., Losch, M., and Goessling, H. F.: Toward a Data Assimilation System for Seamless Sea Ice Prediction Based on the AWI Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001937, <https://doi.org/10.1029/2019MS001937>, 2020.
- 970 OSI SAF: OSI SAF Global Sea Ice Concentration (AMSR-2) (OSI-408-a), [https://doi.org/10.15770/EUM\\_SAF\\_OSI\\_NRT\\_2023](https://doi.org/10.15770/EUM_SAF_OSI_NRT_2023), 2017.
- OSI SAF: OSI SAF Global sea ice concentration climate data record 1978-2020 (3), [https://doi.org/10.15770/EUM\\_SAF\\_OSI\\_0013](https://doi.org/10.15770/EUM_SAF_OSI_0013), 2022.
- 975



- Peng, G., Meier, W. N., Scott, D. J., and Savoie, M. H.: A long-term and reproducible passive microwave sea ice concentration data record for climate studies and monitoring, *Earth System Science Data*, 5, 311–318, <https://doi.org/10.5194/essd-5-311-2013>, 2013.
- 980 Perovich, D., Meier, W., Tschudi, M., Hendricks, S., Petty, A. A., Divine, D., Farrell, S., Gerland, S., Haas, C., Kaleschke, L., Pavlova, O., Ricker, R., Tian-Kunze, X., Webster, M., and Wood, K.: Arctic Report Card 2020: Sea Ice, 2020.
- Petty, A. A., Bagnardi, M., Kurtz, N. T., Tilling, R., Fons, S., Armitage, T., Horvat, C., and Kwok, R.: Assessment of ICESat-2 Sea Ice Surface Classification with Sentinel-2 Imagery: Implications for Freeboard and New Estimates of Lead and Floe Geometry, *Earth and Space Science*, 8, e2020EA001491, <https://doi.org/10.1029/2020EA001491>, 2021.
- 985 Petty, A. A., Kwok, R., Bagnardi, M., Ivanoff, A., Kurtz, N., Lee, J., Wimert, J., and Hancock, D.: ATLAS/ICESat-2 L3B Daily and Monthly Gridded Sea Ice Freeboard (ATL20, Version 4), Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Archive Center [dataset], <https://doi.org/10.5067/ATLAS/ATL20.004>, 2023.
- Reiser, F., Willmes, S., Hausmann, U., and Heinemann, G.: Predominant Sea Ice Fracture Zones Around Antarctica and Their Relation to Bathymetric Features, *Geophysical Research Letters*, 46, 12117–12124, <https://doi.org/10.1029/2019GL084624>, 2019.
- 990 Ricker, R., Hendricks, S., Helm, V., Skourup, H., and Davidson, M.: Sensitivity of CryoSat-2 Arctic sea-ice freeboard and thickness on radar-waveform interpretation, *The Cryosphere*, 8, 1607–1622, <https://doi.org/10.5194/tc-8-1607-2014>, 2014.
- Ricker, R., Hendricks, S., Kaleschke, L., Tian-Kunze, X., King, J., and Haas, C.: A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data, *The Cryosphere*, 11, 1607–1623, <https://doi.org/10.5194/tc-11-1607-2017>, 2017.
- 995 Roach, L. A., Dörr, J., Holmes, C. R., Massonnet, F., Blockley, E. W., Notz, D., Rackow, T., Raphael, M. N., O’Farrell, S. P., Bailey, D. A., and Bitz, C. M.: Antarctic Sea Ice Area in CMIP6, *Geophysical Research Letters*, 47, e2019GL086729, <https://doi.org/10.1029/2019GL086729>, 2020.
- Sallila, H., Farrell, S. L., McCurry, J., and Rinne, E.: Assessment of contemporary satellite sea ice thickness products for Arctic sea ice, *The Cryosphere*, 13, 1187–1213, <https://doi.org/10.5194/tc-13-1187-2019>, 2019.
- 1000 SEALEVEL\_GLO\_PHY\_L3\_NRT\_008\_044: Global Ocean Along Track L 3 Sea Surface Heights NRT, E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). [dataset], <https://doi.org/10.48670/moi-00147>, 2023.
- Shu, Q., Wang, Q., Song, Z., Qiao, F., Zhao, J., Chu, M., and Li, X.: Assessment of Sea Ice Extent in CMIP6 With Comparison to Observations and CMIP5, *Geophysical Research Letters*, 47, e2020GL087965, <https://doi.org/10.1029/2020GL087965>, 2020.
- 1005 Sievers, I., Rasmussen, T. A. S., and Stenseng, L.: Assimilating CryoSat-2 freeboard to improve Arctic sea ice thickness estimates, *The Cryosphere*, 17, 3721–3738, <https://doi.org/10.5194/tc-17-3721-2023>, 2023.
- Spreen, G., Kwok, R., Menemenlis, D., and Nguyen, A. T.: Sea-ice deformation in a coupled ocean–sea-ice model and in satellite remote sensing data, *The Cryosphere*, 11, 1553–1573, <https://doi.org/10.5194/tc-11-1553-2017>, 2017.
- 1010 SST\_GLO\_SST\_L4\_NRT\_OBSERVATIONS\_010\_001: Global Ocean OSTIA Sea Surface Temperature and Sea Ice Analysis, E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). [dataset], <https://doi.org/10.48670/moi-00165>, 2023.



- Sumata, H., Divine, D., and de Steur, L.: Monthly mean sea ice draft from the Fram Strait Arctic Outflow Observatory since 1990, <https://doi.org/10.21334/npolar.2021.5b717274>, 2021.
- 1015 Tietsche, S., Notz, D., Jungclaus, J. H., and Marotzke, J.: Assimilation of sea-ice concentration in a global climate model – physical and statistical aspects, *Ocean Science*, 9, 19–36, <https://doi.org/10.5194/os-9-19-2013>, 2013.
- Tilling, R. L., Ridout, A., and Shepherd, A.: Estimating Arctic sea ice thickness and volume using CryoSat-2 radar altimeter data, *Advances in Space Research*, 62, 1203–1225, <https://doi.org/10.1016/j.asr.2017.10.051>, 2018.
- 1020 Uotila, P., Goosse, H., Haines, K., Chevallier, M., Barthélemy, A., Bricaud, C., Carton, J., Fučkar, N., Garric, G., Iovino, D., Kauker, F., Korhonen, M., Lien, V. S., Marnela, M., Massonnet, F., Mignac, D., Peterson, K. A., Sadikni, R., Shi, L., Tietsche, S., Toyoda, T., Xie, J., and Zhang, Z.: An assessment of ten ocean reanalyses in the polar regions, *Clim Dyn*, 52, 1613–1650, <https://doi.org/10.1007/s00382-018-4242-z>, 2019.
- Vancoppenolle, M., Rousset, C., Blockley, E., Aksenov, Y., Feltham, D., Fichefet, T., Garric, G., Guémas, V., Iovino, D., Keeley, S., Madec, G., Massonnet, F., Ridley, J., Schroeder, D., and Tietsche, S.: SI3, the NEMO Sea Ice Engine, <https://doi.org/10.5281/zenodo.7534900>, 2023.
- 1025 Wang, Q., Danilov, S., Jung, T., Kaleschke, L., and Wernecke, A.: Sea ice leads in the Arctic Ocean: Model assessment, interannual variability and trends, *Geophysical Research Letters*, 43, 7019–7027, <https://doi.org/10.1002/2016GL068696>, 2016.
- Warren, S. G., Rigor, I. G., Untersteiner, N., Radionov, V. F., Bryazgin, N. N., Aleksandrov, Y. I., and Colony, R.: Snow Depth on Arctic Sea Ice, *J. Climate*, 12, 1814–1829, [https://doi.org/10.1175/1520-0442\(1999\)012<1814:SDOASI>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1814:SDOASI>2.0.CO;2), 1999.
- 1030 Williams, N., Byrne, N., Feltham, D., Van Leeuwen, P. J., Bannister, R., Schroeder, D., Ridout, A., and Nerger, L.: The effects of assimilating a sub-grid-scale sea ice thickness distribution in a new Arctic sea ice data assimilation system, *The Cryosphere*, 17, 2509–2532, <https://doi.org/10.5194/tc-17-2509-2023>, 2023.
- Xie, J., Counillon, F., Bertino, L., Tian-Kunze, X., and Kaleschke, L.: Benefits of assimilating thin sea ice thickness from SMOS into the TOPAZ system, *The Cryosphere*, 10, 2745–2761, <https://doi.org/10.5194/tc-10-2745-2016>, 2016.
- 1035 Xie, J., Counillon, F., and Bertino, L.: Impact of assimilating a merged sea-ice thickness from CryoSat-2 and SMOS in the Arctic reanalysis, *The Cryosphere*, 12, 3671–3691, <https://doi.org/10.5194/tc-12-3671-2018>, 2018.
- Zhang, X., Zhao, Q., Gao, G., Zhang, J., Bao, M., and Meng, J.: Impact and Correction of Sea Ice, Snow, and Seawater Density on Arctic Sea-Ice Thickness Retrieval From Ku-Band SAR Altimeters, *IEEE Journal on Miniaturization for Air and Space Systems*, 3, 276–283, <https://doi.org/10.1109/JMASS.2022.3212880>, 2022.
- 1040 Zhang, Y., Cheng, X., Liu, J., and Hui, F.: The potential of sea ice leads as a predictor for summer Arctic sea ice extent, *The Cryosphere*, 12, 3747–3757, <https://doi.org/10.5194/tc-12-3747-2018>, 2018.