

## Extremophiles

November 2007, Volume 11 (6) : Pages 747-757

<http://dx.doi.org/10.1007/s00792-007-0092-z>

© 2007 Springer. Part of Springer Science+Business Media

Archimer, archive institutionnelle de l'Ifremer

<http://www.ifremer.fr/docelec/>

The original publication is available at <http://www.springerlink.com>

---

## Continuous enrichment cultures: insights into prokaryotic diversity and metabolic interactions in deep-sea vent chimneys

Anne Postec<sup>1,3,\*</sup>, Françoise Lesongeur<sup>1</sup>, Patricia Pignet<sup>1</sup>, Bernard Ollivier<sup>2</sup>, Joël Querellou<sup>1</sup> and Anne Godfroy<sup>1</sup>

<sup>1</sup>Laboratoire de Microbiologie des Environnements Extrêmes, UMR 6197, IFREMER, Centre de Brest, BP 70, 29280 Plouzané, France

<sup>2</sup>Laboratoire IRD de Microbiologie des Anaérobies, UR 101, Universités de Provence et de la Méditerranée, CESB-ESIL, case 925, 163 avenue de Luminy, 13288 Marseille, France

<sup>3</sup>Geologisches Institut, ETH Zentrum, Universitätsstrasse 16, 8092 Zürich, Switzerland

\*: Corresponding author : A. Postec, email address : [anne.postec@univ-brest.fr](mailto:anne.postec@univ-brest.fr)

---

### Abstract:

The prokaryotic diversity of culturable thermophilic communities of deep-sea hydrothermal chimneys was analysed using a continuous enrichment culture performed in a gas-lift bioreactor, and compared to classical batch enrichment cultures in vials. Cultures were conducted at 60°C and pH 6.5 using a complex medium containing carbohydrates, peptides and sulphur, and inoculated with a sample of a hydrothermal black chimney collected at the Rainbow field, Mid-Atlantic Ridge, at 2,275 m depth. To assess the relevance of both culture methods, bacterial and archaeal diversity was studied using cloning and sequencing, DGGE, and whole-cell hybridisation of 16S rRNA genes. Sequences of heterotrophic microorganisms belonging to the genera *Marinitoga*, *Thermosipho*, *Caminicella* (Bacteria) and *Thermococcus* (Archaea) were obtained from both batch and continuous enrichment cultures while sequences of the autotrophic bacterial genera *Deferribacter* and *Thermodesulfator* were only detected in the continuous bioreactor culture. It is presumed that over time constant metabolite exchanges will have occurred in the continuous enrichment culture enabling the development of a more diverse prokaryotic community. In particular, CO<sub>2</sub> and H<sub>2</sub> produced by the heterotrophic population would support the growth of autotrophic populations. Therefore, continuous enrichment culture is a useful technique to grow over time environmentally representative microbial communities and obtain insights into prokaryotic species interactions that play a crucial role in deep hydrothermal environments.

**Keywords:** Microbial diversity - Deep-sea hydrothermal vent - Continuous enrichment cultures - Bioreactor - Thermophiles - 16S rRNA gene - Metabolic interactions

## 46 **Introduction**

47

48 The widespread application of 16S rRNA gene based molecular methods to identify microorganisms in natural  
49 samples has revealed an extensive and, in many cases, unexpected microbial diversity. Within deep-sea  
50 hydrothermal environments, the diversity of microbial communities associated with *in situ* colonizers  
51 (McCliment et al. 2006), mats (Moussard et al. 2006), animals (DeChaine et al. 2006), sediments (Inagaki et al.  
52 2006) and chimneys (Kormas et al. 2006) have been reported in recent molecular surveys. The rise of molecular  
53 microbial ecology has resulted in the detection of many microorganisms that have as yet not been cultivated, for  
54 example the widespread Marine *Crenarchaeota* Group I (MGI) and the *Korarchaeota*. To understand the  
55 physiology and ecological significance of these uncultivated microorganisms, an effort has to be made to  
56 improve and develop cultural approaches. Indeed, culture conditions routinely used reveal only a small fraction  
57 of the global microbial community. As an alternative to batch cultures in vials, a gas-lift bioreactor was  
58 developed to grow anaerobic and hyperthermophilic microorganisms in continuous culture (Raven et al. 1992).  
59 Recently, it has been used to study the metabolism of members of the *Thermococcales*, including *Pyrococcus*  
60 *abyssi* (Godfroy et al. 2000), to develop minimal media and to optimize the growth conditions of *Pyrococcus*  
61 *furiosus* and *Thermococcus hydrothermalis* (Raven and Sharp 1997; Postec et al. 2005a). In addition to the study  
62 of pure cultures, the gas-lift bioreactor can also be used to cultivate representative microorganisms from  
63 environmental samples in continuous enrichment culture under controlled conditions. The bioreactor allows the  
64 long-term cultivation of microbes by enabling a continuous substrate supply, the elimination of volatile  
65 metabolic end-products (potentially toxic for microbial growth) by gas sparging and pH and temperature  
66 regulation. These features help to grow less dominant microorganisms having poor representation, long latency  
67 phase and/or slow growth. In a previous study, the gas-lift bioreactor was used to enrich microorganisms from a  
68 black smoker collected at 2275 m depth on the Rainbow hydrothermal field of the Mid-Atlantic Ridge (Postec et  
69 al. 2005b). A fifty-days continuous culture at 90°C on a rich medium containing sulphur under anaerobic  
70 conditions demonstrated a large diversity inside the cultivated community, including (in addition to archaeal  
71 species belonging to the order *Thermococcales*) moderately thermophilic members of the orders *Clostridiales* and  
72 *Thermotogales*, and members of the *Epsilonproteobacteria* that were not detected in vial cultures. In the present  
73 study, the same black smoker chimney was used as inoculum to perform similar experiments on both batch and  
74 continuous enrichment cultures under the same conditions, except the temperature lowered to 60°C. The  
75 microbial diversity of the communities enriched in vials and bioreactor was analysed using the following

76 techniques based on the 16S rRNA genes: cloning, sequencing, denaturing gradient gel electrophoresis (DGGE)  
77 and whole-cell hybridisation. Molecular results were the guidelines for subsequent isolation of microorganisms  
78 from the enrichment cultures.

79

## 80 **Materials and methods**

81

### 82 **Samples**

83

84 During the ATOS cruise (European project VENTOX) on the Rainbow field (36°13'N 33°54W, 2275 m) located  
85 on the Mid-Atlantic Ridge (MAR), an active black smoker was collected by the Remotely Operated Vehicle  
86 (ROV) Victor, and brought to the surface in an insulated box under aseptic conditions. On board, eight fragments  
87 of the chimney were subsampled according to mineral zonations from the inner part to the outer part. The  
88 subsamples were crushed in an anaerobic chamber (La Calhene, France) and stored in sterile serum vials filled  
89 with sterile seawater containing 0.5 mg l<sup>-1</sup> of Na<sub>2</sub>S. All subsamples were pooled to represent the whole chimney  
90 and this suspension was used to inoculate the enrichment cultures.

91

### 92 **Continuous enrichment culture in bioreactor: conditions and monitoring**

93

#### 94 *Medium*

95 The growth medium was the modified SME medium (Sharp and Raven 1997) in which yeast and peptone were  
96 replaced by: 1 g yeast extract (Difco), 0.5 g casaminoacids (Difco), 0.4 g glucose, 0.4 g dextrin (from corn), 0.2  
97 g galactose, 0.2 g dextran, 0.1 g glycogen, 0.2 g pyruvate and 0.1 g acetate (all purchased from Sigma). The  
98 medium was supplemented with 3 g l<sup>-1</sup> colloidal sulphur.

99

#### 100 *Culture conditions*

101 The medium was sterilized by filtration (Sartroban, 0.22 µm) in a 20 litre Nalgene bottle containing the colloidal  
102 sulphur, previously sterilized by heating twice at 100°C for 30 min on two successive days. The culture was  
103 performed at 60°C and pH 6.5 in a 2 litre glass gas-lift bioreactor as previously described (Raven et al. 1992;  
104 Godfroy et al. 2005). The bioreactor was inoculated at 2% (v/v) with the chimney sample suspension. The  
105 temperature was controlled by a heated circulating bath filled with water and monitored with a standard PT100

106 probe covered with Teflon. The pH was monitored using a combination gel pH electrode (Mettler Toledo). Acid  
107 (1N HCl) and base (1N NaOH) were added with peristaltic pumps (Masterflex). Temperature and pH were  
108 controlled with a 4-20 mA controller and AFS Biocommand system from New Brunswick (Nijmegen,  
109 Netherlands). Fresh medium addition and culture withdrawal were performed using peristaltic pumps  
110 (Masterflex). The culture was sparged with  $N_2$  ( $0.1 \text{ v } \bar{v}^{-1} \text{ min}^{-1}$ ) to maintain anaerobic conditions, and to  
111 eliminate volatile metabolic end products that might inhibit the growth of microorganisms (i.e.  $H_2S$ ). The  
112 bioreactor was maintained as a batch culture for the first 34 h to prevent wash-out of the cells before they have  
113 grown. After 34 h, fresh medium was provided by applying a dilution rate of  $0.04 \text{ h}^{-1}$  ( $80 \text{ ml h}^{-1}$ ). To test the  
114 effect of the temperature on the composition of the cultivated microbial community, the temperature was  
115 increased from  $60^\circ\text{C}$  to  $70^\circ\text{C}$  at T31 (day 31) and to  $80^\circ\text{C}$  at T36 (day 36).

116

#### 117 *Culture monitoring and sample preservation*

118 The culture was maintained for 45 days. Culture samples from the bioreactor were collected every 24 hours,  
119 from T0 (day 0) to T45 (day 45). Cell concentration was determined every day by direct cell counting, using a  
120 Thoma chamber (0.02 mm depth) viewed with an Olympus BX60 phase contrast microscope ( $\times 400$ ). For each  
121 sampling, 10 ml of culture were preserved anaerobically at  $4^\circ\text{C}$  in a serum vial, and eight cryotubes containing  
122 each 1.8 ml of culture were frozen at  $-20^\circ\text{C}$  with 5% DMSO (v/v). For DNA extraction, cells were recovered  
123 from 15 ml of culture by centrifugation (20 min at 8,000 g). Cell pellets were washed with  $23 \text{ g l}^{-1}$  sterile NaCl,  
124 then mixed to 5 ml of lysis buffer TE Na 1X (Tris-HCl pH 8, 100 mM ; NaCl, 100 mM, EDTA pH 8, 50 mM),  
125 and stored at  $-20^\circ\text{C}$  until the DNA extraction procedure was undertaken. For whole-cell hybridisation, 12 ml of  
126 culture sample were fixed for two hours with 3% (v/v) formaldehyde. Fixed cells recovered by centrifugation (10  
127 min at 6,000 rpm) were washed with PBS buffer (phosphate-buffered: 8 g NaCl, 0.2 g KCl, 1.44 g  $Na_2HPO_4$ ,  
128 0.24 g  $KH_2PO_4$ , per litre of distilled water, pH 7.4) before storage in 50% (v/v) ethanol in PBS at  $-20^\circ\text{C}$ . For  
129 HPLC analyses, 1.5 ml of culture sample was centrifuged (10 min at 10,000 rpm) and the supernatant was stored  
130 at  $4^\circ\text{C}$  until analysis.

131

#### 132 Batch enrichment cultures in vial: conditions and monitoring

133

134 Cultures were performed anaerobically in 100 ml serum vials (Godfroy et al. 1996), using the medium described  
135 for the enrichment culture in the bioreactor to which  $6.05 \text{ g l}^{-1}$  PIPES was added and colloidal sulphur was

136 replaced by 10 g l<sup>-1</sup> sulphur powder. Inoculation was performed with 2% (v/v) of the chimney sample  
137 suspension. The same temperature and pH (60°C, pH 6.5) as in the bioreactor were used. Cell pellets dedicated  
138 to DNA extractions were recovered from 15 ml of culture after 24 h (sample A1) and 41 h (sample A2) of  
139 incubation. The 24 h culture (A1) was subcultured for 17 h in vial in the same conditions (sample B).

140

#### 141 Nucleic acid extraction

142

143 DNA was extracted from frozen cell pellets in lysis buffer (cells recovered from 15 ml of culture, see procedure  
144 above). A modified version of the protocol described by Alain et al. (2002) was followed combining chemical  
145 and enzymatic lysis. For these culture samples, the applied lysis treatment was 1.5 h and 2.5 h. Afterwards, intact  
146 cells could not be observed by microscopy. Supernatants from the 1.5 h and 2.5 h lysis were pooled and  
147 extracted twice with equal volumes of buffered (pH 8.0) PCI (phenol/chloroform/isoamyl alcohol : 25/24/1) and  
148 once with an equal volume of chloroform. DNA was finally precipitated by addition of 70 % (v/v) isopropanol.  
149 After centrifugation at 11,000×g for 30 min, DNA was air dried before being resuspended in 250 µl TE 1×  
150 buffer (10 mM Tris-HCl, 2 mM EDTA, pH 7.5). The extracted DNA quality was routinely checked using 0.8%  
151 agarose-TAE-1× gels.

152

#### 153 Amplification of the 16S rRNA gene and DGGE analysis

154

155 The variable v3 region of 16S rRNA genes from extracted DNA were amplified using the primers 341F-GC and  
156 907R specific to the bacterial domain (Muyzer et al. 1993; Muyzer and Smalla 1998). The PCR procedure and  
157 the analysis of the fragments by DGGE using the Bio-Rad Dcode apparatus are described in Muyzer et al.  
158 (1993). Electrophoresis conditions, gel staining, DGGE band extraction, DNA reamplification, and PCR product  
159 purification were performed using the conditions described by Postec et al. (2005b).

160

#### 161 Amplification of the 16S rRNA gene and cloning

162

163 Archaeal DNA was amplified using the primer A24F (5'-TTC CGG TTG ATC CTG CCG GA -3') and the  
164 reverse primer 1407R (5'-GAC GGG CGG TGW GTR CAA-3') or alternatively A23SR (5'-CTT TCG GTC  
165 GCC CCT ACT-3', position 257-234 on *Thermococcus celer* 23S rRNA gene sequence). Bacterial DNA was

166 amplified using primer E8F (5' -AGA GTT TGA TCA TGG CTC AG-3') and the reverse primer U1492R (5'-  
167 GTT ACC TTG TTA CGA CTT-3'). PCR reactions were performed on a Robocycler Gradient 96 (Stratagene)  
168 (Wery et al. 2002; Nercessian et al. 2003). PCR products were then checked on a 0.8% (w/v) agarose gel and  
169 directly cloned using the TOPO TA Cloning<sup>®</sup> kit (pCR2.1 vector), according to the manufacturer's instructions  
170 (Invitrogen). Clone libraries were constructed by transforming *E. coli* TOP10F' cells. An archaeal and a bacterial  
171 library were constructed from two culture samples from the enrichment culture in bioreactor named T7 and T28,  
172 collected respectively after 7 and 28 days of culture, and from each sample of enrichment cultures in vials (A1,  
173 A2 and B).

174

175 16S rRNA gene sequencing and phylogenetic analysis

176

177 DNA fragments obtained by DGGE were sequenced by Genome Express S.A. (Grenoble, France). From clone  
178 libraries, each clone was cultivated overnight at 37°C with shaking (320 rpm) on deepwell microplates, in 1 ml  
179 Luria Bertani broth 2X medium containing ampicillin (50 µg ml<sup>-1</sup>). Plasmids were extracted and purified using  
180 Montage Plasmid Miniprep<sub>96</sub> Kits (Millipore) and partially sequenced using the BigDye Terminator chemistry  
181 with an automated capillary sequencer (Applied Biosystem). Sequences were compared to all GenBank, RefSeq  
182 Nucleotides, EMBL, DDBJ and PDB sequences using the BLAST (<http://www.ncbi.nlm.nih.gov/BLAST>)  
183 network service (Altschul et al. 1990) in order to determine phylogenetic affiliations and detect chimeric  
184 sequences. Alignment of 16S rRNA gene sequences was performed using the CLUSTALW program (Thompson  
185 et al. 1994), then refined manually using the SEAVIEW program (Galtier 1996). Sequences displaying more  
186 than 97% similarity were considered to be related and grouped in the same phylotype. Complete 16S rRNA gene  
187 sequences were obtained for the representative clone of each unique phylotype: the related partial sequences  
188 were first assembled using the SEQMAN module of the DNASTAR software (Madison, WI, USA), and the  
189 complete sequences were analysed with the BLAST program.

190

191 Whole-cell hybridisation

192

193 Fixed cells in PBS/ethanol 50% (v/v) (fixation procedure described above) were diluted if necessary and filtered  
194 on a 0.2 µm pore size white polycarbonate filter (Isopore Membrane Filters, Millipore) laying on a nitrocellulose  
195 membrane. After drying at room temperature, cells were hybridised with the archaeal universal probe ARCH915

196 (5'-GTG CTC CCC CGC CAA TTC CT-3') labelled with indocarbocyanin (Cy3) (Eurogentec) as well as with  
197 the universal bacterial probe EUB338 (5'-GCT GCC TCC CGT AGG AGT-3') labelled with fluorescein-  
198 isothiocyanate (FITC) (Eurogentec). Whole-cell hybridisations were carried out at 46°C after addition of 1.5 µl  
199 ARCH915 and 1.5 µl EUB338 each at 50 ng µl<sup>-1</sup> and 12 µl hybridisation solution [NaCl 5 M: 360 µl, Tris-HCl 1  
200 M pH 7.4: 200 µl, formamide: 400 µl, sodium dodecyl sulphate (SDS) 10%: 1 µl, deionised water: 1039 µl] onto  
201 each filter. After 2 h hybridisation, the filters were rinsed up 15 min at 48°C in a wash solution (NaCl 5M: 1.8  
202 ml, Tris-HCl 1 M pH 7.4: 5 ml, SDS 10%: 25 µl, deionised water: 43.2 ml). After the hybridisation procedure,  
203 each sample was stained with 10 µl of 4',6'-diamidino-2-phenylindole (DAPI 1 µg ml<sup>-1</sup>). After addition of  
204 Citifluor (Citifluor, UK) to the filters, the hybridised cells were viewed with an Olympus microscope (BX60)  
205 equipped with a UV lamp and filters for DAPI (excitation 365 nm, emission 397 nm), FITC (excitation 492 nm,  
206 emission 520 nm), or Cy3 (excitation 550 nm, emission 570 nm).

207

208 Analyses of amino acids, organic acids, and glucose

209

210 The HPLC procedure used for analyses of amino acids, organic acids and glucose is in Wery et al. (2001).

211

212 Gas analyses

213

214 The gas outflow from the bioreactor was directly analysed using a MTI M200D micro gas chromatograph  
215 equipped with a thermal conductivity detector. A Molecular Sieve column with argon as the carrier gas was used  
216 at a temperature of 30°C to detect H<sub>2</sub>. CO<sub>2</sub> and H<sub>2</sub>S were detected with a Poraplot U column at 100°C, with  
217 helium as the carrier gas.

218

219 Subcultures and isolations

220

221 The media employed for the subcultures were designed to cultivate heterotrophic as well as autotrophic  
222 microorganisms that might be expected to grow with different electron acceptors (sulphur, nitrate, sulphate).  
223 Isolation of strains whose phylotypes were recovered in the clone libraries was attempted from the enrichment  
224 culture samples T7 and T28 from the bioreactor. Four culture media were used. The enrichment medium was  
225 used as described above and also according to the three following modifications. Vitamins and minerals were

226 preserved and organic substrates were replaced by: (1)  $\text{NH}_4\text{Cl}$   $0.03 \text{ g l}^{-1}$ , acetate  $0.016 \text{ g l}^{-1}$ , yeast extract  $0.01 \text{ g l}^{-1}$ , and sulphur  $5 \text{ g l}^{-1}$  in the DS medium, or (2)  $\text{NaNO}_3$   $0.2 \text{ g l}^{-1}$  in the DN medium, or (3)  $\text{Na}_2\text{SO}_4$   $0.3 \text{ g l}^{-1}$  in the  
227  
228 T medium. The DS, DN and T media were adjusted at pH 6.5. The DS medium was sterilized by tyndallisation  
229 (twice 30 min at  $100^\circ\text{C}$ ) while the DN and T media were autoclaved (20 min at  $121^\circ\text{C}$ ). In the anaerobic  
230 chamber, the media were reduced with  $\text{Na}_2\text{S}$  (final concentration:  $0.05 \text{ g l}^{-1}$ ), then aliquoted into Hungate tubes  
231 or penicillin vials under  $\text{N}_2/\text{H}_2/\text{CO}_2$  (90:5:5). The gas phase was then replaced performing 10 cycles of vacuum  
232 extraction / addition of  $\text{H}_2/\text{CO}_2$  (80/20, v/v, 2 bar). All the incubations were performed at  $60^\circ\text{C}$  and pH6. Strains  
233 were isolated by repeated dilutions-to-extinction cultures or streaking on solidified enrichment medium.

234

235 Nucleotide sequence accession numbers

236

237 The sequence data used in this study have been submitted to the EMBL databases under accession number  
238 AJ874300 to AJ874328.

239

## 240 **Results**

241

242 Monitoring of the continuous enrichment culture in bioreactor: cell morphologies, whole-cell hybridisation,  
243 DGGE, gas chromatography and HPLC

244

245 Starting from  $2.7 \cdot 10^6 \text{ cell ml}^{-1}$  at T0, cell density reached  $7.4 \cdot 10^8 \text{ cell ml}^{-1}$  at T2 and its maximal value  $2.2 \cdot 10^9$   
246  $\text{cell ml}^{-1}$  at T31 (Fig. 1). Coccoid cells single or in pairs were dominant at T2 (Fig. 2a, 2d). Afterwards rods  
247 displaying various morphologies became widely dominant (Fig. 2b, 2e). Short, rod-shaped cells appeared single  
248 or in chains within an outer sheath-like structure, similarly to the specific toga of *Thermotogales*. Long rods  
249 exhibiting a terminal endospore were observed. From T27, coccoid cells, single or in pairs, increased in density  
250 compared to the rod morphologies. At the end of the culture, rod morphologies had nearly disappeared.

251 The relative proportions of archaeal and bacterial cells were determined by whole-cell hybridisation in eleven  
252 culture samples of the bioreactor, from T2 to T41 (Fig. 1). Approximately 2,000 cells were counted on filters for  
253 each sample. 99.0% of the cells detected at T2 belonged to the *Archaea* (Fig. 2a, 2d); *Bacteria* were dominant  
254 from T4 to T31 (between 94.8 to 99.8% until T24, then 60.0% at T28 and 54.6% at T31). At T36 and T41,  
255 *Archaea* became predominant again, representing 61.6% of the cells at T36 and 98.8% of the cells at T41.

256 Temperature was increased from 60°C to 70°C at T31, and from 70°C to 80°C at T36, which was associated  
257 with a significant decrease in the cell density ( $2.2 \cdot 10^9$  cell ml<sup>-1</sup> at T31 and  $1.8 \cdot 10^7$  cell ml<sup>-1</sup> at T41).

258 The forty-five days enrichment culture was further investigated by DGGE analysis. The v3 hypervariable region  
259 of the bacterial 16S rRNA gene was amplified from T1, T4, T9, T13, T16, T24, T28, T31 and T36 culture  
260 samples. No amplification was obtained from samples collected after T36, probably due to the low cell density  
261 corresponding to the temperature increase at T36. Representative DGGE bands migrating at different distances  
262 and originating from various samples were extracted from gels (not shown) and re-amplified directly. The  
263 DGGE sequence types (approximately 500 pb) were affiliated to *Thermosipho* MV1063 (99% identity),  
264 *Marinitoga camini* (96%), *Caminicella sporogenes* (99-98%) and *Deferribacter abyssi* (87%). *Thermosipho* spp.  
265 was detected from T1 to T36, *Marinitoga* spp. at T31, *Caminicella* spp. from T4 to T31 and *Deferribacter* spp.  
266 from T1 to T31.

267 H<sub>2</sub>, CO<sub>2</sub> and H<sub>2</sub>S were detected in T8, T21, T30 and T38 samples by gas chromatography. H<sub>2</sub>S production was  
268 also detected using Zn acetate strips (Lead Acetate, Whatman) from T2 until the end of the culture. From T3  
269 until T41, HPLC analysis of the free amino acids in the culture medium showed that they were all completely  
270 consumed, and might be therefore a limiting factor for microbial growth. Analysis of glucose by HPLC indicated  
271 an initial concentration of 0.440 g l<sup>-1</sup>, between T4 and T32 glucose was not detected at all showing a complete  
272 consumption, and endly detected again and reaching 0.389 g l<sup>-1</sup>. Also organic acids as pyruvate, succinate,  
273 lactate, formate, acetate, propionate, butyrate, isobutyrate and isovalerate were detected and assumed to be  
274 metabolic end products.

275

276 16S rRNA gene libraries from the continuous enrichment culture in bioreactor

277

278 Archaeal and bacterial 16S rRNA genes were amplified from T7 and T28 culture samples from the continuous  
279 enrichment culture in bioreactor. All the archaeal sequences from T7 and T28 were related to the genus  
280 *Thermococcus* (Table 1). The sequence types A704 and A800 were affiliated to *T. siculi*, and shared more than  
281 97% of identity with a large number of 16S rRNA gene sequences related to members of the group *T. siculi* -  
282 *T. celer*, according to the BLAST analysis. Bacterial sequences affiliated with the orders *Clostridiales*,  
283 *Thermotogales* and *Deferribacterales* were retrieved in both T7 and T28 libraries (Table 1, Fig. 3) and were  
284 closely related to the hydrothermal species *Caminicella sporogenes* (97% 16S rRNA gene identity with clones  
285 775 and 813), *Marinitoga camini* (94% identity with the clone 716, 95% with clone 805) and *Deferribacter*

286 *abyssi* (98% identity with clones 737 and 820), respectively. The proportion of clones related to *Deferribacter*  
287 spp. and *Marinitoga* spp. increased slightly at T28, while the number of clones related to *Caminiella* spp.  
288 decreased from 50% to 7% (Fig. 3). Two phlotypes were recovered only in the T28 bacterial library: (i) 12  
289 clones were affiliated to *Thermosipho* spp. and the sequence type 840 shared 99% identity with *Thermosipho*  
290 MV1063, the closest species being *T. melanesiensis* (96% identity) and (ii) 3 clones were affiliated to  
291 *Thermodesulfatator* spp. and the sequence type 850 shared 96% identity with *T. indicus* (Table 1).

292

293 16S rRNA gene libraries from the batch enrichment cultures in vials

294

295 Total DNA was extracted from A1, A2 and B culture samples. Archaeal 16S rRNA gene was amplified by PCR  
296 only from sample A1, corresponding to the shortest incubated enrichment culture (24 h incubation). No archaeal  
297 16S rRNA gene sequences was amplified after longer incubation (A2: 41 h incubation), and after subculturing  
298 from A1 (B). The 55 archaeal clones from the A1 library were all related to the genus *Thermococcus* (Table 1).  
299 The sequence type A254 displayed 97 % identity with the closest described strain *T. barophilus* (AY099172),  
300 also originating from the Mid-Atlantic Ridge (Snake Pit) (Marteinsson et al. 1999) and growing in the range of  
301 48-95°C under atmospheric pressure. Bacterial 16S rRNA gene was amplified by PCR from all three samples.  
302 Sequences related to *Marinitoga* spp. and *Caminiella* spp. were retrieved in each library. Sequences related to  
303 the *Thermosipho* spp. were not recovered in the A1 culture, but only after 41 hours of incubation (A2) and after  
304 17 h subculturing (B). Moreover, a shift was observed in the library compositions; sequences related to  
305 *Marinitoga* spp. were widely dominant in the A1 library whereas the A2 and B libraries were largely dominated  
306 by sequences related to *Caminiella* spp. (Fig. 3). Extending the incubation time or subculturing from the  
307 primary A1 culture resulted in similar changes in the composition of the bacterial libraries.

308

309 Subcultures and isolations

310

311 Several strains were isolated from culture samples (T3, T7 and T28) from the bioreactor: (i) an archaeal strain  
312 *Thermococcus* spp. designated as AT1273 (99% 16S rRNA gene similarity with *Thermococcus siculi*) (ii) a  
313 strain *Thermosipho* spp. (order *Thermotogales*) designated as AT1272 (98% similarity with *Thermosipho*  
314 MV1063, 95% with *T. melanesiensis*) (iii) a new bacterial species of the *Marinitoga* genus (order  
315 *Thermotogales*) named *M. hydrogenitolerans* (Postec et al. 2005c); (iv) using the T medium, a new bacterial

316 species among the *Thermodesulfator* genus capable of sulphate-reduction (96% similarity with the  
317 hydrothermal species *T. indicus*), (v) using the DS and DN media, we isolated a strain closely affiliated to the  
318 hydrothermal species *Deferribacter abyssi* (99% identity).

319

320

## 321 **Discussion**

322

323 In this study, we used an original culture method to enrich thermophilic microorganisms from a hydrothermal  
324 black smoker: a continuous culture was performed in a gas-lift bioreactor during 45 days at 60°C and pH 6.5  
325 under anaerobic conditions. The microbial diversity in continuous culture and classical batch cultures in vials  
326 was compared.

327

328 Methodological considerations

329

330 The molecular inventories gave snapshots of the microbial diversity on a restricted number of samples while  
331 DGGE and whole-cell hybridisation revealed a temporal dynamics in the continuous culture. The use of different  
332 molecular techniques based on 16S rRNA gene analysis gave complementary data. For example, sequences of  
333 *Thermosipho* spp. were detected by DGGE at T7 in the bioreactor but were not evidenced by cloning. Inversely,  
334 *Thermodesulfator* spp. was detected by cloning but not by DGGE. The possible limitations of primer  
335 selectivity and cloning biases (Theron and Cloete 2000) or PCR biases (Suzuki and Giovannoni 1996) are well  
336 established and can explain the variability in results from different methods.

337 Considering the gas-lift bioreactor as a system for continuous cultivation, the potential adhesion of bacteria on  
338 the inner wall and the formation of a biofilm have to be examined, since the bioreactor includes no device to  
339 clean surfaces. From a precipitate sampled on the inner wall, no cells were detected by microscopy observation.

340 Culture attempt in vial and DNA extraction failed as well. X-RD analysis indicated that the precipitate was  
341 mainly composed of sulphurs (data not shown). The microorganisms detected in this study represent therefore  
342 cells in suspension. The dilution rate applied in the bioreactor after 34 h of batch culture was 0.04 h<sup>-1</sup> and  
343 corresponded to a generation time of 17.25 h. Although the continuous culture involves a progressive dilution of  
344 the medium inside the bioreactor, four volume changes, corresponding to 100 h (about 4 days) at a dilution rate  
345 of 0.04 h<sup>-1</sup>, have been considered sufficient to completely renew the culture medium inside the bioreactor (Raven

346 et al. 1992). Thus, microorganisms thriving in the continuous culture from T6 were not washed-out and should  
347 have grown with a minimal growth rate of 0.04 h<sup>-1</sup>.

348

349 *Thermococcales* as early heterotrophic colonizers

350

351 *Thermococcales* at deep-sea hydrothermal vents are widespread and members of the genus *Thermococcus* are  
352 some of the most numerous hyperthermophiles described from deep-sea vents. Investigation of their natural  
353 distribution showed that a viable *Thermococcus* population was present in the surface layers of mature  
354 hydrothermal chimneys (Harmsen et al. 1997; Takai et al. 2001; Schrenk et al. 2003). The early growth of  
355 *Thermococcales* was observed in enrichment cultures from hydrothermal chimney performed in vials and  
356 bioreactor at 90°C (Postec 2005b) and at 60°C (this study). Although all the members of the *Thermococcales* are  
357 hyperthermophilic, their early growth at 60°C both in the bioreactor and in vials is possible since (i) some  
358 *Thermococcus* sp. are able to grow at 60°C (Godfroy et al. 1997), (ii) an absence of latency phase could explain  
359 the early growth (*T. hydrothermalis*; Godfroy, pers. com.), (iii) the number of *Thermococcus*-related cells might  
360 be abundant in the chimney sample used as inoculum. This is also suggested by the detection of *Thermococcus*  
361 spp. in the molecular inventory performed directly on the studied chimney sample, while the bacteria grown in  
362 this study were not detected (Postec 2005d). Their growth at 60°C at the beginning of the enrichment culture and  
363 then when temperature was risen from 60 to 80°C after T31 may confer on *Thermococcales* a great ecological  
364 advantage to colonize new hydrothermal environments and they may be the first heterotrophs colonizing this  
365 ecosystem. This idea is supported by a recent study of nascent vent colonization in which protochimneys were  
366 deployed for short time on hydrothermal vents and heterotrophic groups including *Thermococcales* dominated  
367 the colonization of mineral surfaces after 72h (McCliment et al. 2006). The temporal sequence of colonization  
368 was analysed in the study of *in situ* samplers deployed on hydrothermal vents: the widespread occurrence of  
369 *Thermococcales* was demonstrated in short deployments (4-7 days) and decreased with time, suggesting that this  
370 group is an early surface colonizer (Nercessian et al. 2003).

371

372 Bacterial diversity and insight into microbial interactions

373

374 All the bacteria cultivated in this study were related to microorganisms from deep-sea hydrothermal vents.

375 *Caminiella* spp. and *Marinitoga* spp. were detected in both vials and bioreactor, while the autotrophs

376 *Deferribacter* spp. and *Thermodesulfatator* spp. were only detected in the bioreactor. Compared to batch cultures  
377 in vials, a larger diversity was described in the enriched community in bioreactor continuous supply of nutrients,  
378 gaseous inhibitory by-products removal and and pH regulation over time, what confirms previous results (Postec  
379 et al. 2005b).

380 Guided by the results of the molecular analysis, we succeeded in isolating microorganisms (heterotrophs and  
381 autotrophs) enriched in continuous in the gas-lift bioreactor. Three new species belonging to the genera  
382 *Thermosipho*, *Marinitoga* and *Thermodesulfatator* were obtained in pure culture in vials. They were related to *T.*  
383 *melanesiensis* and *M. camini*, both originating from Atlantic deep sea vents, and *T. indicus*, isolated from the  
384 Central Indian Ridge, respectively. *Marinitoga* sp. nov. was recently characterised and named *M.*  
385 *hydrogenitolerans* because its growth is not inhibited by high hydrogen concentrations (Postec et al. 2005c). The  
386 genus *Thermodesulfatator* is only represented so far by the species *T. indicus*, a thermophilic, anaerobic and  
387 strictly chemolithoautotrophic bacterium growing exclusively with CO<sub>2</sub> as sole carbon source, H<sub>2</sub> as sole  
388 electron donor and sulphate as sole electron acceptor (Moussard et al. 2004). A second chemolithoautotrophic to  
389 mixotrophic strain isolated from the bioreactor shared 99% 16S rRNA gene similarity with *Deferribacter abyssi*  
390 (Miroshnichenko et al. 2003). *D. abyssi* is thermophilic, anaerobic and facultative chemolithoautotrophic using  
391 elemental sulphur or nitrate as electron acceptors, similarly to the strain isolated in this study.

392 In the enrichment culture, growth of *Thermococcales* clearly occurred first. Their fermentative metabolism on  
393 proteinaceous substrates and in smaller extent carbohydrates through sulphur reduction might have generated a  
394 propitious environment for the growth of bacterial heterotrophs (*Caminiella* spp., *Thermosipho* spp. and  
395 *Marinitoga* spp) and then autotrophs (*Deferribacter* spp. and *Thermodesulfatator* spp.). The growth of  
396 *Thermosipho* sp. and *Marinitoga* sp. is correlated with the diminution of the glucose concentration in the  
397 medium and species of these genera are known to be able to use glucose as carbon substrate and to produce  
398 acetate as end metabolic product (Antoine et al. 1997; Alain et al. 2002b; Postec et al. 2005c).

399 Indeed 16S rRNA gene sequences related to the autotrophic microorganisms were detected by molecular  
400 analysis late at T28 but not at T7. Similarly, a temporal study of *in situ* collectors deployed on deep-sea  
401 hydrothermal vents showed that the microbial diversity of the colonizing community increased with time and  
402 that chemolithoautotrophs emerged during late stages (Nercessian et al. 2003).

403 The late growth of chemolithotrophs in co-culture with heterotrophs may be explained by interactions between  
404 species by means of metabolites exchanges. Organic carbon provided by the medium supported the growth of  
405 heterotrophs, afterwards the chemolithotrophs utilized carbon dioxide or acetate as carbon source and hydrogen

406 and acetate as possible electron donor, both compounds being end-products of fermentation. The study of a  
407 natural community showed that acetate and a range of other organic electron donors can be oxidised under  
408 sulphate-reducing conditions in hydrothermal vents at high temperature (90°C) (Tor et al. 2003). It has been  
409 suggested that acetate and hydrogen are the most prevalent organic fermentation products and important  
410 extracellular intermediates in the degradation of organic matter in hot microbial ecosystems, and that cooperative  
411 activity between fermentative microorganisms and sulphate reducers is important for the metabolism of  
412 fermentable compounds. Syntrophic interactions can also have an impact in hyperthermophilic co-cultures, for  
413 example on the metabolism of heterotrophic microorganisms co-cultivated with methanogens (Johnson et al.  
414 2006).

415

#### 416 Conclusions

417

418 A gas-lift bioreactor was used to cultivate in continuous a thermophilic microbial community from a deep sea  
419 hydrothermal chimney, on an organic-rich medium with sulphur under anaerobic conditions. The enrichment  
420 culture was monitored with molecular and chemical analyses. In the long-term running culture, the cultivated  
421 populations were evidenced to continuously evolve with time, instead of reaching a stationary state.  
422 *Thermococcales* dominated in the first hours of the enrichment cultures suggesting that this group early  
423 colonizes hydrothermal edifices, and may represent the first heterotrophic colonizers. A larger diversity was  
424 detected in the enrichment culture in bioreactor compared to culture in vials and most of the microorganisms  
425 enriched in bioreactor, including three new bacterial species, were successfully isolated by subculturing in vials.  
426 Results indicate that the continuous culture in a gas-lift bioreactor, combined with the use of molecular tools,  
427 could be of further use to access the "uncultivated" microbial community. The microorganisms isolated in this  
428 study displayed a phylogenetic and metabolic diversity. They are involved in the sulphur cycle (sulphur- and  
429 sulphate-reduction) and in the carbon cycle (autotrophy and heterotrophy). Autotrophic microorganisms were  
430 enriched in co-culture with heterotrophs, suggesting that prokaryotic species interact by means of metabolite  
431 exchange to support the growth of autotrophs. This emphasizes the importance of microbial interactions with  
432 surrounding microorganisms, animals or minerals within ecological niches. Inter-species interactions should be  
433 further taken into account to attempt the growth of as-yet uncultivated microorganisms and microbial  
434 metabolism need to be examined inside communities rather than extrapolated from pure cultures (Tor et al.  
435 2003). The bioreactor can be considered as a window to investigate *in vitro* interactions between population

436 interactions that may occur *in situ*. Considering that deep-sea hydrothermal vents are extreme environments  
437 encompassing intense thermal and chemical gradient (Karl 1995), the microbial communities inhabiting these  
438 disturbed systems must be strongly affected by environmental changes. The gas-lift bioreactor represents a  
439 promising tool to investigate *in vitro* the effect of physico-chemical perturbations on the microbial community  
440 structure.

441

442 **Acknowledgements** The authors want to thank P.M. Sarradin, chief scientist of the ATOS cruise, as well as the  
443 captain and crew of the RV/Atalante and the Victor team. This work was supported by Ifremer, European  
444 VENTOX Program and Région Bretagne. We also wish to thank Dr C. L. Van Dover and Dr G. Webster for  
445 their helpful reading of the manuscript.

446

447


448 **References**

449

- 450 Alain K, Olagnon M, Desbruyeres D, Page A, Barbier G, Juniper SK, Querellou J, Cambon-Bonavita MA  
451 (2002) Phylogenetic characterization of the bacterial assemblage associated with mucous secretions of the  
452 hydrothermal vent polychaete *Paralvinella palmiformis*. FEMS Microbiol Ecol 42:463-476
- 453 Altschul SF, Gish W, Miller W, Meyers EW, Lipman DJ (1990) Basic Local Alignment Search Tool. J Mol Biol  
454 215:403-410
- 455 DeChaine EG, Bates AE, Shank TM, Cavanaugh CM (2006) Off-axis symbiosis found: characterization and  
456 biogeography of bacterial symbionts of *Bathymodiolus* mussels from Lost City hydrothermal vents. Environ  
457 Microbiol 8:1902-1912
- 458 Galtier N, Gouy M, Gautier C (1996) SEAVIEW and PHYLO\_WIN: two graphic tools for sequence alignment  
459 and molecular phylogeny. CABIOS 12:543-548
- 460 Godfroy A, Meunier JR, Guezennec J, Lesongeur F, Raguénès G, Rimbault A, Barbier G (1996) *Thermococcus*  
461 *fumicolans* sp. nov. a new hyperthermophilic archaeum isolated from deep-sea hydrothermal vent in North  
462 Fiji basin. Int J Syst Bacteriol 46:1113-1119
- 463 Godfroy A, Lesongeur F, Raguénès G, Quérellou J, Antoine E, Meunier JR, Guezennec J, Barbier G (1997)  
464 *Thermococcus hydrothermalis* sp. nov., a new hyperthermophilic archaeon isolated from deep-sea  
465 hydrothermal vent. Int J Syst Bacteriol 47:622-626
- 466 Godfroy A, Raven NDH, Sharp RJ (2000) Physiology and continuous culture of the hyperthermophilic deep-sea  
467 vent archaeon *Pyrococcus abyssi* ST549. FEMS Microbiol Lett 186:127-132
- 468 Godfroy A, Postec A, Raven NDH (2005) Growth of hyperthermophilic microorganisms for physiological and  
469 nutritional studies. In: Rainey FA, Oren A (eds) Methods in Microbiology, Extremophiles. Academic Press,  
470 Oxford, pp 91-106
- 471 Harmsen HJM, Prieur D, Jeanthon C (1997) Distribution of microorganisms in deep-sea hydrothermal vent  
472 chimneys investigated by whole-cell hybridization and enrichment culture of thermophilic subpopulations.  
473 Appl Environ Microbiol 63:2876-2883
- 474 Inagaki F, Kuypers MMM, Tsunogai U, Ishibashi JI, Nakamura KI, Treude T, Ohkubo S, Nakaseama M, Gena  
475 K, Chiba H, Hirayama H, Nunoura T, Takai K, Jorgensen BB, Horikoshi K, Boetius A (2006) From the  
476 Cover: Microbial community in a sediment-hosted CO<sub>2</sub> lake of the southern Okinawa Trough hydrothermal  
477 system. PNAS 103:14164-14169
- 478 Johnson MR, Connors SB, Montero CI, Chou CJ, Shockley KR, Kelly RM (2006) The *Thermotoga maritima*  
479 phenotype is impacted by syntrophic interaction with *Methanococcus jannaschii* in hyperthermophilic  
480 coculture. Appl Environ Microbiol 72:811-818
- 481 Karl DM (1995) Ecology of free-living hydrothermal vent microbial communities. In: Karl DM (eds) The  
482 microbiology of deep-sea hydrothermal vents. CRC Press, pp 35-125
- 483 Kormas KA, Tivey MK, Von Damm K, Teske A (2006) Bacterial and archaeal phylotypes associated with  
484 distinct mineralogical layers of a white smoker spire from a deep-sea hydrothermal vent site (9°N, East  
485 Pacific Rise). Environ Microbiol 8:909-920
- 486 López-García P, Duperron S, Philippot P, Foriel J, Susini S, Moreira D (2003) Bacterial diversity in  
487 hydrothermal sediment and epsilon-proteobacterial dominance in experimental microcolonizers at the Mid-  
488 Atlantic Ridge. Environ Microbiol 5:961-976
- 489 Marteinsson V, Birrien J, Reysenbach A, Vernet M, Marie D, Gambacorta A, Messner P, Sleytr U, Prieur D  
490 (1999) *Thermococcus barophilus* sp. nov., a new barophilic and hyperthermophilic archaeon isolated under  
491 high hydrostatic pressure from a deep-sea hydrothermal vent. Int J Syst Bacteriol 49:351-359
- 492 McCliment EA, Voglesonger KM, O'Day PA, Dunn EE, Holloway JR, Cary SC (2006) Colonization of nascent,  
493 deep-sea hydrothermal vents by a novel archaeal and nanoarchaeal assemblage. Environ Microbiol 8:114-125
- 494 Miroshnichenko ML, Slobodkin AI, Kostrikina NA, L'Haridon S, Nercessian O, Spring S, Stackebrandt E,  
495 Bonch-Osmolovskaya EA, Jeanthon C (2003) *Deferribacter abyssi* sp. nov., an anaerobic thermophile from  
496 deep-sea hydrothermal vents of the Mid-Atlantic Ridge. Int J Syst Evol Microbiol 53:1637-1641
- 497 Moussard H, L'Haridon S, Tindall BJ, Banta A, Schumann P, Stackebrandt E, Reysenbach AL, Jeanthon C  
498 (2004) *Thermodesulfatator indicus* gen. nov., sp. nov., a novel thermophilic chemolithoautotrophic sulfate-  
499 reducing bacterium isolated from the Central Indian Ridge. Int J Syst Evol Microbiol 54:227-233
- 500 Moussard H, Corre E, Cambon-Bonavita MA, Fouquet Y, Jeanthon C (2006) Novel uncultured  
501 *Epsilonproteobacteria* dominate a filamentous sulphur mat from the 13°N hydrothermal vent field, East  
502 Pacific Rise. FEMS Microbiol Ecol 58:449-463
- 503 Muyzer G, Smalla K (1998) Application of denaturing gradient gel electrophoresis (DGGE) and temperature  
504 gradient gel electrophoresis (TGGE) in microbial ecology. Ant van Leeuw 73:127-141

505 Muyzer G, De Waal EC, Uitterlinden AG (1993) Profiling of complex microbial populations by denaturing  
 506 gradient gel electrophoresis analysis of polymerase chain reaction- amplified genes coding for 16S rRNA.  
 507 Appl Environ Microbiol 59:695-700  
 508 Nercessian O, Reysenbach AL, Prieur D, Jeanthon C (2003) Archaeal diversity associated with *in situ* samplers  
 509 deployed on hydrothermal vents on the East Pacific Rise (13°N). Environ Microbiol 5:492-502  
 510 Pace NR, Stahl DA, Lane DJ, Olsen GJ (1986) The analysis of natural microbial populations by ribosomal RNA  
 511 sequences. Adv Microbiol Ecol 9:1-55  
 512 Postec A, Pignet P, Cuff-Gauchard V, Schmitt A, Querellou J, Godfroy A (2005a) Optimisation of growth  
 513 conditions for continuous culture of the hyperthermophilic archaeon *Thermococcus hydrothermalis* and  
 514 development of sulphur-free defined and minimal media. Res Microbiol 156:82-87  
 515 Postec A, Urios L, Lesongeur L, Ollivier B, Querellou J, Godfroy A (2005b) Continuous enrichment culture and  
 516 molecular monitoring to investigate the microbial diversity of thermophiles inhabiting the deep-sea  
 517 hydrothermal ecosystems. Curr microbiol 50:138-144  
 518 Postec A, Le Breton C, Fardeau ML, Lesongeur F, Pignet P, Querellou J, Ollivier B, Godfroy A (2005c)  
 519 *Marinitoga hydrogenitolerans* sp. nov., a novel member of the order *Thermotogales* isolated from a black  
 520 smoker chimney on the Mid-Atlantic Ridge. Int J Syst Evol Microbiol 55:1217-1221  
 521 Postec A (2005d) Diversité de populations microbiennes thermophiles d'une cheminée hydrothermale océanique:  
 522 cultures d'enrichissement en bioréacteur et isolement d'espèces nouvelles. In: Thesis, Université de Provence,  
 523 France  
 524 Raven N, Ladwa N, Sharp R (1992) Continuous culture of the hyperthermophilic archaeum *Pyrococcus furiosus*.  
 525 Appl Microbiol Biotechnol 38:263-267  
 526 Raven NDH, Sharp RJ (1997) Development of defined and minimal media for the growth of the  
 527 hyperthermophilic archaeon *Pyrococcus furiosus* Vc1. FEMS Microbiol Lett 146:135-141  
 528 Schrenk MO, Kelley DS, Delaney JR, Baross JA (2003) Incidence and diversity of microorganisms within the  
 529 walls of an active deep-sea sulfide chimney. Appl Environ Microbiol 69:3580-3592  
 530 Sharp RJ, Raven NDH (1997) Isolation and growth of hyperthermophiles. In: Rhodes PM, Stanbury PF (eds)  
 531 Applied microbial physiology: a practical approach. IRL Press, Oxford, pp 23-51  
 532 Suzuki M, Giovannoni S (1996) Bias caused by template annealing in the amplification of mixtures of  
 533 16S rRNA genes by PCR. Appl Environ Microbiol 62:625-630  
 534 Takai K, Komatsu T, Inagaki F, Horikoshi K (2001) Distribution of *Archaea* in a black smoker chimney  
 535 structure. Appl Environ Microbiol 67:3618-3629  
 536 Theron J, Cloete TE (2000) Molecular techniques for determining microbial diversity and community structure  
 537 in natural environments. Crit Rev Microbiol 26:37-57  
 538 Thompson J, Higgins D, Gibson T (1994) CLUSTAL W: improving the sensitivity of progressive multiple  
 539 sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice.  
 540 Nuc Acids Res 22:4673-4680  
 541 Tor JM, Amend JP, Lovley DR (2003) Metabolism of organic compounds in anaerobic, hydrothermal sulphate-  
 542 reducing marine sediments. Environ Microbiol 5:583-591  
 543 Wery N, Cambon-Bonavita MA, Lesongeur F, Barbier G (2002) Diversity of anaerobic heterotrophic  
 544 thermophiles isolated from deep-sea hydrothermal vents of the Mid-Atlantic Ridge. FEMS Microbiol Ecol  
 545 41:105-114  
 546 Wery N, Lesongeur F, Pignet P, Derennes V, Cambon-Bonavita MA, Godfroy A, Barbier G (2001) *Marinitoga*  
 547 *camini* gen. nov., sp. nov., a rod-shaped bacterium belonging to the order *Thermotogales*, isolated from a  
 548 deep-sea hydrothermal vent. Int J Syst Evol Microbiol 51:495-504  
 549

550




551

552

553 **Fig. 1** Total cell densities of the bioreactor culture from T2 to T41 expressed in cell ml<sup>-1</sup> and *Archaea* and  
554 *Bacteria* ratio expressed in percentages of total cells determined by whole-cell hybridisation using the universal  
555 probes ARC915 and EUB338 respectively. The temperature was increased from 60°C to 70°C at T31 and from  
556 70°C to 80°C at T36.


557



558  
559

560 **Fig. 2** Whole-cell hybridisation of fixed cells from 3 samples of the enrichment culture in bioreactor: T2 (a, d),  
561 T7 (b, e) and T28 (c, f). Cells were stained with DAPI (a, b and c), and hybridised with the FITC-labelled  
562 Eub338 probe (e), or the Cy-3-labeled Arch 915 (d and f). Cells were viewed by epifluorescence microscopy in  
563 which DAPI-, FITC- and Cy-3-specific filters were used. The relative proportions of archaeal and bacterial cells  
564 were determined by counting approximately 2,000 cells on filter and resulted in: 99.0% of the cells detected at  
565 T2 belonged to the *Archaea* (d), 99.0% of the cells detected at T7 belonged to the *Bacteria* (e), 40% of the cells  
566 detected at T28 belonged to the *Archaea* (f). Scale bars = 10  $\mu$ m.

567



569

570

571 **Fig. 3** Composition of the bacterial clone libraries from enrichment cultures in vials (A1: 24 hours culture, A2:  
 572 41 hours culture and B: 17 hours subculture from A1) and from the enrichment culture in bioreactor (T7: 7 days  
 573 culture, T28: 28 days culture). The percentages of clones of each phylogenetic group are indicated on the pie-  
 574 charts.

575

576

577

578 **Table 1.** Distribution and phylogenetic affiliations of archaeal and bacterial 16S rDNA sequences ( $\geq$  97%  
579 similarity in each phylotype) from the enrichment culture in bioreactor and from enrichment cultures in flask  
580 both performed at 60°C. Representative clones were completely sequenced. The sequence types deposited in  
581 GenBank appear in bold.

582

	<b>Culture sample</b>	<b>Phylogenetic affiliation</b>	<b>Representative clones</b>	<b>Number of clones</b>	<b>Closest match organism<sup>a</sup></b>	<b>Identity<sup>a</sup> (%)</b>
<b>Bioreactor</b>	T7	<i>Thermococcales</i>	<b>A704</b> - A710-A712- A715- A730-A732-A737-A739	76	<i>Thermococcus siculi</i> (AY099185)	98
	T28	<i>Thermococcales</i>	<b>A800</b> -A811- A816-A817	71	<i>Thermococcus siculi</i> (AY099185)	98
	T7	<i>Clostridiales</i>	<b>775</b> -700-725-728-750-770-	38	<i>Caminicella sporogenes</i> (AJ320233)	97
		<i>Thermotogales</i>	<b>716</b> -705-706-709-724-768	29	<i>Marinitoga camini</i> (AJ250439)	94
		<i>Deferribacteriales</i>	<b>737</b> -711-740-754-769	9	<i>Deferribacter abyssi</i> (AJ515882)	98
	T28	<i>Clostridiales</i>	<b>813</b> -874	5	<i>Caminicella sporogenes</i> (AJ320233)	97
		<i>Thermotogales</i>	<b>805</b> -802-822-823-832	35	<i>Marinitoga camini</i> (AJ250439)	95
			<b>840</b> -812-821-825-868-884-	12	<i>Thermosipho</i> MV1063 (AJ419874)	99
		<i>Deferribacteriales</i>	893	17	<i>Deferribacter abyssi</i> (AJ515882)	98
		<i>Thermodesulfobacteriales</i>	<b>820</b> -829-856 <b>850</b> -816-858	3	<i>Thermodesulfatator indicus</i> (AF393376)	96
<b>Flask</b>	A1	<i>Thermococcales</i>	<b>A254</b>	55	<i>Thermococcus barophilus</i> (AY099172)	97
	A1	<i>Clostridiales</i>	<b>238</b>	1	<i>Caminicella sporogenes</i> (AJ320233)	97
		<i>Thermotogales</i>	<b>207</b> -245-219-240-255	58	<i>Marinitoga camini</i> (AJ250439)	94
	A2	<i>Clostridiales</i>	<b>413</b> -404-462	47	<i>Caminicella sporogenes</i> (AJ320233)	97
		<i>Thermotogales</i>	<b>436</b>	8	<i>Marinitoga camini</i> (AJ250439)	95
			<b>440<sup>b</sup></b>	1	<i>Thermosipho</i> MV1063 (AJ419874)	98
	B	<i>Clostridiales</i>	<b>608</b> -626-669-635	68	<i>Caminicella sporogenes</i> (AJ320233)	97
		<i>Thermotogales</i>	<b>660<sup>b</sup></b>	1	<i>Marinitoga camini</i> (AJ250439)	93
		<b>609</b>	1	<i>Thermosipho</i> MV1063 (AJ419874)	97	

583 <sup>a</sup> based on BLAST search. GenBank accession numbers are in brackets.

584 <sup>b</sup> partial sequence (600pb)

585